

Department of Pathology and Laboratory Medicine No.201, Sec. 2, Shipai Rd., Beitou District, Taipei City, Taiwan 11217, R.O.C.

Tel: 02-2875-7449

**Date**: 11 Aug 2023 1 of 3

## Sample Information

Patient Name: 胡慧貞 Gender: Female ID No.: A221321454 History No.: 22731960

**Age:** 55

Ordering Doctor: DOC1686E 陳玟均

Ordering REQ.: 0CPMBSY Signing in Date: 2023/08/09

Path No.: M112-00211 MP No.: MY23054

Assay: Oncomine Myeloid Assay Sample Type: Bone Marrow

**Bone Marrow Aspirating Date: 2023/08/08** 

Reporting Doctor: DOC5444B 楊靜芬 (Phone: 8#5444)

Note:

# Sample Cancer Type: Myelodysplastic Syndrome

Table of Contents	Page	Report Highlights	
Variants (Exclude variant in Taiwan BioBank with >1% allele frequency)	1 2	0 Relevant Biomarkers 0 Therapies Available	
Biomarker Descriptions		0 Clinical Trials	

### **Relevant Biomarkers**

No clinically significant biomarkers found in this sample.

Prevalent cancer biomarkers without relevant evidence based on included data sources  $RB1\ p.(R661Q)\ c.1982G>A$ 

# Variants (Exclude variant in Taiwan BioBank with >1% allele frequency)

DNA Sequence Variants											
Gene	Amino Acid Change	Coding	Variant ID	Locus	Allele Frequency	Transcript	Variant Effect	Coverage			
RB1	p.(R661Q)	c.1982G>A	COSM4428412	chr13:49033845	49.75%	NM_000321.2	missense	2000			
CALR	p.(?)	c68C>T		chr19:13049426	49.52%	NM_004343.4	unknown	1999			

## **Biomarker Descriptions**

#### **RB1** (RB transcriptional corepressor 1)

Background: The RB1 gene encodes the retinoblastoma protein (pRB), and is an early molecular hallmark of cancer. RB1 belongs to the family of pocket proteins that also includes p107 and p130, which play a crucial role in the cell proliferation, apoptosis, and differentiation<sup>1,2</sup>. RB1 is well characterized as a tumor suppressor gene that restrains cell cycle progression from G1 phase to S phase<sup>3</sup>. Specifically, RB1 binds and represses the E2F family of transcription factors that regulate the expression of genes involved in the G1/S cell cycle regulation<sup>1,2,4</sup>. Germline mutations in RB1 are associated with retinoblastoma (a rare childhood tumor) as well as other cancer types such as osteosarcoma, soft tissue sarcoma, and melanoma<sup>5</sup>.

Alterations and prevalence: Recurrent somatic alterations in RB1, including mutations and biallelic loss, lead to the inactivation of the RB1 protein. RB1 mutations are observed in urothelial carcinoma (approximately 16%), endometrial cancer (approximately 12%), and sarcomas (approximately 9%)<sup>6</sup>. Similarly, biallelic loss of RB1 is observed in sarcomas (approximately 13%), urothelial carcinoma (approximately 6%), and endometrial cancer (approximately 1%)<sup>6</sup>. Biallelic loss of the RB1 gene is also linked to the activation of chemotherapy-induced acute myeloid leukemia (AML) and acute lymphoblastic leukemia (ALL)<sup>7,8,9</sup>.

Potential relevance: Currently, there are no therapies approved for RB1 aberrations.

### References

- 1. Korenjak et al. E2F-Rb complexes regulating transcription of genes important for differentiation and development. Curr Opin Genet Dev . 2005 Oct;15(5):520-7. doi: 10.1016/j.gde.2005.07.001. PMID: 16081278
- 2. Sachdeva et al. Understanding pRb: toward the necessary development of targeted treatments for retinoblastoma. J. Clin. Invest. 2012 Feb;122(2):425-34. PMID: 22293180
- 3. Dyson. RB1: a prototype tumor suppressor and an enigma. Genes Dev. 2016 Jul 1;30(13):1492-502. PMID: 27401552
- 4. Cobrinik. Pocket proteins and cell cycle control. Oncogene. 2005 Apr 18;24(17):2796-809. PMID: 15838516
- 5. Dommering et al. RB1 mutations and second primary malignancies after hereditary retinoblastoma. Fam. Cancer. 2012 Jun;11(2):225-33. PMID: 22205104
- Cerami et al. The cBio cancer genomics portal: an open platform for exploring multidimensional cancer genomics data. Cancer Discov. 2012 May;2(5):401-4. PMID: 22588877
- 7. Anasua et al. Acute lymphoblastic leukemia as second primary tumor in a patient with retinoblastoma. . Oman J Ophthalmol . May-Aug 2016;9(2):116-8. PMID: 27433042
- 8. Tanaka et al. Frequent allelic loss of the RB, D13S319 and D13S25 locus in myeloid malignancies with deletion/translocation at 13q14 of chromosome 13, but not in lymphoid malignancies. Leukemia. 1999 Sep;13(9):1367-73. PMID: 10482987
- 9. Gombos et al. Secondary acute myelogenous leukemia in patients with retinoblastoma: is chemotherapy a factor?. Ophthalmology. 2007 Jul;114(7):1378-83. PMID: 17613328