

ABOUT THE TEST FoundationOne®Liquid CDx is a next generation sequencing (NGS) assay that identifies clinically relevant genomic alterations in circulating cell-free DNA.

PATIENT	DISEASE Unknown primary adenocarcinoma	PHYSICIAN	ORDERING PHYSICIAN Yeh, Yi-Chen	SPECIMEN	SPECIMEN ID YTC 6/2/1956
	NAME Chou, Yung-Tsung		MEDICAL FACILITY Taipei Veterans General Hospital		SPECIMEN TYPE Blood
	DATE OF BIRTH 02 June 1956		ADDITIONAL RECIPIENT None		DATE OF COLLECTION 12 June 2023
	SEX Male		MEDICAL FACILITY ID 205872		SPECIMEN RECEIVED 14 June 2023
	MEDICAL RECORD # 48929302		PATHOLOGIST Not Provided		

Biomarker Findings

Blood Tumor Mutational Burden - 3 Muts/Mb
Microsatellite status - MSI-High Not Detected
Tumor Fraction - Elevated Tumor Fraction Not Detected

Genomic Findings

For a complete list of the genes assayed, please refer to the Appendix.

BAP1T254fs*3
BRAFD594N
NFE2L2L23P
PBRM1splice site 4132+2T>A

Report Highlights

- Evidence-matched clinical trial options based on this patient's genomic findings: (p. [9](#))

BIOMARKER FINDINGS

Blood Tumor Mutational Burden -
 3 Muts/Mb

Microsatellite status -
 MSI-High Not Detected

Tumor Fraction -
 Elevated Tumor Fraction Not Detected

THERAPY AND CLINICAL TRIAL IMPLICATIONS

No therapies or clinical trials. See Biomarker Findings section

MSI-High not detected. No evidence of microsatellite instability in this sample (see Appendix section).

Tumor fraction is considered elevated when ctDNA levels are high enough that aneuploidy can be detected. The fact that elevated tumor fraction was not detected in this specimen indicates the possibility of lower levels of ctDNA but does not compromise confidence in any reported alterations. However, in the setting of a negative liquid biopsy result, orthogonal testing of a tissue specimen should be considered if clinically indicated (see Biomarker Findings section).

GENOMIC FINDINGS

VAF%

BAP1 - T254fs*3 4.1%
 10 Trials see p. [9](#)

THERAPIES WITH CLINICAL RELEVANCE (IN PATIENT'S TUMOR TYPE)

None

THERAPIES WITH CLINICAL RELEVANCE (IN OTHER TUMOR TYPE)

None

GENOMIC FINDINGS WITH NO REPORTABLE THERAPEUTIC OR CLINICAL TRIAL OPTIONS

For more information regarding biological and clinical significance, including prognostic, diagnostic, germline, and potential chemosensitivity implications, see the Genomic Findings section.

BRAF - D594N p. [7](#) **PBRM1** - splice site 4132+2T>A p. [8](#)
NFE2L2 - L23P p. [8](#)

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

NOTE Genomic alterations detected may be associated with activity of certain approved therapies; however, the therapies listed in this report may have varied clinical evidence in the patient's tumor type. Therapies and the clinical trials listed in this report may not be complete and/or exhaustive. Neither the therapies nor the trials identified are ranked in order of potential or predicted efficacy for this patient, nor are they ranked in order of level of evidence for this patient's tumor type. This report should be regarded and used as a supplementary source of information and not as the single basis for the making of a therapy decision. All treatment decisions remain the full and final responsibility of the treating physician and physicians should refer to approved prescribing information for all therapies. Therapies contained in this report may have been approved by the US FDA or other national authorities; however, they might not have been approved in your respective country. In the appropriate clinical context, germline testing of *APC*, *ATM*, *BAP1*, *BRCA1*, *BRCA2*, *BRIP1*, *CHEK2*, *FH*, *FLCN*, *MEN1*, *MLH1*, *MSH2*, *MSH6*, *MUTYH*, *NF1*, *NF2*, *PALB2*, *PMS2*, *POLE*, *PTEN*, *RAD51C*, *RAD51D*, *RB1*, *RET*, *SDHA*, *SDHB*, *SDHC*, *SDHD*, *SMAD4*, *STK11*, *TGFBR2*, *TP53*, *TSC1*, *TSC2*, *VHL*, and *WT1* is recommended.

Variant Allele Frequency is not applicable for copy number alterations.


Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Irene Shyu, M.D. | 21 June 2023
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1651473-01



FoundationOne®Liquid CDx
21 Jun 2023

HISTORIC PATIENT FINDINGS

ORD-1651473-01
VAF%

Blood Tumor Mutational Burden

3 Muts/Mb

Microsatellite status

MSI-High Not Detected

Tumor Fraction

Elevated Tumor Fraction Not Detected

BAP1	● T254fs*3	4.1%
BRAF	● D594N	0.10%
NFE2L2	● L23P	5.1%
PBRM1	● splice site 4132+2T>A	4.3%

IMPORTANT NOTE This comparison table refers only to genes and biomarkers assayed by prior FoundationOne®Liquid CDx or FoundationOne®CDx tests. Up to five previous tests may be shown.

For some genes in FoundationOne Liquid CDx, only select exons are assayed. Therefore, an alteration found by a previous test may not have been confirmed despite overlapping gene lists. Please refer to the Appendix for the complete list of genes and exons assayed. Variants reported for prior time points reflect reporting practices at the time of the historical test(s). Changes in variant reporting nomenclature, classification, or handling may result in the appearance of discrepancies across time points. The gene and biomarker list will be updated periodically to reflect new knowledge about cancer biology.

As new scientific information becomes available, alterations that had previously been listed as Variants of Unknown Significance (VUS) may become reportable or reportable variants may become VUS.

Tissue Tumor Mutational Burden (TMB) and blood TMB (bTMB) are estimated from the number of synonymous and non-synonymous single-nucleotide variants (SNVs) and insertions and deletions (indels) per area of coding genome sampled, after the removal of known and likely oncogenic driver events and germline SNPs. Tissue TMB is calculated based on variants with an allele frequency of $\geq 5\%$, and bTMB is calculated based on variants with an allele frequency of $\geq 0.5\%$.

Not Tested = not baited, not reported on test, or test preceded addition of biomarker or gene

Not Detected = baited but not detected on test

Detected = present (VAF% is not applicable)

VAF% = variant allele frequency percentage

Cannot Be Determined = Sample is not of sufficient data quality to confidently determine biomarker status

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Irene Shyu, M.D. | 21 June 2023
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1651473-01

Please note that other aspects of this table may have changed from the previous version to reflect the most up-to-date reporting information.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Irene Shyu, M.D. | 21 June 2023
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1651473-01

BIOMARKER FINDINGS

BIOMARKER

Blood Tumor Mutational Burden

RESULT

3 Muts/Mb

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

On the basis of clinical evidence in solid tumors, increased blood tumor mutational burden (bTMB) may be associated with greater sensitivity to immunotherapeutic agents, including anti-PD-L1¹⁻³, anti-PD-1³⁻⁴, anti-PD-1/CTLA4 therapies⁵⁻⁶, anti-PD-L1/CTLA4 therapies⁷⁻¹⁰. A Phase 2 multi-solid-tumor trial showed that bTMB ≥ 16 Muts/Mb (as measured by this assay) was associated with improved survival from treatment with a PD-1 inhibitor alone or in combination with a CTLA-4 inhibitor⁵. In non-small cell lung cancer (NSCLC), multiple clinical trials have shown patients with higher bTMB derive clinical benefit from immune checkpoint inhibitors following single-agent or combination treatments with either CTLA4 inhibitors or chemotherapy, with reported high bTMB cutpoints ranging from 6 Muts/Mb-16 Muts/Mb^{1,8-10}. In head and neck squamous cell

carcinoma (HNSCC), a Phase 3 trial showed that bTMB ≥ 16 Muts/Mb (approximate equivalency ≥ 8 Muts/Mb as measured by this assay) was associated with improved survival from treatment with a PD-L1 inhibitor alone or in combination with a CTLA-4 inhibitor¹¹. In colorectal cancer (CRC), a Phase 2 study showed that bTMB TMB ≥ 28 Muts/Mb (approximate equivalency ≥ 14 Muts/Mb as measured by this assay) was associated with improved OS from a PD-L1 inhibitor alone or in combination with a CTLA-4 inhibitor⁷.

FREQUENCY & PROGNOSIS

Average bTMB levels in solid tumors other than NSCLC have not been evaluated (PubMed, Mar 2023). Published data investigating the prognostic implications of TMB have mainly been investigated in the context of tissue TMB. In patients with NSCLC, increased TMB is associated with higher tumor grade and poor prognosis¹², as well as with a decreased frequency of known driver mutations in EGFR, ALK, ROS1, or MET (1% of high-TMB samples each), but not BRAF (10.3%) or KRAS (9.4%)¹³. Although some studies have reported a lack of association between smoking and increased TMB in NSCLC^{12,14}, several other large studies did find a strong link¹⁵⁻¹⁸. In CRC, elevated TMB is associated with a higher frequency of BRAF V600E driver mutations¹⁹⁻²⁰ and with

microsatellite instability (MSI)²⁰, which in turn has been reported to correlate with better prognosis²¹⁻²⁸. Although increased TMB is associated with increased tumor grade in endometrioid endometrial carcinoma²⁹⁻³² and bladder cancer³³, it is also linked with improved prognosis in patients with these tumor types³⁰.

FINDING SUMMARY

Blood tumor mutational burden (bTMB, also known as mutation load) is a measure of the number of somatic protein-coding base substitution and insertion/deletion mutations from circulating tumor DNA in blood. TMB is affected by a variety of causes, including exposure to mutagens such as ultraviolet light in melanoma³⁴⁻³⁵ and cigarette smoke in lung cancer³⁶⁻³⁷, treatment with temozolomide-based chemotherapy in glioma³⁸⁻³⁹, mutations in the proofreading domains of DNA polymerases encoded by the POLE and POLD1 genes^{19,30,40-42}, and microsatellite instability (MSI)^{19,30,42}. High bTMB levels were not detected in this sample. It is unclear whether the bTMB levels in this sample would be predicted to be associated with sensitivity to PD-1- or PD-L1-targeting immune checkpoint inhibitors, alone or in combination with other agents¹⁻²⁴. Depending on the clinical context, TMB testing of an alternate sample or by another methodology could be considered.

BIOMARKER

Tumor Fraction

RESULT

Elevated Tumor Fraction Not Detected

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

Specimens with elevated tumor fraction values have high circulating-tumor DNA (ctDNA) content, and thus higher sensitivity for identifying genomic alterations. Such specimens are at a lower risk of false negative results. However, if elevated tumor fraction is not detected, it does not exclude the presence of disease burden or compromise the confidence of reported alterations. Tumor fraction levels currently have limited implications for diagnosis, surveillance, or therapy and should not

be overinterpreted or compared from one blood draw to another. There are currently no targeted approaches to address specific tumor fraction levels. In the research setting, changes in tumor fraction estimates have been associated with treatment duration and clinical response and may be a useful indicator for future cancer management⁴³⁻⁴⁸.

FREQUENCY & PROGNOSIS

Detectable ctDNA levels have been reported in a variety of tumor types, with higher tumor fraction levels reported for patients with metastatic (Stage 4) tumors compared with patients with localized disease (Stages 1 to 3)⁴⁹. Elevated tumor fraction levels have been reported to be associated with worse prognosis in a variety of cancer types, including pancreatic cancer⁵⁰, Ewing sarcoma and osteosarcoma⁵¹, prostate cancer⁴⁶, breast cancer⁵², leiomyosarcoma⁵³, esophageal cancer⁵⁴, colorectal

cancer⁵⁵, and gastrointestinal cancer⁵⁶.

FINDING SUMMARY

Tumor fraction provides an estimate of the percentage of ctDNA present in a cell-free DNA (cfDNA) sample. The tumor fraction estimate for this sample is based on the observed level of aneuploid instability. The tumor fraction algorithm utilized for FoundationOne Liquid CDx uses the allele frequencies of approximately 1,000 single-nucleotide polymorphism (SNP) sites across the genome. Unlike the maximum somatic allele frequency (MSAF) method of estimating ctDNA content⁵⁷, the tumor fraction metric does not take into account the allele frequency of individual variants but rather produces a more holistic estimate of ctDNA content using data from across the genome. The amount of ctDNA detected may correlate with disease burden and response to therapy⁵⁸⁻⁵⁹.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Irene Shyu, M.D. | 21 June 2023
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1651473-01

GENOMIC FINDINGS

GENE

BAP1

ALTERATION

T254fs*3

HGVS VARIANT

NM_004656.2:c.759del (p.T254Qfs*3)

VARIANT CHROMOSOMAL POSITION

chr3:52440292-52440293

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

Clinical⁶⁰ and preclinical⁶¹ evidence in the context of mesothelioma suggests that tumors with BAP1 inactivation may be sensitive to EZH2 inhibitors such as tazemetostat. In a Phase 1 study of the EZH2 inhibitor CPI-0209, 67% (4/6) of patients with mesothelioma had BAP1 loss or alterations, 1 of whom achieved PR⁶². Clinical⁶³⁻⁶⁶ and preclinical⁶⁷⁻⁷⁰ studies suggest that BAP1 inactivation might be associated with sensitivity to PARP inhibitors. Phase 2, retrospective, and case studies have reported PR or SD for patients with BAP1-deficient cholangiocarcinoma, uveal melanoma, mesothelioma, and clear cell renal cell carcinoma treated with olaparib, rucaparib, niraparib, or veliparib⁶³⁻⁶⁶. One preclinical study suggests that histone deacetylase inhibitors may be

beneficial in BAP1-mutated uveal melanoma; however, it is unclear if these inhibitors are effective in other BAP1-mutated cancers⁷¹.

FREQUENCY & PROGNOSIS

Mutations in BAP1 have been reported in a variety of tumor types, but most frequently in mesothelioma (21%), cholangiocarcinoma (19%), uveal melanoma (16%), kidney renal clear cell carcinoma (9.5%), and uterine corpus endometrial carcinoma (5.6%) (cBioPortal, Feb 2023)⁷²⁻⁷³. Studies reported in the literature confirm the relatively high frequency of BAP1 mutation in clear cell renal cell carcinoma and malignant pleural mesothelioma⁷⁴⁻⁷⁵. The chromosomal region where BAP1 is located is subject to frequent deletion in non-small cell lung cancer (NSCLC) and other cancers⁷⁶. High BAP1 protein expression in patients with NSCLC has been associated with increased median survival time as compared to patients with low expression⁷⁷; however, decreases in the expression BAP1 mRNA and protein have been reported in colorectal cancers and have been associated with poor prognosis⁷⁸. In other tumor types, such as clear cell renal cell carcinoma, BAP1 mutations have been associated with high tumor grade, worse cancer-specific survival, and shorter overall survival^{69,79-80}.

FINDING SUMMARY

BAP1 (BRCA1 associated protein-1) encodes a ubiquitin hydrolase, a protein involved in regulating the availability of target proteins for the ubiquitin-proteasome protein degradation pathway; BAP1 is located on chromosome 3p21.3, in a region of frequent loss of heterozygosity (LOH) in breast and lung cancer, and has been postulated to be a tumor suppressor^{76,81}. Alterations such as seen here may disrupt BAP1 function or expression⁸¹⁻⁹⁰.

POTENTIAL GERMLINE IMPLICATIONS

BAP1 germline inactivating alterations, including mutations and deletions, are associated with BAP1 tumor predisposition syndrome (BAP1-TPDS), an autosomal-dominant syndrome characterized by early onset of benign melanocytic skin tumors^{84,91-92}. An estimated 2% of patients with BAP1-inactivated melanocytic tumors display germline BAP1 mutations⁹³. Later in life, patients have an increased risk of cancers such as uveal melanoma, mesothelioma, clear cell renal cell carcinoma, basal cell carcinoma, and meningioma^{83-87,94}. In small studies, the prevalence of pathogenic germline BAP1 mutation has been reported as 22% in familial uveal melanoma and 4.4% in mesothelioma⁹⁵⁻⁹⁶. In the appropriate clinical context, germline testing of BAP1 is recommended.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Irene Shyu, M.D. | 21 June 2023
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1651473-01

GENOMIC FINDINGS

GENE

BRAF

ALTERATION

D594N

HGVS VARIANT

NM_004333.4:c.1780G>A (p.D594N)

VARIANT CHROMOSOMAL POSITION

chr7:140453155

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

Clinical outcomes for patients with activating BRAF alterations treated with BRAF and MEK inhibitors are most extensive at the V600 codon; outcomes are more limited for BRAF class 3 kinase-impaired or inactivating mutations such as one or more of the alterations seen here. A retrospective study of immunotherapies in NSCLC reported a 78% DCR (7/9) for patients with BRAF class 3 mutations⁹⁷. MEK inhibitors alone or in combination with BRAF inhibitors may be efficacious in these alterations; a basket trial of single-agent MEK inhibitor trametinib reported 1 PR, 8 SDs, and 9 PDs for these patients⁹⁸, and

combination therapies reported individual responses in other basket trials⁹⁹⁻¹⁰⁰. A retrospective analysis in BRAF-mutated melanoma reported PD as the best response in BRAF class 3 alterations for 2 patients treated with MEK inhibitors and 3 patients treated with RAF inhibitors¹⁰¹. Single-agent BRAF inhibitor vemurafenib was not effective in a Phase 2 trial in NSCLC, which reported no responses for 6 patients with class 3 BRAF alterations¹⁰²; a basket trial of vemurafenib also observed no responses for these patients (n=3)¹⁰³. Investigational BRAF¹⁰⁴ and ERK¹⁰⁵ inhibitors are also in development; a basket trial of ulixertinib reported no responses and 3 SDs for patients across class 3-mutated tumors¹⁰⁵.

FREQUENCY & PROGNOSIS

BRAF mutation has been most extensively studied in melanoma, where it has been reported in 37-66% of cases¹⁰⁶⁻¹⁰⁹. BRAF mutation also occurs at high frequencies in patients with papillary craniopharyngiomas (95%)¹¹⁰, metanephric kidney adenomas (90%)¹¹¹, and papillary thyroid carcinoma (45%)¹¹²⁻¹¹⁴, and has also been reported in lung adenocarcinoma (10%)¹¹⁵ and colorectal cancer (9%)¹⁹. Studies on the effect of BRAF alteration on prognosis are conflicting with reports of

association with poor prognosis in cholangiocarcinoma¹¹⁶⁻¹¹⁸ and colorectal cancer¹¹⁹⁻¹²⁶, improved prognosis in ovarian cancer¹²⁷, or no association in NSCLC¹²⁸⁻¹²⁹ and pancreatic ductal adenocarcinoma¹³⁰. BRAF mutation in papillary thyroid carcinoma have been reported to correlate with poor prognosis in some studies^{112,114,131-135}, but not in other studies¹³⁶⁻¹³⁷. There are similarly conflicting reports regarding the prognostic significance of BRAF mutation in the context of melanoma¹³⁸⁻¹⁴¹.

FINDING SUMMARY

BRAF encodes a member of the RAF family of protein kinases, which includes ARAF, BRAF, and CRAF. These kinases function downstream of RAS as part of the MAPK (RAF-MEK-ERK) signaling cascade that facilitates cell proliferation, survival and transformation¹⁴²⁻¹⁴³. BRAF mutations have been reported in up to 20% of all cancers, with the majority of mutations occurring at the V600 position^{106,144}. Alterations such as the class 3 mutation seen here have been shown to require concomitant upstream RAS activity in contrast with independently activating BRAF V600 or class 2 alterations¹⁴⁵⁻¹⁵⁸, and may activate the MEK-ERK signaling pathway via CRAF^{145-147,154,159}.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Irene Shyu, M.D. | 21 June 2023
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1651473-01

GENOMIC FINDINGS

GENE

NFE2L2

ALTERATION

L23P

HGVS VARIANT

NM_006164.4:c.68T>C (p.L23P)

VARIANT CHROMOSOMAL POSITION

chr2:178098977

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

There are no approved therapies that directly target NFE2L2 alterations; however, activating alterations in NFE2L2 may indicate sensitivity to mTORC1/2 inhibitors¹⁶⁰⁻¹⁶¹. A Phase 2 study of the mTORC1/2 inhibitor sapanisertib for patients with advanced or recurrent lung squamous cell carcinoma (SCC) reported a 29% (2/7) ORR and a 100% (7/7) DCR for patients with NFE2L2 alterations¹⁶². A study of patients with localized non-small cell lung cancer (NSCLC) identified pathogenic KEAP1 and NFE2L2 mutations as predictors of local recurrence following radiotherapy but not surgery; limited

preclinical data also showed that treatment with a glutaminase inhibitor sensitized KEAP1-mutated NSCLC cells to radiation¹⁶³. In other preclinical studies, treatment with AKT inhibitors sensitized lung cancer cells harboring KEAP1 or NFE2L2 mutations to both chemotherapy and radiation therapy¹⁶⁴⁻¹⁶⁵.

FREQUENCY & PROGNOSIS

NFE2L2 mutations have been reported in several cancer types, including tumors affecting the liver (3%-5%), lung (4%-8%), and esophagus (2%-6%); 1.4% of samples with lung cancer, 1.2% of samples with liver cancer, and 1.2% of samples with esophageal cancer had a copy number alteration^{72-73,166-173} (cBioPortal, COSMIC, PubMed, Jan 2023). In the context of lung cancer, NFE2L2 mutations are most prevalent in smokers and patients with squamous cell carcinomas (SCC); NFE2L2 mutations also correlate with poor survival in multivariate analysis (mean survival 55 months vs. 81 months)¹⁷⁴⁻¹⁷⁵. NRF2 protein expression, especially nuclear expression, correlates with poor patient prognosis in lung cancer¹⁷⁶, colorectal cancer¹⁷⁷, gastric cancer¹⁷⁸⁻¹⁸⁰, esophageal SCC¹⁸¹⁻¹⁸², and osteosarcoma¹⁸³⁻¹⁸⁴, among others.

Additionally, cancers with increased NRF2 activity show increased resistance to chemotherapy and radiotherapy^{178-179,181,185}. Patients with non-small cell lung cancer (NSCLC) harboring mutation in both NFE2L2 and KEAP1 experienced significantly shorter OS and poorer responses to both atezolizumab and docetaxel than patients' wildtype for these genes in a retrospective analysis of the Phase 2 POPLAR and Phase 3 OAK trials¹⁸⁶, and concurrent mutation of NFE2L2 and KEAP1 has been proposed as a distinct subtype of NSCLC associated with increased rates of progression¹⁸⁷.

FINDING SUMMARY

NFE2L2 encodes nuclear factor E2-related factor 2 (NRF2), a transcription factor that plays a critical role in responses to oxidative and electrophilic stress¹⁸⁸. NRF2 activity is antagonized by KEAP1, which binds NRF2 to promote its ubiquitination and subsequent degradation¹⁸⁹. NRF2 missense mutations tend to cluster within the KEAP1 binding domain (especially amino acids 21-36 and 74-86); these mutations prevent degradation and lead to high levels of NRF2 protein and constitutive expression of NRF2 target genes¹⁷⁴. These mutations are considered activating.

GENE

PBRM1

ALTERATION

splice site 4132+2T>A

HGVS VARIANT

NM_018313.4:c.4132+2T>A (p.?)

VARIANT CHROMOSOMAL POSITION

chr3:52595781

POTENTIAL TREATMENT STRATEGIES

— Targeted Therapies —

On the basis of significant clinical data from prospective studies, PBRM1 inactivation may predict benefit from PD-1-targeting immune checkpoint inhibitors, such as nivolumab, pembrolizumab, cemiplimab, or dostarlimab, for patients with clear cell renal cell carcinoma and prior anti-angiogenic therapy¹⁹⁰⁻¹⁹². However, multiple retrospective analyses report that PBRM1 mutation status is not associated with clinical benefit from various immune checkpoint inhibitors

in other solid tumor types, including non-small cell lung cancer, urothelial carcinoma, melanoma, or esophagogastric cancer, suggesting that the impact of PBRM1 loss of function may depend on tumor type¹⁹³⁻¹⁹⁶.

FREQUENCY & PROGNOSIS

Somatic mutations in PBRM1 are prevalent in clear cell renal cell carcinomas (ccRCC) (41%)¹⁹⁷, intrahepatic cholangiocarcinomas (9-13%)¹⁹⁸⁻²⁰¹, and bladder urothelial carcinomas (6-14%)²⁰²⁻²⁰⁴. PBRM1 mutations are detected in other tumor types, including in 37% (11/30) of papillary meningiomas and 4% (2/54) of thymic carcinomas²⁰⁵⁻²⁰⁶ and in tumors of the skin (7.2%), stomach (5.7%), large intestine (4.8%), lung (2.7%), and soft tissue (2.4%) (COSMIC, Jan 2023)¹⁷³. Preclinical studies have shown that PBRM1 loss increases the proliferation of clear cell renal cell carcinoma (ccRCC) cell lines¹⁹⁷. PBRM1 protein loss or mutations are correlated with late tumor stage, low differentiation grade, and/or poor patient prognosis in ccRCC^{80,207-208}, extrahepatic cholangiocarcinoma²⁰⁰, and pancreatic cancer²⁰⁹.

However, 1 ccRCC study reported no correlation between PBRM1 mutations and cancer-specific survival²¹⁰. In ccRCC, PBRM1 alterations are generally observed to be mutually exclusive with BAP1 alterations^{69,197}; a retrospective analysis of 145 primary ccRCCs found a decreased median OS for patients with mutations in both BAP1 and PBRM1 compared with patients having either mutated gene alone⁷⁹. A trend toward worse survival was also seen for patients with intrahepatic cholangiocarcinoma harboring mutations in chromatin modifiers, including BAP1, ARID1A, or PBRM1¹⁹⁹.

FINDING SUMMARY

PBRM1 (Polybromo-1), also known as BAF180, encodes a subunit of ATP-dependent chromatin-remodeling complexes and a required cofactor for ligand-dependent transactivation by nuclear hormone receptors²¹¹. Mutation, loss, or inactivation of PBRM1 has been reported in several cancers, suggesting PBRM1 is a tumor suppressor^{197,199,212}. Alterations such as seen here may disrupt PBRM1 function or expression²¹³⁻²¹⁸.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Irene Shyu, M.D. | 21 June 2023
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1651473-01

CLINICAL TRIALS

IMPORTANT Clinical trials are ordered by gene and prioritized by: age range inclusion criteria for pediatric patients, proximity to ordering medical facility, later trial phase, and verification of trial information within the last two months. While every effort is made to ensure the accuracy of the information contained below, the information available in the public domain is continually updated and should be investigated by the physician or

research staff. This is not a comprehensive list of all available clinical trials. There may also be compassionate use or early access programs available, which are not listed in this report. Foundation Medicine displays a subset of trial options and ranks them in this order of descending priority: Qualification for pediatric trial → Geographical proximity → Later trial phase. Clinical trials are not ranked in order of potential or predicted efficacy for this patient or

in order of level of evidence for this patient's tumor type. Clinical trials listed here may have additional enrollment criteria that may require medical screening to determine final eligibility. For additional information about listed clinical trials or to conduct a search for additional trials, please see clinicaltrials.gov. However, clinicaltrials.gov does not list all clinical trials that might be available.

GENE
BAP1
RATIONALE

BAP1 inactivating alterations may predict sensitivity to PARP inhibitors.

ALTERATION

T254fs*3

NCT04434482
PHASE 1

IMP4297 in Combination With Temozolomide in Patients With Advanced Solid Tumors and Small Cell Lung Cancer

TARGETS
 PARP

LOCATIONS: Taipei (Taiwan), Taichung (Taiwan), Tainan (Taiwan), Gyeonggi-do (Korea, Republic of), Orange (Australia), Blacktown (Australia), Albury (Australia)

NCT03742895
PHASE 2

Efficacy and Safety of Olaparib (MK-7339) in Participants With Previously Treated, Homologous Recombination Repair Mutation (HRRm) or Homologous Recombination Deficiency (HRD) Positive Advanced Cancer (MK-7339-002 / LYNK-002)

TARGETS
 PARP

LOCATIONS: Seongnam-si (Korea, Republic of), Seoul (Korea, Republic of), Darlinghurst (Australia), Adana (Turkey), Jerusalem (Israel), Konya (Turkey), Ramat Gan (Israel), Istanbul (Turkey), Antalya (Turkey), Brasov (Romania)

NCT02264678
PHASE 1/2

Ascending Doses of AZD6738 in Combination With Chemotherapy and/or Novel Anti Cancer Agents

TARGETS
 ATR, PARP, PD-L1

LOCATIONS: Seoul (Korea, Republic of), Goyang-si (Korea, Republic of), Cambridge (United Kingdom), Withington (United Kingdom), Manchester (United Kingdom), London (United Kingdom), Coventry (United Kingdom), Sutton (United Kingdom), Oxford (United Kingdom), Villejuif (France)

NCT05021367
PHASE 1

A Clinical Study of TQB3823 in Patients With Advanced Malignant Tumor

TARGETS
 PARP

LOCATIONS: Guangzhou (China)

NCT04170153
PHASE 1

M1774 in Participants With Metastatic or Locally Advanced Unresectable Solid Tumors

TARGETS
 ATR, PARP

LOCATIONS: Beijing (China), Chuo-ku (Japan), Kashiwa-shi (Japan), Newcastle upon Tyne (United Kingdom), Cambridge (United Kingdom), Manchester (United Kingdom), Sutton (United Kingdom), Barcelona (Spain), Valencia (Spain), Madrid (Spain)

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Irene Shyu, M.D. | 21 June 2023
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
 Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
 Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1651473-01

CLINICAL TRIALS
NCT05035745
PHASE 1/2

Selinexor & Talazoparib in Advanced Refractory Solid Tumors; Advanced/Metastatic Triple Negative Breast Cancer (START)

TARGETS
 XPO1, PARP

LOCATIONS: Singapore (Singapore)

NCT03772561
PHASE 1

Phase I Study of AZD5363 + Olaparib + Durvalumab in Patients With Advanced or Metastatic Solid Tumor Malignancies

TARGETS
 PARP, AKTs, PD-L1

LOCATIONS: Singapore (Singapore)

NCT03297606
PHASE 2

Canadian Profiling and Targeted Agent Utilization Trial (CAPTUR)

TARGETS
 VEGFRs, ABL, SRC, ALK, ROS1, AXL, TRKA, MET, TRKC, DDR2, KIT, EGFR, PD-1, CTLA-4, PARP, CDK4, CDK6, FLT3, CSF1R, RET, mTOR, ERBB2, MEK, BRAF, SMO

LOCATIONS: Vancouver (Canada), Kelowna (Canada), Edmonton (Canada), Saskatoon (Canada), Regina (Canada), Ottawa (Canada), Montreal (Canada), Toronto (Canada), Kingston (Canada), London (Canada)

NCT05327010
PHASE 2

Testing the Combination of the Anti-cancer Drugs ZEN003694 (ZEN-3694) and Talazoparib in Patients With Advanced Solid Tumors, The ComBET Trial

TARGETS
 PARP, BRD4, BRDT, BRD2, BRD3

LOCATIONS: Colorado, Illinois, Texas, North Carolina, Georgia

NCT02769962
PHASE 1/2

Trial of CRLX101, a Nanoparticle Camptothecin With Olaparib in People With Relapsed/Refractory Small Cell Lung Cancer

TARGETS
 PARP, TOP1

LOCATIONS: Maryland

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

 Electronically signed by Irene Shyu, M.D. | 21 June 2023
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
 Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
 Foundation Medicine, Inc. | www.rochefoundationmedicine.com

 Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1651473-01

APPENDIX
Variants of Unknown Significance

NOTE One or more variants of unknown significance (VUS) were detected in this patient's tumor. These variants may not have been adequately characterized in the scientific literature at the time this report was issued, and/or the genomic context of these alterations makes their significance unclear. We choose to include them here in the event that they become clinically meaningful in the future.

EGFR

NM_005228.3: c.104G>A
(p.S35N)
chr7:55209994 and
NM_005228.3: c.776C>T
(p.T259M)
chr7:55221732

HGF

NM_000601.4: c.1859C>T
(p.T620I)
chr7:81334968

JAK1

NM_002227.2: c.1554C>G
(p.S518R)
chr1:65321286

KMT2D (MLL2)

NM_003482.4: c.794G>A
(p.R265H)
chr12:49447304 and
NM_003482.4: c.5285G>A
(p.S1762N)
chr12:49437685

NBN

NM_002485.4:
c.102_104delinsAAC (p.I35T)
chr8:90995017-90995019

NF1

NM_001042492.2: c.4726C>T
(p.H1576Y)
chr17:29592248

PIK3C2B

NM_002646.3: c.388A>G
(p.I130V)
chr1:204438543

TGFBR2

NM_003242.5: c.1189G>A
(p.D397N)
chr3:30713864

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Irene Shyu, M.D. | 21 June 2023
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1651473-01

APPENDIX

Genes assayed in FoundationOne®Liquid CDx

FoundationOne Liquid CDx interrogates 324 genes, including 309 genes with complete exonic (coding) coverage and 15 genes with only select non-coding coverage (indicated with an *); 75 genes (indicated in bold) are captured with increased sensitivity and have complete exonic (coding) coverage unless otherwise noted.

ABL1 Exons 4-9	ACVR1B	AKT1 Exon 3	AKT2	AKT3	ALK Exons 20-29, Introns 18, 19	ALOX12B	AMER1 (FAM123B or WTX)	APC
AR	ARAF Exons 4, 5, 7, 11, 13, 15, 16	ARFRP1	ARID1A	ASXL1	ATM	ATR	ATRX	AURKA
AURKB	AXIN1	AXL	BAP1	BARD1	BCL2	BCL2L1	BCL2L2	BCL6
BCOR	BCORL1	BCR* Introns 8, 13, 14	BRAF Exons 11-18, Introns 7-10	BRCA1 Introns 2, 7, 8, 12, 16, 19, 20	BRCA2 Intron 2	BRD4	BRIP1	BTG1
BTG2	BTK Exons 2, 15	CALR	CARD11	CASP8	CBFB	CBL	CCND1	CCND2
CCND3	CCNE1	CD22	CD70	CD74* Introns 6-8	CD79A	CD79B	CD274 (PD-L1)	CDC73
CDH1	CDK12	CDK4	CDK6	CDK8	CDKN1A	CDKN1B	CDKN2A	CDKN2B
CDKN2C	CEBPA	CHEK1	CHEK2	CIC	CREBBP	CRKL	CSF1R	CSF3R
CTCF	CTNNA1	CTNNB1 Exon 3	CUL3	CUL4A	CXCR4	CYP17A1	DAXX	DDR1
DDR2 Exons 5, 17, 18	DIS3	DNMT3A	DOT1L	EED	EGFR Introns 7, 15, 24-27	EMSY (C11orf30)	EP300	EPHA3
EPHB1	EPHB4	ERBB2	ERBB3 Exons 3, 6, 7, 8, 10, 12, 20, 21, 23, 24, 25	ERBB4	ERCC4	ERG	ERRF1	ESR1 Exons 4-8
ETV4* Intron 8	ETV5* Introns 6, 7	ETV6* Introns 5, 6	EWSR1* Introns 7-13	EZH2 Exons 4, 16, 17, 18	EZR* Introns 9-11	FANCA	FANCC	FANCG
FANCL	FAS	FBXW7	FGF10	FGF12	FGF14	FGF19	FGF23	FGF3
FGF4	FGF6	FGFR1 Introns 1, 5, Intron 17	FGFR2 Intron 1, Intron 17	FGFR3 Exons 7, 9 (alternative designation exon 10), 14, 18, Intron 17	FGFR4	FH	FLCN	FLT1
FLT3 Exons 14, 15, 20	FOXL2	FUBP1	GABRA6	GATA3	GATA4	GATA6	GID4 (C17orf39)	GNA11 Exons 4, 5
GNA13	GNAQ Exons 4, 5	GNAS Exons 1, 8	GRM3	GSK3B	H3-3A (H3F3A)	HDAC1	HGF	HNFI1A
HRAS Exons 2, 3	HSD3B1	ID3	IDH1 Exon 4	IDH2 Exon 4	IGF1R	IKBKE	IKZF1	INPP4B
IRF2	IRF4	IRS2	JAK1	JAK2 Exon 14	JAK3 Exons 5, 11, 12, 13, 15, 16	JUN	KDM5A	KDM5C
KDM6A	KDR	KEAP1	KEL	KIT Exons 8, 9, 11, 12, 13, 17, Intron 16	KLHL6	KMT2A (MLL) Introns 6, 8-11, Intron 7	KMT2D (MLL2)	KRAS

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

ORDERED TEST # ORD-1651473-01

APPENDIX

Genes assayed in FoundationOne® Liquid CDx

FoundationOne Liquid CDx interrogates 324 genes, including 309 genes with complete exonic (coding) coverage and 15 genes with only select non-coding coverage (indicated with an *); 75 genes (indicated in bold) are captured with increased sensitivity and have complete exonic (coding) coverage unless otherwise noted.

LTK	LYN	MAF	MAP2K1 (MEK1) Exons 2, 3	MAP2K2 (MEK2) Exons 2-4, 6, 7	MAP2K4	MAP3K1	MAP3K13	MAPK1
MCL1	MDM2	MDM4	MED12	MEF2B	MEN1	MERTK	MET	MITF
MKNK1	MLH1	MPL Exon 10	MRE11 (MRE11A)	MSH2 Intron 5	MSH3	MSH6	MST1R	MTAP
MTOR Exons 19, 30, 39, 40, 43-45, 47, 48, 53, 56	MUTYH	MYB* Intron 14	MYC Intron 1	MYCL (MYCL1)	MYCN	MYD88 Exon 4	NBN	NF1
NF2	NFE2L2	NFKBIA	NKX2-1	NOTCH1	NOTCH2 Intron 26	NOTCH3	NPM1 Exons 4-6, 8, 10	NRAS Exons 2, 3
NSD2 (WHSC1 or MMSET)	NSD3 (WHSC1L1)	NT5C2	NTRK1 Exons 14, 15, Introns 8-11	NTRK2 Intron 12	NTRK3 Exons 16, 17	NUTM1* Intron 1	P2RY8	PALB2
PARP1	PARP2	PARP3	PAX5	PBRM1	PDCD1 (PD-1)	PDCD1LG2 (PD-L2)	PDGFRA Exons 12, 18, Introns 7, 9, 11	PDGFRB Exons 12-21, 23 9, 11
PDK1	PIK3C2B	PIK3C2G	PIK3CA Exons 2, 3, 5-8, 10, 14, 19, 21 (Coding Exons 1, 2, 4-7, 9, 13, 18, 20)	PIK3CB	PIK3R1	PIM1	PMS2	POLD1
POLE	PPARG	PPP2R1A	PPP2R2A	PRDM1	PRKAR1A	PRKCI	PRKN (PARK2)	PTCH1
PTEN	PTPN11	PTPRO	QKI	RAC1	RAD21	RAD51	RAD51B	RAD51C
RAD51D	RAD52	RAD54L	RAF1 Exons 3, 4, 6, 7, 10, 14, 15, 17, Introns 4-8	RARA Intron 2	RB1	RBM10	REL	RET Introns 7, 8, Exons 11, 13-16, Introns 9-11
RICTOR	RNF43	ROS1 Exons 31, 36-38, 40, Introns 31-35	RPTOR	RSP02* Intron 1	SDC4* Intron 2	SDHA	SDHB	SDHC
SDHD	SETD2	SF3B1	SGK1	SLC34A2* Intron 4	SMAD2	SMAD4	SMARCA4	SMARCB1
SMO	SNCAIP	SOCS1	SOX2	SOX9	SPEN	SPOP	SRC	STAG2
STAT3	STK11	SUFU	SYK	TBX3	TEK	TENT5C (FAM46C)	TERC* ncRNA	TERT* Promoter
TET2	TGFBR2	TIPARP	TMPRSS2* Introns 1-3	TNFAIP3	TNFRSF14	TP53	TSC1	TSC2
TYRO3	U2AF1	VEGFA	VHL	WT1	XPO1	XRCC2	ZNF217	ZNF703

ADDITIONAL ASSAYS: FOR THE DETECTION OF SELECT CANCER BIOMARKERS

Microsatellite (MS) status
Blood Tumor Mutational Burden (bTMB)
Tumor Fraction

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Irene Shyu, M.D. | 21 June 2023
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1651473-01

APPENDIX

About FoundationOne® Liquid CDx

FoundationOne Liquid CDx fulfills the requirements of the European Directive 98/79 EC for in vitro diagnostic medical devices and is registered as a CE-IVD product by Foundation Medicine's EU Authorized Representative, Qarad b.v.b.a, Ciplastraat 3, 2440 Geel, Belgium. The CE-IVD regulatory status of FoundationOne Liquid CDx is applicable in countries that accept and/or recognize the CE mark.



ABOUT FOUNDATIONONE LIQUID CDx

FoundationOne Liquid CDx was developed and its performance characteristics determined by Foundation Medicine, Inc. (Foundation Medicine). FoundationOne Liquid CDx may be used for clinical purposes and should not be regarded as purely investigational or for research only. Foundation Medicine's clinical reference laboratories are qualified to perform high-complexity clinical testing.

Please refer to technical information for performance specification details.

INTENDED USE

FoundationOne Liquid CDx is a next generation sequencing based *in vitro* diagnostic device that analyzes 324 genes. Substitutions and insertion and deletion alterations (indels) are reported in 311 genes, copy number alterations (CNAs) are reported in 310 genes, and gene rearrangements are reported in 324 genes. The test also detects the genomic signatures blood tumor mutational burden (bTMB), microsatellite instability (MSI), and tumor fraction. FoundationOne Liquid CDx utilizes circulating cell-free DNA (cfDNA) isolated from plasma derived from the anti-coagulated peripheral whole blood of cancer patients. The test is intended to be used as a companion diagnostic to identify patients who may benefit from treatment with targeted therapies in accordance with the approved therapeutic product labeling. Additionally, FoundationOne Liquid CDx is intended to provide tumor mutation profiling to be used by qualified health care professionals in accordance with professional guidelines in oncology for patients with malignant neoplasms.

TEST PRINCIPLES

The FoundationOne Liquid CDx assay is performed exclusively as a laboratory service using circulating cell-free DNA (cfDNA) isolated from plasma derived from anti-coagulated peripheral whole blood from patients with solid malignant neoplasms. The assay employs a single DNA extraction method to obtain cfDNA from plasma from whole blood. Extracted

cfDNA undergoes whole-genome shotgun library construction and hybridization-based capture of 324 cancer-related genes including coding exons and select introns of 309 genes, as well as only select intronic regions or non-coding regions of 15 genes. Hybrid-capture selected libraries are sequenced with deep coverage using the NovaSeq® 6000 platform. Sequence data are processed using a customized analysis pipeline designed to accurately detect genomic alterations, including base substitutions, indels, select copy number variants, and select genomic rearrangements. Substitutions and insertion and deletion alterations (indels) are reported in 311 genes, copy number alterations (CNAs) are reported in 310 genes, and gene rearrangements are reported in 324 genes. The assay also reports tumor fraction, and genomic signatures including MSI and bTMB. A subset of targeted regions in 75 genes is baited for increased sensitivity.

THE REPORT

Incorporates analyses of peer-reviewed studies and other publicly available information identified by Foundation Medicine; these analyses and information may include associations between a molecular alteration (or lack of alteration) and one or more drugs with potential clinical benefit (or potential lack of clinical benefit), including drug candidates that are being studied in clinical research. *Note:* A finding of biomarker alteration does not necessarily indicate pharmacologic effectiveness (or lack thereof) of any drug or treatment regimen; a finding of no biomarker alteration does not necessarily indicate lack of pharmacologic effectiveness (or effectiveness) of any drug or treatment regimen.

QUALIFIED ALTERATION CALLS (EQUIVOCAL)

All equivocal calls, regardless of alteration type, imply that there is adequate evidence to call the alteration with confidence. However, the repeatability of equivocal calls may be lower than non-equivocal calls.

RANKING OF THERAPIES AND CLINICAL TRIALS

Ranking of Therapies in Summary Table

Therapies are ranked based on the following criteria: Therapies with clinical benefit (ranked alphabetically within each evidence category), followed by therapies associated with resistance (when applicable).

Ranking of Clinical Trials

Pediatric trial qualification → Geographical proximity → Later trial phase.

LIMITATIONS

1. For *in vitro* diagnostic use.
2. For prescription use only. This test must be ordered by a qualified medical professional in accordance with clinical laboratory regulations.
3. A negative result does not rule out the presence of a mutation below the limits of detection of the assay. Patients for whom no companion diagnostic alterations are detected should be considered for confirmation with an appropriately validated tumor tissue test, if available.
4. The FoundationOne Liquid CDx assay does not detect heterozygous deletions.
5. The test is not intended to provide information on cancer predisposition.
6. Performance has not been validated for cfDNA input below the specified minimum input.
7. Tissue TMB and blood TMB (bTMB) are estimated from the number of synonymous and nonsynonymous single-nucleotide variants (SNVs) and insertions and deletions (indels) per area of coding genome sampled, after the removal of known and likely oncogenic driver events and germline SNPs. Tissue TMB is calculated based on variants with an allele frequency of $\geq 5\%$, and bTMB is calculated based on variants with an allele frequency of $\geq 0.5\%$.
8. Tumor fraction is the percentage of circulating tumor DNA (ctDNA) present in a cell-free DNA (cfDNA) sample. The tumor fraction estimate is computationally derived from the observed level of aneuploidy in the sample. Tumor fraction is considered elevated when ctDNA levels are high enough that aneuploidy can be detected and is significantly distinct from that typically found in non-tumor samples.
9. Microsatellite instability (MSI) is a condition of genetic hypermutability that generates excessive amounts of short insertion/deletion mutations in the tumor genome; it generally occurs at microsatellite DNA sequences and is caused by a deficiency in DNA mismatch repair (MMR) in the tumor. The MSI algorithm is based on genome wide analysis of 1765 microsatellite loci and not based on the 5 or 7 MSI loci described in current clinical practice guidelines for solid tissue testing.
10. Genomic findings from circulating cell-free DNA (cfDNA) may originate from circulating tumor DNA fragments, germline alterations, or non-tumor somatic alterations, such as clonal hematopoiesis of indeterminate potential (CHIP). Genes with alterations that may be derived from CHIP include, but are not limited to: *ASXL1*, *ATM*, *CBL*, *CHEK2*, *DNMT3A*, *JAK2*,

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

ORDERED TEST # ORD-1651473-01

APPENDIX

About FoundationOne® Liquid CDx

KMT2D (MLL2), MPL, MYD88, SF3B1, TET2, TP53, and U2AF1.

11. Alterations reported may include somatic (not inherited) or germline (inherited) alterations; however, the test does not distinguish between germline and somatic alterations. If a reported alteration is suspected to be germline, confirmatory testing should be considered in the appropriate clinical context.
12. The test is not intended to replace germline testing or to provide information about cancer predisposition.

REPORT HIGHLIGHTS

The Report Highlights includes select genomic and therapeutic information with potential impact on patient care and treatment that is specific to the genomics and tumor type of the sample analyzed. This section may highlight information including targeted therapies with potential sensitivity or resistance; evidence-matched clinical trials; and variants with potential diagnostic, prognostic, nontargeted treatment, germline, or clonal hematopoiesis implications. Information included in the Report Highlights is expected to evolve with advances in scientific and clinical research. Findings included in the Report Highlights should be considered in the context of all other information in this report and other relevant patient information. Decisions on patient care and treatment are the responsibility of the treating physician.

VARIANTS TO CONSIDER FOR FOLLOW-UP GERMLINE TESTING

The variants indicated for consideration of follow-up germline testing are 1) limited to reportable short variants with a protein effect listed in the ClinVar genomic database (Landrum et al., 2018; 29165669) as Pathogenic, Pathogenic/Likely Pathogenic, or Likely Pathogenic (by an expert panel or multiple submitters), 2) associated with hereditary cancer-predisposing disorder(s), 3) detected at an allele frequency of >30%, and 4) in select genes reported by the ESMO Precision Medicine Working Group (Mandelker et al., 2019; 31050713) to have a greater than 10% probability of germline origin if identified during tumor sequencing. The selected genes are *ATM, BAP1, BRCA1, BRCA2, BRIP1, CHEK2, FH, FLCN, MLH1, MSH2, MSH6, MUTYH, PALB2, PMS2, POLE, RAD51C, RAD51D, RET, SDHA, SDHB, SDHC, SDHD, TSC2, and VHL*, and are not inclusive of all cancer susceptibility genes. The content in this report should not substitute for genetic counseling or follow-up germline testing, which is needed to distinguish whether a finding in this patient's

tumor sequencing is germline or somatic. Interpretation should be based on clinical context.

VARIANTS THAT MAY REPRESENT CLONAL HEMATOPOIESIS

Variants that may represent clonal hematopoiesis (CH) are limited to select reportable short variants in defined genes identified in solid tumors only. Variant selection was determined based on gene tumor-suppressor or oncogene status, known role in solid tumors versus hematological malignancies, and literature prevalence. The defined genes are *ASXL1, ATM, CBL, CHEK2, DNMT3A, IDH2, JAK2, KMT2D (MLL2), MPL, MYD88, SF3B1, TET2, and U2AF1* and are not inclusive of all CH genes. The content in this report should not substitute for dedicated hematological workup. Comprehensive genomic profiling of solid tumors detects nontumor alterations that are due to CH. Patient-matched peripheral blood mononuclear cell sequencing is required to conclusively determine if this alteration is present in tumor or is secondary to CH. Interpretation should be based on clinical context.

NATIONAL COMPREHENSIVE CANCER NETWORK® (NCCN®) CATEGORIZATION

Biomarker and genomic findings detected may be associated with certain entries within the NCCN Drugs & Biologics Compendium® (NCCN Compendium®) (www.nccn.org). The NCCN Categories of Evidence and Consensus indicated reflect the highest possible category for a given therapy in association with each biomarker or genomic finding. Please note, however, that the accuracy and applicability of these NCCN categories within a report may be impacted by the patient's clinical history, additional biomarker information, age, and/or co-occurring alterations. For additional information on the NCCN categories, please refer to the NCCN Compendium®. Referenced with permission from the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®). © National Comprehensive Cancer Network, Inc. 2023. All rights reserved. To view the most recent and complete version of the guidelines, go online to NCCN.org. NCCN makes no warranties of any kind whatsoever regarding their content, use or application and disclaims any responsibility for their application or use in any way.

LEVEL OF EVIDENCE NOT PROVIDED

Drugs with potential clinical benefit (or potential lack of clinical benefit) are not evaluated for source or level of published evidence.

NO GUARANTEE OF CLINICAL BENEFIT

This report makes no promises or guarantees that a particular drug will be effective in the treatment of

disease in any patient. This report also makes no promises or guarantees that a drug with potential lack of clinical benefit will in fact provide no clinical benefit.

NO GUARANTEE OF REIMBURSEMENT

Foundation Medicine makes no promises or guarantees that a healthcare provider, insurer or other third party payor, whether private or governmental, will reimburse a patient for the cost of FoundationOne Liquid CDx.

TREATMENT DECISIONS ARE THE RESPONSIBILITY OF PHYSICIAN

Drugs referenced in this Report may not be suitable for a particular patient. The selection of any, all or none of the drugs associated with potential clinical benefit (or potential lack of clinical benefit) resides entirely within the discretion of the treating physician. Indeed, the information in this Report must be considered in conjunction with all other relevant information regarding a particular patient, before the patient's treating physician recommends a course of treatment. Decisions on patient care and treatment must be based on the independent medical judgment of the treating physician, taking into consideration all applicable information concerning the patient's condition, such as patient and family history, physical examinations, information from other diagnostic tests, and patient preferences, in accordance with the standard of care in a given community. A treating physician's decisions should not be based on a single test, such as this test or the information contained in this report.

Certain sample of variant characteristics may result in reduced sensitivity. These include: low sample quality, deletions and insertions >40bp, or repetitive/high homology sequences. FoundationOne Liquid CDx is performed using cell-free DNA, and as such germline events may not be reported.

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Irene Shyu, M.D. | 21 June 2023
Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # ORD-1651473-01

APPENDIX

About FoundationOne®Liquid CDx

SELECT ABBREVIATIONS

ABBREVIATION	DEFINITION
CR	Complete response
DCR	Disease control rate
DNMT	DNA methyltransferase
HR	Hazard ratio
ITD	Internal tandem duplication
MMR	Mismatch repair
Muts/Mb	Mutations per megabase
NOS	Not otherwise specified
ORR	Objective response rate
OS	Overall survival
PD	Progressive disease
PFS	Progression-free survival
PR	Partial response
SD	Stable disease
TKI	Tyrosine kinase inhibitor

REFERENCE SEQUENCE INFORMATION

Sequence data is mapped to the human genome, Genome Reference Consortium Human Build 37 (GRCh37), also known as hg19.

MR Suite Version (RG) 7.9.0

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Irene Shyu, M.D. | 21 June 2023
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
 Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
 Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
 Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531
 Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 · CLIA: 22D2027531

ORDERED TEST # **ORD-1651473-01**
APPENDIX References

1. Gandara DR, et al. *Nat. Med.* (2018) PMID: 30082870
2. Wang Z, et al. *JAMA Oncol* (2019) PMID: 30816954
3. Sturgill EG, et al. *Oncologist* (2022) PMID: 35274716
4. Aggarwal C, et al. *Clin. Cancer Res.* (2020) PMID: 32102950
5. Schenker et al., 2022; AACR Abstract CT022
6. Saori et al., 2021; ESMO Abstract 80P
7. Chen EX, et al. *JAMA Oncol* (2020) PMID: 32379280
8. Rizvi NA, et al. *JAMA Oncol* (2020) PMID: 32271377
9. Si H, et al. *Clin. Cancer Res.* (2021) PMID: 33355200
10. Leighi NB, et al. *J Thorac Oncol* (2022) PMID: 34800700
11. Li et al., 2020; ASCO Abstract 6511
12. Xiao D, et al. *Oncotarget* (2016) PMID: 27009843
13. Spigel et al., 2016; ASCO Abstract 9017
14. Shim HS, et al. *J Thorac Oncol* (2015) PMID: 26200269
15. Govindan R, et al. *Cell* (2012) PMID: 22980976
16. Ding L, et al. *Nature* (2008) PMID: 18948947
17. Imielinski M, et al. *Cell* (2012) PMID: 22980975
18. Kim Y, et al. *J. Clin. Oncol.* (2014) PMID: 24323028
19. *Nature* (2012) PMID: 22810696
20. Stadler ZK, et al. *J. Clin. Oncol.* (2016) PMID: 27022117
21. Samowitz WS, et al. *Cancer Epidemiol. Biomarkers Prev.* (2001) PMID: 11535541
22. Elsaleh H, et al. *Clin Colorectal Cancer* (2001) PMID: 12445368
23. Brueckl WM, et al. *Anticancer Res.* () PMID: 12820457
24. Guidoboni M, et al. *Am. J. Pathol.* (2001) PMID: 11438476
25. Gryfe R, et al. *N. Engl. J. Med.* (2000) PMID: 10631274
26. Sinicrope FA, et al. *Gastroenterology* (2006) PMID: 16952542
27. Guastadisegni C, et al. *Eur. J. Cancer* (2010) PMID: 20627535
28. Laghi L, et al. *Dig Dis* (2012) PMID: 22722556
29. Mehnert JM, et al. *J. Clin. Invest.* (2016) PMID: 27159395
30. Cancer Genome Atlas Research Network, et al. *Nature* (2013) PMID: 23636398
31. Hussein YR, et al. *Mod. Pathol.* (2015) PMID: 25394778
32. Church DN, et al. *Hum. Mol. Genet.* (2013) PMID: 23528559
33. Cazier JB, et al. *Nat Commun* (2014) PMID: 24777035
34. Pfeifer GP, et al. *Mutat. Res.* (2005) PMID: 15748635
35. Hill VK, et al. *Annu Rev Genomics Hum Genet* (2013) PMID: 23875803
36. Pfeifer GP, et al. *Oncogene* (2002) PMID: 12379884
37. Rizvi NA, et al. *Science* (2015) PMID: 25765070
38. Johnson BE, et al. *Science* (2014) PMID: 24336570
39. Choi S, et al. *Neuro-oncology* (2018) PMID: 29452419
40. Briggs S, et al. *J. Pathol.* (2013) PMID: 23447401
41. Heitzner E, et al. *Curr. Opin. Genet. Dev.* (2014) PMID: 24583393
42. Roberts SA, et al. *Nat. Rev. Cancer* (2014) PMID: 25568919
43. Bronkhorst AJ, et al. *Biomol Detect Quantif* (2019) PMID: 30923679
44. Raja R, et al. *Clin. Cancer Res.* (2018) PMID: 30093454
45. Hrebien S, et al. *Ann. Oncol.* (2019) PMID: 30860573
46. Choudhury AD, et al. *JCI Insight* (2018) PMID: 30385733
47. Goodall J, et al. *Cancer Discov* (2017) PMID: 28450425
48. Goldberg SB, et al. *Clin. Cancer Res.* (2018) PMID: 29330207
49. Bettgowda C, et al. *Sci Transl Med* (2014) PMID: 24553385
50. Lapin M, et al. *J Transl Med* (2018) PMID: 30400802
51. Shulman DS, et al. *Br. J. Cancer* (2018) PMID: 30131550
52. Stover DG, et al. *J. Clin. Oncol.* (2018) PMID: 29298117
53. Hemming ML, et al. *JCO Precis Oncol* (2019) PMID: 30793095
54. Egyud M, et al. *Ann. Thorac. Surg.* (2019) PMID: 31059681
55. Fan G, et al. *PLoS ONE* (2017) PMID: 28187169
56. Vu et al., 2020; DOI: 10.1200/PO.19.00204
57. Li G, et al. *J Gastrointest Oncol* (2019) PMID: 31602320
58. Zhang EW, et al. *Cancer* (2020) PMID: 32757294
59. Butler TM, et al. *Cold Spring Harb Mol Case Stud* (2019) PMID: 30833418
60. Zauderer et al., 2018; ASCO Abstract 8515
61. LaFave LM, et al. *Nat. Med.* (2015) PMID: 26437366
62. Papadopoulos et al., 2022; ENA Abstract 188
63. Dudnik E, et al. *Front Oncol* (2021) PMID: 33777745
64. Fennell DA, et al. *Lancet Respir Med* (2021) PMID: 33515503
65. Tomczak A, et al. *Br J Cancer* (2022) PMID: 35986087
66. Lian BJ, et al. *Front Oncol* (2022) PMID: 35875073
67. Yu H, et al. *Proc. Natl. Acad. Sci. U.S.A.* (2014) PMID: 24347639
68. Ismail IH, et al. *Cancer Res.* (2014) PMID: 24894717
69. Peña-Llopis S, et al. *Nat. Genet.* (2012) PMID: 22683710
70. Nishi R, et al. *Nat. Cell Biol.* (2014) PMID: 25194926
71. Landreville S, et al. *Clin. Cancer Res.* (2012) PMID: 22038994
72. Cerami E, et al. *Cancer Discov* (2012) PMID: 22588877
73. Gao J, et al. *Sci Signal* (2013) PMID: 23550210
74. Gossage L, et al. *Genes Chromosomes Cancer* (2014) PMID: 24166983
75. Zauderer MG, et al. *J Thorac Oncol* (2013) PMID: 24128712
76. Jensen DE, et al. *Oncogene* (1998) PMID: 9528852
77. Fan LH, et al. *Clin Invest Med* (2012) PMID: 22863556
78. Tang J, et al. *Med. Oncol.* (2013) PMID: 23526420
79. Kapur P, et al. *Lancet Oncol.* (2013) PMID: 23333114
80. Hakimi AA, et al. *Eur. Urol.* (2013) PMID: 23036577
81. Ventii KH, et al. *Cancer Res.* (2008) PMID: 18757409
82. Chan-On W, et al. *Nat. Genet.* (2013) PMID: 24185513
83. Abdel-Rahman MH, et al. *J. Med. Genet.* (2011) PMID: 21941004
84. Testa JR, et al. *Nat. Genet.* (2011) PMID: 21874000
85. Wiesner T, et al. *Nat. Genet.* (2011) PMID: 21874003
86. Farley MN, et al. *Mol. Cancer Res.* (2013) PMID: 23709298
87. Aoude LG, et al. *PLoS ONE* (2013) PMID: 23977234
88. Peng H, et al. *Cancer Res* (2018) PMID: 29284740
89. Nishikawa H, et al. *Cancer Res* (2009) PMID: 19117993
90. Zhang Y, et al. *Nat Cell Biol* (2018) PMID: 30202049
91. Walpole S, et al. *J Natl Cancer Inst* (2018) PMID: 30517737
92. Boru G, et al. *Genes Chromosomes Cancer* (2019) PMID: 30883995
93. Garfield EM, et al. *J Am Acad Dermatol* (2018) PMID: 29753057
94. Shankar GM, et al. *Neuro Oncol* (2017) PMID: 28170043
95. Rai K, et al. *Genes Chromosomes Cancer* (2017) PMID: 27718540
96. Zauderer MG, et al. *J Thorac Oncol* (2019) PMID: 31323388
97. Marin-Acevedo et al., 2021; ASCO Abstract e21016
98. Johnson DB, et al. *Clin Cancer Res* (2020) PMID: 31924734
99. Nikanjam M, et al. *Mol Cancer Ther* (2021) PMID: 33722853
100. Nebhan CA, et al. *Oncologist* (2021) PMID: 33861486
101. Kim DW, et al. *Cancer* (2017) PMID: 27911979
102. Mazieres J, et al. *Ann. Oncol.* (2020) PMID: 31959346
103. Hainsworth JD, et al. *J. Clin. Oncol.* (2018) PMID: 29320312
104. Janku et al., 2021; AACR Abstract CT212
105. Sullivan RJ, et al. *Cancer Discov* (2018) PMID: 29247021
106. Davies H, et al. *Nature* (2002) PMID: 12068308
107. Hodis E, et al. *Cell* (2012) PMID: 22817889
108. Krauthammer M, et al. *Nat. Genet.* (2012) PMID: 22842228
109. Greaves WO, et al. *J Mol Diagn* (2013) PMID: 23273605
110. Brastianos PK, et al. *Nat. Genet.* (2014) PMID: 24413733
111. Choueiri TK, et al. *Eur. Urol.* (2012) PMID: 22727996
112. Tufano RP, et al. *Medicine (Baltimore)* (2012) PMID: 22932786
113. *Nat. Rev. Cancer* (2013) PMID: 23429735
114. Hsiao SJ, et al. *Endocr. Relat. Cancer* (2014) PMID: 24829266
115. *Nature* (2014) PMID: 25079552
116. Sia D, et al. *Gastroenterology* (2013) PMID: 23295441
117. Andersen JB, et al. *Gastroenterology* (2012) PMID: 22178589
118. Robertson S, et al. *Hum. Pathol.* (2013) PMID: 24139215
119. Van Cutsem E, et al. *J. Clin. Oncol.* (2011) PMID: 21502544
120. Bokemeyer C, et al. *Eur. J. Cancer* (2012) PMID: 22446022
121. Gavin PG, et al. *Clin. Cancer Res.* (2012) PMID: 23045248
122. Laurent-Puig P, et al. *J. Clin. Oncol.* (2009) PMID: 19884556
123. Ogino S, et al. *Clin. Cancer Res.* (2012) PMID: 22147942
124. Roth AD, et al. *J. Clin. Oncol.* (2010) PMID: 20008640
125. Douillard JY, et al. *N. Engl. J. Med.* (2013) PMID: 24024839
126. Hsu HC, et al. *Oncotarget* (2016) PMID: 26989027
127. Grisham RN, et al. *Cancer* (2013) PMID: 22930283
128. An et al., 2013; ASCO Abstract 8101
129. Chen D, et al. *PLoS ONE* (2014) PMID: 24979348
130. Schultz NA, et al. *Pancreas* (2012) PMID: 22699145
131. Elisei R, et al. *J. Clin. Endocrinol. Metab.* (2012) PMID: 23066120
132. Li C, et al. *J. Clin. Endocrinol. Metab.* (2012) PMID: 23055546
133. Zeiger MA, et al. *Ann. Surg. Oncol.* (2013) PMID: 22941167
134. Xing M, et al. *JAMA* (2013) PMID: 23571588
135. Xing M, et al. *J. Clin. Oncol.* (2015) PMID: 25332244
136. Gouveia C, et al. *JAMA Otolaryngol Head Neck Surg* (2013) PMID: 24030686
137. Zoghbi A, et al. *Eur Ann Otorhinolaryngol Head Neck Dis* (2014) PMID: 23845288
138. Long GV, et al. *J. Clin. Oncol.* (2011) PMID: 21343559
139. El-Osta H, et al. *PLoS ONE* (2011) PMID: 22039425
140. Ekedahl H, et al. *Br. J. Dermatol.* (2013) PMID: 23855428
141. Egberts F, et al. *Ann. Oncol.* (2014) PMID: 24276025
142. Holderfield M, et al. *Nat. Rev. Cancer* (2014) PMID: 24957944
143. Burotto M, et al. *Cancer* (2014) PMID: 24948110
144. Kandath C, et al. *Nature* (2013) PMID: 24132290
145. Wan PT, et al. *Cell* (2004) PMID: 15035987
146. Noeparast A, et al. *Oncotarget* (2017) PMID: 28947956
147. Smalley KS, et al. *Oncogene* (2009) PMID: 18794803
148. Smalley KS, et al. *Cancer Res.* (2009) PMID: 19351826
149. Ikenoue T, et al. *Cancer Res.* (2004) PMID: 15150094
150. Sen B, et al. *Sci Transl Med* (2012) PMID: 22649091
151. Kamata T, et al. *Cancer Res.* (2010) PMID: 20978199

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

ORDERED TEST # **ORD-1651473-01**
APPENDIX
References

152. Cardarella S, et al. Clin. Cancer Res. (2013) pmid: 23833300
153. Niihori T, et al. Nat. Genet. (2006) pmid: 16474404
154. Moretti S, et al. Biochim. Biophys. Acta (2009) pmid: 19735675
155. Rodriguez-Viciana P, et al. Meth. Enzymol. (2008) pmid: 18413255
156. Anastasaki C, et al. Dis Model Mech (2012) pmid: 22301711
157. Yao Z, et al. Nature (2017) pmid: 28783719
158. Sheikine Y, et al. JCO Precis Oncol (2018) pmid: 32913992
159. Garnett MJ, et al. Mol. Cell (2005) pmid: 16364920
160. Shibata T, et al. Cancer Res. (2010) pmid: 21062981
161. Bendavit G, et al. J. Biol. Chem. (2016) pmid: 27784786
162. Paik et al., 2019; ASCO Abstract 9085
163. Binkley MS, et al. Cancer Discov (2020) pmid: 33071215
164. Chowdhry S, et al. Oncogene (2013) pmid: 22964642
165. Abazeed ME, et al. Cancer Res. (2013) pmid: 23980093
166. Ahn SM, et al. Hepatology (2014) pmid: 24798001
167. Schulze K, et al. Nat. Genet. (2015) pmid: 25822088
168. Harding JJ, et al. Clin. Cancer Res. (2018) pmid: 30373752
169. Rizvi H, et al. J. Clin. Oncol. (2018) pmid: 29337640
170. Dulak AM, et al. Nat. Genet. (2013) pmid: 23525077
171. Cancer Genome Atlas Research Network, et al. Nature (2017) pmid: 28052061
172. Lin DC, et al. Nat. Genet. (2014) pmid: 24686850
173. Tate JG, et al. Nucleic Acids Res. (2019) pmid: 30371878
174. Shibata T, et al. Proc. Natl. Acad. Sci. U.S.A. (2008) pmid: 18757741
175. Sasaki H, et al. Int. J. Mol. Med. (2013) pmid: 23545629
176. Yang H, et al. Clin Lung Cancer (2011) pmid: 21663859
177. Ji L, et al. Int J Clin Exp Pathol (2014) pmid: 24695690
178. Hu XF, et al. Asian Pac. J. Cancer Prev. (2013) pmid: 24175806
179. Kawasaki Y, et al. BMC Cancer (2015) pmid: 25588809
180. Yu S, et al. Anticancer Agents Med Chem (2018) pmid: 29110625
181. Kawasaki Y, et al. Ann. Surg. Oncol. (2014) pmid: 24599410
182. Jiang X, et al. Mod Pathol (2022) pmid: 35194221
183. Park JY, et al. Pathology (2012) pmid: 23172081
184. Zhang J, et al. Biochem Biophys Res Commun (2016) pmid: 26987716
185. Wang XJ, et al. Carcinogenesis (2008) pmid: 18413364
186. Zhu H, et al. Front Oncol (2021) pmid: 34381706
187. Goeman F, et al. J Thorac Oncol (2019) pmid: 31323387
188. Leinonen HM, et al. Adv. Cancer Res. (2014) pmid: 24974185
189. Kobayashi A, et al. Mol. Cell. Biol. (2004) pmid: 15282312
190. Braun DA, et al. JAMA Oncol (2019) pmid: 31486842
191. Braun DA, et al. Nat Med (2020) pmid: 32472114
192. Miao D, et al. Science (2018) pmid: 29301960
193. Abou Alaiwi S, et al. Cancer Immunol Res (2020) pmid: 32321774
194. Hakimi AA, et al. Nat Commun (2020) pmid: 32820162
195. Miao D, et al. Nat. Genet. (2018) pmid: 30150660
196. Yang Q, et al. Ann Transl Med (2021) pmid: 33850862
197. Varela I, et al. Nature (2011) pmid: 21248752
198. Fujimoto A, et al. Nat Commun (2015) pmid: 25636086
199. Jiao Y, et al. Nat. Genet. (2013) pmid: 24185509
200. Churi CR, et al. PLoS ONE (2014) pmid: 25536104
201. Simbolo M, et al. Oncotarget (2014) pmid: 24867389
202. Nature (2014) pmid: 24476821
203. Robertson AG, et al. Cell (2017) pmid: 28988769
204. Pietzak EJ, et al. Eur. Urol. (2017) pmid: 28583311
205. Williams EA, et al. Acta Neuropathol. (2020) pmid: 32405805
206. Petrini I, et al. Nat. Genet. (2014) pmid: 24974848
207. da Costa WH, et al. BJU Int. (2014) pmid: 24053427
208. Pawłowski R, et al. Int. J. Cancer (2013) pmid: 22949125
209. Numata M, et al. Int. J. Oncol. (2013) pmid: 23229642
210. Hakimi AA, et al. Clin. Cancer Res. (2013) pmid: 23620406
211. Lemon B, et al. Nature () pmid: 11780067
212. Xia W, et al. Cancer Res. (2008) pmid: 18339845
213. Hopson S, et al. ACS Chem. Biol. (2017) pmid: 28921948
214. Porter EG, et al. J. Biol. Chem. (2017) pmid: 28053089
215. Niimi A, et al. Mutat. Res. (2015) pmid: 26117423
216. Brownlee PM, et al. Cell Rep (2014) pmid: 24613357
217. Kakarougkas A, et al. Mol. Cell (2014) pmid: 25066234
218. Gao W, et al. Proc. Natl. Acad. Sci. U.S.A. (2017) pmid: 28082722

Disclaimer: Foundation Medicine Inc. only provides PDF report as an official issuance of the test result. Any other transformed format is not an "official / formal solution" and not guarantee the accuracy of this conversion. It is suggested the hospital to verify the outputs and validate the suitability of use.

© 2023 Foundation Medicine, Inc. All rights reserved.

Electronically signed by Irene Shyu, M.D. | 21 June 2023
 Julia Elvin, M.D., Ph.D., Laboratory Director CLIA: 22D2027531
 Nimesh R. Patel, M.D., Laboratory Director CLIA: 34D2044309
 Foundation Medicine, Inc. | www.rochefoundationmedicine.com

Sample Preparation: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Sample Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531
Post-Sequencing Analysis: 150 Second St., 1st Floor, Cambridge, MA 02141 • CLIA: 22D2027531