



Sample Information

Patient Name: 張家恩  
Gender: Male  
ID No.: P120343983  
History No.: 41643612  
Age: 53

Ordering Doctor: DOC1326K 簡聖軒  
Ordering REQ.: H489M84  
Signing in Date: 2023/11/10

Path No.: M112-00289  
MP No.: MY23071  
Assay: Oncomine Myeloid Assay  
Sample Type: Bone Marrow  
Bone Marrow Aspirating Date: 2023/11/02

Reporting Doctor: DOC5444B 楊靜芬 (Phone: 8#5444)

Note:

Sample Cancer Type: Acute Myeloid Leukemia

Table of Contents	Page	Report Highlights
Variants (Exclude variant in Taiwan BioBank with >1% allele frequency)	1	1 Relevant Biomarkers
Biomarker Descriptions	2	5 Therapies Available
Relevant Therapy Summary	2	0 Clinical Trials
Relevant Therapy Details	3	

Relevant Biomarkers

Tier	Genomic Alteration	Relevant Therapies (In this cancer type)	Relevant Therapies (In other cancer type)	Clinical Trials
IA	IDH2 p.(R172K) c.515G>A isocitrate dehydrogenase (NADP(+)) 2 Allele Frequency: 21.70%	enasidenib <sup>1</sup> azacitidine decitabine enasidenib + chemotherapy venetoclax + chemotherapy	None	0

Public data sources included in relevant therapies: FDA<sup>1</sup>, NCCN, EMA<sup>2</sup>, ESMO

Variants (Exclude variant in Taiwan BioBank with >1% allele frequency)

DNA Sequence Variants								
Gene	Amino Acid Change	Coding	Variant ID	Locus	Allele Frequency	Transcript	Variant Effect	Coverage
IDH2	p.(R172K)	c.515G>A	COSM33733	chr15:90631838	21.70%	NM_002168.4	missense	1995

Disclaimer: The data presented here is from a curated knowledgebase of publicly available information, but may not be exhaustive. The data version is 2023.10(004). The content of this report has not been evaluated or approved by FDA, EMA or other regulatory agencies.

## Biomarker Descriptions

### IDH2 (isocitrate dehydrogenase (NADP(+)) 2)

**Background:** The IDH1 and IDH2 genes encode homologous isocitrate dehydrogenase enzymes that catalyze the conversion of isocitrate to  $\alpha$ -ketoglutarate ( $\alpha$ -KG)<sup>1</sup>. The IDH1 gene encodes the NADP+ dependent cytoplasmic isocitrate dehydrogenase enzyme; IDH2 encodes the mitochondrial isoform.

**Alterations and prevalence:** Recurrent somatic mutations in IDH1 and IDH2 are mutually exclusive and observed in several malignancies including glioma, chondrosarcoma, intrahepatic cholangiocarcinoma, acute myeloid leukemia (AML), and myelodysplastic syndrome (MDS)<sup>2</sup>. Recurrent IDH2 variants include predominately R140Q and R172K plus other substitutions at lower frequencies. These gain of function variants confer neomorphic enzyme activity<sup>3</sup>. Although wild-type enzymatic activity is ablated, recurrent IDH2 variants catalyze the conversion of  $\alpha$ -KG to D-2-hydroxyglutarate, an oncometabolite with diverse effects on cellular metabolism, epigenetic regulation, redox states, and DNA repair<sup>1,4</sup>. Recurrent IDH2 mutations are present in 10-20% of patients with AML and 5% of patients with MDS<sup>5,6,7</sup>.

**Potential relevance:** Enasidenib<sup>8</sup> is FDA approved (2017) for the treatment of AML patients with IDH2 R140G/L/Q/W and R172G/K/M/S/W mutations. In AML, acquired resistance to enasidenib has been associated with the emergence of Q316E or I319M mutations<sup>9</sup>. IDH2 R172 and R140Q variants are associated with poor prognosis in MDS but have been shown to confer improved prognosis in lower grade gliomas<sup>10,11,12</sup>. Additionally, IDH2 mutations are associated with inferior overall survival in polycythemia vera (PV) and essential thrombocythemia (ET) as well as inferior leukemia-free survival in primary myelofibrosis (PMF)<sup>13</sup>. Mutations in IDH2 are diagnostic of astrocytoma IDH-mutant and oligodendroglioma IDH-mutant and 1p/19q-codeleted subtypes of central nervous system (CNS) tumors<sup>14</sup>.

## Relevant Therapy Summary

☒ In this cancer type
 ☐ In other cancer type
 ☒ In this cancer type and other cancer types
 ☒ No evidence

### IDH2 p.(R172K) c.515G>A

Relevant Therapy	FDA	NCCN	EMA	ESMO	Clinical Trials*
enasidenib	●	●	×	●	×
azacitidine	×	●	×	×	×
decitabine	×	●	×	×	×
enasidenib + azacitidine	×	●	×	×	×
venetoclax + azacitidine	×	●	×	×	×
venetoclax + cytarabine	×	●	×	×	×
venetoclax + decitabine	×	●	×	×	×

## Relevant Therapy Details

### Current FDA Information

☒ In this cancer type    ☐ In other cancer type    ☒ In this cancer type and other cancer types

FDA information is current as of 2023-09-13. For the most up-to-date information, search [www.fda.gov](https://www.fda.gov).

#### IDH2 p.(R172K) c.515G>A

##### ☒ enasidenib

**Cancer type:** Acute Myeloid Leukemia

**Label as of:** 2020-11-24

**Variant class:** IDH2 R172K mutation

**Indications and usage:**

IDHIFA® is an isocitrate dehydrogenase-2 inhibitor indicated for the treatment of adult patients with relapsed or refractory acute myeloid leukemia (AML) with an isocitrate dehydrogenase-2 (IDH2) mutation as detected by an FDA-approved test.

**Reference:**

[https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2020/209606s004lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2020/209606s004lbl.pdf)

## Current NCCN Information

☒ In this cancer type    ☐ In other cancer type    ☒ In this cancer type and other cancer types

NCCN information is current as of 2023-09-01. For the most up-to-date information, search [www.nccn.org](http://www.nccn.org). For NCCN International Adaptations & Translations, search [www.nccn.org/global/what-we-do/international-adaptations](http://www.nccn.org/global/what-we-do/international-adaptations).

Some variant specific evidence in this report may be associated with a broader set of alterations from the NCCN Guidelines. Specific variants listed in this report were sourced from approved therapies or scientific literature. These therapeutic options are appropriate for certain population segments with cancer. Refer to the NCCN Guidelines® for full recommendation.

### IDH2 p.(R172K) c.515G>A

#### ☒ venetoclax + azacitidine

Cancer type: Acute Myeloid Leukemia

Variant class: IDH2 R172 mutation

NCCN Recommendation category: 1

Population segment (Line of therapy):

- (Induction therapy); Preferred intervention

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 4.2023]

#### ☒ enasidenib

Cancer type: Acute Myeloid Leukemia

Variant class: IDH2 R172 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Relapsed, Refractory (Line of therapy not specified)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 4.2023]

#### ☒ venetoclax + decitabine

Cancer type: Acute Myeloid Leukemia

Variant class: IDH2 R172 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Induction therapy); Preferred intervention

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 4.2023]

#### ☒ azacitidine

Cancer type: Acute Myeloid Leukemia

Variant class: IDH2 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Induction therapy); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 4.2023]

**IDH2 p.(R172K) c.515G>A (continued)****● decitabine**

Cancer type: Acute Myeloid Leukemia

Variant class: IDH2 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Induction therapy); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 4.2023]

**● enasidenib**

Cancer type: Acute Myeloid Leukemia

Variant class: IDH2 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Induction therapy); Preferred intervention

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 4.2023]

**● venetoclax + cytarabine**

Cancer type: Acute Myeloid Leukemia

Variant class: IDH2 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Induction therapy); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 4.2023]

**● enasidenib + azacitidine**

Cancer type: Acute Myeloid Leukemia

Variant class: IDH2 mutation

NCCN Recommendation category: 2B

Population segment (Line of therapy):

- (Induction therapy); Useful in certain circumstances

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 4.2023]

## Current ESMO Information

☒ In this cancer type    ☐ In other cancer type    ☐ In this cancer type and other cancer types

ESMO information is current as of 2023-09-01. For the most up-to-date information, search [www.esmo.org](http://www.esmo.org).

### IDH2 p.(R172K) c.515G>A

#### ☒ enasidenib

Cancer type: Acute Myeloid Leukemia

Variant class: IDH2 mutation

ESMO Level of Evidence/Grade of Recommendation: IV / B

Population segment (Line of therapy):

- Relapsed, Refractory (Second-line therapy)

Reference: ESMO Clinical Practice Guidelines - ESMO-Acute Myeloblastic Leukaemia in Adult Patients [Ann Oncol (2020); 31(6): 697-712.]

## References

1. Molenaar et al. Wild-type and mutated IDH1/2 enzymes and therapy responses. *Oncogene*. 2018 Apr;37(15):1949-1960. PMID: 29367755
2. Yan et al. IDH1 and IDH2 mutations in gliomas. *N. Engl. J. Med.* 2009 Feb 19;360(8):765-73. PMID: 19228619
3. Dang et al. Cancer-associated IDH1 mutations produce 2-hydroxyglutarate. *Nature*. 2009 Dec 10;462(7274):739-44. PMID: 19935646
4. Ward et al. The common feature of leukemia-associated IDH1 and IDH2 mutations is a neomorphic enzyme activity converting alpha-ketoglutarate to 2-hydroxyglutarate. *Cancer Cell*. 2010 Mar 16;17(3):225-34. PMID: 20171147
5. Paschka et al. IDH1 and IDH2 mutations are frequent genetic alterations in acute myeloid leukemia and confer adverse prognosis in cytogenetically normal acute myeloid leukemia with NPM1 mutation without FLT3 internal tandem duplication. *J. Clin. Oncol.* 2010 Aug 1;28(22):3636-43. PMID: 20567020
6. Chou et al. The prognostic impact and stability of Isocitrate dehydrogenase 2 mutation in adult patients with acute myeloid leukemia. *Leukemia*. 2011 Feb;25(2):246-53. PMID: 21079611
7. Marcucci et al. IDH1 and IDH2 gene mutations identify novel molecular subsets within de novo cytogenetically normal acute myeloid leukemia: a Cancer and Leukemia Group B study. *J. Clin. Oncol.* 2010 May 10;28(14):2348-55. PMID: 20368543
8. [https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2020/209606s004lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2020/209606s004lbl.pdf)
9. Intlekofer et al. Acquired resistance to IDH inhibition through trans or cis dimer-interface mutations. *Nature*. 2018 Jul;559(7712):125-129. PMID: 29950729
10. NCCN Guidelines® - NCCN-Myelodysplastic Syndromes [Version 1.2023]
11. Cancer Genome Atlas Research Network. Comprehensive, Integrative Genomic Analysis of Diffuse Lower-Grade Gliomas. *N Engl J Med*. 2015 Jun 25;372(26):2481-98. doi: 10.1056/NEJMoa1402121. Epub 2015 Jun 10. PMID: 26061751
12. Houillier et al. IDH1 or IDH2 mutations predict longer survival and response to temozolomide in low-grade gliomas. *Neurology*. 2010 Oct 26;75(17):1560-6. PMID: 20975057
13. NCCN Guidelines® - NCCN-Myeloproliferative Neoplasms [Version 2.2023]
14. Louis et al. The 2021 WHO Classification of Tumors of the Central Nervous System: a summary. *Neuro Oncol*. 2021 Aug 2;23(8):1231-1251. PMID: 34185076