



Sample Information

Patient Name: 何瑞源  
Gender: Male  
ID No.: A110972818  
History No.: 50117469  
Age: 65  
  
Ordering Doctor: DOC8131B 呂錯東  
Ordering REQ.: 0CVDXTC  
Signing in Date: 2023/12/28

Path No.: M112-00341  
MP No.: MY23086  
Assay: Oncomine Myeloid Assay  
Sample Type: Bone Marrow  
Bone Marrow Aspirating Date: 2023/12/25

Reporting Doctor: DOC5444B 楊靜芬 (Phone: 8#5444)

Note:

Sample Cancer Type: Acute Myeloid Leukemia

Table of Contents	Page	Report Highlights
Variants (Exclude variant in Taiwan BioBank with >1% allele frequency)	2	4 Relevant Biomarkers
Biomarker Descriptions	3	15 Therapies Available
Relevant Therapy Summary	4	0 Clinical Trials
Relevant Therapy Details	6	
Alert Details	21	

Relevant Biomarkers

Tier	Genomic Alteration	Relevant Therapies (In this cancer type)	Relevant Therapies (In other cancer type)	Clinical Trials
IA	IDH1 p.(R132C) c.394C>T, IDH1 p.(R132L) c.395G>T isocitrate dehydrogenase (NADP(+)) 1 Allele Frequency: 4.10%, 9.51% (2 variants)	ivosidenib 1 ivosidenib + chemotherapy 1,2 olutasidenib 1 azacitidine decitabine venetoclax + chemotherapy	ivosidenib 1 ivosidenib + chemotherapy 1	0

Public data sources included in relevant therapies: FDA1, NCCN, EMA2, ESMO

## Relevant Biomarkers (continued)

Tier	Genomic Alteration	Relevant Therapies (In this cancer type)	Relevant Therapies (In other cancer type)	Clinical Trials
IA	<b><i>RUNX1</i> p.(R204*) c.610C&gt;T</b> RUNX family transcription factor 1 Allele Frequency: 12.85%	allogeneic stem cells azacitidine cytarabine cytarabine + daunorubicin cytarabine + daunorubicin + etoposide cytarabine + etoposide + idarubicin cytarabine + fludarabine + idarubicin + filgrastim cytarabine + idarubicin cytarabine + mitoxantrone decitabine liposomal cytarabine-daunorubicin CPX-351 venetoclax + chemotherapy	None	0
	<b><i>ZRSR2</i> p.(R30Vfs*8) c.88delC</b> zinc finger CCCH-type, RNA binding motif and serine/arginine rich 2 Allele Frequency: 10.47%	allogeneic stem cells azacitidine cytarabine cytarabine + daunorubicin cytarabine + daunorubicin + etoposide cytarabine + etoposide + idarubicin cytarabine + fludarabine + idarubicin + filgrastim cytarabine + idarubicin cytarabine + mitoxantrone decitabine liposomal cytarabine-daunorubicin CPX-351 venetoclax + chemotherapy	None	0

Public data sources included in relevant therapies: FDA1, NCCN, EMA2, ESMO

### Prevalent cancer biomarkers without relevant evidence based on included data sources

*PHF6* p.(K241\*) c.721A>T

## Variants (Exclude variant in Taiwan BioBank with >1% allele frequency)

### DNA Sequence Variants

Gene	Amino Acid Change	Coding	Variant ID	Locus	Allele Frequency	Transcript	Variant Effect	Coverage
IDH1	p.(R132L)	c.395G>T	COSM28750	chr2:209113112	9.51%	NM_005896.3	missense	1998
IDH1	p.(R132C)	c.394C>T	COSM28747	chr2:209113113	4.10%	NM_005896.3	missense	1998
RUNX1	p.(R204*)	c.610C>T	.	chr21:36231774	12.85%	NM_001754.4	nonsense	2000
ZRSR2	p.(R30Vfs*8)	c.88delC	.	chrX:15809102	10.47%	NM_005089.3	frameshift Deletion	1996
PHF6	p.(K241*)	c.721A>T	.	chrX:133547988	28.20%	NM_032458.3	nonsense	2000
EZH2	p.(Y741C)	c.2222A>G	.	chr7:148504772	20.35%	NM_004456.5	missense	2000
ETV6	p.(N90_L96dup)	c.288_289insAATGG CAAAGCTCTCCTGCT G	.	chr12:11992175	13.67%	NM_001987.5	nonframeshift Insertion	1990

## Variants (Exclude variant in Taiwan BioBank with >1% allele frequency) (continued)

### DNA Sequence Variants (continued)

Gene	Amino Acid Change	Coding	Variant ID	Locus	Allele Frequency	Transcript	Variant Effect	Coverage
ETV6	p.(R103=)	c.309C>T	.	chr12:11992219	14.45%	NM_001987.5	synonymous	2000

## Biomarker Descriptions

### IDH1 p.(R132C) c.394C>T, IDH1 p.(R132L) c.395G>T

*isocitrate dehydrogenase (NADP(+)) 1*

**Background:** The IDH1 and IDH2 genes encode homologous isocitrate dehydrogenase enzymes that catalyze the conversion of isocitrate to  $\alpha$ -ketoglutarate ( $\alpha$ -KG)<sup>1</sup>. The IDH1 gene encodes the NADP+ dependent cytoplasmic isocitrate dehydrogenase enzyme; IDH2 encodes the mitochondrial isoform.

**Alterations and prevalence:** Recurrent somatic mutations in IDH1 and IDH2 are mutually exclusive and observed in several malignancies including glioma, chondrosarcoma, intrahepatic cholangiocarcinoma, acute myeloid leukemia (AML), and myelodysplastic syndrome (MDS)<sup>2</sup>. Recurrent IDH1 variants include predominately R132H/C plus other substitutions at lower frequencies. These gain of function variants confer neomorphic enzyme activity<sup>3</sup>. Although wild-type enzymatic activity is ablated, recurrent IDH1 variants catalyze the conversion of  $\alpha$ -KG to D-2-hydroxyglutarate, an oncometabolite with diverse effects on cellular metabolism, epigenetic regulation, redox states, and DNA repair<sup>1,4</sup>. Recurrent IDH1 mutations are present in 5-10% of patients with AML and 5% of patients with MDS<sup>5,6,7</sup>. Recurrent IDH1 mutations are present in nearly 80% of lower grade gliomas<sup>8,9</sup>.

**Potential relevance:** The IDH1 inhibitor, olutasidenib<sup>10</sup> is approved (2022) for the treatment of IDH1 R132C/G/H/L/S variants in AML. Ivosidenib<sup>11</sup> is also FDA approved (2018) for the treatment of AML or cholangiocarcinoma patients with IDH1 R132C/G/H/L/S variants<sup>12</sup>. Ivosidenib was granted breakthrough therapy designation (2020) for the treatment of IDH1 mutated relapsed or refractory myelodysplastic syndrome (MDS)<sup>13</sup>. IDH1 mutations are associated with inferior leukemia-free survival in primary myelofibrosis (PMF) and inferior overall survival in polycythemia vera (PV) but have been shown to confer improved prognosis in lower grade gliomas<sup>14,15,16</sup>. Mutations in IDH1 are diagnostic of astrocytoma IDH-mutant and oligodendroglioma IDH-mutant and 1p/19q-codeleted subtypes of central nervous system (CNS) tumors<sup>17</sup>.

### RUNX1 p.(R204\*) c.610C>T

*RUNX family transcription factor 1*

**Background:** The RUNX1 gene encodes the runt-related transcription factor (RUNX) 1, part of the RUNX family of transcription factors which also includes RUNX2 and RUNX3<sup>18</sup>. All RUNX proteins share several conserved regions with similar functionality including a highly conserved N-terminal 'runt' domain responsible for binding DNA, a C-terminal region composed of an activation domain, inhibitory domain, protein interacting motifs, and a nuclear matrix targeting signal<sup>19</sup>. Each of these proteins are capable of interacting with core binding factor beta (CBF $\beta$ ) to form the core binding factor (CBF) complex. Consequently, RUNX1, RUNX2, and RUNX3 are collectively known as core binding factor alpha (CBF $\alpha$ ) since they can each function as the alpha subunit of CBF. Specifically, CBF $\beta$  binds to the 'runt' domain of RUNX1 leading to RUNX1 stabilization and increased affinity of the CBF complex for promoters involved in hematopoietic differentiation and cell cycle regulation<sup>20,21</sup>. RUNX1 is frequently mutated in various hematological malignancies<sup>21</sup>. Germline mutations in RUNX1 result in a rare autosomal dominant condition known as familial platelet disorder, with predisposition to acute myeloid leukemia (FPD/AML)<sup>22,23</sup>. Somatic mutations and chromosomal translocations in RUNX1 are often observed in myelodysplastic syndrome (MDS), acute myeloid leukemia (AML), acute lymphoblastic leukemia (ALL), and chronic myelomonocytic leukemia (CMML)<sup>21</sup>.

**Alterations and prevalence:** RUNX1 is frequently rearranged in hematological malignancies with over 50 different observed translocations<sup>24</sup>. The most recurrent translocation, t(12;21)(q34;q11), results in ETV6-RUNX1 fusion and is observed in 20-25% of childhood ALL<sup>25,26,27</sup>. This translocation is also observed in adult ALL at a lower frequency (2%)<sup>26,27</sup>. Another recurrent translocation, t(8;21)(q22;q22), results in RUNX1-RUNX1T1 fusion and is observed in 5-10% of AML<sup>28</sup>. The RUNX1-RUNX1T1 fusion, consists of the RHD domain of RUNX1 and the majority of RUNX1T1, which promotes oncogenesis by altering transcriptional regulation of RUNX1 target genes<sup>21,28</sup>. Somatic mutations in RUNX1 include missense, nonsense, and frameshift mutations resulting in loss of function or dominant negative effects<sup>21</sup>. RUNX1 mutations are reported in approximately 10% of de novo AML as well as 10-15% of MDS<sup>8,21,29,30</sup>.

**Potential relevance:** AML with RUNX1-RUNX1T1 fusions is considered a distinct molecular subtype by the World Health Organization (WHO)<sup>29,31</sup>. Translocations involving RUNX1, specifically t(8;21)(q22;q22)/RUNX1-RUNX1T1 in AML and t(12;21)(q34;q11)/ETV6-

## Biomarker Descriptions (continued)

RUNX1 in ALL, are associated with favorable risk<sup>32,33</sup>. On the other hand, mutations in RUNX1 confer poor prognosis in AML, MDS, and systemic mastocytosis (SM)<sup>30,32,34</sup>.

### ZRSR2 p.(R30Vfs\*8) c.88delC

*zinc finger CCCH-type, RNA binding motif and serine/arginine rich 2*

**Background:** The ZRSR2 gene encodes the zinc finger CCCH-type, RNA binding motif and serine/arginine-rich 2 protein, a component of the spliceosome. Specifically, ZRSR2 encodes a splicing factor that is involved in the recognition of the 3' intron splice site<sup>35</sup>. ZRSR2 interacts with components of the pre-spliceosome assembly including SRSF2 and U2AF2/U2AF1 heterodimer<sup>35,36</sup>. Mutations in ZRSR2 can lead to deregulated global and alternative mRNA splicing, nuclear-cytoplasm export, and unspliced mRNA degradation while concurrently altering the expression of multiple genes<sup>35,37</sup>.

**Alterations and prevalence:** ZRSR2 alterations including nonsense and frameshift mutations are observed in 5-10% of myelodysplastic syndromes (MDS) and 4% of uterine cancer. ZRSR2 deletions are observed in 4% of diffuse large B-cell lymphoma (DLBCL), 3% of head and neck and esophageal cancers<sup>9,30</sup>.

**Potential relevance:** Mutation of ZRSR2 is associated with poor prognosis in myelodysplastic syndromes as well as poor/adverse risk in acute myeloid leukemia (AML)<sup>29,30,32</sup>.

### PHF6 p.(K241\*) c.721A>T

*PHD finger protein 6*

**Background:** The PHF6 gene encodes the plant homeodomain (PHD) finger protein 6 which contains four nuclear localization signals and two imperfect PHD zinc finger domains. PHF6 is a tumor suppressor that interacts with the nucleosome remodeling deacetylase (NuRD) complex, which regulates nucleosome positioning and transcription of genes involved in development and cell-cycle progression<sup>38,39</sup>.

**Alterations and prevalence:** The majority of PHF6 aberrations are nonsense, frameshift (70%), or missense (30%) mutations, which result in complete loss of protein expression<sup>38,40,41,42</sup>. Truncating or missense mutations in PHF6 are observed in 38% of adult and 16% of pediatric T-cell acute lymphoblastic leukemia (T-ALL), 20-25% of mixed phenotype acute leukemias (MPAL), and 3% of AML, and 2.6% of hepatocellular carcinoma (HCC)<sup>40,42</sup>. Missense mutations recurrently involve codon C215 and the second zinc finger domain of PHF6<sup>40</sup>. PHF6 mutations are frequently observed in hematologic malignancies from male patients<sup>38,40</sup>.

**Potential relevance:** Somatic mutations in PHF6 are associated with reduced overall survival in AML patients treated with high-dose induction chemotherapy<sup>43</sup>.

## Relevant Therapy Summary

☒ In this cancer type
 ☐ In other cancer type
 ☒ In this cancer type and other cancer types
 ☒ No evidence

### IDH1 p.(R132C) c.394C>T, IDH1 p.(R132L) c.395G>T

Relevant Therapy	FDA	NCCN	EMA	ESMO	Clinical Trials*
ivosidenib + azacitidine	<input checked="" type="radio"/>	<input checked="" type="radio"/>	<input checked="" type="radio"/>	<input checked="" type="radio"/>	<input checked="" type="radio"/>
ivosidenib	<input checked="" type="radio"/>	<input checked="" type="radio"/>	<input checked="" type="radio"/>	<input checked="" type="radio"/>	<input checked="" type="radio"/>
olutasidenib	<input checked="" type="radio"/>	<input checked="" type="radio"/>	<input checked="" type="radio"/>	<input checked="" type="radio"/>	<input checked="" type="radio"/>
azacitidine	<input checked="" type="radio"/>	<input checked="" type="radio"/>	<input checked="" type="radio"/>	<input checked="" type="radio"/>	<input checked="" type="radio"/>
decitabine	<input checked="" type="radio"/>	<input checked="" type="radio"/>	<input checked="" type="radio"/>	<input checked="" type="radio"/>	<input checked="" type="radio"/>
venetoclax + azacitidine	<input checked="" type="radio"/>	<input checked="" type="radio"/>	<input checked="" type="radio"/>	<input checked="" type="radio"/>	<input checked="" type="radio"/>
venetoclax + cytarabine	<input checked="" type="radio"/>	<input checked="" type="radio"/>	<input checked="" type="radio"/>	<input checked="" type="radio"/>	<input checked="" type="radio"/>

## Relevant Therapy Summary (continued)

☒ In this cancer type    
 ☐ In other cancer type    
 ☒ In this cancer type and other cancer types    
 ✕ No evidence

### IDH1 p.(R132C) c.394C>T, IDH1 p.(R132L) c.395G>T (continued)

Relevant Therapy	FDA	NCCN	EMA	ESMO	Clinical Trials*
venetoclax + decitabine	✕	●	✕	✕	✕

### RUNX1 p.(R204\*) c.610C>T

Relevant Therapy	FDA	NCCN	EMA	ESMO	Clinical Trials*
Allogeneic hematopoietic stem cell transplantation	✕	●	✕	✕	✕
azacitidine	✕	●	✕	✕	✕
cytarabine	✕	●	✕	✕	✕
cytarabine + daunorubicin	✕	●	✕	✕	✕
cytarabine + daunorubicin + etoposide	✕	●	✕	✕	✕
cytarabine + etoposide + idarubicin	✕	●	✕	✕	✕
cytarabine + fludarabine + idarubicin + filgrastim	✕	●	✕	✕	✕
cytarabine + idarubicin	✕	●	✕	✕	✕
cytarabine + mitoxantrone	✕	●	✕	✕	✕
decitabine	✕	●	✕	✕	✕
liposomal cytarabine-daunorubicin CPX-351	✕	●	✕	✕	✕
venetoclax + azacitidine	✕	●	✕	✕	✕
venetoclax + cytarabine	✕	●	✕	✕	✕
venetoclax + cytarabine + fludarabine + idarubicin + filgrastim	✕	●	✕	✕	✕
venetoclax + decitabine	✕	●	✕	✕	✕

### ZRSR2 p.(R30Vfs\*8) c.88delC

Relevant Therapy	FDA	NCCN	EMA	ESMO	Clinical Trials*
Allogeneic hematopoietic stem cell transplantation	✕	●	✕	✕	✕
azacitidine	✕	●	✕	✕	✕
cytarabine	✕	●	✕	✕	✕
cytarabine + daunorubicin	✕	●	✕	✕	✕
cytarabine + daunorubicin + etoposide	✕	●	✕	✕	✕
cytarabine + etoposide + idarubicin	✕	●	✕	✕	✕
cytarabine + fludarabine + idarubicin + filgrastim	✕	●	✕	✕	✕

## Relevant Therapy Summary (continued)

☒ In this cancer type
 ☐ In other cancer type
 ☒ In this cancer type and other cancer types
 ☒ No evidence

### ZRSR2 p.(R30Vfs\*8) c.88delC (continued)

Relevant Therapy	FDA	NCCN	EMA	ESMO	Clinical Trials*
cytarabine + idarubicin	×	●	×	×	×
cytarabine + mitoxantrone	×	●	×	×	×
decitabine	×	●	×	×	×
liposomal cytarabine-daunorubicin CPX-351	×	●	×	×	×
venetoclax + azacitidine	×	●	×	×	×
venetoclax + cytarabine	×	●	×	×	×
venetoclax + cytarabine + fludarabine + idarubicin + filgrastim	×	●	×	×	×
venetoclax + decitabine	×	●	×	×	×

## Relevant Therapy Details

### Current FDA Information

☒ In this cancer type
 ☐ In other cancer type
 ☒ In this cancer type and other cancer types

FDA information is current as of 2023-11-15. For the most up-to-date information, search [www.fda.gov](https://www.fda.gov).

### IDH1 p.(R132C) c.394C>T

#### ☒ ivosidenib, ivosidenib + azacitidine

**Cancer type:** Acute Myeloid Leukemia, Myelodysplastic Syndrome

**Label as of:** 2023-10-24

**Variant class:** IDH1 R132C mutation

#### Indications and usage:

TIBSOVO® is an isocitrate dehydrogenase-1 (IDH1) inhibitor indicated for the treatment of adult patients with a susceptible IDH1 mutation as detected by an FDA-approved test with:

#### Newly Diagnosed Acute Myeloid Leukemia (AML)

- In combination with azacitidine or as monotherapy for the treatment of newly diagnosed AML in adults 75 years or older, or who have comorbidities that preclude use of intensive induction chemotherapy.

#### Relapsed or refractory AML

- For the treatment of adult patients with relapsed or refractory AML.

#### Relapsed or refractory Myelodysplastic Syndromes (MDS)

- For the treatment of adult patients with relapsed or refractory myelodysplastic syndromes.

#### Locally Advanced or Metastatic Cholangiocarcinoma

- For the treatment of adult patients with locally advanced or metastatic cholangiocarcinoma who have been previously treated.

#### Reference:

[https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2023/211192s011lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2023/211192s011lbl.pdf)

**IDH1 p.(R132C) c.394C>T (continued)****● olutasidenib****Cancer type:** Acute Myeloid Leukemia**Label as of:** 2022-12-01**Variant class:** IDH1 R132C mutation**Indications and usage:**

REZLIDHIA™ is an isocitrate dehydrogenase-1 (IDH1) inhibitor indicated for the treatment of adult patients with relapsed or refractory acute myeloid leukemia (AML) with a susceptible IDH1 mutation as detected by an FDA-approved test.

**Reference:**

[https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2022/215814s000lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2022/215814s000lbl.pdf)

**IDH1 p.(R132L) c.395G>T****① ivosidenib, ivosidenib + azacitidine****Cancer type:** Acute Myeloid Leukemia,  
Myelodysplastic Syndrome**Label as of:** 2023-10-24**Variant class:** IDH1 R132L mutation**Indications and usage:**

TIBSOVO® is an isocitrate dehydrogenase-1 (IDH1) inhibitor indicated for the treatment of adult patients with a susceptible IDH1 mutation as detected by an FDA-approved test with:

Newly Diagnosed Acute Myeloid Leukemia (AML)

- In combination with azacitidine or as monotherapy for the treatment of newly diagnosed AML in adults 75 years or older, or who have comorbidities that preclude use of intensive induction chemotherapy.

Relapsed or refractory AML

- For the treatment of adult patients with relapsed or refractory AML.

Relapsed or refractory Myelodysplastic Syndromes (MDS)

- For the treatment of adult patients with relapsed or refractory myelodysplastic syndromes.

Locally Advanced or Metastatic Cholangiocarcinoma

- For the treatment of adult patients with locally advanced or metastatic cholangiocarcinoma who have been previously treated.

**Reference:**

[https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2023/211192s011lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2023/211192s011lbl.pdf)

**● olutasidenib****Cancer type:** Acute Myeloid Leukemia**Label as of:** 2022-12-01**Variant class:** IDH1 R132L mutation**Indications and usage:**

REZLIDHIA™ is an isocitrate dehydrogenase-1 (IDH1) inhibitor indicated for the treatment of adult patients with relapsed or refractory acute myeloid leukemia (AML) with a susceptible IDH1 mutation as detected by an FDA-approved test.

**Reference:**

[https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2022/215814s000lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2022/215814s000lbl.pdf)

## Current NCCN Information

- ☒ In this cancer type    ☐ In other cancer type    ☒ In this cancer type and other cancer types

NCCN information is current as of 2023-11-01. To view the most recent and complete version of the guideline, go online to NCCN.org.

For NCCN International Adaptations & Translations, search [www.nccn.org/global/what-we-do/international-adaptations](http://www.nccn.org/global/what-we-do/international-adaptations).

Some variant specific evidence in this report may be associated with a broader set of alterations from the NCCN Guidelines. Specific variants listed in this report were sourced from approved therapies or scientific literature. These therapeutic options are appropriate for certain population segments with cancer. Refer to the NCCN Guidelines® for full recommendation.

All guidelines cited below are referenced with permission from the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®) National Comprehensive Cancer Network, Inc. 2023. All rights reserved. NCCN makes no warranties regarding their content.

### IDH1 p.(R132C) c.394C>T, IDH1 p.(R132L) c.395G>T

#### ☒ ivosidenib + azacitidine

Cancer type: Acute Myeloid Leukemia

Variant class: IDH1 R132 mutation

NCCN Recommendation category: 1

Population segment (Line of therapy):

- (Induction therapy); Preferred intervention

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 6.2023]

#### ☒ venetoclax + azacitidine

Cancer type: Acute Myeloid Leukemia

Variant class: IDH1 R132 mutation

NCCN Recommendation category: 1

Population segment (Line of therapy):

- (Induction therapy); Preferred intervention

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 6.2023]

#### ☒ ivosidenib

Cancer type: Acute Myeloid Leukemia

Variant class: IDH1 R132 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Relapsed, Refractory (Line of therapy not specified)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 6.2023]



**IDH1 p.(R132C) c.394C>T, IDH1 p.(R132L) c.395G>T (continued)****olutasidenib**

Cancer type: Acute Myeloid Leukemia

Variant class: IDH1 R132 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Relapsed, Refractory (Line of therapy not specified)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 6.2023]

**venetoclax + decitabine**

Cancer type: Acute Myeloid Leukemia

Variant class: IDH1 R132 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Induction therapy); Preferred intervention

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 6.2023]

**azacitidine**

Cancer type: Acute Myeloid Leukemia

Variant class: IDH1 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Induction therapy); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 6.2023]

**decitabine**

Cancer type: Acute Myeloid Leukemia

Variant class: IDH1 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Induction therapy); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 6.2023]

**ivosidenib**

Cancer type: Acute Myeloid Leukemia

Variant class: IDH1 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Induction therapy); Preferred intervention

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 6.2023]

**IDH1 p.(R132C) c.394C>T, IDH1 p.(R132L) c.395G>T (continued)****● venetoclax + cytarabine**

Cancer type: Acute Myeloid Leukemia

Variant class: IDH1 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Induction therapy); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 6.2023]

**RUNX1 p.(R204\*) c.610C>T****● azacitidine**

Cancer type: Acute Myeloid Leukemia

Variant class: RUNX1 mutation

NCCN Recommendation category: 1

Population segment (Line of therapy):

- (Maintenance therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 6.2023]

**● cytarabine + daunorubicin**

Cancer type: Acute Myeloid Leukemia

Variant class: RUNX1 mutation

NCCN Recommendation category: 1

Population segment (Line of therapy):

- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 6.2023]

**● cytarabine + daunorubicin + etoposide**

Cancer type: Acute Myeloid Leukemia

Variant class: RUNX1 mutation

NCCN Recommendation category: 1

Population segment (Line of therapy):

- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 6.2023]

**RUNX1 p.(R204\*) c.610C>T (continued)****● cytarabine + etoposide + idarubicin**

Cancer type: Acute Myeloid Leukemia

Variant class: RUNX1 mutation

NCCN Recommendation category: 1

Population segment (Line of therapy):

- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 6.2023]

**● cytarabine + idarubicin**

Cancer type: Acute Myeloid Leukemia

Variant class: RUNX1 mutation

NCCN Recommendation category: 1

Population segment (Line of therapy):

- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 6.2023]

**● Allogeneic hematopoietic stem cell transplantation**

Cancer type: Acute Myeloid Leukemia

Variant class: RUNX1 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Consolidation therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 6.2023]

**● azacitidine**

Cancer type: Acute Myeloid Leukemia

Variant class: RUNX1 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Maintenance therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 6.2023]

**● cytarabine**

Cancer type: Acute Myeloid Leukemia

Variant class: RUNX1 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Consolidation therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 6.2023]

**RUNX1 p.(R204\*) c.610C>T (continued)****● cytarabine + mitoxantrone**

Cancer type: Acute Myeloid Leukemia

Variant class: RUNX1 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 6.2023]

**● decitabine**

Cancer type: Acute Myeloid Leukemia

Variant class: RUNX1 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 6.2023]

**● liposomal cytarabine-daunorubicin CPX-351**

Cancer type: Acute Myeloid Leukemia

Variant class: RUNX1 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Consolidation therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 6.2023]

**● venetoclax + azacitidine**

Cancer type: Acute Myeloid Leukemia

Variant class: RUNX1 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Consolidation therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 6.2023]

**● venetoclax + azacitidine**

Cancer type: Acute Myeloid Leukemia

Variant class: RUNX1 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 6.2023]

**RUNX1 p.(R204\*) c.610C>T (continued)****● venetoclax + cytarabine**

Cancer type: Acute Myeloid Leukemia

Variant class: RUNX1 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 6.2023]

**● venetoclax + decitabine**

Cancer type: Acute Myeloid Leukemia

Variant class: RUNX1 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Consolidation therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 6.2023]

**● venetoclax + decitabine**

Cancer type: Acute Myeloid Leukemia

Variant class: RUNX1 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 6.2023]

**● azacitidine**

Cancer type: Acute Myeloid Leukemia

Variant class: RUNX1 mutation

NCCN Recommendation category: 2B

Population segment (Line of therapy):

- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 6.2023]

**● cytarabine + daunorubicin + etoposide**

Cancer type: Acute Myeloid Leukemia

Variant class: RUNX1 mutation

NCCN Recommendation category: 2B

Population segment (Line of therapy):

- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 6.2023]

**RUNX1 p.(R204\*) c.610C>T (continued)****● cytarabine + etoposide + idarubicin**

Cancer type: Acute Myeloid Leukemia

Variant class: RUNX1 mutation

NCCN Recommendation category: 2B

Population segment (Line of therapy):

- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 6.2023]

**● cytarabine + fludarabine + idarubicin + filgrastim**

Cancer type: Acute Myeloid Leukemia

Variant class: RUNX1 mutation

NCCN Recommendation category: 2B

Population segment (Line of therapy):

- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 6.2023]

**● decitabine**

Cancer type: Acute Myeloid Leukemia

Variant class: RUNX1 mutation

NCCN Recommendation category: 2B

Population segment (Line of therapy):

- (Maintenance therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 6.2023]

**● venetoclax + cytarabine + fludarabine + idarubicin + filgrastim**

Cancer type: Acute Myeloid Leukemia

Variant class: RUNX1 mutation

NCCN Recommendation category: 3

Population segment (Line of therapy):

- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 6.2023]

**● venetoclax + decitabine**

Cancer type: Acute Myeloid Leukemia

Variant class: RUNX1 mutation

NCCN Recommendation category: 3

Population segment (Line of therapy):

- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 6.2023]

**ZRSR2 p.(R30Vfs\*8) c.88delC****● azacitidine**

Cancer type: Acute Myeloid Leukemia

Variant class: ZRSR2 mutation

NCCN Recommendation category: 1

Population segment (Line of therapy):

- (Maintenance therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 6.2023]

**● cytarabine + daunorubicin**

Cancer type: Acute Myeloid Leukemia

Variant class: ZRSR2 mutation

NCCN Recommendation category: 1

Population segment (Line of therapy):

- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 6.2023]

**● cytarabine + daunorubicin + etoposide**

Cancer type: Acute Myeloid Leukemia

Variant class: ZRSR2 mutation

NCCN Recommendation category: 1

Population segment (Line of therapy):

- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 6.2023]

**● cytarabine + etoposide + idarubicin**

Cancer type: Acute Myeloid Leukemia

Variant class: ZRSR2 mutation

NCCN Recommendation category: 1

Population segment (Line of therapy):

- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 6.2023]

**● cytarabine + idarubicin**

Cancer type: Acute Myeloid Leukemia

Variant class: ZRSR2 mutation

NCCN Recommendation category: 1

Population segment (Line of therapy):

- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 6.2023]

**ZRSR2 p.(R30Vfs\*8) c.88delC (continued)****● Allogeneic hematopoietic stem cell transplantation**

Cancer type: Acute Myeloid Leukemia

Variant class: ZRSR2 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Consolidation therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 6.2023]

**● azacitidine**

Cancer type: Acute Myeloid Leukemia

Variant class: ZRSR2 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Maintenance therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 6.2023]

**● cytarabine**

Cancer type: Acute Myeloid Leukemia

Variant class: ZRSR2 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Consolidation therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 6.2023]

**● cytarabine + mitoxantrone**

Cancer type: Acute Myeloid Leukemia

Variant class: ZRSR2 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 6.2023]

**● decitabine**

Cancer type: Acute Myeloid Leukemia

Variant class: ZRSR2 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 6.2023]



**ZRSR2 p.(R30Vfs\*8) c.88delC (continued)****● liposomal cytarabine-daunorubicin CPX-351**

Cancer type: Acute Myeloid Leukemia

Variant class: ZRSR2 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Consolidation therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 6.2023]

**● venetoclax + azacitidine**

Cancer type: Acute Myeloid Leukemia

Variant class: ZRSR2 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Consolidation therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 6.2023]

**● venetoclax + azacitidine**

Cancer type: Acute Myeloid Leukemia

Variant class: ZRSR2 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 6.2023]

**● venetoclax + cytarabine**

Cancer type: Acute Myeloid Leukemia

Variant class: ZRSR2 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 6.2023]

**● venetoclax + decitabine**

Cancer type: Acute Myeloid Leukemia

Variant class: ZRSR2 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Consolidation therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 6.2023]

**ZRSR2 p.(R30Vfs\*8) c.88delC (continued)****● venetoclax + decitabine**

Cancer type: Acute Myeloid Leukemia

Variant class: ZRSR2 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 6.2023]

**● azacitidine**

Cancer type: Acute Myeloid Leukemia

Variant class: ZRSR2 mutation

NCCN Recommendation category: 2B

Population segment (Line of therapy):

- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 6.2023]

**● cytarabine + daunorubicin + etoposide**

Cancer type: Acute Myeloid Leukemia

Variant class: ZRSR2 mutation

NCCN Recommendation category: 2B

Population segment (Line of therapy):

- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 6.2023]

**● cytarabine + etoposide + idarubicin**

Cancer type: Acute Myeloid Leukemia

Variant class: ZRSR2 mutation

NCCN Recommendation category: 2B

Population segment (Line of therapy):

- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 6.2023]

**● cytarabine + fludarabine + idarubicin + filgrastim**

Cancer type: Acute Myeloid Leukemia

Variant class: ZRSR2 mutation

NCCN Recommendation category: 2B

Population segment (Line of therapy):

- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 6.2023]

**ZRSR2 p.(R30Vfs\*8) c.88delC (continued)****● decitabine**

Cancer type: Acute Myeloid Leukemia

Variant class: ZRSR2 mutation

NCCN Recommendation category: 2B

Population segment (Line of therapy):

- (Maintenance therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 6.2023]

**● venetoclax + cytarabine + fludarabine + idarubicin + filgrastim**

Cancer type: Acute Myeloid Leukemia

Variant class: ZRSR2 mutation

NCCN Recommendation category: 3

Population segment (Line of therapy):

- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 6.2023]

**● venetoclax + decitabine**

Cancer type: Acute Myeloid Leukemia

Variant class: ZRSR2 mutation

NCCN Recommendation category: 3

Population segment (Line of therapy):

- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 6.2023]

## Current EMA Information

☒ In this cancer type    ☐ In other cancer type    ☒ In this cancer type and other cancer types

EMA information is current as of 2023-11-15. For the most up-to-date information, search [www.ema.europa.eu/ema](https://www.ema.europa.eu/ema).

### IDH1 p.(R132C) c.394C>T

#### ☒ ivosidenib + azacitidine

Cancer type: Acute Myeloid Leukemia

Label as of: 2023-05-12

Variant class: IDH1 R132C mutation

Reference:

[https://www.ema.europa.eu/en/documents/product-information/tibsovo-epar-product-information\\_en.pdf](https://www.ema.europa.eu/en/documents/product-information/tibsovo-epar-product-information_en.pdf)

### IDH1 p.(R132L) c.395G>T

#### ☒ ivosidenib + azacitidine

Cancer type: Acute Myeloid Leukemia

Label as of: 2023-05-12

Variant class: IDH1 R132L mutation

Reference:

[https://www.ema.europa.eu/en/documents/product-information/tibsovo-epar-product-information\\_en.pdf](https://www.ema.europa.eu/en/documents/product-information/tibsovo-epar-product-information_en.pdf)

## Current ESMO Information

☒ In this cancer type    ☐ In other cancer type    ☐ In this cancer type and other cancer types

ESMO information is current as of 2023-11-01. For the most up-to-date information, search [www.esmo.org](http://www.esmo.org).

### IDH1 p.(R132C) c.394C>T, IDH1 p.(R132L) c.395G>T

#### ivosidenib

Cancer type: Acute Myeloid Leukemia

Variant class: IDH1 mutation

ESMO Level of Evidence/Grade of Recommendation: IV / B

Population segment (Line of therapy):

- Relapsed, Refractory (Second-line therapy)

Reference: ESMO Clinical Practice Guidelines - ESMO-Acute Myeloblastic Leukaemia in Adult Patients [Ann Oncol (2020); 31(6): 697-712.]

## Alerts Informed By Public Data Sources

### Current FDA Information

 Contraindicated     Not recommended     Resistance     Breakthrough     Fast Track

FDA information is current as of 2023-11-15. For the most up-to-date information, search [www.fda.gov](http://www.fda.gov).

### IDH1 p.(R132C) c.394C>T

#### ivosidenib

Cancer type: Myelodysplastic Syndrome

Variant class: IDH1 mutation

Supporting Statement:

The FDA has granted Breakthrough Designation to the isocitrate dehydrogenase-1 inhibitor, ivosidenib, for the treatment of adult patients with relapsed or refractory myelodysplastic syndrome (MDS) with a susceptible IDH1 mutation as detected by an FDA-approved test.

Reference:

<https://investor.agios.com/news-releases/news-release-details/agios-receives-fda-breakthrough-therapy-designation-tibsovor-0>

### IDH1 p.(R132L) c.395G>T

#### ivosidenib

Cancer type: Myelodysplastic Syndrome

Variant class: IDH1 mutation

Supporting Statement:

The FDA has granted Breakthrough Designation to the isocitrate dehydrogenase-1 inhibitor, ivosidenib, for the treatment of adult patients with relapsed or refractory myelodysplastic syndrome (MDS) with a susceptible IDH1 mutation as detected by an FDA-approved test.

Reference:

<https://investor.agios.com/news-releases/news-release-details/agios-receives-fda-breakthrough-therapy-designation-tibsovor-0>

## References

1. Molenaar et al. Wild-type and mutated IDH1/2 enzymes and therapy responses. *Oncogene*. 2018 Apr;37(15):1949-1960. PMID: 29367755
2. Yan et al. IDH1 and IDH2 mutations in gliomas. *N. Engl. J. Med.* 2009 Feb 19;360(8):765-73. PMID: 19228619
3. Dang et al. Cancer-associated IDH1 mutations produce 2-hydroxyglutarate. *Nature*. 2009 Dec 10;462(7274):739-44. PMID: 19935646
4. Ward et al. The common feature of leukemia-associated IDH1 and IDH2 mutations is a neomorphic enzyme activity converting alpha-ketoglutarate to 2-hydroxyglutarate. *Cancer Cell*. 2010 Mar 16;17(3):225-34. PMID: 20171147
5. Paschka et al. IDH1 and IDH2 mutations are frequent genetic alterations in acute myeloid leukemia and confer adverse prognosis in cytogenetically normal acute myeloid leukemia with NPM1 mutation without FLT3 internal tandem duplication. *J. Clin. Oncol.* 2010 Aug 1;28(22):3636-43. PMID: 20567020
6. Chou et al. The prognostic impact and stability of Isocitrate dehydrogenase 2 mutation in adult patients with acute myeloid leukemia. *Leukemia*. 2011 Feb;25(2):246-53. PMID: 21079611
7. Marcucci et al. IDH1 and IDH2 gene mutations identify novel molecular subsets within de novo cytogenetically normal acute myeloid leukemia: a Cancer and Leukemia Group B study. *J. Clin. Oncol.* 2010 May 10;28(14):2348-55. PMID: 20368543
8. Weinstein et al. The Cancer Genome Atlas Pan-Cancer analysis project. *Nat. Genet.* 2013 Oct;45(10):1113-20. PMID: 24071849
9. Cerami et al. The cBio cancer genomics portal: an open platform for exploring multidimensional cancer genomics data. *Cancer Discov.* 2012 May;2(5):401-4. PMID: 22588877
10. [https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2022/215814s000lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2022/215814s000lbl.pdf)
11. [https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2023/211192s011lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2023/211192s011lbl.pdf)
12. Abou et al. The role of enasidenib in the treatment of mutant IDH2 acute myeloid leukemia. *Ther Adv Hematol*. 2018 Jul;9(7):163-173. PMID: 30013764
13. <https://investor.agios.com/news-releases/news-release-details/agios-receives-fda-breakthrough-therapy-designation-tibsovor-0>
14. NCCN Guidelines® - NCCN-Myeloproliferative Neoplasms [Version 3.2023]
15. Cancer Genome Atlas Research Network. Comprehensive, Integrative Genomic Analysis of Diffuse Lower-Grade Gliomas. *N Engl J Med*. 2015 Jun 25;372(26):2481-98. doi: 10.1056/NEJMoa1402121. Epub 2015 Jun 10. PMID: 26061751
16. Houillier et al. IDH1 or IDH2 mutations predict longer survival and response to temozolomide in low-grade gliomas. *Neurology*. 2010 Oct 26;75(17):1560-6. PMID: 20975057
17. Louis et al. The 2021 WHO Classification of Tumors of the Central Nervous System: a summary. *Neuro Oncol*. 2021 Aug 2;23(8):1231-1251. PMID: 34185076
18. de et al. Runx transcription factors in the development and function of the definitive hematopoietic system. *Blood*. 2017 Apr 13;129(15):2061-2069. PMID: 28179276
19. Chuang et al. RUNX family: Regulation and diversification of roles through interacting proteins. *Int. J. Cancer*. 2013 Mar 15;132(6):1260-71. PMID: 23180629
20. Jung et al. Prognostic factor analysis in core-binding factor-positive acute myeloid leukemia. *Anticancer Res*. 2014 Feb;34(2):1037-45. PMID: 24511052
21. Sood et al. Role of RUNX1 in hematological malignancies. *Blood*. 2017 Apr 13;129(15):2070-2082. PMID: 28179279
22. Béri-Dexheimer et al. Clinical phenotype of germline RUNX1 haploinsufficiency: from point mutations to large genomic deletions. *Eur. J. Hum. Genet.* 2008 Aug;16(8):1014-8. PMID: 18478040
23. Hayashi et al. Myeloid neoplasms with germ line RUNX1 mutation. *Int. J. Hematol*. 2017 Aug;106(2):183-188. PMID: 28534116
24. De et al. RUNX1 translocations and fusion genes in malignant hemopathies. *Future Oncol*. 2011 Jan;7(1):77-91. PMID: 21174539
25. De et al. ETV6 fusion genes in hematological malignancies: a review. *Leuk. Res*. 2012 Aug;36(8):945-61. PMID: 22578774
26. Pui et al. Acute lymphoblastic leukemia. *N. Engl. J. Med.* 2004 Apr 8;350(15):1535-48. PMID: 15071128
27. NCCN Guidelines® - Acute Lymphoblastic Leukemia [Version 2.2019]. 2019 May 15
28. Huret et al. Atlas of genetics and cytogenetics in oncology and haematology in 2013. *Nucleic Acids Res*. 2013 Jan;41(Database issue):D920-4. PMID: 23161685
29. NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 6.2023]
30. NCCN Guidelines® - NCCN-Myelodysplastic Syndromes [Version 2.2023]
31. Khoury et al. The 5th edition of the World Health Organization Classification of Haematolymphoid Tumours: Myeloid and Histiocytic/Dendritic Neoplasms. *Leukemia*. 2022 Jul;36(7):1703-1719. PMID: 35732831

## References (continued)

32. Döhner et al. Diagnosis and management of AML in adults: 2022 recommendations from an international expert panel on behalf of the ELN. *Blood*. 2022 Sep 22;140(12):1345-1377. PMID: 35797463
33. NCCN Guidelines® - NCCN-Acute Lymphoblastic Leukemia [Version 3.2023]
34. NCCN Guidelines® - NCCN-Systemic Mastocytosis [Version 1.2020]
35. Madan et al. Aberrant splicing of U12-type introns is the hallmark of ZRSR2 mutant myelodysplastic syndrome. *Nat Commun*. 2015 Jan 14;6:6042. doi: 10.1038/ncomms7042. PMID: 25586593
36. Tronchère et al. A protein related to splicing factor U2AF35 that interacts with U2AF65 and SR proteins in splicing of pre-mRNA. *Nature*. 1997 Jul 24;388(6640):397-400. PMID: 9237760
37. Chesnais et al. Spliceosome mutations in myelodysplastic syndromes and chronic myelomonocytic leukemia. *Oncotarget*. 2012 Nov;3(11):1284-93. PMID: 23327988
38. Wendorff et al. Phf6 Loss Enhances HSC Self-Renewal Driving Tumor Initiation and Leukemia Stem Cell Activity in T-ALL. *Cancer Discov*. 2019 Mar;9(3):436-451. PMID: 30567843
39. Lower et al. Mutations in PHF6 are associated with Börjeson-Forssman-Lehmann syndrome. *Nat. Genet*. 2002 Dec;32(4):661-5. PMID: 12415272
40. Van et al. PHF6 mutations in T-cell acute lymphoblastic leukemia. *Nat. Genet*. 2010 Apr;42(4):338-42. PMID: 20228800
41. Van et al. PHF6 mutations in adult acute myeloid leukemia. *Leukemia*. 2011 Jan;25(1):130-4. PMID: 21030981
42. Yoo et al. Somatic mutation of PHF6 gene in T-cell acute lymphoblastic leukemia, acute myelogenous leukemia and hepatocellular carcinoma. *Acta Oncol*. 2012 Jan;51(1):107-11. PMID: 21736506
43. Patel et al. Prognostic relevance of integrated genetic profiling in acute myeloid leukemia. *N. Engl. J. Med*. 2012 Mar 22;366(12):1079-89. PMID: 22417203