






REPORTING	PHYSICIAN	 <i>Complete Tumor Response Map on page 2</i>
Report Date: JAN-28-2024	Chih-Hsueh Chen	
Receipt Date: JAN-23-2024	Account: Genconn Biotech Co., LTD	
Collection Date: JAN-22-2024	Address: F15., No 207-5 Sec 3, Beixin Rd, Xindian Dist, New Taipei City, 23143, Taiwan	
Specimen: Blood	Ph: +886 963 820 633 Fax: N/A	
Status: FINAL	Additional Recipient: N/A	

Summary of Detected Somatic Alterations, Immunotherapy Biomarkers & Associated Treatment Options

KEY  Approved in indication  Approved in other indication  Lack of response

Detected Alteration(s) / Biomarker(s)	Associated FDA-approved therapies	Clinical trial availability (see page 3)	% cfDNA or Amplification
FGFR3-TACC3 Fusion	 Erdafitinib	Yes	2.0%

Comments


Reported by: DO

Additional Biomarkers

Biomarker	Additional Details
MSI-High	NOT DETECTED

Guardant360 Tumor Response Map

The Guardant360 Tumor Response Map illustrates the variant allele fraction (% cfDNA) of observed somatic variants at each sample submission. Amplifications are not plotted, and only the first and last five test dates are plotted. Please see the Physician Portal (portal.guardanthealth.com) for the Tumor Response Map with all test dates.



Detected Alteration(s) / Biomarker(s)	% cfDNA or Amp
FGFR3-TACC3 Fusion	2.0%

The table above annotates the variant allele fraction (% cfDNA) detected in this sample, listed in descending order.
§ See definitions section for more detail

Available Clinical Trials (within the same state as the ordering physician)

There may be additional trials not listed here. Visit: portal.guardanthealth.com or email clientservices@guardanthealth.com with A0949052 in the subject line of the email, for additional trials.

Alteration	Trial ID / Contact	Title	Phase	Site(s)
FGFR3-TACC3 Fusion	NCT04008797 Eisai Inquiry Service,eisai-chicken_hotline@hnc.eisai.co.jp	A Study of E7386 in Combination With Other Anticancer Drug in Participants With Solid Tumor	Phase 1	Tainan, Taiwan Taoyuan, Taiwan Kao-Hsiung, Taiwan Taipei, Taiwan (2) Additional trial sites available

Visit portal.guardanthealth.com for trials not within the same state as the physician's office

More clinical trial options available at portal.guardanthealth.com

Definitions

Fusion: Fusion events are gene rearrangements that fuse two previously distinct genes into a single genetic unit. Guardant360 detects fusions in the genes listed in Table 1.

Interpretation

Somatic alterations were detected in the circulating cell-free DNA isolated from this patient's blood specimen. These genomic alterations are cancer-associated somatic variants, some of which have been associated with either increased or reduced clinical response to specific treatments. The percentage of altered cell-free DNA circulating (% cfDNA) in blood is related to the unique tumor biology of each patient. Factors that may affect the % cfDNA of detected somatic alterations include tumor growth, turn over, size, heterogeneity, vascularization, disease progression, and treatment.

Method and Limitations

Guardant360 sequences 74 cancer-associated genes to identify somatic alterations. Cell-free DNA (cfDNA) is extracted from plasma, enriched for targeted regions, and sequenced using the Illumina platform and hg19 as the reference genome. All exons are sequenced in some genes; only clinically significant exons are sequenced in other genes. The types of genomic alterations detected by Guardant360 include single nucleotide variants, gene amplifications, fusions, short insertions/deletions (longest detected, 70 base pairs), and splice site disrupting events (see Table 1). Microsatellite Instability (MSI) is assessed for all cancer types by evaluating somatic changes in the length of repetitive sequences on the Guardant360 panel. A "Not Detected" result in samples where the highest % cfDNA is < 0.2% is an inconclusive result because it does not preclude MSI-High status in tissue. MSI status is currently not reported for earlier panel versions. This version of the Guardant360 test is not validated for the detection of other types of genomic alterations, such as complex rearrangements or gene deletions. Certain sample or variant characteristics, such as low cfDNA concentration, may result in reduced analytic sensitivity. Guardant360 cannot discern the source of circulating cfDNA, and for some variants in the range of ~40 to 60% cfDNA, the test cannot easily distinguish germline variants from somatic alterations. Guardant360 is not validated for the detection of germline or de novo variants that are associated with hereditary cancer risk. Tissue genotyping should be considered when plasma genotyping is negative, if clinically appropriate.

Table 1: Genes on the Guardant360 Panel

Guardant360 reports single nucleotide variants, splice site mutations, and insertion and deletion variants (indels) in all clinically relevant exons in 74 genes and reports other variant types in select genes as indicated below.

AKT1	ALK #	APC	AR †	ARAF	ARID1A	ATM	BRAF †	BRCA1
BRCA2	CCND1 †	CCND2 †	CCNE1 †	CDH1	CDK12	CDK4 †	CDK6 †	CDKN2A
CTNNB1	DDR2	EGFR †	ERBB2 †	ESR1	EZH2	FBXW7	FGFR1 †	FGFR2 # †
FGFR3 #	GATA3	GNA11	GNAQ	GNAS	HNF1A	HRAS	IDH1	IDH2
JAK2	JAK3	KIT †	KRAS †	MAP2K1	MAP2K2	MAPK1	MAPK3	MET †
MLH1	MPL	MTOR	MYC †	NF1	NFE2L2	NOTCH1	NPM1	NRAS
NTRK1 #	NTRK3	PDGFRA †	PIK3CA †	PTEN	PTPN11	RAF1 †	RB1	RET #
RHEB	RHOA	RIT1	ROS1 #	SMAD4	SMO	STK11	TERT ‡	TP53
TSC1	VHL							

‡ Guardant360 reports alterations in the promoter region of this gene.
Guardant360 reports fusion events involving this gene.
† Guardant360 reports amplifications of this gene.

About the Test

The Guardant360 assay was developed and its performance characteristics were determined by Guardant Health, Inc. This test has not been cleared or approved by the U.S. Food and Drug Administration (FDA). The FDA has determined that such clearance or approval is not necessary. This test may be used for clinical purposes and should not be regarded as investigational or for research only. Guardant Health's clinical reference laboratory is certified under the Clinical Laboratory Improvement Amendments of 1988 (CLIA) as qualified to perform high complexity clinical laboratory testing. The laboratory report should be interpreted in the context of other clinical information and laboratory, pathology, and imaging studies by a qualified medical professional prior to initiating or changing a patient's treatment plan. The selection of any, all, or none of the drugs associated with potential clinical benefit (or potential lack of clinical benefit) is entirely at the discretion of the treating medical professional. Drug and trial information are based on the diagnosis written on the submitted test request form; this information is not based on any supplemental information provided by the requesting medical professional, including pathology reports or other molecular studies. Some drugs listed in this report may not be approved or cleared by the FDA for the indicated use. Guardant Health makes no endorsement, express or implied, of any product, physician, or procedure contained in this report. This report makes no promises or guarantees that a particular medication will affect (or not affect) the clinical outcome of any patient.

Testing Performed at: Guardant Health

Laboratory Director: Martina Lefterova, MD PhD | CLIA ID: 05D2070300 | CAP #: 8765297 | 505 Penobscot Drive, Redwood City, CA, 94063, USA

Additional information is available

Any therapeutic annotations are based on publicly available information. This information is described in the "Detailed Therapy Results" and "Relevance of Detected Alterations" sections.

Visit portal.guardanthealth.com or email clientservices@guardanthealth.com with A0949052 in the subject line of the email for:

- Additional clinical trials
- Detailed Therapy Results
- Relevance of Detected Alterations
- References

If you would like to receive this additional information with every Guardant360 report, please call client services at [855.698.8887](tel:855.698.8887) to opt-in.

Additional information begins on the next page.

List of Available Clinical Trials

Alteration	Trial ID / Contact	Title	Phase	Site (number in parenthesis is count of trial sites)
FGFR3-TACC3 Fusion	NCT04008797 Eisai Inquiry Service,eisai-chicken_hotline@hhc.eisai.co.jp	A Study of E7386 in Combination With Other Anticancer Drug in Participants With Solid Tumor	Phase 1	Los Angeles, CA; Houston, TX; Oklahoma City, OK; West Palm Beach, FL; La Jolla, CA; New York, NY; Aurora, CO; Sarasota, FL; Nashville, TN; Charleston, SC; Santa Monica, CA; Kansas City, MO; Dallas, TX (2); Japan (12); Taiwan (6); Korea, Republic of (5); France (5)
	NCT04565275 Olivia Yang,olivia.yang@INNOCAREPHARMA.COM,+1 (609) 524-0684	A Study of ICP-192 in Patients With Advanced Solid Tumors	Phase 1 /Phase 2	Tucson, AZ; Houston, TX; Minneapolis, MN; La Jolla, CA; Columbus, OH; Lake Success, NY; Aurora, CO; Bronx, NY; Orange City, FL; Australia (7)
	NCT04601857 Taiho Oncology, INC, clinicaltrialinfo@taihooncology.com,1-609-250-7336	Futibatinib and Pembrolizumab Combination in the Treatment of Advanced or Metastatic Urothelial Carcinoma	Phase 2	Las Vegas, NV; Detroit, MI; San Francisco, CA; France (6); Spain (8)
	NCT05544552 Jennifer M Davis,TyraClinicalTrials@tyra.bio,(619)728-4805	Safety and Preliminary Anti-Tumor Activity of TYRA-300 in Advanced Urothelial Carcinoma and Other Solid Tumors With FGFR3 Gene Alterations	Phase 1 /Phase 2	Seattle, WA; Boston, MA; New York, NY; Worcester, MA; Nashville, TN; Durham, NC; France (3); Australia (6); Spain (3)
	NCT05614739 Patient Advocacy, clinicaltrials@loxooncology.com,855-569-6305	A Study of LOXO-435 in Participants With Cancer With a Change in a Gene Called FGFR3	Phase 1	Orlando, FL; Oklahoma City, OK; Philadelphia, PA; Saint Louis, MO; Chicago, IL; Baltimore, MD; Myrtle Beach, SC; Santa Monica, CA; Houston, TX; Duarte, CA; Detroit, MI; Chapel Hill, NC; Boston, MA; Atlanta, GA; Pittsburgh, PA; Dallas, TX; Salt Lake City, UT; Nashville, TN; New York, NY (3); Norway; United Kingdom; France; Canada (2); Japan (4); Korea, Republic of (4); Australia (2); Germany (2); Spain (2)
	NCT05627063 May Litt,may.litt@abbisko.com,+1 5033091243	A Study to Assess Safety, Tolerability, and Pharmacokinetics of ABSK121-NX in Patients With Advanced Solid Tumors	Phase 1	Las Vegas, NV; Detroit, MI; Canton, OH; China (6)

Detailed Therapy Results

Alteration	Drug	Trade Name	Target	Current Status
FGFR3-TACC3 Fusion	ABSK121-NX		Pan-Fgfr inhibitor.	Phase 1 (Solid Tumor)
	AL8326		Aurora kinase B/VEGFRs/Fgfr multi-kinase inhibitor.	Phase 1 (Solid Tumor) Phase 2 (Small cell lung carcinoma (SCLC))
	Anlotinib		Multi-tyrosine kinase inhibitor of VEGFR-1/2/3, Fgfr1-4, Pdgfr-beta, and Kit.	Phase 1 (Solid Tumor) Phase 3 (Gastric carcinoma, Non-small cell lung carcinoma (NSCLC), Thyroid carcinoma, Colorectal carcinoma (CRC), Soft tissue sarcoma)
	AZD4547		Pan-Fgfr inhibitor.	Phase 2 (Urothelial carcinoma) Phase 2 (Gastric carcinoma, Glioma, Non-small cell lung carcinoma (NSCLC), Gastroesophageal junction carcinoma, Esophageal carcinoma, Breast carcinoma, Multiple myeloma (MM), Lung cancer, Upper gastrointestinal carcinoma, Mesothelioma)
	B-701		Anti-Fgfr3 antibody.	Phase 2 (Urothelial carcinoma)
	Debio 1347		Fgfr1/2/3 inhibitor.	Phase 1 (Solid Tumor) Phase 2 (Breast carcinoma)
	Derazantinib		Fgfr1/2/3 inhibitor.	Phase 1 (Solid Tumor) Phase 2 (Cholangiocarcinoma)
	E7090		Fgfr1/2/3 kinase inhibitor.	Phase 1 (Solid Tumor) Phase 2 (Cholangiocarcinoma)
	Erdafitinib	Balversa	Pan-Fgfr inhibitor.	Phase 3 (Urothelial carcinoma) FDA Approved in other indications (Urothelial carcinoma with a susceptible FGFR3 alteration)
	Futibatinib	Lytgobi	Covalent pan-Fgfr kinase inhibitor.	Phase 2 (Urothelial carcinoma) FDA Approved in other indications (FGFR2-rearranged cholangiocarcinoma)
	Gunagratinib		Covalent pan-Fgfr inhibitor.	Phase 2 (Urothelial carcinoma) Phase 2 (Solid Tumor, Brain and Central Nervous System Tumors)
	HMPL-453		Fgfr1/2/3 inhibitor.	Phase 2 (Solid Tumor) Phase 2 (Brain and Central Nervous System Tumors)
	KIN-3248		Irreversible pan-Fgfr inhibitor.	Phase 1 (Urothelial carcinoma) Phase 1 (Solid Tumor, Brain and Central Nervous System Tumors)
	Lenvatinib	Lenvima	Multi-kinase inhibitor targeting VEGF, FGF, and SCF receptors.	Phase 3 (Urothelial carcinoma) FDA Approved in other indications (Thyroid carcinoma, well differentiated, Hepatocellular carcinoma (HCC))
	LOXO-435		Fgfr3-specific inhibitor.	Phase 1 (Urothelial carcinoma) Phase 1 (Solid Tumor)
	Lucitanib		Oral inhibitor of Fgfr1,2,3, VEGFR-1,2,3, Pdgfr-alpha, and Pdgfr-beta.	Phase 2 (Solid Tumor) Phase 2 (Small cell lung carcinoma (SCLC))
	LY2874455		Pan-Fgfr inhibitor.	Phase 1 (Solid Tumor) Phase 1 (Acute myeloid leukemia (AML))
	LY3076226		Anti-Fgfr3 antibody drug conjugate.	Phase 1 (Solid Tumor) Phase 1 (Bladder urothelial (transitional cell) carcinoma)
	Nintedanib	Vargatef	Multitargeted kinase inhibitor (VEGFRs, Fgfrs, Pdgfrs, Lyn, Lck, Src, Ret, Flt3).	Phase 2 (Urothelial carcinoma) FDA Approved in other indications (Idiopathic pulmonary fibrosis)
	ODM-203		Fgfr/VEGFR inhibitor.	Phase 1 (Solid Tumor)

Detailed Therapy Results

Alteration	Drug	Trade Name	Target	Current Status
	Pazopanib	Votrient	VEGFR-1,2,3/Kit/Pdgfrs/Fgfrs small molecule kinase inhibitor.	Phase 2 (Urothelial carcinoma) FDA Approved in other indications (Renal cell carcinoma, Soft tissue sarcoma)
	Pemigatinib	Pemazyre	Fgfr1/2/3 inhibitor.	Phase 2 (Urothelial carcinoma) FDA Approved in other indications (FGFR2-rearranged cholangiocarcinoma, FGFR1-rearranged myeloid /lymphoid neoplasm)
	Ponatinib	Iclusig	Bcr-Abl/VEGFR-1,2,3/Fgfrs/Kit/Tie-2/Flt3 kinase inhibitor.	Phase 2 (Solid Tumor) FDA Approved in other indications (Chronic myeloid leukemia, BCR-ABL1 positive, Acute lymphoblastic leukemia (Ph+) (ALL-Ph+))
	PRN1371		Pan-Fgfr inhibitor.	Phase 1 (Solid Tumor)
	Rogaratinib		Fgfr1/2/3 inhibitor.	Phase 3 (Urothelial carcinoma) Phase 2 (Lung squamous cell carcinoma)
	TT-00434		Fgfr1/2/3 inhibitor.	Phase 1 (Solid Tumor)
	TYRA-300		Fgfr3-specific inhibitor.	Phase 2 (Solid Tumor)

Relevance of Detected Alterations

Alteration	Role in Disease	Effect on Drug Sensitivity	Effect on Drug Resistance
FGFR3-TACC3 Fusion	Gain of function mutations in FGFRs have been reported in several cancer types. ^(1,2) . The presence of an FGFR3 abnormality (mutation, amplification, or overexpression) can result in an overabundance or overactivity of Fgfr3 protein, which may drive tumorigenesis. ⁽³⁻⁵⁾ . FGFR3 mutation has been associated with Fgfr3 overexpression in bladder urothelial carcinoma samples, although Fgfr3 overexpression without FGFR3 mutation has also been reported in muscle-invasive tumors. ⁽⁶⁻⁸⁾ . FGFR3 mutations have been reported to occur early during bladder urothelial transformation, and to be more frequent in low stage and low grade bladder lesions. ⁽⁸⁻¹⁶⁾ . FGFR3 mutations have been significantly correlated with TERT promoter mutations, and shorter relative telomere lengths as compared with non-mutated samples, in a cohort of 327 bladder urothelial carcinoma samples. ⁽¹⁷⁾ .	FGFR3 amplification or mutations may lead to activation of Fgfr3 and may therefore confer sensitivity to Fgfr family inhibitors. ^(3,18) . Erdafitinib is a pan-FGFR inhibitor that has been FDA-approved for the treatment of patients with locally advanced or metastatic urothelial carcinoma who experienced disease progression during or after at least one line of systemic therapy and have a susceptible FGFR3 alteration, as determined by an FDA-approved companion diagnostic test. Erdafitinib is not recommended for the treatment of patients who are eligible for and have not received prior PD-1/PD-L1 inhibitor therapy ⁽¹⁹⁾ . Several multi-kinase inhibitors that target Fgfrs, including pazopanib, ponatinib, and lenvatinib, have been approved for certain indications and continue to be studied in clinical trials. ⁽²⁰⁻²⁴⁾ . Additional agents that target Fgfrs are also being studied in clinical trials. ⁽²⁵⁻³¹⁾ .	Preclinical and clinical studies suggest that FGFR3-TACC3 fusion may result in resistance to Egfr-targeted therapies. ^(32,33) .

References

1. Eswarakumar V, Lax I, Schlessinger J "Cellular signaling by fibroblast growth factor receptors." *Cytokine & growth factor reviews*(2005): 139-49
2. Wesche J, Haglund K, Haugsten E "Fibroblast growth factors and their receptors in cancer." *The Biochemical journal*(2011): 199-213
3. Turner N, Grose R "Fibroblast growth factor signalling: from development to cancer." *Nature reviews. Cancer*(2010): 116-29
4. Plowright E, Li Z, Bergsagel P, Chesi M, Barber D, Branch D, Hawley R, Stewart A "Ectopic expression of fibroblast growth factor receptor 3 promotes myeloma cell proliferation and prevents apoptosis." *Blood*(2000): 992-8
5. Li Z, Zhu Y, Plowright E, Bergsagel P, Chesi M, Patterson B, Hawley T, Hawley R, Stewart A "The myeloma-associated oncogene fibroblast growth factor receptor 3 is transforming in hematopoietic cells." *Blood*(2001): 2413-9
6. Kang H, Kim Y, Jeong P, Park C, Kim W, Ryu D, Cha E, Ha Y, Kim T, Kwon T, Moon S, Choi Y, Yun S, Kim W "Expression levels of FGFR3 as a prognostic marker for the progression of primary pT1 bladder cancer and its association with mutation status." *Oncology letters*(2017): 3817-3824
7. Tomlinson D, Baldo O, Harnden P, Knowles M "FGFR3 protein expression and its relationship to mutation status and prognostic variables in bladder cancer." *The Journal of pathology*(2007): 91-8
8. Poyet C, Hermanns T, Zhong Q, Drescher E, Eberli D, Burger M, Hofstaedter F, Hartmann A, Stöhr R, Zwarthoff E, Sulser T, Wild P "Positive fibroblast growth factor receptor 3 immunoreactivity is associated with low-grade non-invasive urothelial bladder cancer." *Oncology letters*(2015): 2753-2760
9. Nassar A, Umeton R, Kim J, Lundgren K, Harshman L, Van Allen E, Preston M, Dong F, Bellmunt J, Mouw K, Choueiri T, Sonpavde G, Kwiatkowski D "Mutational Analysis of 472 Urothelial Carcinoma Across Grades and Anatomic Sites." *Clinical cancer research : an official journal of the American Association for Cancer Research*(2019): 2458-2470
10. van Oers J, Adam C, Denzinger S, Stoehr R, Bertz S, Zaak D, Stief C, Hofstaedter F, Zwarthoff E, van der Kwast T, Knuechel R, Hartmann A "Chromosome 9 deletions are more frequent than FGFR3 mutations in flat urothelial hyperplasias of the bladder." *International journal of cancer*(2006): 1212-5
11. van Rhijn B, Montironi R, Zwarthoff E, Jöbsis A, van der Kwast T "Frequent FGFR3 mutations in urothelial papilloma." *The Journal of pathology*(2002): 245-51
12. Hernández S, López-Knowles E, Lloreta J, Kogevinas M, Amorós A, Tardón A, Carrato A, Serra C, Malats N, Real F "Prospective study of FGFR3 mutations as a prognostic factor in nonmuscle invasive urothelial bladder carcinomas." *Journal of clinical oncology : official journal of the American Society of Clinical Oncology*(2006): 3664-71
13. Billerey C, Chopin D, Aubriot-Lorton M, Ricol D, Gil Diez de Medina S, Van Rhijn B, Bralet M, Lefrere-Belda M, Lahaye J, Abbou C, Bonaventure J, Zafrani E, van der Kwast T, Thiery J, Radvanyi F "Frequent FGFR3 mutations in papillary non-invasive bladder (pTa) tumors." *The American journal of pathology*(2001): 1955-9
14. Kimura T, Suzuki H, Ohashi T, Asano K, Kiyota H, Eto Y "The incidence of thanatophoric dysplasia mutations in FGFR3 gene is higher in low-grade or superficial bladder carcinomas." *Cancer*(2001): 2555-61
15. Pandith A, Hussain A, Khan M, Shah Z, Wani M, Siddiqi M "Oncogenic Activation of Fibroblast Growth Factor Receptor-3 and RAS Genes as Non-Overlapping Mutual Exclusive Events in Urinary Bladder Cancer." *Asian Pacific journal of cancer prevention : APJCP*(2016): 2787-93
16. Bagrodia A, Cha E, Sfakianos J, Zabor E, Bochner B, Al-Ahmadie H, Solit D, Coleman J, Iyer G, Scott S, Shah R, Ostrovnya I, Lee B, Desai N, Ren Q, Rosenberg J, Dalbagni G, Bajorin D, Reuter V, Berger M "Genomic Biomarkers for the Prediction of Stage and Prognosis of Upper Tract Urothelial Carcinoma." *The Journal of urology* (2016): 1684-1689
17. Hosen I, Rachakonda P, Heidenreich B, de Verdier P, Ryk C, Steineck G, Hemminki K, Kumar R "Mutations in TERT promoter and FGFR3 and telomere length in bladder cancer." *International journal of cancer*(2015): 1621-9
18. Socinski M "Multitargeted receptor tyrosine kinase inhibition: an antiangiogenic strategy in non-small cell lung cancer." *Cancer treatment reviews*(2011): 611-7
19. Llorca Y, Matsubara N, Park S, Huddart R, Burgess E, Houede N, Banek S, Guadalupe V, Ku J, Valderrama B, Tran B, Triantos S, Kean Y, Akapame S, Deprince K, Mukhopadhyay S, Stone N, Siefker-Radtke A "Erdafitinib or Chemotherapy in Advanced or Metastatic Urothelial Carcinoma." *The New England journal of medicine*(2023): 1961-1971
20. Schlumberger M, Tahara M, Wirth L, Robinson B, Brose M, Elisei R, Habra M, Newbold K, Shah M, Hoff A, Gianoukakis A, Kiyota N, Taylor M, Kim S, Krzyzanowska M, Dutcus C, de las Heras B, Zhu J, Sherman S "Lenvatinib versus placebo in radioiodine-refractory thyroid cancer." *The New England journal of medicine*(2015): 621-30
21. van der Graaf W, Blay J, Chawla S, Kim D, Bui-Nguyen B, Casali P, Schöffski P, Aglietta M, Staddon A, Beppu Y, Le Cesne A, Gelderblom H, Judson I, Araki N, Ouali M, Marreud S, Hodge R, Dewji M, Coens C, Demetri G, Fletcher C, Dei Tos A, Hohenberger P "Pazopanib for metastatic soft-tissue sarcoma (PALETTE): a randomised, double-blind, placebo-controlled phase 3 trial." *Lancet (London, England)*(2012): 1879-86
22. Sternberg C, Davis I, Mardiyah I, Szczyluk C, Lee E, Wagstaff J, Barrios C, Salman P, Gladkov O, Kavina A, Zarbá J, Chen M, McCann L, Pandite L, Roychowdhury D, Hawkins R "Pazopanib in locally advanced or metastatic renal cell carcinoma: results of a randomized phase III trial." *Journal of clinical oncology : official journal of the American Society of Clinical Oncology*(2010): 1061-8
23. Motzer R, Hutson T, Glen H, Michaelson M, Molina A, Eisen T, Jassem J, Zolnieriek J, Maroto J, Mellado B, Melichar B, Tomasek J, Kremer A, Kim H, Wood K, Dutcus C, Larkin J "Lenvatinib, everolimus, and the combination in patients with metastatic renal cell carcinoma: a randomised, phase 2, open-label, multicentre trial." *The Lancet. Oncology*(2015): 1473-1482
24. Cortes J, Kim D, Pinilla-Ibarz J, le Coutre P, Paquette R, Chuah C, Nicolini F, Apperley J, Khoury H, Talpaz M, DiPersio J, DeAngelo D, Abruzzese E, Rea D, Baccarani M, Müller M, Gambacorti-Passerini C, Wong S, Lustgarten S, Rivera V, Clackson T, Turner C, Haluska F, Guilhot F, Deininger M, Hochhaus A, Hughes T, Goldman J, Shah N, Kantarjian H "A phase 2 trial of ponatinib in Philadelphia chromosome-positive leukemias." *The New England journal of medicine*(2013): 1783-96
25. Nogova L, Sequist L, Perez Garcia J, Andre F, Delord J, Hidalgo M, Schellens J, Cassier P, Camidge D, Schuler M, Vaishampayan U, Burris H, Tian G, Campone M, Wainberg Z, Lim W, LoRusso P, Shapiro G, Parker K, Chen X, Choudhury S, Ringelsen F, Graus-Porta D, Porter D, Isaacs R, Buettner R, Wolf J "Evaluation of BGJ398, a Fibroblast Growth Factor Receptor 1-3 Kinase Inhibitor, in Patients With Advanced Solid Tumors Harboring Genetic Alterations in Fibroblast Growth Factor Receptors: Results of a Global Phase I, Dose-Escalation and Dose-Expansion Study." *Journal of clinical oncology : official journal of the American Society of Clinical Oncology*(2017): 157-165
26. Javle M, Lowery M, Shroff R, Weiss K, Springfield C, Borad M, Ramanathan R, Goyal L, Sadeghi S, Macarulla T, El-Khoueiry A, Kelley R, Borbath I, Choo S, Oh D, Philip P, Chen L, Reungwetwattana T, Van Cutsem E, Yeh K, Ciombor K, Finn R, Patel A, Sen S, Porter D, Isaacs R, Zhu A, Abou-Alfa G, Bekaii-Saab T "Phase II Study of BGJ398 in Patients With FGFR-Altered Advanced Cholangiocarcinoma." *Journal of clinical oncology : official journal of the American Society of Clinical Oncology*(2018): 276-282

References

27. Van Cutsem E, Bang Y, Mansoor W, Petty R, Chao Y, Cunningham D, Ferry D, Smith N, Frewer P, Ratnayake J, Stockman P, Kilgour E, Landers D "A randomized, open-label study of the efficacy and safety of AZD4547 monotherapy versus paclitaxel for the treatment of advanced gastric adenocarcinoma with FGFR2 polysomy or gene amplification." *Annals of oncology : official journal of the European Society for Medical Oncology*(2017): 1316-1324
28. Smyth EC, Turner NC, Peckitt C, et al. "Phase II multicenter proof of concept study of AZD4547 in FGFR amplified tumours." *J Clin Oncol*(2015): Abstract 2508
29. Tabernero J, Bahleda R, Dienstmann R, Infante J, Mita A, Italiano A, Calvo E, Moreno V, Adamo B, Gazzah A, Zhong B, Platero S, Smit J, Stuyckens K, Chatterjee-Kishore M, Rodon J, Peddareddigari V, Luo F, Soria J "Phase I Dose-Escalation Study of JNJ-42756493, an Oral Pan-Fibroblast Growth Factor Receptor Inhibitor, in Patients With Advanced Solid Tumors." *Journal of clinical oncology : official journal of the American Society of Clinical Oncology*(2015): 3401-8
30. Reck M, Kaiser R, Mellemaard A, Douillard J, Orlov S, Krzakowski M, von Pawel J, Gottfried M, Bondarenko I, Liao M, Gann C, Barrueco J, Gaschler-Markefski B, Novello S "Docetaxel plus nintedanib versus docetaxel plus placebo in patients with previously treated non-small-cell lung cancer (LUME-Lung 1): a phase 3, double-blind, randomised controlled trial." *The Lancet. Oncology*(2014): 143-55
31. Babina I, Turner N "Advances and challenges in targeting FGFR signalling in cancer." *Nature reviews. Cancer*(2017): 318-332
32. Daly C, Castanaro C, Zhang W, Zhang Q, Wei Y, Ni M, Young T, Zhang L, Burova E, Thurston G "FGFR3-TACC3 fusion proteins act as naturally occurring drivers of tumor resistance by functionally substituting for EGFR/ERK signaling." *Oncogene*(2017): 471-481
33. Ou S, Horn L, Cruz M, Vafai D, Lovly C, Spradlin A, Williamson M, Dagogo-Jack I, Johnson A, Miller V, Gadgeel S, Ali S, Schrock A "Emergence of FGFR3-TACC3 fusions as a potential by-pass resistance mechanism to EGFR tyrosine kinase inhibitors in EGFR mutated NSCLC patients." *Lung cancer (Amsterdam, Netherlands)*(2017): 61-64