Taipei Veterans General Hospital

Department of Pathology and Laboratory Medicine
No.201, Sec. 2, Shipai Rd., Beitou District, Taipei City, Taiwan
11217, R.O.C.
Tel: 02-2875-7449

Sample Information

Patient Name: 彭冉妹
Gender: Female
ID No.: K201667841
History No.: 49589426
Age: 68

Ordering Doctor: DOC8023E 李品臻
Ordering REQ.: 0CNUCRC
Signing in Date: 2023/07/22

Path No.: M112-00195
MP No.: MY23049
Assay: Oncomine Myeloid Assay
Sample Type: Bone Marrow
Bone Marrow Aspirating Date: 2023/07/24

Reporting Doctor: DOC5444B 楊靜芬 (Phone: 8#5444)

Note:

Sample Cancer Type: Myelodysplastic/Myeloproliferative Neoplasm

Table of Contents

	Page
Variants (Exclude variant in Taiwan BioBank with >1% allele frequency)	2
Biomarker Descriptions	2
Relevant Therapy Summary	4
Relevant Therapy Details	5

Report Highlights

2 Relevant Biomarkers
12 Therapies Available
0 Clinical Trials

Relevant Myelodysplastic/Myeloproliferative Neoplasm Variants

Gene	Finding
PDGFRA	None detected
PDGFRB	None detected

Relevant Biomarkers

Tier	Genomic Alteration	Relevant Therapies (In this cancer type)	Relevant Therapies (In other cancer type)	Clinical Trials
IIC	*ASXL1 p.(W1037Lfs*7)* *c.3110_3119delGGAACCAATC* ASXL transcriptional regulator 1 Allele Frequency: 48.03%	None	allogeneic stem cells azacitidine cytarabine cytarabine + daunorubicin + etoposide cytarabine + etoposide + idarubicin	0

Public data sources included in relevant therapies: FDA[1], NCCN, **EMA**[2], ESMO

Disclaimer: The data presented here is from a curated knowledgebase of publicly available information, but may not be exhaustive. The data version is 2023.06(005). The content of this report has not been evaluated or approved by FDA, EMA or other regulatory agencies.

Relevant Biomarkers (continued)

Tier	Genomic Alteration	Relevant Therapies (In this cancer type)	Relevant Therapies (In other cancer type)	Clinical Trials
			cytarabine + fludarabine + idarubicin + filgrastim daunorubicin decitabine idarubicin liposomal cytarabine-daunorubicin CPX-351 mitoxantrone venetoclax + chemotherapy	
IIC	*SRSF2 p.(P95L) c.284C>T* serine and arginine rich splicing factor 2 Allele Frequency: 28.13%	None	allogeneic stem cells azacitidine cytarabine cytarabine + daunorubicin + etoposide cytarabine + etoposide + idarubicin cytarabine + fludarabine + idarubicin + filgrastim daunorubicin decitabine idarubicin liposomal cytarabine-daunorubicin CPX-351 mitoxantrone venetoclax + chemotherapy	0

Public data sources included in relevant therapies: FDA[1], NCCN, **EMA**[2], ESMO

Prevalent cancer biomarkers without relevant evidence based on included data sources

TET2 p.(Q933) c.2797C>T*

Variants (Exclude variant in Taiwan BioBank with >1% allele frequency)

DNA Sequence Variants

Gene	Amino Acid Change	Coding	Variant ID	Locus	Allele Frequency	Transcript	Variant Effect	Coverage
TET2	p.(Q933*)	c.2797C>T	.	chr4:106157896	47.57%	NM_001127208.2	nonsense	1999
SRSF2	p.(P95L)	c.284C>T	COSM146288	chr17:74732959	28.13%	NM_003016.4	missense	672
ASXL1	p.(W1037Lfs*7)	c.3110_3119delGGAA CCAATC	.	chr20:31023624	48.03%	NM_015338.6	frameshift Deletion	1978
CBL	p.(E366K)	c.1096G>A	.	chr11:119148876	97.20%	NM_005188.4	missense	2000

Biomarker Descriptions

ASXL1 (ASXL transcriptional regulator 1)

Background: The ASXL1 gene encodes the ASXL transcriptional regulator 1 protein, a ligand-dependent co-activator and epigenetic scaffolding protein involved in transcriptional regulation[1,2]. ASXL1 belongs to the ASXL gene family, which also includes ASXL2 and ASXL3[2]. ASXL proteins contain a conserved c-terminal plant homeodomain (PHD) which facilitates interaction with DNA and histones[2,3]. ASXL1 influences chromatin remodeling and transcription through interaction with BAP1 and polycomb repressive complex (PRC) proteins, as well as other transcriptional activators and repressors[2,4]. In cancer, ASXL1 is the target of somatic mutations which

Biomarker Descriptions (continued)

often result in a truncated ASXL1 protein and loss of its PHD[5,6,7]. Such mutations can lead to impaired protein function and consequent upregulation of HOXA gene expression, supporting a tumor suppressor role for ASXL1[8].

Alterations and prevalence: Missense, nonsense, and frameshift mutations in ASXL1 are reported in 3-6% of de novo acute myeloid leukemia (AML), up to 36% of secondary AML, approximately 15% of myelodysplastic syndromes (MDS), up to 23% of myeloproliferative neoplasms (MPN), up to 30% of systemic mastocytosis (SM), and approximately 45% of chronic myelomonocytic leukemia (CMML)[4,9,10,11,12,13,14,15,16]. The ASXL1 G646Wfs*12 mutation accounts for over 50% of ASXL1 mutated cases in myeloid malignancies[6,11,17]. This mutation results from a single nucleotide expansion that occurs within an eight base pair guanine repeat that extends from c.1927 to c.1934. It is proposed that the high prevalence of the G646Wfs*12 variant is due to replication slippage which can occur in areas of repetitive sequence[18]. As a consequence, detection of G646Wfs*12 may result as an artifact of PCR and/or sequencing[19]. However, multiple studies observe an increase in the frequency of G646Wfs*12 in myeloid cancer relative to normal suggesting that G646Wfs*12 is a bona fide somatic mutation[9,18,20].

Potential relevance: The majority of frameshift and nonsense mutations in ASXL1 that result in protein truncation and removal of the PHD domain are considered pathogenic[21]. Mutations in ASXL1 confer poor/adverse risk in AML[16,22]. Additionally, ASXL1 nonsense or frameshift mutations are independently associated with poor prognosis in MDS and CMML[23]. Moreover, ASXL1 mutations are independently associated with inferior overall survival (OS) in patients with MPN or SM[24,25].

SRSF2 (serine and arginine rich splicing factor 2)

Background: The SRFS2 gene encodes the serine/arginine (SR)-rich splicing factor 2, a member of the SR-rich family of pre-mRNA splicing factors which make up part of the spliceosome. SRFS2 contains an RNA recognition motif (RRM) that recognizes and binds exonic splicing enhancers (ESE) in a sequence-specific manner[26]. SR proteins are essential regulators of alternative RNA splicing due to their ability to bind RNA and interact with other splicing factors. These proteins can influence the exclusion of cassette exons, a form of alternative splicing also known as exon skipping, which allows for the production of different protein isoforms[26,27]. SRSF2 is the target of somatic missense mutations and in-frame deletions in hematological malignancies, particularly myelodysplastic syndromes (MDS), chronic myelomonocytic leukemia (CMML), and myeloproliferative neoplasms (MPN)[28,29,30]. Such mutations in SRSF2 result in a differential gain of function which influences cassette exon exclusion, thereby supporting an oncogenic role in cancer[31].

Alterations and prevalence: Mutations in SRSF2 are observed in approximately 10% of MDS cases and 30-40% of CMML[29,32,33]. Missense mutations at P95 are most recurrent, which leads to an amino acid change from proline to histidine (H), leucine (L), or arginine (R)[33]. Specifically, the P95H substitution alters SRSF2 affinity for ESEs and drives preferential recognition of cassette exons containing C- versus G-rich ESEs[30,31]. Although less prevalent, recurrent in-frame deletions (P95H_R102del) are observed in primary myelofibrosis (PMF)[34]. This mutation results in the deletion of 8 amino acids which has been shown to exhibit greater variation of splicing events relative to the P95 missense mutation alone[35].

Potential relevance: Mutation of SRSF2 considered is one of the molecular abnormalities that defines acute myeloid leukemia, myelodysplasia related (AML-MR) according to the World Health Organization (WHO)[36]. SRSF2 mutations confer poor prognosis and risk in MDS, systemic mastocytosis (SM), and acute myeloid leukemia (AML) and are associated with decreased overall survival (OS)[16,22,23,25,37]. In MPN, SRSF2 mutations are considered high-risk mutations and are independently associated with inferior OS as well as leukemia-free survival[24,38]. Additionally, SRSF2 mutations are predictive of leukemic transformation in patients with PMF[24].

TET2 (tet methylcytosine dioxygenase 2)

Background: TET2 encodes the tet methylcytosine dioxygenase 2 protein and belongs to a family of ten-eleven translocation (TET) proteins that also includes TET1 and TET3[39]. TET2 is involved in DNA methylation, specifically in the conversion of 5-methylcytosine to 5-hydroxymethylcytosine[40,41]. The TET proteins contain a C-terminal core catalytic domain that contains a cysteine-rich domain and a double stranded ß-helix domain (DSBH)[42]. TET2 is a tumor suppressor gene. Loss of function mutations in TET2 are associated with loss of catalytic activity and transformation to hematological malignancies[39,40,41]

Alterations and prevalence: Somatic TET2 mutations, including nonsense, frameshift, splice site, and missense, are observed in 20-25% of myelodysplastic syndrome (MDS) associated diseases, including 40%-60% chronic myelomonocytic leukemia (CMML)[23]. TET2 mutations at H1881 and R1896 are frequently observed in myeloid malignancies[40,43]. TET2 mutations are also observed in 9% of uterine, 8% of melanoma and acute myeloid leukemia (AML), as well as 6% of diffuse large B-cell lymphoma (DLBCL).

Potential relevance: The presence of TET2 mutations may be used as one of the major diagnostic criteria in pre-primary myelofibrosis (pre-PMF) and overt PMF in the absence of JAK2/CALR/MPL mutations[24]. TET2 mutations are associated with poor prognosis in PMF and increased rate of transformation to leukemia[24,44]

Disclaimer: The data presented here is from a curated knowledgebase of publicly available information, but may not be exhaustive. The data version is 2023.06(005).

Relevant Therapy Summary

● In this cancer type ○ In other cancer type ◑ In this cancer type and other cancer types ✖ No evidence

ASXL1 p.(W1037Lfs*7) c.3110_3119delGGAACCAATC

Relevant Therapy	FDA	NCCN	EMA	ESMO	Clinical Trials*
Allogeneic hematopoietic stem cell transplantation	✖	○	✖	✖	✖
azacitidine	✖	○	✖	✖	✖
cytarabine	✖	○	✖	✖	✖
cytarabine + daunorubicin + etoposide	✖	○	✖	✖	✖
cytarabine + etoposide + idarubicin	✖	○	✖	✖	✖
cytarabine + fludarabine + idarubicin + filgrastim	✖	○	✖	✖	✖
daunorubicin	✖	○	✖	✖	✖
decitabine	✖	○	✖	✖	✖
idarubicin	✖	○	✖	✖	✖
liposomal cytarabine-daunorubicin CPX-351	✖	○	✖	✖	✖
mitoxantrone	✖	○	✖	✖	✖
venetoclax + azacitidine	✖	○	✖	✖	✖
venetoclax + cytarabine	✖	○	✖	✖	✖
venetoclax + cytarabine + fludarabine + idarubicin + filgrastim	✖	○	✖	✖	✖
venetoclax + decitabine	✖	○	✖	✖	✖

SRSF2 p.(P95L) c.284C>T

Relevant Therapy	FDA	NCCN	EMA	ESMO	Clinical Trials*
Allogeneic hematopoietic stem cell transplantation	✖	○	✖	✖	✖
azacitidine	✖	○	✖	✖	✖
cytarabine	✖	○	✖	✖	✖
cytarabine + daunorubicin + etoposide	✖	○	✖	✖	✖
cytarabine + etoposide + idarubicin	✖	○	✖	✖	✖
cytarabine + fludarabine + idarubicin + filgrastim	✖	○	✖	✖	✖
daunorubicin	✖	○	✖	✖	✖
decitabine	✖	○	✖	✖	✖
idarubicin	✖	○	✖	✖	✖
liposomal cytarabine-daunorubicin CPX-351	✖	○	✖	✖	✖
mitoxantrone	✖	○	✖	✖	✖

Disclaimer: The data presented here is from a curated knowledgebase of publicly available information, but may not be exhaustive. The data version is 2023.06(005).

Relevant Therapy Summary (continued)

● In this cancer type ○ In other cancer type ◑ In this cancer type and other cancer types ✖ No evidence

SRSF2 p.(P95L) c.284C>T (continued)

Relevant Therapy	FDA	NCCN	EMA	ESMO	Clinical Trials*
venetoclax + azacitidine	✖	○	✖	✖	✖
venetoclax + cytarabine	✖	○	✖	✖	✖
venetoclax + cytarabine + fludarabine + idarubicin + filgrastim	✖	○	✖	✖	✖
venetoclax + decitabine	✖	○	✖	✖	✖

Relevant Therapy Details

Current NCCN Information

● In this cancer type ○ In other cancer type ◑ In this cancer type and other cancer types

NCCN information is current as of 2023-05-01. For the most up-to-date information, search www.nccn.org.
For NCCN International Adaptations & Translations, search www.nccn.org/global/what-we-do/international-adaptations.

Some variant specific evidence in this report may be associated with a broader set of alterations from the NCCN Guidelines. Specific variants listed in this report were sourced from approved therapies or scientific literature. These therapeutic options are appropriate for certain population segments with cancer. Refer to the NCCN Guidelines® for full recommendation.

ASXL1 p.(W1037Lfs*7) c.3110_3119delGGAACCAATC

○ **azacitidine**

Cancer type: Acute Myeloid Leukemia **Variant class:** ASXL1 mutation

NCCN Recommendation category: 1

Population segment (Line of therapy):
- (Maintenance therapy); Preferred intervention
- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2023]

○ **cytarabine + daunorubicin + etoposide**

Cancer type: Acute Myeloid Leukemia **Variant class:** ASXL1 mutation

NCCN Recommendation category: 1

Population segment (Line of therapy):
- (Maintenance therapy); Preferred intervention
- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2023]

Disclaimer: The data presented here is from a curated knowledgebase of publicly available information, but may not be exhaustive. The data version is 2023.06(005).

ASXL1 p.(W1037Lfs*7) c.3110_3119delGGAACCAATC (continued)

cytarabine + etoposide + idarubicin

Cancer type: Acute Myeloid Leukemia **Variant class:** ASXL1 mutation

NCCN Recommendation category: 1

Population segment (Line of therapy):
- (Maintenance therapy); Preferred intervention
- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2023]

daunorubicin

Cancer type: Acute Myeloid Leukemia **Variant class:** ASXL1 mutation

NCCN Recommendation category: 1

Population segment (Line of therapy):
- (Maintenance therapy); Preferred intervention
- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2023]

idarubicin

Cancer type: Acute Myeloid Leukemia **Variant class:** ASXL1 mutation

NCCN Recommendation category: 1

Population segment (Line of therapy):
- (Maintenance therapy); Preferred intervention
- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2023]

Allogeneic hematopoietic stem cell transplantation

Cancer type: Acute Myeloid Leukemia **Variant class:** ASXL1 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):
- (Maintenance therapy)
- (Consolidation therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2023]

Disclaimer: The data presented here is from a curated knowledgebase of publicly available information, but may not be exhaustive. The data version is 2023.06(005).

ASXL1 p.(W1037Lfs*7) c.3110_3119delGGAACCAATC (continued)

azacitidine

Cancer type: Acute Myeloid Leukemia **Variant class:** ASXL1 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):
- (Maintenance therapy)
- (Consolidation therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2023]

cytarabine

Cancer type: Acute Myeloid Leukemia **Variant class:** ASXL1 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):
- (Maintenance therapy)
- (Consolidation therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2023]

decitabine

Cancer type: Acute Myeloid Leukemia **Variant class:** ASXL1 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):
- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2023]

liposomal cytarabine-daunorubicin CPX-351

Cancer type: Acute Myeloid Leukemia **Variant class:** ASXL1 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):
- (Maintenance therapy)
- (Consolidation therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2023]

Disclaimer: The data presented here is from a curated knowledgebase of publicly available information, but may not be exhaustive. The data version is 2023.06(005).

ASXL1 p.(W1037Lfs*7) c.3110_3119delGGAACCAATC (continued)

mitoxantrone

Cancer type: Acute Myeloid Leukemia **Variant class:** ASXL1 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):
- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2023]

venetoclax + azacitidine

Cancer type: Acute Myeloid Leukemia **Variant class:** ASXL1 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):
- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2023]

venetoclax + azacitidine

Cancer type: Acute Myeloid Leukemia **Variant class:** ASXL1 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):
- (Maintenance therapy)
- (Consolidation therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2023]

venetoclax + cytarabine

Cancer type: Acute Myeloid Leukemia **Variant class:** ASXL1 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):
- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2023]

venetoclax + decitabine

Cancer type: Acute Myeloid Leukemia **Variant class:** ASXL1 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):
- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2023]

Disclaimer: The data presented here is from a curated knowledgebase of publicly available information, but may not be exhaustive. The data version is 2023.06(005).

ASXL1 p.(W1037Lfs*7) c.3110_3119delGGAACCAATC (continued)

venetoclax + decitabine

Cancer type: Acute Myeloid Leukemia **Variant class:** ASXL1 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):
- (Maintenance therapy)
- (Consolidation therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2023]

azacitidine

Cancer type: Acute Myeloid Leukemia **Variant class:** ASXL1 mutation

NCCN Recommendation category: 2B

Population segment (Line of therapy):
- (Maintenance therapy)
- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2023]

cytarabine + daunorubicin + etoposide

Cancer type: Acute Myeloid Leukemia **Variant class:** ASXL1 mutation

NCCN Recommendation category: 2B

Population segment (Line of therapy):
- (Maintenance therapy)
- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2023]

cytarabine + etoposide + idarubicin

Cancer type: Acute Myeloid Leukemia **Variant class:** ASXL1 mutation

NCCN Recommendation category: 2B

Population segment (Line of therapy):
- (Maintenance therapy)
- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2023]

Disclaimer: The data presented here is from a curated knowledgebase of publicly available information, but may not be exhaustive. The data version is 2023.06(005).

ASXL1 p.(W1037Lfs*7) c.3110_3119delGGAACCAATC (continued)

cytarabine + fludarabine + idarubicin + filgrastim

Cancer type: Acute Myeloid Leukemia **Variant class:** ASXL1 mutation

NCCN Recommendation category: 2B

Population segment (Line of therapy):
- (Maintenance therapy)
- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2023]

decitabine

Cancer type: Acute Myeloid Leukemia **Variant class:** ASXL1 mutation

NCCN Recommendation category: 2B

Population segment (Line of therapy):
- (Maintenance therapy)
- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2023]

venetoclax + cytarabine + fludarabine + idarubicin + filgrastim

Cancer type: Acute Myeloid Leukemia **Variant class:** ASXL1 mutation

NCCN Recommendation category: 3

Population segment (Line of therapy):
- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2023]

venetoclax + decitabine

Cancer type: Acute Myeloid Leukemia **Variant class:** ASXL1 mutation

NCCN Recommendation category: 3

Population segment (Line of therapy):
- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2023]

Disclaimer: The data presented here is from a curated knowledgebase of publicly available information, but may not be exhaustive. The data version is 2023.06(005).

SRSF2 p.(P95L) c.284C>T

azacitidine

Cancer type: Acute Myeloid Leukemia **Variant class:** SRSF2 mutation

NCCN Recommendation category: 1

Population segment (Line of therapy):
- (Maintenance therapy); Preferred intervention
- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2023]

cytarabine + daunorubicin + etoposide

Cancer type: Acute Myeloid Leukemia **Variant class:** SRSF2 mutation

NCCN Recommendation category: 1

Population segment (Line of therapy):
- (Maintenance therapy); Preferred intervention
- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2023]

cytarabine + etoposide + idarubicin

Cancer type: Acute Myeloid Leukemia **Variant class:** SRSF2 mutation

NCCN Recommendation category: 1

Population segment (Line of therapy):
- (Maintenance therapy); Preferred intervention
- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2023]

daunorubicin

Cancer type: Acute Myeloid Leukemia **Variant class:** SRSF2 mutation

NCCN Recommendation category: 1

Population segment (Line of therapy):
- (Maintenance therapy); Preferred intervention
- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2023]

Disclaimer: The data presented here is from a curated knowledgebase of publicly available information, but may not be exhaustive. The data version is 2023.06(005).

SRSF2 p.(P95L) c.284C>T (continued)

idarubicin

Cancer type: Acute Myeloid Leukemia **Variant class:** SRSF2 mutation

NCCN Recommendation category: 1

Population segment (Line of therapy):
- (Maintenance therapy); Preferred intervention
- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2023]

Allogeneic hematopoietic stem cell transplantation

Cancer type: Acute Myeloid Leukemia **Variant class:** SRSF2 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):
- (Maintenance therapy)
- (Consolidation therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2023]

azacitidine

Cancer type: Acute Myeloid Leukemia **Variant class:** SRSF2 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):
- (Maintenance therapy)
- (Consolidation therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2023]

cytarabine

Cancer type: Acute Myeloid Leukemia **Variant class:** SRSF2 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):
- (Maintenance therapy)
- (Consolidation therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2023]

Disclaimer: The data presented here is from a curated knowledgebase of publicly available information, but may not be exhaustive. The data version is 2023.06(005).

SRSF2 p.(P95L) c.284C>T (continued)

decitabine

Cancer type: Acute Myeloid Leukemia **Variant class:** SRSF2 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):
- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2023]

liposomal cytarabine-daunorubicin CPX-351

Cancer type: Acute Myeloid Leukemia **Variant class:** SRSF2 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):
- (Maintenance therapy)
- (Consolidation therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2023]

mitoxantrone

Cancer type: Acute Myeloid Leukemia **Variant class:** SRSF2 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):
- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2023]

venetoclax + azacitidine

Cancer type: Acute Myeloid Leukemia **Variant class:** SRSF2 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):
- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2023]

venetoclax + azacitidine

Cancer type: Acute Myeloid Leukemia **Variant class:** SRSF2 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):
- (Maintenance therapy)
- (Consolidation therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2023]

Disclaimer: The data presented here is from a curated knowledgebase of publicly available information, but may not be exhaustive. The data version is 2023.06(005).

SRSF2 p.(P95L) c.284C>T (continued)

venetoclax + cytarabine

Cancer type: Acute Myeloid Leukemia **Variant class:** SRSF2 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):
- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2023]

venetoclax + decitabine

Cancer type: Acute Myeloid Leukemia **Variant class:** SRSF2 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):
- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2023]

venetoclax + decitabine

Cancer type: Acute Myeloid Leukemia **Variant class:** SRSF2 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):
- (Maintenance therapy)
- (Consolidation therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2023]

azacitidine

Cancer type: Acute Myeloid Leukemia **Variant class:** SRSF2 mutation

NCCN Recommendation category: 2B

Population segment (Line of therapy):
- (Maintenance therapy)
- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2023]

Disclaimer: The data presented here is from a curated knowledgebase of publicly available information, but may not be exhaustive. The data version is 2023.06(005).

SRSF2 p.(P95L) c.284C>T (continued)

cytarabine + daunorubicin + etoposide

Cancer type: Acute Myeloid Leukemia **Variant class:** SRSF2 mutation

NCCN Recommendation category: 2B

Population segment (Line of therapy):
- (Maintenance therapy)
- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2023]

cytarabine + etoposide + idarubicin

Cancer type: Acute Myeloid Leukemia **Variant class:** SRSF2 mutation

NCCN Recommendation category: 2B

Population segment (Line of therapy):
- (Maintenance therapy)
- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2023]

cytarabine + fludarabine + idarubicin + filgrastim

Cancer type: Acute Myeloid Leukemia **Variant class:** SRSF2 mutation

NCCN Recommendation category: 2B

Population segment (Line of therapy):
- (Maintenance therapy)
- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2023]

decitabine

Cancer type: Acute Myeloid Leukemia **Variant class:** SRSF2 mutation

NCCN Recommendation category: 2B

Population segment (Line of therapy):
- (Maintenance therapy)
- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2023]

Disclaimer: The data presented here is from a curated knowledgebase of publicly available information, but may not be exhaustive. The data version is 2023.06(005).

SRSF2 p.(P95L) c.284C>T (continued)

venetoclax + cytarabine + fludarabine + idarubicin + filgrastim

Cancer type: Acute Myeloid Leukemia **Variant class:** SRSF2 mutation

NCCN Recommendation category: 3

Population segment (Line of therapy):
- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2023]

venetoclax + decitabine

Cancer type: Acute Myeloid Leukemia **Variant class:** SRSF2 mutation

NCCN Recommendation category: 3

Population segment (Line of therapy):
- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2023]

Disclaimer: The data presented here is from a curated knowledgebase of publicly available information, but may not be exhaustive. The data version is 2023.06(005).

References

1. O'Leary et al. Reference sequence (RefSeq) database at NCBI: current status, taxonomic expansion, and functional annotation. Nucleic Acids Res. 2016 Jan 4;44(D1):D733-45. PMID: 26553804
2. Katoh. Functional and cancer genomics of ASXL family members. Br. J. Cancer. 2013 Jul 23;109(2):299-306. PMID: 23736028
3. Gelsi-Boyer et al. Mutations of polycomb-associated gene ASXL1 in myelodysplastic syndromes and chronic myelomonocytic leukaemia. Br. J. Haematol. 2009 Jun;145(6):788-800. PMID: 19388938
4. Gelsi-Boyer et al. Mutations in ASXL1 are associated with poor prognosis across the spectrum of malignant myeloid diseases. J Hematol Oncol. 2012 Mar 21;5:12. doi: 10.1186/1756-8722-5-12. PMID: 22436456
5. Larsson et al. The changing mutational landscape of acute myeloid leukemia and myelodysplastic syndrome. Mol. Cancer Res. 2013 Aug;11(8):815-27. PMID: 23645565
6. Alvarez et al. ASXL1 mutations in myeloid neoplasms: pathogenetic considerations, impact on clinical outcomes and survival. Curr Med Res Opin. 2018 May;34(5):757-763. PMID: 28027687
7. Yang et al. Gain of function of ASXL1 truncating protein in the pathogenesis of myeloid malignancies. Blood. 2018 Jan 18;131(3):328-341. PMID: 29113963
8. Abdel-Wahab et al. ASXL1 mutations promote myeloid transformation through loss of PRC2-mediated gene repression. Cancer Cell. 2012 Aug 14;22(2):180-93. PMID: 22897849
9. Alberti et al. Discriminating a common somatic ASXL1 mutation (c.1934dup; p.G646Wfs*12) from artifact in myeloid malignancies using NGS. Leukemia. 2018 Aug;32(8):1874-1878. PMID: 29959414
10. Kakosaiou et al. ASXL1 mutations in AML are associated with specific clinical and cytogenetic characteristics. Leuk. Lymphoma. 2018 Oct;59(10):2439-2446. PMID: 29411666
11. Paschka et al. ASXL1 mutations in younger adult patients with acute myeloid leukemia: a study by the German-Austrian Acute Myeloid Leukemia Study Group. Haematologica. 2015 Mar;100(3):324-30. PMID: 25596267
12. Weinstein et al. The Cancer Genome Atlas Pan-Cancer analysis project. Nat. Genet. 2013 Oct;45(10):1113-20. PMID: 24071849
13. Jawhar et al. The clinical and molecular diversity of mast cell leukemia with or without associated hematologic neoplasm. Haematologica. 2017 Jun;102(6):1035-1043. PMID: 28255023
14. Jawhar et al. KIT D816 mutated/CBF-negative acute myeloid leukemia: a poor-risk subtype associated with systemic mastocytosis. Leukemia. 2019 May;33(5):1124-1134. PMID: 30635631
15. Damaj et al. ASXL1 but not TET2 mutations adversely impact overall survival of patients suffering systemic mastocytosis with associated clonal hematologic non-mast-cell diseases. PLoS ONE. 2014;9(1):e85362. PMID: 24465546
16. NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2023]
17. Boultwood et al. Frequent mutation of the polycomb-associated gene ASXL1 in the myelodysplastic syndromes and in acute myeloid leukemia. Leukemia. 2010 May;24(5):1062-5. doi: 10.1038/leu.2010.20. Epub 2010 Feb 25. PMID: 20182461
18. Yannakou et al. ASXL1 c.1934dup;p.Gly646Trpfs*12-a true somatic alteration requiring a new approach. Blood Cancer J. 2017 Dec 15;7(12):656. doi: 10.1038/s41408-017-0025-8. PMID: 29242575
19. Abdel-Wahab et al. The most commonly reported variant in ASXL1 (c.1934dupG;p.Gly646TrpfsX12) is not a somatic alteration. Leukemia. 2010 Sep;24(9):1656-7. doi: 10.1038/leu.2010.144. Epub 2010 Jul 1. PMID: 20596031
20. Montes-Moreno et al. Clinical molecular testing for ASXL1 c.1934dupG p.Gly646fs mutation in hematologic neoplasms in the NGS era. PLoS ONE. 2018;13(9):e0204218. PMID: 30222780
21. Landrum et al. ClinVar: improving access to variant interpretations and supporting evidence. Nucleic Acids Res. 2018 Jan 4;46(D1):D1062-D1067. PMID: 29165669
22. Döhner et al. Diagnosis and management of AML in adults: 2022 recommendations from an international expert panel on behalf of the ELN. Blood. 2022 Sep 22;140(12):1345-1377. PMID: 35797463
23. NCCN Guidelines® - NCCN-Myelodysplastic Syndromes [Version 1.2023]
24. NCCN Guidelines® - NCCN-Myeloproliferative Neoplasms [Version 3.2022]
25. NCCN Guidelines® - NCCN-Systemic Mastocytosis [Version 1.2020]
26. Liang et al. SRSF2 mutations drive oncogenesis by activating a global program of aberrant alternative splicing in hematopoietic cells. Leukemia. 2018 Dec;32(12):2659-2671. PMID: 29858584
27. Cui et al. Comparative Analysis and Classification of Cassette Exons and Constitutive Exons. Biomed Res Int. 2017;2017:7323508. doi: 10.1155/2017/7323508. Epub 2017 Dec 4. PMID: 29349080
28. Meggendorfer et al. The mutational landscape of 18 investigated genes clearly separates four subtypes of myelodysplastic/myeloproliferative neoplasms. Haematologica. 2018 May;103(5):e192-e195. PMID: 29700173

Disclaimer: The data presented here is from a curated knowledgebase of publicly available information, but may not be exhaustive. The data version is 2023.06(005).

References (continued)

29. Arbab et al. Prognostic significance of SRSF2 mutations in myelodysplastic syndromes and chronic myelomonocytic leukemia: a meta-analysis. Hematology. 2018 Dec;23(10):778-784. PMID: 29757120
30. Kim et al. SRSF2 Mutations Contribute to Myelodysplasia by Mutant-Specific Effects on Exon Recognition. Cancer Cell. 2015 May 11;27(5):617-30. PMID: 25965569
31. Zhang et al. Disease-associated mutation in SRSF2 misregulates splicing by altering RNA-binding affinities. Proc. Natl. Acad. Sci. U.S.A. 2015 Aug 25;112(34):E4726-34. PMID: 26261309
32. Ethan et al. AACR Project GENIE: Powering Precision Medicine through an International Consortium. Cancer Discov. 2017 Aug;7(8):818-831. PMID: 28572459
33. Thol et al. Frequency and prognostic impact of mutations in SRSF2, U2AF1, and ZRSR2 in patients with myelodysplastic syndromes. Blood. 2012 Apr 12;119(15):3578-84. PMID: 22389253
34. Lasho et al. SRSF2 mutations in primary myelofibrosis: significant clustering with IDH mutations and independent association with inferior overall and leukemia-free survival. Blood. 2012 Nov 15;120(20):4168-71. PMID: 22968464
35. Komeno et al. SRSF2 Is Essential for Hematopoiesis, and Its Myelodysplastic Syndrome-Related Mutations Dysregulate Alternative Pre-mRNA Splicing. Mol. Cell. Biol. 2015 Sep 1;35(17):3071-82. PMID: 26124281
36. Khoury et al. The 5th edition of the World Health Organization Classification of Haematolymphoid Tumours: Myeloid and Histiocytic/Dendritic Neoplasms. Leukemia. 2022 Jul;36(7):1703-1719. PMID: 35732831
37. Jawhar et al. Splenomegaly, elevated alkaline phosphatase and mutations in the SRSF2/ASXL1/RUNX1 gene panel are strong adverse prognostic markers in patients with systemic mastocytosis. Leukemia. 2016 Dec;30(12):2342-2350. PMID: 27416984
38. Vannucchi et al. Mutations and prognosis in primary myelofibrosis. Leukemia. 2013 Sep;27(9):1861-9. PMID: 23619563
39. Pan et al. The TET2 interactors and their links to hematological malignancies. IUBMB Life. 2015 Jun;67(6):438-45. PMID: 26099018
40. Ko et al. Impaired hydroxylation of 5-methylcytosine in myeloid cancers with mutant TET2. Nature. 2010 Dec 9;468(7325):839-43. PMID: 21057493
41. Solary et al. The Ten-Eleven Translocation-2 (TET2) gene in hematopoiesis and hematopoietic diseases. Leukemia. 2014 Mar;28(3):485-96. PMID: 24220273
42. An et al. TET family dioxygenases and DNA demethylation in stem cells and cancers. Exp. Mol. Med. 2017 Apr 28;49(4):e323. PMID: 28450733
43. Kosmider et al. TET2 mutation is an independent favorable prognostic factor in myelodysplastic syndromes (MDSs). Blood. 2009 Oct 8;114(15):3285-91. PMID: 19666869
44. Lundberg et al. Clonal evolution and clinical correlates of somatic mutations in myeloproliferative neoplasms. Blood. 2014 Apr 3;123(14):2220-8. PMID: 24478400

Disclaimer: The data presented here is from a curated knowledgebase of publicly available information, but may not be exhaustive. The data version is 2023.06(005).