Taipei Veterans General Hospital

Department of Pathology and Laboratory Medicine
No.201, Sec. 2, Shipai Rd., Beitou District, Taipei City, Taiwan
11217, R.O.C.
Tel: 02-2875-7449

Sample Information

Patient Name: 俞新昌
Gender: Male
ID No.: A102335347
History No.: 48947893
Age: 78

Ordering Doctor: DOC1751J 蕭樑材
Ordering REQ.: 0CNWMQX
Signing in Date: 2023/07/22

Path No.: M112-00196
MP No.: MY23050
Assay: Oncomine Myeloid Assay
Sample Type: Bone Marrow
Bone Marrow Aspirating Date: 2023/07/24

Reporting Doctor: DOC5444B 楊靜芬 (Phone: 8#5444)

Note:

Sample Cancer Type: Myelodysplastic Syndrome

Table of Contents

	Page
Variants (Exclude variant in Taiwan BioBank with >1% allele frequency)	2
Biomarker Descriptions	3
Relevant Therapy Summary	4
Relevant Therapy Details	5

Report Highlights

2 Relevant Biomarkers
12 Therapies Available
0 Clinical Trials

Relevant Myelodysplastic Syndrome Variants

Gene	Finding	Gene	Finding
ASXL1	None detected	NPM1	None detected
BCOR	***BCOR p.(R1217*) c.3649C>T***	NRAS	None detected
CBL	None detected	NUP214	None detected
CREBBP	None detected	RUNX1	None detected
ETV6	None detected	SF3B1	None detected
EZH2	None detected	SRSF2	None detected
FLT3	None detected	STAG2	None detected
GATA2	None detected	TP53	None detected
IDH2	None detected	U2AF1	***U2AF1 p.(S34F) c.101C>T***
KIT	None detected	WT1	None detected
KMT2A	None detected	ZRSR2	None detected
MECOM	None detected		

Disclaimer: The data presented here is from a curated knowledgebase of publicly available information, but may not be exhaustive. The data version is 2023.06(005). The content of this report has not been evaluated or approved by FDA, EMA or other regulatory agencies.

Relevant Biomarkers

Tier	Genomic Alteration	Relevant Therapies (In this cancer type)	Relevant Therapies (In other cancer type)	Clinical Trials
IIC	*BCOR p.(R1217*) c.3649C>T* BCL6 corepressor Allele Frequency: 78.71%	None	allogeneic stem cells azacitidine cytarabine cytarabine + daunorubicin + etoposide cytarabine + etoposide + idarubicin cytarabine + fludarabine + idarubicin + filgrastim daunorubicin decitabine idarubicin liposomal cytarabine-daunorubicin CPX-351 mitoxantrone venetoclax + chemotherapy	0
IIC	*U2AF1 p.(S34F) c.101C>T* U2 small nuclear RNA auxiliary factor 1 Allele Frequency: 37.05%	None	allogeneic stem cells azacitidine cytarabine cytarabine + daunorubicin + etoposide cytarabine + etoposide + idarubicin cytarabine + fludarabine + idarubicin + filgrastim daunorubicin decitabine idarubicin liposomal cytarabine-daunorubicin CPX-351 mitoxantrone venetoclax + chemotherapy	0

Public data sources included in relevant therapies: FDA[1], NCCN, **EMA**[2], ESMO

Prevalent cancer biomarkers without relevant evidence based on included data sources

TET2 p.(Q917) c.2749C>T*

Variants (Exclude variant in Taiwan BioBank with >1% allele frequency)

DNA Sequence Variants

Gene	Amino Acid Change	Coding	Variant ID	Locus	Allele Frequency	Transcript	Variant Effect	Coverage
TET2	p.(Q917*)	c.2749C>T	.	chr4:106157848	4.60%	NM_001127208.2	nonsense	2000
U2AF1	p.(S34F)	c.101C>T	COSM166866	chr21:44524456	37.05%	NM_006758.2	missense	2000
BCOR	p.(R1217*)	c.3649C>T	.	chrX:39923059	78.71%	NM_001123385.2	nonsense	1996
CEBPA	p.(H195_P196dup)	c.589_590insACCCGC	.	chr19:33792731	36.53%	NM_004364.4	nonframeshift Insertion	1656

Disclaimer: The data presented here is from a curated knowledgebase of publicly available information, but may not be exhaustive. The data version is 2023.06(005).

Biomarker Descriptions

BCOR (BCL6 corepressor)

Background: The BCOR gene encodes the B-cell CLL/lymphoma 6 (BCL6) corepressor protein which potentiates transcriptional repression by BCL6[1,2]. BCOR also associates with class I and II histone deacetylases (HDACs) suggesting an alternate mechanism for BCOR mediated transcriptional repression independent of BCL6[2]. Genetic alterations in BCOR result in protein dysfunction which suggests BCOR functions as a tumor suppressor gene[3,4,5].

Alterations and prevalence: Genetic alterations in BCOR include missense, nonsense, and frameshift mutations that result in loss of function and have been observed in up to 5% of myelodysplastic syndromes (MDS), 5-10% of chronic myelomonocytic leukemia (CMML), and 1-5% of acute myeloid leukemia (AML)[6,7,8,9]. Higher mutational frequencies are reported in some solid tumors, including up to 15% of uterine cancer and 5-10% of colorectal cancer, stomach cancer, cholangiocarcinoma, and melanoma. Although less common, BCOR fusions and internal tandem duplications (ITDs) have been reported in certain rare cancer types[10,11,12]. Specifically, BCOR-CCNB3 rearrangements define a particular subset of sarcomas with Ewing sarcoma-like morphology known as BCOR-CCNB3 sarcomas (BCS)[13,14]. In clear cell carcinoma of the kidney, a rare pediatric renal malignant tumor, one study described the presence of BCOR ITDs in more than 90% of cases[10].

Potential relevance: BCOR rearrangement, including inv(X)(p11.4p11.22) resulting in BCOR-CCNB3 fusion, is diagnostic of sarcoma with BCOR genetic alterations, a subset of undifferentiated round cell sarcomas[15,16]. Additionally, translocation t(x;22)(p11;q13) resulting in ZC3H7B-BCOR fusion is a useful ancillary diagnostic marker of high-grade endometrial stromal sarcoma[15]. Somatic mutation in BCOR is one of the possible molecular abnormality requirements for the diagnosis of myelodysplasia-related AML (AML-MR) and is associated with poor prognosis in AML and MDS[6,7,17,18,19]. In FLT3-ITD negative AML patients under 65 with intermediate cytogenetic prognosis, mutations in BCOR confer inferior overall survival (OS) as well as relapse free survival (RFS) compared to those without BCOR abnormalities (OS= 13.6% vs. 55%; RFS= 14.3% vs. 44.5%)[9].

TET2 (tet methylcytosine dioxygenase 2)

Background: TET2 encodes the tet methylcytosine dioxygenase 2 protein and belongs to a family of ten-eleven translocation (TET) proteins that also includes TET1 and TET3[20]. TET2 is involved in DNA methylation, specifically in the conversion of 5-methylcytosine to 5-hydroxymethylcytosine[21,22]. The TET proteins contain a C-terminal core catalytic domain that contains a cysteine-rich domain and a double stranded ß-helix domain (DSBH)[23]. TET2 is a tumor suppressor gene. Loss of function mutations in TET2 are associated with loss of catalytic activity and transformation to hematological malignancies[20,21,22]

Alterations and prevalence: Somatic TET2 mutations, including nonsense, frameshift, splice site, and missense, are observed in 20-25% of myelodysplastic syndrome (MDS) associated diseases, including 40%-60% chronic myelomonocytic leukemia (CMML)[6]. TET2 mutations at H1881 and R1896 are frequently observed in myeloid malignancies[21,24]. TET2 mutations are also observed in 9% of uterine, 8% of melanoma and acute myeloid leukemia (AML), as well as 6% of diffuse large B-cell lymphoma (DLBCL).

Potential relevance: The presence of TET2 mutations may be used as one of the major diagnostic criteria in pre-primary myelofibrosis (pre-PMF) and overt PMF in the absence of JAK2/CALR/MPL mutations[25]. TET2 mutations are associated with poor prognosis in PMF and increased rate of transformation to leukemia[25,26]

U2AF1 (U2 small nuclear RNA auxiliary factor 1)

Background: The U2AF1 gene encodes the U2 small nuclear RNA auxiliary factor 1 protein that belongs to the splicing factor SR family of genes involved in RNA splicing[27,28]. U2AF1, also known as U2AF35, mediates the recruitment of the U2AF complex to the 3' end of that pre-mRNA that is being spliced[29]. U2AF1 is the smaller subunit of the U2 auxiliary factor and along with the larger subunit, U2AF65 regulates the removal of introns from pre-mRNAs to produce mature mRNAs for translation during protein synthesis[30]. Mutations in U2AF1 alter the differential splicing of genes that are involved in various biological pathways, including DNMT3B in DNA methylation, ATR along with FANCA in DNA damage response, and H2AFY in X-chromosome inactivation[31]. Spliceosomal genes such as U2AF1 are common targets of somatic mutations in myelodysplastic syndrome (MDS) and are associated with the progression of MDS to acute myeloid leukemia (AML)[31,32,33].

Alterations and prevalence: Recurrent mutations in U2AF1 occur at S34 and Q157 and are observed in 8-12% of MDS[6]. Somatic mutations in U2AF1 are also observed in 10% of uterine carcinoma, 4% of AML, as well as 2% of lung adenocarcinoma and stomach adenocarcinoma[34].

Potential relevance: U2AF1 mutations including S34 and Q157 are associated with poor prognosis in MDS[6]. U2AF1 mutations are associated with inferior overall survival and adverse risk in primary myelofibrosis (PMF) and AML[17,18,25]. Specifically, the Q157 mutation is associated with a significantly shorter overall survival than U2AF1 S34 mutated and U2AF1 unmutated myeloproliferative neoplasms (MPN)[25].

Disclaimer: The data presented here is from a curated knowledgebase of publicly available information, but may not be exhaustive. The data version is 2023.06(005).

Relevant Therapy Summary

● In this cancer type ○ In other cancer type ◑ In this cancer type and other cancer types ✖ No evidence

BCOR p.(R1217*) c.3649C>T

Relevant Therapy	FDA	NCCN	EMA	ESMO	Clinical Trials*
Allogeneic hematopoietic stem cell transplantation	✖	○	✖	✖	✖
azacitidine	✖	○	✖	✖	✖
cytarabine	✖	○	✖	✖	✖
cytarabine + daunorubicin + etoposide	✖	○	✖	✖	✖
cytarabine + etoposide + idarubicin	✖	○	✖	✖	✖
cytarabine + fludarabine + idarubicin + filgrastim	✖	○	✖	✖	✖
daunorubicin	✖	○	✖	✖	✖
decitabine	✖	○	✖	✖	✖
idarubicin	✖	○	✖	✖	✖
liposomal cytarabine-daunorubicin CPX-351	✖	○	✖	✖	✖
mitoxantrone	✖	○	✖	✖	✖
venetoclax + azacitidine	✖	○	✖	✖	✖
venetoclax + cytarabine	✖	○	✖	✖	✖
venetoclax + cytarabine + fludarabine + idarubicin + filgrastim	✖	○	✖	✖	✖
venetoclax + decitabine	✖	○	✖	✖	✖

U2AF1 p.(S34F) c.101C>T

Relevant Therapy	FDA	NCCN	EMA	ESMO	Clinical Trials*
Allogeneic hematopoietic stem cell transplantation	✖	○	✖	✖	✖
azacitidine	✖	○	✖	✖	✖
cytarabine	✖	○	✖	✖	✖
cytarabine + daunorubicin + etoposide	✖	○	✖	✖	✖
cytarabine + etoposide + idarubicin	✖	○	✖	✖	✖
cytarabine + fludarabine + idarubicin + filgrastim	✖	○	✖	✖	✖
daunorubicin	✖	○	✖	✖	✖
decitabine	✖	○	✖	✖	✖
idarubicin	✖	○	✖	✖	✖
liposomal cytarabine-daunorubicin CPX-351	✖	○	✖	✖	✖
mitoxantrone	✖	○	✖	✖	✖

Disclaimer: The data presented here is from a curated knowledgebase of publicly available information, but may not be exhaustive. The data version is 2023.06(005).

Relevant Therapy Summary (continued)

● In this cancer type ○ In other cancer type ◑ In this cancer type and other cancer types ✖ No evidence

U2AF1 p.(S34F) c.101C>T (continued)

Relevant Therapy	FDA	NCCN	EMA	ESMO	Clinical Trials*
venetoclax + azacitidine	✖	○	✖	✖	✖
venetoclax + cytarabine	✖	○	✖	✖	✖
venetoclax + cytarabine + fludarabine + idarubicin + filgrastim	✖	○	✖	✖	✖
venetoclax + decitabine	✖	○	✖	✖	✖

Relevant Therapy Details

Current NCCN Information

● In this cancer type ○ In other cancer type ◑ In this cancer type and other cancer types

NCCN information is current as of 2023-05-01. For the most up-to-date information, search www.nccn.org.
For NCCN International Adaptations & Translations, search www.nccn.org/global/what-we-do/international-adaptations.

Some variant specific evidence in this report may be associated with a broader set of alterations from the NCCN Guidelines. Specific variants listed in this report were sourced from approved therapies or scientific literature. These therapeutic options are appropriate for certain population segments with cancer. Refer to the NCCN Guidelines® for full recommendation.

BCOR p.(R1217*) c.3649C>T

○ azacitidine

Cancer type: Acute Myeloid Leukemia **Variant class:** BCOR mutation

NCCN Recommendation category: 1

Population segment (Line of therapy):
- (Maintenance therapy); Preferred intervention
- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2023]

○ cytarabine + daunorubicin + etoposide

Cancer type: Acute Myeloid Leukemia **Variant class:** BCOR mutation

NCCN Recommendation category: 1

Population segment (Line of therapy):
- (Maintenance therapy); Preferred intervention
- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2023]

Disclaimer: The data presented here is from a curated knowledgebase of publicly available information, but may not be exhaustive. The data version is 2023.06(005).

BCOR p.(R1217*) c.3649C>T (continued)

cytarabine + etoposide + idarubicin

Cancer type: Acute Myeloid Leukemia **Variant class:** BCOR mutation

NCCN Recommendation category: 1

Population segment (Line of therapy):
- (Maintenance therapy); Preferred intervention
- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2023]

daunorubicin

Cancer type: Acute Myeloid Leukemia **Variant class:** BCOR mutation

NCCN Recommendation category: 1

Population segment (Line of therapy):
- (Maintenance therapy); Preferred intervention
- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2023]

idarubicin

Cancer type: Acute Myeloid Leukemia **Variant class:** BCOR mutation

NCCN Recommendation category: 1

Population segment (Line of therapy):
- (Maintenance therapy); Preferred intervention
- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2023]

Allogeneic hematopoietic stem cell transplantation

Cancer type: Acute Myeloid Leukemia **Variant class:** BCOR mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):
- (Maintenance therapy)
- (Consolidation therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2023]

Disclaimer: The data presented here is from a curated knowledgebase of publicly available information, but may not be exhaustive. The data version is 2023.06(005).

BCOR p.(R1217*) c.3649C>T (continued)

azacitidine

Cancer type: Acute Myeloid Leukemia **Variant class:** BCOR mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):
- (Maintenance therapy)
- (Consolidation therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2023]

cytarabine

Cancer type: Acute Myeloid Leukemia **Variant class:** BCOR mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):
- (Maintenance therapy)
- (Consolidation therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2023]

decitabine

Cancer type: Acute Myeloid Leukemia **Variant class:** BCOR mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):
- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2023]

liposomal cytarabine-daunorubicin CPX-351

Cancer type: Acute Myeloid Leukemia **Variant class:** BCOR mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):
- (Maintenance therapy)
- (Consolidation therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2023]

Disclaimer: The data presented here is from a curated knowledgebase of publicly available information, but may not be exhaustive. The data version is 2023.06(005).

BCOR p.(R1217*) c.3649C>T (continued)

mitoxantrone

Cancer type: Acute Myeloid Leukemia **Variant class:** BCOR mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):
- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2023]

---

venetoclax + azacitidine

Cancer type: Acute Myeloid Leukemia **Variant class:** BCOR mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):
- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2023]

---

venetoclax + azacitidine

Cancer type: Acute Myeloid Leukemia **Variant class:** BCOR mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):
- (Maintenance therapy)
- (Consolidation therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2023]

---

venetoclax + cytarabine

Cancer type: Acute Myeloid Leukemia **Variant class:** BCOR mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):
- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2023]

---

venetoclax + decitabine

Cancer type: Acute Myeloid Leukemia **Variant class:** BCOR mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):
- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2023]

Disclaimer: The data presented here is from a curated knowledgebase of publicly available information, but may not be exhaustive. The data version is 2023.06(005).

BCOR p.(R1217*) c.3649C>T (continued)

venetoclax + decitabine

Cancer type: Acute Myeloid Leukemia **Variant class:** BCOR mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):
- (Maintenance therapy)
- (Consolidation therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2023]

azacitidine

Cancer type: Acute Myeloid Leukemia **Variant class:** BCOR mutation

NCCN Recommendation category: 2B

Population segment (Line of therapy):
- (Maintenance therapy)
- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2023]

cytarabine + daunorubicin + etoposide

Cancer type: Acute Myeloid Leukemia **Variant class:** BCOR mutation

NCCN Recommendation category: 2B

Population segment (Line of therapy):
- (Maintenance therapy)
- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2023]

cytarabine + etoposide + idarubicin

Cancer type: Acute Myeloid Leukemia **Variant class:** BCOR mutation

NCCN Recommendation category: 2B

Population segment (Line of therapy):
- (Maintenance therapy)
- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2023]

Disclaimer: The data presented here is from a curated knowledgebase of publicly available information, but may not be exhaustive. The data version is 2023.06(005).

BCOR p.(R1217*) c.3649C>T (continued)

cytarabine + fludarabine + idarubicin + filgrastim

Cancer type: Acute Myeloid Leukemia **Variant class:** BCOR mutation

NCCN Recommendation category: 2B

Population segment (Line of therapy):
- (Maintenance therapy)
- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2023]

decitabine

Cancer type: Acute Myeloid Leukemia **Variant class:** BCOR mutation

NCCN Recommendation category: 2B

Population segment (Line of therapy):
- (Maintenance therapy)
- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2023]

venetoclax + cytarabine + fludarabine + idarubicin + filgrastim

Cancer type: Acute Myeloid Leukemia **Variant class:** BCOR mutation

NCCN Recommendation category: 3

Population segment (Line of therapy):
- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2023]

venetoclax + decitabine

Cancer type: Acute Myeloid Leukemia **Variant class:** BCOR mutation

NCCN Recommendation category: 3

Population segment (Line of therapy):
- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2023]

Disclaimer: The data presented here is from a curated knowledgebase of publicly available information, but may not be exhaustive. The data version is 2023.06(005).

U2AF1 p.(S34F) c.101C>T

azacitidine

Cancer type: Acute Myeloid Leukemia **Variant class:** U2AF1 mutation

NCCN Recommendation category: 1

Population segment (Line of therapy):
- (Maintenance therapy); Preferred intervention
- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2023]

cytarabine + daunorubicin + etoposide

Cancer type: Acute Myeloid Leukemia **Variant class:** U2AF1 mutation

NCCN Recommendation category: 1

Population segment (Line of therapy):
- (Maintenance therapy); Preferred intervention
- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2023]

cytarabine + etoposide + idarubicin

Cancer type: Acute Myeloid Leukemia **Variant class:** U2AF1 mutation

NCCN Recommendation category: 1

Population segment (Line of therapy):
- (Maintenance therapy); Preferred intervention
- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2023]

daunorubicin

Cancer type: Acute Myeloid Leukemia **Variant class:** U2AF1 mutation

NCCN Recommendation category: 1

Population segment (Line of therapy):
- (Maintenance therapy); Preferred intervention
- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2023]

Disclaimer: The data presented here is from a curated knowledgebase of publicly available information, but may not be exhaustive. The data version is 2023.06(005).

U2AF1 p.(S34F) c.101C>T (continued)

idarubicin

Cancer type: Acute Myeloid Leukemia **Variant class:** U2AF1 mutation

NCCN Recommendation category: 1

Population segment (Line of therapy):
- (Maintenance therapy); Preferred intervention
- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2023]

Allogeneic hematopoietic stem cell transplantation

Cancer type: Acute Myeloid Leukemia **Variant class:** U2AF1 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):
- (Maintenance therapy)
- (Consolidation therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2023]

azacitidine

Cancer type: Acute Myeloid Leukemia **Variant class:** U2AF1 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):
- (Maintenance therapy)
- (Consolidation therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2023]

cytarabine

Cancer type: Acute Myeloid Leukemia **Variant class:** U2AF1 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):
- (Maintenance therapy)
- (Consolidation therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2023]

Disclaimer: The data presented here is from a curated knowledgebase of publicly available information, but may not be exhaustive. The data version is 2023.06(005).

U2AF1 p.(S34F) c.101C>T (continued)

decitabine

Cancer type: Acute Myeloid Leukemia **Variant class:** U2AF1 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):
- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2023]

liposomal cytarabine-daunorubicin CPX-351

Cancer type: Acute Myeloid Leukemia **Variant class:** U2AF1 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):
- (Maintenance therapy)
- (Consolidation therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2023]

mitoxantrone

Cancer type: Acute Myeloid Leukemia **Variant class:** U2AF1 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):
- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2023]

venetoclax + azacitidine

Cancer type: Acute Myeloid Leukemia **Variant class:** U2AF1 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):
- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2023]

venetoclax + azacitidine

Cancer type: Acute Myeloid Leukemia **Variant class:** U2AF1 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):
- (Maintenance therapy)
- (Consolidation therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2023]

Disclaimer: The data presented here is from a curated knowledgebase of publicly available information, but may not be exhaustive. The data version is 2023.06(005).

U2AF1 p.(S34F) c.101C>T (continued)

venetoclax + cytarabine

Cancer type: Acute Myeloid Leukemia **Variant class:** U2AF1 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):
- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2023]

venetoclax + decitabine

Cancer type: Acute Myeloid Leukemia **Variant class:** U2AF1 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):
- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2023]

venetoclax + decitabine

Cancer type: Acute Myeloid Leukemia **Variant class:** U2AF1 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):
- (Maintenance therapy)
- (Consolidation therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2023]

azacitidine

Cancer type: Acute Myeloid Leukemia **Variant class:** U2AF1 mutation

NCCN Recommendation category: 2B

Population segment (Line of therapy):
- (Maintenance therapy)
- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2023]

Disclaimer: The data presented here is from a curated knowledgebase of publicly available information, but may not be exhaustive. The data version is 2023.06(005).

U2AF1 p.(S34F) c.101C>T (continued)

cytarabine + daunorubicin + etoposide

Cancer type: Acute Myeloid Leukemia **Variant class:** U2AF1 mutation

NCCN Recommendation category: 2B

Population segment (Line of therapy):
- (Maintenance therapy)
- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2023]

cytarabine + etoposide + idarubicin

Cancer type: Acute Myeloid Leukemia **Variant class:** U2AF1 mutation

NCCN Recommendation category: 2B

Population segment (Line of therapy):
- (Maintenance therapy)
- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2023]

cytarabine + fludarabine + idarubicin + filgrastim

Cancer type: Acute Myeloid Leukemia **Variant class:** U2AF1 mutation

NCCN Recommendation category: 2B

Population segment (Line of therapy):
- (Maintenance therapy)
- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2023]

decitabine

Cancer type: Acute Myeloid Leukemia **Variant class:** U2AF1 mutation

NCCN Recommendation category: 2B

Population segment (Line of therapy):
- (Maintenance therapy)
- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2023]

Disclaimer: The data presented here is from a curated knowledgebase of publicly available information, but may not be exhaustive. The data version is 2023.06(005).

U2AF1 p.(S34F) c.101C>T (continued)

venetoclax + cytarabine + fludarabine + idarubicin + filgrastim

Cancer type: Acute Myeloid Leukemia **Variant class:** U2AF1 mutation

NCCN Recommendation category: 3

Population segment (Line of therapy):
- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2023]

venetoclax + decitabine

Cancer type: Acute Myeloid Leukemia **Variant class:** U2AF1 mutation

NCCN Recommendation category: 3

Population segment (Line of therapy):
- (Induction therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2023]

Disclaimer: The data presented here is from a curated knowledgebase of publicly available information, but may not be exhaustive. The data version is 2023.06(005).

References

1. Gearhart et al. Polycomb group and SCF ubiquitin ligases are found in a novel BCOR complex that is recruited to BCL6 targets. Mol. Cell. Biol. 2006 Sep;26(18):6880-9. PMID: 16943429
2. Huynh et al. BCoR, a novel corepressor involved in BCL-6 repression. Genes Dev. 2000 Jul 15;14(14):1810-23. PMID: 10898795
3. Kelly et al. Bcor loss perturbs myeloid differentiation and promotes leukaemogenesis. Nat Commun. 2019 Mar 22;10(1):1347. PMID: 30902969
4. Cao et al. BCOR regulates myeloid cell proliferation and differentiation. Leukemia. 2016 May;30(5):1155-65. PMID: 26847029
5. Yamamoto et al. Clarifying the impact of polycomb complex component disruption in human cancers. Mol. Cancer Res. 2014 Apr;12(4):479-84. PMID: 24515802
6. NCCN Guidelines® - NCCN-Myelodysplastic Syndromes [Version 1.2023]
7. Damm et al. BCOR and BCORL1 mutations in myelodysplastic syndromes and related disorders. Blood. 2013 Oct 31;122(18):3169-77. PMID: 24047651
8. Weinstein et al. The Cancer Genome Atlas Pan-Cancer analysis project. Nat. Genet. 2013 Oct;45(10):1113-20. PMID: 24071849
9. Terada et al. Usefulness of BCOR gene mutation as a prognostic factor in acute myeloid leukemia with intermediate cytogenetic prognosis. Genes Chromosomes Cancer. 2018 Aug;57(8):401-408. PMID: 29663558
10. Wong et al. Clear cell sarcomas of the kidney are characterised by BCOR gene abnormalities, including exon 15 internal tandem duplications and BCOR-CCNB3 gene fusion. Histopathology. 2018 Jan;72(2):320-329. PMID: 28833375
11. Cramer et al. Successful Treatment of Recurrent Primitive Myxoid Mesenchymal Tumor of Infancy With BCOR Internal Tandem Duplication. J Natl Compr Canc Netw. 2017 Jul;15(7):868-871. PMID: 28687574
12. Peters et al. BCOR-CCNB3 fusions are frequent in undifferentiated sarcomas of male children. Mod. Pathol. 2015 Apr;28(4):575-86. PMID: 25360585
13. Puls et al. BCOR-CCNB3 (Ewing-like) sarcoma: a clinicopathologic analysis of 10 cases, in comparison with conventional Ewing sarcoma. Am. J. Surg. Pathol. 2014 Oct;38(10):1307-18. PMID: 24805859
14. Kao et al. BCOR-CCNB3 Fusion Positive Sarcomas: A Clinicopathologic and Molecular Analysis of 36 Cases With Comparison to Morphologic Spectrum and Clinical Behavior of Other Round Cell Sarcomas. Am. J. Surg. Pathol. 2018 May;42(5):604-615. PMID: 29300189
15. NCCN Guidelines® - NCCN-Soft Tissue Sarcoma [Version 2.2023]
16. NCCN Guidelines® - NCCN-Bone Cancer [Version 3.2023]
17. NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 3.2023]
18. Döhner et al. Diagnosis and management of AML in adults: 2022 recommendations from an international expert panel on behalf of the ELN. Blood. 2022 Sep 22;140(12):1345-1377. PMID: 35797463
19. Khoury et al. The 5th edition of the World Health Organization Classification of Haematolymphoid Tumours: Myeloid and Histiocytic/Dendritic Neoplasms. Leukemia. 2022 Jul;36(7):1703-1719. PMID: 35732831
20. Pan et al. The TET2 interactors and their links to hematological malignancies. IUBMB Life. 2015 Jun;67(6):438-45. PMID: 26099018
21. Ko et al. Impaired hydroxylation of 5-methylcytosine in myeloid cancers with mutant TET2. Nature. 2010 Dec 9;468(7325):839-43. PMID: 21057493
22. Solary et al. The Ten-Eleven Translocation-2 (TET2) gene in hematopoiesis and hematopoietic diseases. Leukemia. 2014 Mar;28(3):485-96. PMID: 24220273
23. An et al. TET family dioxygenases and DNA demethylation in stem cells and cancers. Exp. Mol. Med. 2017 Apr 28;49(4):e323. PMID: 28450733
24. Kosmider et al. TET2 mutation is an independent favorable prognostic factor in myelodysplastic syndromes (MDSs). Blood. 2009 Oct 8;114(15):3285-91. PMID: 19666869
25. NCCN Guidelines® - NCCN-Myeloproliferative Neoplasms [Version 3.2022]
26. Lundberg et al. Clonal evolution and clinical correlates of somatic mutations in myeloproliferative neoplasms. Blood. 2014 Apr 3;123(14):2220-8. PMID: 24478400
27. O'Leary et al. Reference sequence (RefSeq) database at NCBI: current status, taxonomic expansion, and functional annotation. Nucleic Acids Res. 2016 Jan 4;44(D1):D733-45. PMID: 26553804
28. Long et al. The SR protein family of splicing factors: master regulators of gene expression. Biochem. J. 2009 Jan 1;417(1):15-27. PMID: 19061484
29. Zuo et al. The splicing factor U2AF35 mediates critical protein-protein interactions in constitutive and enhancer-dependent splicing. Genes Dev. 1996 Jun 1;10(11):1356-68. PMID: 8647433

Disclaimer: The data presented here is from a curated knowledgebase of publicly available information, but may not be exhaustive. The data version is 2023.06(005).

References (continued)

30. Ruskin et al. A factor, U2AF, is required for U2 snRNP binding and splicing complex assembly. Cell. 1988 Jan 29;52(2):207-19. PMID: 2963698
31. Ilagan et al. U2AF1 mutations alter splice site recognition in hematological malignancies. Genome Res. 2015 Jan;25(1):14-26. PMID: 25267526
32. Graubert et al. Recurrent mutations in the U2AF1 splicing factor in myelodysplastic syndromes. Nat. Genet. 2011 Dec 11;44(1):53-7. PMID: 22158538
33. Papaemmanuil et al. Clinical and biological implications of driver mutations in myelodysplastic syndromes. Blood. 2013 Nov 21;122(22):3616-27; quiz 3699. PMID: 24030381
34. Cerami et al. The cBio cancer genomics portal: an open platform for exploring multidimensional cancer genomics data. Cancer Discov. 2012 May;2(5):401-4. PMID: 22588877

Disclaimer: The data presented here is from a curated knowledgebase of publicly available information, but may not be exhaustive. The data version is 2023.06(005).