

# Guardant360 基因檢測服務報告

醫師姓名：江起陸

受檢者姓名：袁金蘭

送檢編號：A1173211

檢測次數：第一次

## 提醒

基因數據乃屬個人隱私，切勿輕易向任何個人、團體或非您的授權者透漏本報告內容。若您有任何疑慮，歡迎來電洽詢，我們很樂意為您提供更詳細的諮詢服務。若因郵遞錯誤收此檔，請予銷毀，多謝合作。

詳細資訊



## 康誠生技股份有限公司 客戶服務中心

諮詢時間 | 週一～週五 9:00～17:00（國定假日除外）



諮詢專線 | 02-55696099

客服信箱 | [service.gb@healthconn.com](mailto:service.gb@healthconn.com)

REPORTING	PHYSICIAN	 <div>Complete Tumor Response Map on page 2</div>
Report Date: OCT-14-2024	Chi-Lu Chiang	
Receipt Date: OCT-10-2024	Account: Genconn Biotech Co., LTD	
Collection Date: OCT-09-2024	Address: 5F., No. 54, Sec. 1, Jhongsiao E. Rd., Zhongzheng Dist., Beixin Rd, Xindian Dist, Taipei City, 100, Taiwan	
Specimen: Blood	Ph: +886 963 820 633   Fax: N/A	
Status: FINAL	Additional Recipient: N/A	

Summary of Detected Somatic Alterations, Immunotherapy Biomarkers & Associated Treatment Options

KEY  Approved in indication  Approved in other indication  Lack of response

Detected Alteration(s) / Biomarker(s)	Associated FDA-approved therapies	Clinical trial availability (see page 3)	% cfDNA or Amplification
EGFR E746_A750del (Exon 19 deletion)	 Afatinib, Amivantamab, Amivantamab+lazertinib, Dacomitinib, Erlotinib, Erlotinib+ramucirumab, Gefitinib, Osimertinib	Yes	13.8%
PTEN Q149Gs*8	 Capivasertib	Yes	22.6%
CTNNB1 S45C	None	Yes	11.3%
KRAS L19F	None	Yes	0.1%
TP53 R175H	None	No	11.7%

Comments  
Reported by: NT3

Additional Biomarkers

Biomarker	Additional Details
MSI-High	NOT DETECTED

We evaluated this sample for 74 genes, including the following guideline-recommended genes for NSCLC

EGFR(T790M and others)

ALK

ROS1

BRAF

MET

ERBB2(HER2)


RET

NTRK

KRAS

Guardant360 Tumor Response Map

The Guardant360 Tumor Response Map illustrates the variant allele fraction (% cfDNA) of observed somatic variants at each sample submission. Amplifications are not plotted, and only the first and last five test dates are plotted. Please see the Physician Portal ([portal.guardanthealth.com](https://portal.guardanthealth.com)) for the Tumor Response Map with all test dates.



Detected Alteration(s) / Biomarker(s)	% cfDNA or Amp
PTEN Q149Gfs*8	22.6%
EGFR E746_A750del (Exon 19 deletion)	13.8%
TP53 R175H	11.7%
CTNNB1 S45C	11.3%
KRAS L19F	0.1%

The table above annotates the variant allele fraction (% cfDNA) detected in this sample, listed in descending order.  
§ See definitions section for more detail

## Available Clinical Trials (within the same state as the ordering physician)

There may be additional trials not listed here. Visit: [portal.guardanthealth.com](https://portal.guardanthealth.com) or email [clientservices@guardanthealth.com](mailto:clientservices@guardanthealth.com) with A1173211 in the subject line of the email, for additional trials.

Alteration	Trial ID / Contact	Title	Phase	Site(s)
EGFR E746_A750del	NCT04077463 Study Contact,Participate-In-This-Study@its.jnj.com,844-434-4210	A Study of Lazertinib as Monotherapy or in Combination With Amivantamab in Participants With Advanced Non-small Cell Lung Cancer	Phase 1	Kaohsiung, Taiwan Taipei City, Taiwan Tainan, Taiwan Taichung, Taiwan
	NCT05215548 Jin-Shing Chen, M.D., Ph.D.,chenjs@ntu.edu.tw,886-2-2322-0322	Primary Tumor Resection With EGFR TKI for Stage IV NSCLC	Phase 2	Taipei, Taiwan (2)
	NCT05442060 Anna Hu,annahu@obipharma.com,886-2-27866589 x104	To Evaluate OBI-833/OBI-821 in Combination With First-Line Erlotinib in Patients With EGFR-Mutated, Globo H-Positive, Locally Advanced or Metastatic Non-Small Cell Lung Cancer	Phase 2	New Taipei City, Taiwan Taoyuan, Taiwan Taichung, Taiwan Taipei, Taiwan (4)
	NCT05801029 AstraZeneca Clinical Study Information Center,information.center@astrazeneca.com,1-877-240-9479	A Study to Investigate Safety and Efficacy of Osimertinib and Amivantamab in Participants With Non-small Cell Lung Cancer With Common Epidermal Growth Factor Receptor Mutations	Phase 2	Yunlin, Taiwan Taipei City, Taiwan Kaohsiung, Taiwan (2) Taipei, Taiwan (2)  Additional trial sites available
	NCT06120140 Study Contact,Participate-In-This-Study@its.jnj.com,844-434-4210	Enhanced Dermatological Care to Reduce Rash and Paronychia in Epidermal Growth Factor Receptor (EGFR)-Mutated Non-Small Cell Lung Cancer (NSCLC) Treated First-line With Amivantamab Plus Lazertinib	Phase 2	Kaohsiung City, Taiwan Taipei, Taiwan Taoyuan City, Taiwan Taichung City, Taiwan  Additional trial sites available
	Visit <a href="https://portal.guardanthealth.com">portal.guardanthealth.com</a> for trials not within the same state as the physician's office			
CTNNB1 S45C	NCT04008797 Eisai Inquiry Service,eisai-chiken_hotline@hcc.eisai.co.jp	A Study of E7386 in Combination With Other Anticancer Drug(s) in Participants With Solid Tumor	Phase 1 /Phase 2	Tainan, Taiwan Taoyuan, Taiwan Kao-Hsiung, Taiwan Taipei, Taiwan (2)  Additional trial sites available
Visit <a href="https://portal.guardanthealth.com">portal.guardanthealth.com</a> for trials not within the same state as the physician's office				
PTEN Q149Gfs*8	Visit <a href="https://portal.guardanthealth.com">portal.guardanthealth.com</a> for trials not within the same state as the physician's office			
KRAS L19F	Visit <a href="https://portal.guardanthealth.com">portal.guardanthealth.com</a> for trials not within the same state as the physician's office			

More clinical trial options available at [portal.guardanthealth.com](https://portal.guardanthealth.com)

Definitions

**Deletion (Del):** The following alteration was detected in this patient: *EGFR* E746\_A750del; *PTEN* Q149Gfs\*8. Guardant360 detects short deletions in exons of certain genes (see Table 1), including potential splice site-disrupting events.

Interpretation

Somatic alterations were detected in the circulating cell-free DNA isolated from this patient's blood specimen. These genomic alterations are cancer-associated somatic variants, some of which have been associated with either increased or reduced clinical response to specific treatments. The percentage of altered cell-free DNA circulating (% cfDNA) in blood is related to the unique tumor biology of each patient. Factors that may affect the % cfDNA of detected somatic alterations include tumor growth, turn over, size, heterogeneity, vascularization, disease progression, and treatment.

Method and Limitations

Guardant360 sequences 74 cancer-associated genes to identify somatic alterations. Cell-free DNA (cfDNA) is extracted from plasma, enriched for targeted regions, and sequenced using the Illumina platform and hg19 as the reference genome. All exons are sequenced in some genes; only clinically significant exons are sequenced in other genes. The types of genomic alterations detected by Guardant360 include single nucleotide variants, gene amplifications, fusions, short insertions/deletions (longest detected, 70 base pairs), and splice site disrupting events (see Table 1). Microsatellite Instability (MSI) is assessed for all cancer types by evaluating somatic changes in the length of repetitive sequences on the Guardant360 panel. A "Not Detected" result in samples where the highest % cfDNA is < 0.2% is an inconclusive result because it does not preclude MSI-High status in tissue. MSI status is currently not reported for earlier panel versions. This version of the Guardant360 test is not validated for the detection of other types of genomic alterations, such as complex rearrangements or gene deletions. Certain sample or variant characteristics, such as low cfDNA concentration, may result in reduced analytic sensitivity. Guardant360 cannot discern the source of circulating cfDNA, and for some variants in the range of ~40 to 60% cfDNA, the test cannot easily distinguish germline variants from somatic alterations. Guardant360 is not validated for the detection of germline or de novo variants that are associated with hereditary cancer risk. Tissue genotyping should be considered when plasma genotyping is negative, if clinically appropriate.

Table 1: Genes on the Guardant360 Panel

Guardant360 reports single nucleotide variants, splice site mutations, and insertion and deletion variants (indels) in all clinically relevant exons in 74 genes and reports other variant types in select genes as indicated below.

AKT1	ALK #	APC	AR †	ARAF	ARID1A	ATM	BRAF †	BRCA1
BRCA2	CCND1 †	CCND2 †	CCNE1 †	CDH1	CDK12	CDK4 †	CDK6 †	CDKN2A
CTNNB1	DDR2	EGFR †	ERBB2 †	ESR1	EZH2	FBXW7	FGFR1 †	FGFR2 † #
FGFR3 #	GATA3	GNA11	GNAQ	GNAS	HNF1A	HRAS	IDH1	IDH2
JAK2	JAK3	KIT †	KRAS †	MAP2K1	MAP2K2	MAPK1	MAPK3	MET †
MLH1	MPL	MTOR	MYC †	NF1	NFE2L2	NOTCH1	NPM1	NRAS
NTRK1 #	NTRK3	PDGFRA †	PIK3CA †	PTEN	PTPN11	RAF1 †	RB1	RET #
RHEB	RHOA	RIT1	ROS1 #	SMAD4	SMO	STK11	TERT ‡	TP53
TSC1	VHL							

‡ Guardant360 reports alterations in the promoter region of this gene.  
# Guardant360 reports fusion events involving this gene.  
† Guardant360 reports amplifications of this gene.

About the Test

The Guardant360 assay was developed and its performance characteristics were determined by Guardant Health, Inc. This test has not been cleared or approved by the U.S. Food and Drug Administration (FDA). The FDA has determined that such clearance or approval is not necessary. This test may be used for clinical purposes and should not be regarded as investigational or for research only. Guardant Health's clinical reference laboratory is certified under the Clinical Laboratory Improvement Amendments of 1988 (CLIA) as qualified to perform high complexity clinical laboratory testing. The laboratory report should be interpreted in the context of other clinical information and laboratory, pathology, and imaging studies by a qualified medical professional prior to initiating or changing a patient's treatment plan. The selection of any, all, or none of the drugs associated with potential clinical benefit (or potential lack of clinical benefit) is entirely at the discretion of the treating medical professional. Drug and trial information are based on the diagnosis written on the submitted test request form; this information is not based on any supplemental information provided by the requesting medical professional, including pathology reports or other molecular studies. Some drugs listed in this report may not be approved or cleared by the FDA for the indicated use. Guardant Health makes no endorsement, express or implied, of any product, physician, or procedure contained in this report. This report makes no promises or guarantees that a particular medication will affect (or not affect) the clinical outcome of any patient.

Testing Performed at: Guardant Health

Laboratory Director: Martina Lefterova, MD PhD | CLIA ID: 05D2070300 | CAP #: 8765297 | 505 Penobscot Drive, Redwood City, CA, 94063, USA

### Additional information is available

Any therapeutic annotations are based on publicly available information. This information is described in the "Detailed Therapy Results" and "Relevance of Detected Alterations" sections.

Visit [portal.guardanthealth.com](https://portal.guardanthealth.com) or email [clientservices@guardanthealth.com](mailto:clientservices@guardanthealth.com) with A1173211 in the subject line of the email for:

- Additional clinical trials
- Detailed Therapy Results
- Relevance of Detected Alterations
- References

If you would like to receive this additional information with every Guardant360 report, please call client services at [855.698.8887](tel:855.698.8887) to opt-in.

Additional information begins on the next page.



## List of Available Clinical Trials

Alteration	Trial ID / Contact	Title	Phase	Site (number in parenthesis is count of trial sites)
EGFR E746_A750del	NCT03574402 Yi-Long Wu, Professor, syylwu@live.cn, 862083827812	Phase II Umbrella Study Directed by Next Generation Sequencing	Phase 2	China
	NCT04077463 Study Contact, Participate-In-This-Study@its.jnj.com, 844-434-4210	A Study of Lazertinib as Monotherapy or in Combination With Amivantamab in Participants With Advanced Non-small Cell Lung Cancer	Phase 1	Seattle, WA; Detroit, MI; Saint Louis, MO; Philadelphia, PA; Portland, OR; Salt Lake City, UT; Tampa, FL; Fairfax, VA; Boston, MA (2); New York, NY (2); CA (5); Japan (7); China (9); Taiwan (4); Korea, Republic of (4); Italy (5); France (7)
	NCT04922138 Baohui Han, doctor, 18930858216@163.com, 18930858216	Aumolertinib Adjuvant Therapy of Resectable Stage I EGFRm+ NSCLC With High-grade Patterns	Phase 2	China (2)
	NCT05163028 John Ning, MD, PhD, FAIC, jning@huyabio.com, 858-280-1866	A Dose Escalation Study of SHP2 Inhibitor in Patients With Solid Tumors Harboring KRAS of EGFR Mutations	Phase 1	Orlando, FL; Tyler, TX; Plantation, FL; Canton, OH; Fairfax, VA; CA (6); Puerto Rico
	NCT05215548 Jin-Shing Chen, M.D., Ph.D., chenjs@ntu.edu.tw, 886-2-2322-0322	Primary Tumor Resection With EGFR TKI for Stage IV NSCLC	Phase 2	Taiwan (2)
	NCT05442060 Anna Hu, annahu@obipharma.com, 886-2-27866589 x104	To Evaluate OBI-833/OBI-821 in Combination With First-Line Erlotinib in Patients With EGFR-Mutated, Globo H-Positive, Locally Advanced or Metastatic Non-Small Cell Lung Cancer	Phase 2	Taiwan (7)
	NCT05801029 AstraZeneca Clinical Study Information Center, information.center@astrazeneca.com, 1-877-240-9479	A Study to Investigate Safety and Efficacy of Osimertinib and Amivantamab in Participants With Non-small Cell Lung Cancer With Common Epidermal Growth Factor Receptor Mutations	Phase 2	Glendale, CA; New York, NY; Washington, DC; Sacramento, CA; Beverly Hills, CA; Fairfax, VA; Canada (2); Singapore (3); Hong Kong (4); Taiwan (9); Korea, Republic of (7); Malaysia (6); Thailand (5)
	NCT06041776 Shun Lu, M.D., shunlu@sjtu.edu.cn, +86 21 2220 0000	Adjuvant Befotertinib in Stage IB-IIIB Non-small Cell Lung Cancer With Positive EGFR Sensitive Mutations	Phase 3	China (3)
	NCT06043973 Degan Lu, Professor, deganlu@126.com, 18753157623	Almonertinib Combined With Anlotinib as First-line Treatment for Advanced Non-small Cell Lung Cancer	Phase 3	China
CTNNB1 S45C	NCT06120140 Study Contact, Participate-In-This-Study@its.jnj.com, 844-434-4210	Enhanced Dermatological Care to Reduce Rash and Paronychia in Epidermal Growth Factor Receptor (EGFR)-Mutated Non-Small Cell Lung Cancer (NSCLC) Treated First-line With Amivantamab Plus Lazertinib	Phase 2	Renton, WA; Hinsdale, IL; Westbury, NY; Cleveland, OH; W. Salem, WI; Athens, GA; Chandler, AZ; Reno, NV; Springfield, MO; Fairfax, VA; Flemington, NJ; Wilson, NC; CA (16); Argentina (5); Turkey (8); China (11); Taiwan (5); Korea, Republic of (3); Brazil (11); Malaysia (4); France (3); Germany (6); Spain (9)
	NCT04008797 Eisai Inquiry Service, eisai-chiken_hotline@hcc.eisai.co.jp	A Study of E7386 in Combination With Other Anticancer Drug(s) in Participants With Solid Tumor	Phase 1 /Phase 2	Oklahoma City, OK; Pasadena, CA; Santa Monica, CA; Los Angeles, CA; Houston, TX; West Palm Beach, FL; La Jolla, CA; Aurora, CO; Sarasota, FL; Nashville, TN; Charleston, SC; Kansas City, MO; New York, NY (2); Dallas, TX (2); Japan (12); Taiwan (6); Korea, Republic of (5); France (21)
	NCT04681248 Leap Therapeutics, Inc.,	Expanded Access Use of DKN-01 for the Treatment of Advanced Solid Tumors		Los Angeles, CA; Milwaukee, WI; Madison, WI; Chicago, IL;

## List of Available Clinical Trials

Alteration	Trial ID / Contact	Title	Phase	Site (number in parenthesis is count of trial sites)
	EarlyAccess@leaptx.com,617-714-0360			Birmingham, AL
	NCT04780568 The Ohio State University Comprehensive Cancer Center, OSUCCCClinicaltrials@osumc.edu,800-293-5066	Osimertinib and Tegavivint as First-Line Therapy for the Treatment of Metastatic EGFR-Mutant Non-small Cell Lung Cancer	Phase 1	Columbus, OH
	NCT04851119 See <a href="https://clinicaltrials.gov/ct2/show/NCT04851119">https://clinicaltrials.gov/ct2/show/NCT04851119</a>	Tegavivint for the Treatment of Recurrent or Refractory Solid Tumors, Including Lymphomas and Desmoid Tumors	Phase 1 /Phase 2	Philadelphia, PA; Saint Louis, MO; Minneapolis, MN; New York, NY; Chicago, IL; Orange, CA; San Francisco, CA; Memphis, TN; Los Angeles, CA; Seattle, WA; Houston, TX; Boston, MA; Indianapolis, IN; Atlanta, GA; Pittsburgh, PA; Cincinnati, OH; Washington, DC; Birmingham, AL; Aurora, CO; Ann Arbor, MI
	NCT05919264 Clinical Trial Inquiries, clinicaltrials@fogpharma.com,(857) 259-6305	FOG-001 in Locally Advanced or Metastatic Solid Tumors	Phase 1 /Phase 2	Houston, TX; Saint Louis, MO; Boston, MA; Scottsdale, AZ; New York, NY; New Haven, CT; Portland, OR; San Antonio, TX; Nashville, TN
PTEN Q149Gfs*8	NCT02264678 AstraZeneca Clinical Study Information Center,information.center@astrazeneca.com,1-877-240-9479	Ascending Doses of Ceralasertib in Combination With Chemotherapy and/or Novel Anti Cancer Agents	Phase 1 /Phase 2	Duarte, CA; Philadelphia, PA; Irvine, CA; Los Angeles, CA (2); United Kingdom (5); France (4)
	NCT02407509 Taleen Shakouri, PhD,DDU3808@icr.ac.uk,02034376629	Phase I Trial of VS-6766 Alone and in Combination With Everolimus	Phase 1	United Kingdom (3)
	NCT03065062 Geoffrey Shapiro, MD, Geoffrey_S Shapiro@dfci.harvard.edu,617-632-4942	Study of the CDK4/6 Inhibitor Palbociclib (PD-0332991) in Combination With the PI3K/mTOR Inhibitor Gedatolisib (PF-05212384) for Patients With Advanced Squamous Cell Lung, Pancreatic, Head & Neck and Other Solid Tumors	Phase 1	Boston, MA (3)
	NCT03739710 US GSK Clinical Trials Call Center, GSKClinicalSupportHD@gsk.com,877-379-3718	Platform Trial of Novel Regimens Versus Standard of Care (SoC) in Participants With Non-small Cell Lung Cancer (NSCLC)	Phase 2	Los Angeles, CA; Nashville, TN; Chattanooga, TN; Bronx, NY (2); Germany; Canada (3); Italy (4); France (4); Spain (3)
	NCT05327010 See <a href="https://clinicaltrials.gov/ct2/show/NCT05327010">https://clinicaltrials.gov/ct2/show/NCT05327010</a>	Testing the Combination of the Anti-cancer Drugs ZEN003694 (ZEN-3694) and Talazoparib in Patients With Advanced Solid Tumors, The ComBET Trial	Phase 2	New Haven, CT; Chicago, IL; Sacramento, CA; Houston, TX; Lee's Summit, MO; Duarte, CA; Chapel Hill, NC; North Kansas City, MO; Atlanta, GA; Pittsburgh, PA; La Jolla, CA; Irvine, CA; Aurora, CO; Gainesville, FL; Galveston, TX; Salt Lake City, UT; Trumbull, CT; Lexington, KY; Charlottesville, VA; Kansas City, MO; Bethesda, MD (2); KS (5)
KRAS L19F	NCT02407509 Taleen Shakouri, PhD,DDU3808@icr.ac.uk,02034376629	Phase I Trial of VS-6766 Alone and in Combination With Everolimus	Phase 1	United Kingdom (3)
	NCT05163028 John Ning, MD,PhD,FAIC,jning@huyabio.com,858-280-1866	A Dose Escalation Study of SHP2 Inhibitor in Patients With Solid Tumors Harboring KRAS of EGFR Mutations	Phase 1	Orlando, FL; Tyler, TX; Plantation, FL; Canton, OH; Fairfax, VA; CA (6); Puerto Rico
	NCT05375084 Navire Clinical Operations,NAV1004ct.gov@bridgebio.com,650-391-9740	SHP2 Inhibitor BBP-398 in Combination With Nivolumab in Patients With Advanced Non-Small Cell Lung Cancer With a KRAS Mutation	Phase 1	Philadelphia, PA; Baltimore, MD; Portland, OR; Buffalo, NY; Santa Rosa, CA; Detroit, MI; Cleveland, OH; Springdale, AR; La Jolla, CA;

List of Available Clinical Trials

Alteration	Trial ID / Contact	Title	Phase	Site (number in parenthesis is count of trial sites)
				Hollywood, FL; Tampa, FL; Charleston, SC; Fairfax, VA; Houston, TX (2)
	NCT05585320 IMM1104-101 Study Team, clinicaltrials@immuneering.com,(860) 321-1302	A Phase 1/2a Study of IMM-1-104 in Participants With Previously Treated, RAS-Mutant, Advanced or Metastatic Solid Tumors	Phase 1 /Phase 2	Denver, CO; New York, NY; Chicago, IL; Jacksonville, FL; Santa Monica, CA; Durham, NC; Houston, TX; Duarte, CA; Madison, WI; Rochester, MN; Boston, MA; San Diego, CA; Scottsdale, AZ; Lake Mary, FL; East Syracuse, NY; Nashville, TN; Fairfax, VA
	NCT06162221 Revolution Medicines,CT- inquiries@RevMed.com,(650)779-2300	Study of RAS(ON) Inhibitor Combinations in Patients with Advanced RAS-mutated NSCLC	Phase 1 /Phase 2	Houston, TX; Westbury, NY; Cleveland, OH; Irving, TX; Plantation, FL; New York, NY; Washington, DC; Sacramento, CA; Irvine, CA; Billings, MT; San Francisco, CA; Fairfax, VA; Netherlands; Germany; Italy (2); France (7); Spain (7)

## Detailed Therapy Results

Alteration	Drug	Trade Name	Target	Current Status
EGFR E746_A750del (Exon 19 deletion)	ABT-101		Egfr/Her2 inhibitor.	Phase 1 (Non-small cell lung carcinoma (NSCLC)) Phase 1 (Head and neck squamous cell carcinoma (HNSCC))
	Afatinib	Gilotrif	Irreversible pan-ErbB kinase inhibitor.	Phase 3 (Non-small cell lung carcinoma (NSCLC)) FDA Approved in other indications (Lung squamous cell carcinoma, EGFR-mutant NSCLC)
	Amivantamab	Rybrevant	Bispecific anti-Met/Egfr antibody.	Phase 3 (Non-small cell lung carcinoma (NSCLC)) FDA Approved in other indications (NSCLC with EGFR exon 20 insertion, NSCLC with EGFR exon 19 del /L858R)
	Amivantamab+lazertinib	Rybrevant+Lazcluze	Bispecific anti-Met/Egfr antibody+Egfr tyrosine kinase inhibitor.	Phase 3 (Non-small cell lung carcinoma (NSCLC)) FDA Approved in other indications (NSCLC with EGFR exon 19 del/L858R)
	Aumolertinib		Egfr T790M inhibitor.	Phase 3 (Non-small cell lung carcinoma (NSCLC))
	Avitinib		Irreversible mutation-specific Egfr kinase inhibitor.	Phase 2 (Non-small cell lung carcinoma (NSCLC)) Phase 1 (Non-Hodgkin lymphoma (NHL))
	BAY2927088		Egfr/Her2 kinase inhibitor.	Phase 1 (Non-small cell lung carcinoma (NSCLC))
	BBP-398		Shp-2 inhibitor.	Phase 1 (Non-small cell lung carcinoma (NSCLC)) Phase 1 (Solid Tumor)
	BDTX-1535		Irreversible brain-penetrant fourth generation Egfr inhibitor.	Phase 1 (Non-small cell lung carcinoma (NSCLC)) Phase 1 (Glioma)
	Befotertinib		Third generation mutation-specific (T790M, L858R, exon 19 deletion) Egfr tyrosine kinase inhibitor.	Phase 3 (Non-small cell lung carcinoma (NSCLC))
	BLU-945		Fourth generation Egfr inhibitor targeting T790M and T790M/C797S mutations.	Phase 2 (Non-small cell lung carcinoma (NSCLC))
	BPI-361175		Fourth generation Egfr inhibitor targeting T790M and T790M/C797S mutations.	Phase 2 (Non-small cell lung carcinoma (NSCLC))
	CLN-081		Covalent mutation-specific (L858R, T790M, exon 19 deletion, exon 20 insertion) Egfr tyrosine kinase inhibitor.	Phase 2 (Non-small cell lung carcinoma (NSCLC))
	CM93		Third generation mutation-specific (T790M, L858R, exon 19 deletion) Egfr tyrosine kinase inhibitor.	Phase 1 (Glioblastoma)
	Dacomitinib	Vizimpro	Pan-ErbB family tyrosine kinase inhibitor.	Phase 3 (Non-small cell lung carcinoma (NSCLC)) FDA Approved in other indications (EGFR-mutant lung carcinoid, EGFR-mutant lung large cell neuroendocrine carcinoma, EGFR-mutant NSCLC)

## Detailed Therapy Results

Alteration	Drug	Trade Name	Target	Current Status
	ERAS-601		Shp-2 inhibitor.	Phase 2 (Solid Tumor)
	Erlotinib	Tarceva	Egfr tyrosine kinase inhibitor.	Phase 3 (Non-small cell lung carcinoma (NSCLC)) FDA Approved in other indications (EGFR-mutant lung carcinoid, EGFR-mutant lung large cell neuroendocrine carcinoma, NSCLC with EGFR exon 19 del/L858R, Pancreatic carcinoma, EGFR-mutant NSCLC)
	Erlotinib+bevacizumab	Tarceva+Avastin	Egfr tyrosine kinase inhibitor + anti-VEGF-A monoclonal antibody combination.	Phase 3 (Non-small cell lung carcinoma (NSCLC)) Phase 3 (Lung cancer)
	Erlotinib+ramucirumab	Tarceva+Cyramza	Egfr tyrosine kinase inhibitor + anti-VEGFR-2 monoclonal antibody combination.	Phase 3 (Non-small cell lung carcinoma (NSCLC)) FDA Approved in other indications (NSCLC with EGFR exon 19 del/L858R)
	ET0038		Shp-2 inhibitor.	Phase 1 (Solid Tumor)
	Furmonertinib		Third generation mutation-specific (T790M, L858R, exon 19 deletion) Egfr tyrosine kinase inhibitor.	Phase 2 (Non-small cell lung carcinoma (NSCLC))
	FWD1509		Egfr/Her2 kinase inhibitor.	Phase 2 (Non-small cell lung carcinoma (NSCLC))
	Gefitinib	Iressa	Egfr tyrosine kinase inhibitor.	Phase 3 (Non-small cell lung carcinoma (NSCLC)) FDA Approved in other indications (EGFR-mutant lung carcinoid, EGFR-mutant lung large cell neuroendocrine carcinoma, EGFR-mutant NSCLC)
	H002		Fourth generation Egfr inhibitor targeting exon 19del /L858R, T790M, and C797S mutations.	Phase 2 (Non-small cell lung carcinoma (NSCLC))
	HBI-2376		Shp-2 inhibitor.	Phase 1 (Solid Tumor)
	Hemay022		Egfr tyrosine kinase inhibitor.	Phase 1 (Breast carcinoma (HER2+))
	Icotinib	Conmana	Egfr inhibitor.	Phase 3 (Non-small cell lung carcinoma (NSCLC)) Phase 2 (Esophageal carcinoma)
	JIN-A02		Fourth generation Egfr inhibitor targeting T790M and T790M/C797S mutations.	Phase 2 (Non-small cell lung carcinoma (NSCLC))
	Lazertinib	Lazcluze	Third generation mutation-specific Egfr tyrosine kinase inhibitor.	Phase 2 (Non-small cell lung carcinoma (NSCLC))
	Lifirafenib		Dual Braf/Egfr inhibitor.	Phase 1 (Non-small cell lung carcinoma (NSCLC)) Phase 1 (Solid Tumor, Brain and Central Nervous System Tumors)
	MCLA-129		Anti-EGFR/c-Met bispecific antibody.	Phase 2 (Non-small cell lung carcinoma (NSCLC)) Phase 2 (Gastric carcinoma, Head and neck squamous cell carcinoma (HNSCC), Esophageal squamous cell carcinoma)

## Detailed Therapy Results

Alteration	Drug	Trade Name	Target	Current Status
	Mobocertinib	Exkivity	Mutation-specific Egfr/Her2 inhibitor.	Phase 3 (Non-small cell lung carcinoma (NSCLC)) Phase 3 (NSCLC with EGFR exon 20 insertion, Lung cancer)
	Modotuximab		Anti-EGFR antibody.	Phase 1 (Gastric carcinoma, Colorectal carcinoma (CRC))
	Nazartinib		Third generation EGFR mutant-specific (T790M, L858R, exon 19 deletion) tyrosine kinase inhibitor.	Phase 2 (Non-small cell lung carcinoma (NSCLC))
	Neratinib	Nerlynx	Egfr/Her2/ErbB4 inhibitor.	Phase 2 (Non-small cell lung carcinoma (NSCLC)) FDA Approved in other indications (Breast carcinoma (HER2+))
	NX-019		Egfr inhibitor.	Phase 1 (Solid Tumor)
	Olafertinib		Third generation mutation-specific (T790M, L858R, exon 19 deletion) Egfr tyrosine kinase inhibitor.	Phase 2 (Non-small cell lung carcinoma (NSCLC)) Phase 1 (Solid Tumor)
	Osimertinib	Tagrisso	Egfr T790M inhibitor.	Phase 3 (Non-small cell lung carcinoma (NSCLC)) FDA Approved in other indications (EGFR-mutant lung carcinoid, EGFR-mutant lung large cell neuroendocrine carcinoma, EGFR-mutant NSCLC)
	Pirotinib		ErbB family inhibitor.	Phase 1 (Solid Tumor)
	Pozotinib		Egfr/Her2/ErbB4 kinase inhibitor.	Phase 2 (Non-small cell lung carcinoma (NSCLC)) Phase 2 (Gastric carcinoma, Head and neck squamous cell carcinoma (HNSCC), Breast carcinoma, Esophageal squamous cell carcinoma, Colorectal carcinoma (CRC))
	Pyrotinib		Egfr/Her2 kinase inhibitor.	Phase 2 (Non-small cell lung carcinoma (NSCLC)) Phase 3 (Breast carcinoma)
	Rezivertinib		Egfr T790M inhibitor.	Phase 3 (Non-small cell lung carcinoma (NSCLC))
	SKLB1028		Egfr/Flt3/c-Abl inhibitor.	Phase 2 (Acute myeloid leukemia (AML))
	Sunvozertinib		Bispecific anti-Egfr/Her2 monoclonal antibody.	Phase 2 (Non-small cell lung carcinoma (NSCLC)) Phase 2 (Non-Hodgkin lymphoma (NHL))
	TAS2940		Egfr/Her2 kinase inhibitor.	Phase 1 (Solid Tumor)
	TAS3351		Fourth generation Egfr inhibitor targeting T790M and T790M/C797S mutations.	Phase 2 (Non-small cell lung carcinoma (NSCLC))
	TAVO412		Anti-c-Met/anti-EGFR/anti-VEGF trispecific antibody.	Phase 1 (Solid Tumor)
	Varlitinib		Egfr/Her2 kinase inhibitor.	Phase 2 (Gastric carcinoma, Hepatocellular carcinoma (HCC), Pancreatic carcinoma, Cholangiocarcinoma)
	WSD0922-FU		Blood-brain barrier penetrable EGFR/EGFRvIII	Phase 1 (Non-small cell lung carcinoma (NSCLC)) Phase 1 (Glioblastoma,

## Detailed Therapy Results

Alteration	Drug	Trade Name	Target	Current Status
			inhibitor.	Anaplastic astrocytoma)
	ZN-e4		Egfr T790M inhibitor.	Phase 1 (Non-small cell lung carcinoma (NSCLC))
	Zorifertinib		Egfr tyrosine kinase inhibitor.	Phase 2 (Non-small cell lung carcinoma (NSCLC))
PTEN Q149Gfs*8	ABTL0812		Inhibitor of mTORC1/mTORC2 /Dhfr.	Phase 2 (Non-small cell lung carcinoma (NSCLC)) Phase 2 (Pancreatic carcinoma, Endometrial carcinoma)
	Alpelisib	Piqray	p110-alpha inhibitor.	Phase 2 (Non-small cell lung carcinoma (NSCLC)) FDA Approved in other indications (Breast carcinoma (HR+, HER2-) with a PIK3CA mutation, as determined by a validated test)
	Archexin	Archexin	Akt1 antisense oligonucleotide.	Phase 1 (Solid Tumor) Phase 2 (Pancreatic carcinoma, Renal cell carcinoma)
	ARQ 751		Akt inhibitor.	Phase 1 (Solid Tumor)
	AZD8186		p110-beta and p110-delta small molecule inhibitor.	Phase 1 (Solid Tumor) Phase 1 (Breast carcinoma (triple negative), Lung squamous cell carcinoma)
	Buparlisib		PI3K inhibitor.	Phase 2 (Non-small cell lung carcinoma (NSCLC)) Phase 3 (Head and neck squamous cell carcinoma (HNSCC), Breast carcinoma)
	Capivasertib	Truqap	Akt inhibitor.	Phase 2 (Non-small cell lung carcinoma (NSCLC)) FDA Approved in other indications (Breast carcinoma (HR+, HER2-) with a PIK3CA/AKT1 /PTEN mutation)
	CC-115		DNA-PK/dual mTORC1/2 kinase inhibitor.	Phase 1 (Solid Tumor) Phase 2 (Glioblastoma, Glioma)
	Copanlisib	Aliqopa	PI3K inhibitor.	Phase 2 (Non-small cell lung carcinoma (NSCLC)) Phase 3 (Non-Hodgkin lymphoma (NHL))
	CYH33		p110-alpha-specific small molecule inhibitor.	Phase 1 (Solid Tumor)
	Everolimus	Afinitor	mTOR inhibitor, immunosuppressant.	Phase 2 (Non-small cell lung carcinoma (NSCLC)) FDA Approved in other indications (TSC associated renal angiomyolipoma and subependymal giant cell astrocytoma, Renal cell carcinoma, Gastrointestinal neuroendocrine carcinoma, Lung carcinoid, Breast carcinoma (hormone receptor +, HER2-), Subependymal giant cell astrocytoma)
	Gedatolisib		Dual PI3K/mTOR inhibitor.	Phase 2 (Non-small cell lung carcinoma (NSCLC)) Phase 3 (Breast carcinoma)
	GSK2636771		p110-beta small molecule inhibitor.	Phase 2 (Solid Tumor) Phase 1 (Prostate carcinoma)
	HS-10352		p110-alpha-specific small molecule inhibitor.	Phase 1 (Breast carcinoma)
	Inavolisib	Itovebi	PI3K inhibitor.	Phase 1 (Solid Tumor) Phase 3 (Cancer, Breast carcinoma)
	Ipatasertib		Akt inhibitor.	Phase 1 (Solid Tumor) Phase 3 (Prostate carcinoma, Breast carcinoma)



## Detailed Therapy Results

Alteration	Drug	Trade Name	Target	Current Status
	ME-344		Dual mTORC1/mTORC2 inhibitor.	Phase 1 (Solid Tumor) Phase 2 (Ovarian carcinoma, Small cell lung carcinoma (SCLC))
	MEN1611		PI3K inhibitor.	Phase 2 (Metaplastic breast carcinoma, Colorectal carcinoma (CRC))
	Miransertib		Akt inhibitor.	Phase 1 (Non-small cell lung carcinoma (NSCLC)) Phase 1 (Solid Tumor)
	MK-2206		Akt inhibitor.	Phase 2 (Non-small cell lung carcinoma (NSCLC)) Phase 2 (Gastric carcinoma, Gallbladder carcinoma, Hepatocellular carcinoma (HCC), Pancreatic carcinoma, Hodgkin lymphoma (HL), Lymphoma, Melanoma, Carcinoid tumor, Prostate carcinoma, Endometrial carcinoma, Renal cell carcinoma, Ovarian carcinoma, Small cell lung carcinoma (SCLC), Adenoid cystic carcinoma, Head and neck carcinoma, Gastroesophageal junction carcinoma, Gastrointestinal neuroendocrine carcinoma, Mucinous colon adenocarcinoma, Breast carcinoma, Chronic lymphocytic leukemia (CLL), Acute myeloid leukemia (AML), Cholangiocarcinoma, Thymic neoplasm, Peritoneal papillary serous carcinoma, Lung cancer, Colorectal carcinoma (CRC), Diffuse large B-cell lymphoma (DLBCL))
	MSC2363318A		Akt1, Akt3, and p70S6K inhibitor.	Phase 1 (Solid Tumor) Phase 1 (Lymphoma)
	Niraparib	Zejula	PARP inhibitor.	Phase 3 (Non-small cell lung carcinoma (NSCLC)) FDA Approved in other indications (Prostate carcinoma with BRCA1/2 mutation, Ovarian carcinoma with BRCA1/2 mutation)
	Olaparib	Lynparza	PARP inhibitor.	Phase 2 (Lung adenocarcinoma) FDA Approved in other indications (Pancreatic adenocarcinoma with germline BRCA1/2 mutation, Prostate cancer with HRR gene mutation, Ovarian carcinoma with BRCA1/2 mutation, Breast carcinoma with germline BRCA1/2 mutation)
	Onatasertib		Dual mTORC1/mTORC2 inhibitor.	Phase 1 (Non-small cell lung carcinoma (NSCLC)) Phase 2 (Solid Tumor, Bladder neuroendocrine carcinoma, Multiple myeloma (MM), Diffuse large B-cell lymphoma (DLBCL))
	Paxalisib		Dual PI3K/mTOR inhibitor.	Phase 2 (Glioblastoma, Breast carcinoma)
	RLY-2608		p110-alpha-specific small molecule inhibitor.	Phase 1 (Solid Tumor)
	RLY-5836		p110-alpha-specific small molecule inhibitor.	Phase 1 (Solid Tumor)
	RMC-5552		mTORC1-specific inhibitor.	Phase 1 (Solid Tumor)
	Samotolisib		Dual PI3K/mTOR inhibitor.	Phase 2 (Non-small cell lung carcinoma (NSCLC)) Phase 2 (Pancreatic carcinoma, Prostate carcinoma)
	Sapanisertib		Dual mTORC1/mTORC2 inhibitor.	Phase 2 (Non-small cell lung carcinoma (NSCLC)) Phase 2 (Uterine carcinosarcoma, Hepatocellular carcinoma (HCC), Lymphoma, Merkel cell carcinoma, Anaplastic thyroid carcinoma, Pancreatic neuroendocrine tumor, Prostate carcinoma, Endometrial carcinoma, Renal cell



## Detailed Therapy Results

Alteration	Drug	Trade Name	Target	Current Status
				carcinoma, Ovarian carcinoma, Urothelial carcinoma, Bladder carcinoma, Breast carcinoma, Multiple myeloma (MM), Non-Hodgkin lymphoma (NHL), Lung cancer, Sarcoma, Acute lymphoblastic leukemia (ALL))
	Serabelisib		PI3K inhibitor.	Phase 1 (Solid Tumor) Phase 2 (Endometrial carcinoma, Renal cell carcinoma, Breast carcinoma (triple negative))
	Talazoparib	Talzenna	PARP inhibitor.	Phase 2 (Non-small cell lung carcinoma (NSCLC)) FDA Approved in other indications (Prostate cancer with HRR gene mutation, Breast carcinoma with germline BRCA1/2 mutation)
	TAS-117		Akt inhibitor.	Phase 2 (Solid Tumor)
	TAS0612		Akt/p70S6K/p90RSK1 multikinase inhibitor.	Phase 1 (Solid Tumor)
	Temsirolimus	Torisel	mTOR inhibitor.	Phase 2 (Non-small cell lung carcinoma (NSCLC)) FDA Approved in other indications (Renal cell carcinoma)
	TOS-358		p110-alpha-specific small molecule inhibitor.	Phase 1 (Non-small cell lung carcinoma (NSCLC)) Phase 1 (Gastric carcinoma, Endometrial carcinoma, Head and neck squamous cell carcinoma (HNSCC), Ovarian carcinoma, Urothelial carcinoma, Breast carcinoma, Cervical carcinoma, Colorectal carcinoma (CRC))
	Triciribine		DNA synthesis and Akt inhibitor.	Phase 1 (Solid Tumor) Phase 2 (Ovarian carcinoma, Breast carcinoma)
	Uprosertib		Akt inhibitor.	Phase 1 (Solid Tumor) Phase 2 (Melanoma, Uveal melanoma, Breast carcinoma, Cervical carcinoma, Acute myeloid leukemia (AML), Multiple myeloma (MM))
	Vistusertib		Dual mTORC1/mTORC2 inhibitor.	Phase 2 (Non-small cell lung carcinoma (NSCLC)) Phase 2 (Meningioma, Small cell lung carcinoma (SCLC), Solid Tumor, Diffuse large B-cell lymphoma (DLBCL))
	WGI-0301		Nanoliposomal Archexin, an Akt1 antisense oligonucleotide.	Phase 1 (Solid Tumor)
KRAS L19F	ASN007		ERK1/2 kinase inhibitor.	Phase 1 (Solid Tumor)
	ASTX029		ERK1/2 kinase inhibitor.	Phase 2 (Solid Tumor)
	Avutometinib		Dual Raf/MEK kinase inhibitor.	Phase 2 (Non-small cell lung carcinoma (NSCLC)) Phase 2 (Uveal melanoma, Ovarian carcinoma)
	BBP-398		Shp-2 inhibitor.	Phase 1 (Non-small cell lung carcinoma (NSCLC)) Phase 1 (Solid Tumor)
	BDTX-4933		Braf class 1, 2, and 3 inhibitor.	Phase 1 (Solid Tumor)
	BI 1701963		Pan-K-Ras inhibitor targeting the interaction of K-Ras and SOS-1.	Phase 1 (Non-small cell lung carcinoma (NSCLC)) Phase 1 (Solid Tumor, Brain and Central Nervous System Tumors)
	BI 3011441		MEK1,2 inhibitor.	Phase 1 (Solid Tumor)
	Binimetinib	Mektovi	MEK1,2 inhibitor.	Phase 2 (Non-small cell lung carcinoma (NSCLC)) FDA Approved in other indications (Melanoma

## Detailed Therapy Results

Alteration	Drug	Trade Name	Target	Current Status
				with BRAF V600E/K mutation)
	BMF-219		Covalent menin inhibitor.	Phase 1 (Non-small cell lung carcinoma (NSCLC)) Phase 2 (Diabetes)
	Brimarafenib		Braf class 1, 2, and 3 inhibitor.	Phase 2 (Solid Tumor) Phase 1 (Pancreatic ductal adenocarcinoma, Colorectal carcinoma (CRC))
	Cobimetinib	Cotellic	MEK1,2 inhibitor.	Phase 3 (Non-small cell lung carcinoma (NSCLC)) FDA Approved in other indications (Melanoma with BRAF V600E/K mutation, Histiocytic and dendritic cell neoplasms)
	Defactinib		Focal adhesion kinase (FAK) inhibitor, upstream of MAPK pathway.	Phase 2 (Non-small cell lung carcinoma (NSCLC)) Phase 2 (Lymphoma, Solid Tumor, Multiple myeloma (MM))
	E6201		MEK1,2, Mekk1, and Flt3 inhibitor.	Phase 1 (Solid Tumor) Phase 1 (Acute myeloid leukemia (AML))
	ERAS-601		Shp-2 inhibitor.	Phase 2 (Solid Tumor)
	ET0038		Shp-2 inhibitor.	Phase 1 (Solid Tumor)
	HBI-2376		Shp-2 inhibitor.	Phase 1 (Solid Tumor)
	HMPL-295		ERK1/2 kinase inhibitor.	Phase 1 (Solid Tumor)
	IMM-1-104		MEK1,2 inhibitor.	Phase 2 (Non-small cell lung carcinoma (NSCLC)) Phase 2 (Melanoma, Pancreatic ductal adenocarcinoma)
	JAB-3068		Shp-2 inhibitor.	Phase 2 (Non-small cell lung carcinoma (NSCLC)) Phase 2 (Head and neck squamous cell carcinoma (HNSCC), Esophageal carcinoma)
	JAB-3312		Shp-2 inhibitor.	Phase 1 (Solid Tumor) Phase 1 (Brain and Central Nervous System Tumors)
	JSI-1187		ERK1/2 kinase inhibitor.	Phase 1 (Solid Tumor)
	KO-2806		Farnesyl transferase inhibitor.	Phase 1 (Solid Tumor)
	LTT462		ERK1/2 kinase inhibitor.	Phase 1 (Non-small cell lung carcinoma (NSCLC)) Phase 1 (Solid Tumor)
	LUNA18		Ras peptide inhibitor.	Phase 1 (Solid Tumor)
	Mirdametinib		MEK1,2 inhibitor.	Phase 2 (Non-small cell lung carcinoma (NSCLC)) Phase 2 (Glioma, Neurofibroma, Breast carcinoma, Glioneuronal tumor, Neurofibromatosis type 1, Lung cancer, Colorectal carcinoma (CRC))
	MK-8353		ERK1/2 kinase inhibitor.	Phase 1 (Solid Tumor)
	MRTX0902		Pan-K-Ras inhibitor targeting the interaction of K-Ras and SOS-1.	Phase 1 (Solid Tumor)
	PF-07284892		Shp-2 inhibitor.	Phase 1 (Non-small cell lung carcinoma (NSCLC)) Phase 1 (Solid Tumor, Brain and Central Nervous System Tumors)
	PF-07799544		MEK Brain Penetrant Inhibitor.	Phase 1 (Non-small cell lung carcinoma (NSCLC)) Phase 1 (Glioma, Melanoma, Thyroid carcinoma, Colorectal carcinoma (CRC))

## Detailed Therapy Results

Alteration	Drug	Trade Name	Target	Current Status
	Pimasertib		MEK1,2 inhibitor.	Phase 1 (Solid Tumor) Phase 2 (Melanoma, Pancreatic ductal adenocarcinoma, Ovarian carcinoma, Hematologic malignancies, Colorectal carcinoma (CRC))
	RMC-6236		Multispecific Ras inhibitor.	Phase 1 (Solid Tumor)
	RSC-1255		Ras inhibitor.	Phase 1 (Solid Tumor)
	Selumetinib	Koselugo	MEK1,2 inhibitor.	Phase 3 (Non-small cell lung carcinoma (NSCLC)) FDA Approved in other indications (NF1-related plexiform neurofibroma)
	SHR7390		MEK1,2 inhibitor.	Phase 1 (Solid Tumor)
	Temuterkib		ERK1/2 kinase inhibitor.	Phase 1 (Non-small cell lung carcinoma (NSCLC)) Phase 1 (Pancreatic ductal adenocarcinoma, Solid Tumor)
	Tizaterkib		ERK1/2 kinase inhibitor.	Phase 1 (Solid Tumor) Phase 1 (Hematologic malignancies)
	TNO155		Shp-2 inhibitor.	Phase 2 (Non-small cell lung carcinoma (NSCLC)) Phase 2 (Colorectal carcinoma (CRC))
	Trametinib	Mekinist	MEK1,2 inhibitor.	Phase 2 (Non-small cell lung carcinoma (NSCLC)) FDA Approved in other indications (Melanoma with BRAF V600 mutation)
	Ulixertinib		ERK1/2 kinase inhibitor.	Phase 1 (Solid Tumor) Phase 2 (Gastric carcinoma, Melanoma, Histiocytic and dendritic cell neoplasms, Langerhans cell histiocytosis (LCH), Uveal melanoma, Acute myeloid leukemia (AML), Gastrointestinal carcinoma, Non-Hodgkin lymphoma (NHL), Myelodysplastic Syndrome (MDS))
CTNNB1 S45C	Vociprotafib		Shp-2 inhibitor.	Phase 2 (Non-small cell lung carcinoma (NSCLC)) Phase 2 (Solid Tumor)
	DKN-01		Anti-DKK-1 monoclonal antibody.	Phase 1 (Solid Tumor)
	E7386		CBP/beta-catenin inhibitor.	Phase 1 (Solid Tumor) Phase 2 (Hepatocellular carcinoma (HCC), Colorectal carcinoma (CRC))
	FOG-001		Beta-catenin polypeptide inhibitor.	Phase 2 (Non-small cell lung carcinoma (NSCLC)) Phase 2 (Gastric carcinoma, Gastroesophageal junction carcinoma, Colorectal carcinoma (CRC))
	Tegavivint		Wnt/beta-catenin pathway inhibitor.	Phase 1 (Non-small cell lung carcinoma (NSCLC)) Phase 2 (Hepatocellular carcinoma (HCC), Wilms tumor, Ewing sarcoma, Desmoid fibromatosis)

Relevance of Detected Alterations

Alteration	Role in Disease	Effect on Drug Sensitivity	Effect on Drug Resistance
EGFR E746_A750del	The presence of an EGFR abnormality (mutation, amplification, or overexpression) can result in an overabundance or overactivity of Egfr protein, which can lead to excessive proliferation. <sup>(1)</sup> .	The presence of a sensitizing EGFR mutation in a tumor is the strongest biological predictor of sensitivity to an Egfr tyrosine kinase inhibitor (TKI). Compared with conventional chemotherapy, Egfr TKIs have been shown to improve progression-free survival in non-small cell lung cancer patients whose tumors harbor EGFR mutations. <sup>(2-5)</sup> . The Egfr TKIs erlotinib, afatinib, gefitinib, osimertinib, and dacomitinib, as well as the combination of amivantamab plus lazertinib, have been approved by the FDA for the treatment of non-small cell lung cancer (NSCLC) with exon 19 deletion or L858R EGFR mutations; osimertinib has additionally been approved for the treatment of NSCLC with EGFR T790M. <sup>(2,5-14)</sup> . Afatinib has additionally been FDA-approved for the treatment of NSCLC with S768I, L861Q, and/or G719X mutations. <sup>(15)</sup> . The combination of erlotinib and ramucirumab as well as osimertinib plus platinum-based chemotherapy have been FDA-approved for the treatment of metastatic NSCLC patients with tumors harboring an EGFR exon 19 deletion or the exon 21 L858R mutation. <sup>(16-18)</sup> . Amivantamab in combination with carboplatin and pemetrexed has been FDA-approved for the treatment of adult patients with locally advanced or metastatic NSCLC harboring EGFR Exon 19 deletions or Exon 21 L858R substitution mutations whose disease has progressed on or after treatment with an EGFR TKI. <sup>(12,19)</sup> . Amivantamab has also been approved by the FDA for NSCLC patients with EGFR exon 20 insertions, whose disease has progressed on or after platinum-based chemotherapy and as frontline therapy in combination with carboplatin and pemetrexed. The accelerated FDA approval of mobocertinib for NSCLC patients with EGFR exon 20 insertions has been withdrawn due to lack of progression-free survival benefit in the confirmatory Phase 3 trial. <sup>(20-23)</sup> . Studies have reported non-squamous NSCLC patients with metastatic disease and tumors harboring an EGFR exon 19 deletion or L858R mutation to be sensitive to osimertinib, erlotinib, afatinib, gefitinib, dacomitinib, and the combination of erlotinib plus ramucirumab. <sup>(2,5-7,11,16,24)</sup> . Less	Some patients with EGFR-mutant NSCLC exhibit resistance to Egfr inhibition; resistance has been associated with insertions in EGFR exon 20, the T790M mutation in EGFR, and amplification of either MET or ERBB2. <sup>(26-30)</sup> . Third generation irreversible Egfr TKIs that target the EGFR T790M mutation have shown efficacy in T790M-mutant NSCLC, including osimertinib, which has received approval by the FDA, EMA, and PMDA for the treatment of EGFR T790M-mutant metastatic NSCLC. <sup>(31-35)</sup> . Several studies have reported that resistance to Egfr TKIs in NSCLC is mediated by the transformation of NSCLC cell types to those of SCLC with neuroendocrine features. <sup>(36-39)</sup> .

Relevance of Detected Alterations

Alteration	Role in Disease	Effect on Drug Sensitivity	Effect on Drug Resistance
		common activating EGFR mutations have variable sensitivity to EGFR tyrosine kinase inhibitors. <sup>(25)</sup>	
<i>PTEN</i> Q149Gfs*8	Loss of PTEN (through mutation or deletion) can lead to uncontrolled cell growth and suppression of apoptosis. <sup>(40)</sup> . PTEN haploinsufficiency has been associated with tumorigenesis in some tumor types, including astrocytoma and prostate cancer. <sup>(41-44)</sup> . PTEN germline mutations are found in several cancer-predisposition syndromes, such as Cowden syndrome and Proteus syndrome. <sup>(45)</sup> . Loss of Pten protein expression has been associated with higher stage, lymph node metastases, and poorly differentiated disease in NSCLC patients. <sup>(46-51)</sup> . Deletion of PTEN has been shown to lead to the formation of lung squamous cell carcinoma tumors in preclinical studies. <sup>(52,53)</sup>	Because PTEN negatively regulates the PI3K/Akt/mTOR pathway, PTEN loss or mutation leads to activation of the PI3K pathway and may therefore predict sensitivity to inhibitors of the PI3K/Akt /mTOR pathway. <sup>(54)</sup> . The PI3K inhibitors alpelisib and copanlisib, the Akt inhibitor capivasertib, and the mTOR inhibitors everolimus and temsirolimus have been approved in specific cancer indications. These and other PI3K, Akt, and mTOR inhibitors, as well as dual PI3K/mTOR inhibitors are also currently in clinical trials, alone or in combination with other therapies. <sup>(55-59)</sup> . The Akt inhibitor capivasertib has been approved by the FDA in combination with fulvestrant for the treatment of adult patients with hormone receptor positive, Her2 negative locally advanced or metastatic breast cancer with PIK3CA /AKT1/PTEN alterations as detected by an FDA-approved test following progression on at least one endocrine-based regimen in the metastatic setting or recurrence on or within 12 months of completing adjuvant therapy. <sup>(60)</sup> . In addition, preclinical studies have shown that PTEN-deficient tumors may be sensitive to PARP inhibitors, and PARP inhibitors are in clinical trials for patients with PTEN-deficient tumors. <sup>(61-68)</sup>	Preclinical studies have associated decreased expression of Pten with gefitinib and erlotinib resistance in EGFR-mutant NSCLC cell lines. <sup>(69-71)</sup> . Mutation of PTEN has been reported to be associated with a low objective response rate (ORR) to Egr tyrosine kinase inhibitors in lung adenocarcinomas harboring EGFR mutations, with 7.3% ORR in PTEN-mutated cases compared to 70.9% ORR in wild-type PTEN lung adenocarcinomas. <sup>(72)</sup>
<i>CTNNB1</i> S45C	CTNNB1 can act as an oncogene and altered expression of beta-catenin can lead to abnormal signaling in various diseases, including modulation of gene transcription to drive cancer initiation, progression, survival, and relapse. <sup>(73)</sup> . Beta-catenin expression has been reported to be a marker for epithelial-to-mesenchymal transition (EMT) in non-small cell lung carcinoma (NSCLC) cells. <sup>(74)</sup> . In addition, one study analyzing 143 NSCLC cases reported that CTNNB1 promoter hypermethylation was correlated with loss of beta-catenin expression, positive lymph node metastasis, and higher TNM staging. <sup>(75)</sup>	CTNNB1 mutations, particularly those in exon 3, have been associated with increased beta-catenin protein stability and activation of the Wnt pathway. <sup>(76-81)</sup> . At the present time, there are no approved therapies that target CTNNB1 mutation, however, beta-catenin/Wnt pathway inhibitors are under preclinical and clinical investigation. <sup>(82-87)</sup>	Beta-catenin activation has been associated with immune evasion and resistance to checkpoint inhibitors in preclinical cancer models. <sup>(88-93)</sup> . Hyperactivation of the Wnt/beta-catenin pathway may confer resistance to inhibitors of PI3K and Akt. <sup>(94)</sup> . Activation or increased expression of beta-catenin has been suggested to play a role in resistance to a variety of agents in NSCLC cell lines, including gemcitabine, cisplatin, or gefitinib. <sup>(74,96)</sup>
<i>KRAS</i> L19F	The KRAS gene is one of the most commonly mutated genes in human	Many of the current attempts to target K-Ras are directed against its	In some cancer types, such as colorectal cancer (CRC) and non-small

Relevance of Detected Alterations

Alteration	Role in Disease	Effect on Drug Sensitivity	Effect on Drug Resistance
	<p>malignancies, with high incidences in pancreatic, colorectal, and lung cancers. <sup>(97-99)</sup>. KRAS mutation, particularly G12C, has been associated with smoking in NSCLC patients; additionally, KRAS mutations have also been associated with adenocarcinoma histology and are generally mutually exclusive with EGFR mutations and ALK rearrangements. <sup>(100-112)</sup>. Studies analyzing KRAS mutation association with PD-L1 expression in NSCLC patients have reported mixed results; while two older large meta-analyses have reported no association or negative association, smaller studies and one newer large meta-analysis have reported a positive association. <sup>(105,112-127)</sup>.</p>	<p>downstream signaling pathways, Raf /MEK/ERK and PI3K/Akt/mTOR. <sup>(128,129)</sup>. Clinical studies have suggested limited efficacy of MEK inhibitors in KRAS mutant tumors; however, combinations of MEK inhibitors with other targeted therapies may still be relevant. <sup>(130-139)</sup>. Other clinical approaches are being investigated preclinically and clinically in the context of KRAS-mutant tumors, including FAK and Shp-2 inhibitors. <sup>(140-145)</sup>. In addition, inhibitors specifically targeting KRAS G12C and cell-based therapies targeting KRAS G12V and G12D are being investigated clinically and preclinically. <sup>(146-149)</sup>. Sotorasib and adagrasib have been FDA-approved in patients with locally advanced or metastatic non-small cell lung carcinoma harboring a KRAS G12C mutation, as determined by an FDA-approved test, following treatment with at least one prior systemic therapy. <sup>(150-154)</sup>. In addition, combinations of adagrasib or sotorasib with cetuximab or panitumumab have been reported to provide clinical benefit in CRC patients with KRAS G12C mutation. <sup>(155-158)</sup>. Adagrasib plus cetuximab has been FDA-approved for treatment of adults with KRAS G12C-mutated locally advanced or metastatic colorectal cancer, as determined by an FDA-approved test, who have received prior treatment with fluoropyrimidine-, oxaliplatin-, and irinotecan-based chemotherapy. <sup>(157,159)</sup>. Studies analyzing KRAS mutation association with clinical benefit of PD-1/PD-L1 inhibitors to NSCLC patients have reported mixed results, with some studies reporting no significant association while other studies have reported that KRAS mutation correlated with improved clinical outcome of NSCLC patients treated with PD-1/PD-L1 inhibitors. <sup>(115,160-163)</sup>.</p>	<p>cell lung cancer (NSCLC), activating KRAS mutations and KRAS amplification have been associated with resistance to Egfr-targeted therapies. <sup>(164-172)</sup>. While some studies have suggested that KRAS mutation status may predict lack of response to the Egfr inhibitors erlotinib and gefitinib in NSCLC patients, a retrospective study suggests that there is no significant difference in response to Egfr tyrosine kinase inhibitors among NSCLC patients with KRAS wild-type and KRAS mutation, when EGFR mutation status is included in the analysis. <sup>(173-177)</sup>. In addition, KRAS mutations have been reported in 1-3% of EGFR-mutant NSCLC patients following emergence of resistance to osimertinib. Treatment with K-ras inhibitors sotorasib or adagrasib in combination with osimertinib may be an effective strategy in patients with KRAS G12C mutation. <sup>(178-180)</sup>. Case studies of NSCLC patients harboring ALK mutations or EML4-ALK fusions have reported the emergence of KRAS alterations upon acquired resistance to crizotinib, demonstrating a role for KRAS in crizotinib resistance in NSCLC. <sup>(181-185)</sup>. Secondary KRAS mutations have also been reported in case studies of NSCLC patients with KRAS G12C following emergence of resistance to sotorasib or adagrasib. <sup>(186-189)</sup>.</p>
TP53 R175H	<p>Loss of tumor suppressor p53, which is encoded by the TP53 gene, is common in aggressive advanced cancers. <sup>(190)</sup>. Carriers of a germline mutation in TP53 have Li-Fraumeni Syndrome, an inherited cancer syndrome resulting in multiple tumors in early adulthood, including breast cancer, brain tumors, and leukemias. <sup>(191-193)</sup>. Expression of p53 in normal</p>	<p>At present, there are no approved therapies targeting TP53 alterations, despite their high prevalence in cancer. Therapeutic approaches under investigation include gene therapy for TP53 and (dendritic cell-based) TP53 vaccines. <sup>(208-210)</sup>. Inhibition of components of the DNA damage checkpoint, including Wee1, has been reported to enhance the activity of</p>	<p>Mutations in TP53 may increase resistance to ionizing radiation therapy. <sup>(220,221)</sup>.</p>



Relevance of Detected Alterations

Alteration	Role in Disease	Effect on Drug Sensitivity	Effect on Drug Resistance
	cells is low; however, TP53 alterations, including those that result in loss of p53 tumor suppressor function, may lead to stabilization and increased expression of p53, particularly in the nucleus, and several studies have shown that it may have oncogenic gain-of-function effects. (194-198). TP53 alterations are believed to be early events in NSCLC, preceding lymph node metastasis. (199). TP53 mutation and expression of p53 have been correlated with the lung squamous cell carcinoma subtype, and p53 expression in lung squamous cell carcinoma has also been associated with disease stage and higher grade tumors. (200-203). TP53 mutation has been associated with PD-L1 expression and T-cell infiltration in lung adenocarcinoma samples. (121,126,204-206). TP53 mutations have been significantly associated with the development of distant metastases after diagnosis in early-stage NSCLC in a cohort of 759 patients. (207).	DNA-damaging agents in preclinical cancer models with deficiency of p53 function. (211-213). Clinical trials of the Wee1 inhibitor adavosertib (MK-1775) are currently underway for patients with solid tumors and hematologic malignancies. Studies have reported Aurora kinase A to be activated in cells harboring TP53 mutation, and Aurora kinase A and B inhibitors have been reported to activate wild-type p53 in cellular assays; thus, tumors retaining a wild-type TP53 allele may benefit from Aurora kinase inhibitors. (214-219).	

## References

1. Ciardiello F, Tortora G "EGFR antagonists in cancer treatment." *The New England journal of medicine*(2008): 1160-74
2. Mok T, Wu Y, Thongprasert S, Yang C, Chu D, Saijo N, Sunpaweravong P, Han B, Margono B, Ichinose Y, Nishiwaki Y, Ohe Y, Yang J, Chewaskulyong B, Jiang H, Duffield E, Watkins C, Armour A, Fukuoka M "Gefitinib or carboplatin-paclitaxel in pulmonary adenocarcinoma." *The New England journal of medicine*(2009): 947-57
3. Rosell R, Moran T, Queralt C, Porta R, Cardenal F, Camps C, Majem M, Lopez-Vivanco G, Isla D, Provencio M, Insa A, Massuti B, Gonzalez-Larriba J, Paz-Ares L, Bover I, Garcia-Campelo R, Moreno M, Catot S, Rolfo C, Reguart N, Palmero R, Sánchez J, Bastus R, Mayo C, Bertran-Alamillo J, Molina M, Sanchez J, Taron M "Screening for epidermal growth factor receptor mutations in lung cancer." *The New England journal of medicine*(2009): 958-67
4. Tsao M, Sakurada A, Cutz J, Zhu C, Kamel-Reid S, Squire J, Lorimer I, Zhang T, Liu N, Daneshmand M, Marrano P, da Cunha Santos G, Lagarde A, Richardson F, Seymour L, Whitehead M, Ding K, Pater J, Shepherd F "Erlotinib in lung cancer - molecular and clinical predictors of outcome." *The New England journal of medicine* (2005): 133-44
5. Rosell R, Carcereny E, Gervais R, Vergnenegre A, Massuti B, Felip E, Palmero R, Garcia-Gomez R, Pallares C, Sanchez J, Porta R, Cobo M, Garrido P, Longo F, Moran T, Insa A, De Marinis F, Corre R, Bover I, Illiano A, Dansin E, de Castro J, Milella M, Reguart N, Altavilla G, Jimenez U, Provencio M, Moreno M, Terrasa J, Muñoz-Langa J, Valdivia J, Isla D, Domine M, Molinier O, Mazieres J, Baize N, Garcia-Campelo R, Robinet G, Rodriguez-Abreu D, Lopez-Vivanco G, Gebbia V, Ferrera-Delgado L, Bombardieri P, Bernabe R, Bearz A, Arta A, Cortesi E, Rolfo C, Sanchez-Ronco M, Drozdowski A, Queralt C, de Aguirre I, Ramirez J, Sanchez J, Molina M, Taron M, Paz-Ares L "Erlotinib versus standard chemotherapy as first-line treatment for European patients with advanced EGFR mutation-positive non-small-cell lung cancer (EORTC): a multicentre, open-label, randomised phase 3 trial." *The Lancet. Oncology*(2012): 239-46
6. Soria J, Ohe Y, Vansteenkiste J, Reungwetwattana T, Chewaskulyong B, Lee K, Dechaphunkul A, Imamura F, Nogami N, Kurata T, Okamoto I, Zhou C, Cho B, Cheng Y, Cho E, Voon P, Planchard D, Su W, Gray J, Lee S, Hodge R, Marotti M, Rukazekov Y, Ramalingam S "Osimertinib in Untreated EGFR-Mutated Advanced Non-Small-Cell Lung Cancer." *The New England journal of medicine*(2018): 113-125
7. Wu Y, Cheng Y, Zhou X, Lee K, Nakagawa K, Niho S, Tsui F, Linke R, Rosell R, Corral J, Migliorino M, Pluzanski A, Sbar E, Wang T, White J, Nadanaciva S, Sandin R, Mok T "Dacomitinib versus gefitinib as first-line treatment for patients with EGFR-mutation-positive non-small-cell lung cancer (ARCHER 1050): a randomised, open-label, phase 3 trial." *The Lancet. Oncology*(2017): 1454-1466
8. Mok T, Cheng Y, Zhou X, Lee K, Nakagawa K, Niho S, Lee M, Linke R, Rosell R, Corral J, Migliorino M, Pluzanski A, Sbar E, Wang T, White J, Wu Y "Improvement in Overall Survival in a Randomized Study That Compared Dacomitinib With Gefitinib in Patients With Advanced Non-Small-Cell Lung Cancer and EGFR-Activating Mutations." *Journal of clinical oncology : official journal of the American Society of Clinical Oncology*(2018): 2244-2250
9. Shepherd F, Rodrigues Pereira J, Ciuleanu T, Tan E, Hirsh V, Thongprasert S, Campos D, Maoleekoonpiroj S, Smylie M, Martins R, van Kooten M, Dediu M, Findlay B, Tu D, Johnston D, Bezjak A, Clark G, Santabárbara P, Seymour L "Erlotinib in previously treated non-small-cell lung cancer." *The New England journal of medicine*(2005): 123-32
10. Sequist L, Yang J, Yamamoto N, O'Byrne K, Hirsh V, Mok T, Geater S, Orlov S, Tsai C, Boyer M, Su W, Bannouna J, Kato T, Gorbunova V, Lee K, Shah R, Massey D, Zazulina V, Shahidi M, Schuler M "Phase III study of afatinib or cisplatin plus pemetrexed in patients with metastatic lung adenocarcinoma with EGFR mutations." *Journal of clinical oncology : official journal of the American Society of Clinical Oncology*(2013): 3327-34
11. Douillard J, Ostoros G, Cobo M, Ciuleanu T, McCormack R, Webster A, Milenkova T "First-line gefitinib in Caucasian EGFR mutation-positive NSCLC patients: a phase-IV, open-label, single-arm study." *British journal of cancer*(2014): 55-62
12. Passaro A, Wang J, Wang Y, Lee S, Melosky B, Shih J, Wang J, Azuma K, Juan-Vidal O, Cobo M, Felip E, Girard N, Cortot A, Califano R, Cappuzzo F, Owen S, Popat S, Tan J, Salinas J, Tomasini P, Gentzler R, William W, Reckamp K, Takahashi T, Ganguly S, Kowalski D, Bearz A, MacKean M, Barala P, Bourla A, Girvin A, Greger J, Millington D, Withelder M, Xie J, Sun T, Shah S, Diorio B, Knoblauch R, Baum J, Campelo R, Cho B "Amivantamab plus chemotherapy with and without lazertinib in EGFR-mutant advanced NSCLC after disease progression on osimertinib: primary results from the phase III MARIPOSA-2 study." *Annals of oncology : official journal of the European Society for Medical Oncology*(2024): 77-90
13. "A Phase III, Randomized, Double-blind, Placebo-controlled, Multicenter, International Study of Osimertinib as Maintenance Therapy in Patients With Locally Advanced, Unresectable EGFR Mutation-positive Non-Small Cell Lung Cancer (Stage III) Whose Disease Has Not Progressed Following Definitive Platinum-based Chemoradiation Therapy (LAURA)." (2024)
14. Lu S, Kato T, Dong X, Ahn M, Quang L, Soparattanapaisarn N, Inoue T, Wang C, Huang M, Yang J, Cobo M, Özgüroğlu M, Casarini I, Khiem D, Sriuranpong V, Cronemberger E, Takahashi T, Runglodvatana Y, Chen M, Huang X, Grainger E, Ghiorgiu D, van der Gronde T, Ramalingam S "Osimertinib after Chemoradiotherapy in Stage III EGFR-Mutated NSCLC." *The New England journal of medicine*(2024): 585-597
15. Yang J, Sequist L, Geater S, Tsai C, Mok T, Schuler M, Yamamoto N, Yu C, Ou S, Zhou C, Massey D, Zazulina V, Wu Y "Clinical activity of afatinib in patients with advanced non-small-cell lung cancer harbouring uncommon EGFR mutations: a combined post-hoc analysis of LUX-Lung 2, LUX-Lung 3, and LUX-Lung 6." *The Lancet. Oncology*(2015): 830-8
16. Nakagawa K, Garon E, Seto T, Nishio M, Ponce Aix S, Paz-Ares L, Chiu C, Park K, Novello S, Nadal E, Imamura F, Yoh K, Shih J, Au K, Moro-Sibilot D, Enatsu S, Zimmermann A, Frimodt-Moller B, Visseren-Grul C, Reck M "Ramucicirumab plus erlotinib in patients with untreated, EGFR-mutated, advanced non-small-cell lung cancer (RELAY): a randomised, double-blind, placebo-controlled, phase 3 trial." *The Lancet. Oncology*(2019): 1655-1669
17. Planchard D, Jänne P, Cheng Y, Yang J, Yanagitani N, Kim S, Sugawara S, Yu Y, Fan Y, Geater S, Laktionov K, Lee C, Valdiviezo N, Ahmed S, Maurel J, Andrasina I, Goldman J, Ghiorgiu D, Rukazekov Y, Todd A, Kobayashi K "Osimertinib with or without Chemotherapy in EGFR-Mutated Advanced NSCLC." *The New England journal of medicine*(2023): 1935-1948
18. "A Phase III, Open-label, Randomized Study of Osimertinib With or Without Platinum Plus Pemetrexed Chemo, as First-line Treatment in Patients With Epidermal Growth Factor Receptor (EGFR) Mutation Positive, Locally Advanced or Metastatic Non-small Cell Lung Cancer (FLAURA2)." (2024)
19. "A Phase 3, Open-Label, Randomized Study of Amivantamab and Lazertinib in Combination With Platinum-Based Chemotherapy Compared With Platinum-Based Chemotherapy in Patients With EGFR-Mutated Locally Advanced or Metastatic Non-Small Cell Lung Cancer After Osimertinib Failure" (2024)
20. Cho B, Lee K, Cho E, et al. "Amivantamab (JNJ-61186372), an EGFR-MET bispecific antibody, in combination with lazertinib, a 3rd-generation tyrosine kinase inhibitor (TKI), in advanced EGFR NSCLC" *Annals of Oncology*(2020): Abstract 1258O
21. Park K, John T, Kim S, et al. "Amivantamab (JNJ-61186372), an anti-EGFR-MET bispecific antibody, in patients with EGFR exon 20 insertion (exon20ins)-mutated non-small cell lung cancer (NSCLC)." *Journal of Clinical Oncology*(2020)



## References

22. Riely G, Neal J, Camidge D, Spira A, Piotrowska Z, Costa D, Tsao A, Patel J, Gadgeel S, Bazhenova L, Zhu V, West H, Mekhail T, Gentzler R, Nguyen D, Vincent S, Zhang S, Lin J, Bunn V, Jin S, Li S, Jänne P "Activity and Safety of Mobocertinib (TAK-788) in Previously Treated Non-Small Cell Lung Cancer with EGFR Exon 20 Insertion Mutations from a Phase I/II Trial." *Cancer discovery*(2021): 1688-1699
23. Zhou C, Ramalingam S, Li B, et al. "Mobocertinib in NSCLC With EGFR Exon 20 Insertions: Results From EXCLAIM and Pooled Platinum-Pretreated Patient Populations" *Journal of Thoracic Oncology*(2021)
24. Yang J, Wu Y, Schuler M, Sebastian M, Popat S, Yamamoto N, Zhou C, Hu C, O'Byrne K, Feng J, Lu S, Huang Y, Geater S, Lee K, Tsai C, Gorbunova V, Hirsh V, Bennouna J, Orlov S, Mok T, Boyer M, Su W, Lee K, Kato T, Massey D, Shahidi M, Zazulina V, Sequist L "Afatinib versus cisplatin-based chemotherapy for EGFR mutation-positive lung adenocarcinoma (LUX-Lung 3 and LUX-Lung 6): analysis of overall survival data from two randomised, phase 3 trials." *The Lancet. Oncology* (2015): 141-51
25. O'Kane G, Bradbury P, Feld R, Leigh N, Liu G, Pisters K, Kamel-Reid S, Tsao M, Shepherd F "Uncommon EGFR mutations in advanced non-small cell lung cancer." *Lung cancer* (Amsterdam, Netherlands)(2017): 137-144
26. Engelman J, Zejnullahu K, Mitsudomi T, Song Y, Hyland C, Park J, Lindeman N, Gale C, Zhao X, Christensen J, Kosaka T, Holmes A, Rogers A, Cappuzzo F, Mok T, Lee C, Johnson B, Cantley L, Jänne P "MET amplification leads to gefitinib resistance in lung cancer by activating ERBB3 signaling." *Science* (New York, N.Y.)(2007): 1039-43
27. Greulich H, Chen T, Feng W, Jänne P, Alvarez J, Zappaterra M, Bulmer S, Frank D, Hahn W, Sellers W, Meyerson M "Oncogenic transformation by inhibitor-sensitive and -resistant EGFR mutants." *PLoS medicine*(2005): e313
28. Kwak E, Sordella R, Bell D, Godin-Heymann N, Okimoto R, Brannigan B, Harris P, Driscoll D, Fidias P, Lynch T, Rabindran S, McGinnis J, Wissner A, Sharma S, Isselbacher K, Settleman J, Haber D "Irreversible inhibitors of the EGF receptor may circumvent acquired resistance to gefitinib." *Proceedings of the National Academy of Sciences of the United States of America*(2005): 7665-70
29. Takezawa K, Pirazzoli V, Arcila M, Nebhan C, Song X, de Stanchina E, Ohashi K, Janjigian Y, Spitzler P, Melnick M, Riely G, Kris M, Miller V, Ladanyi M, Politi K, Pao W "HER2 amplification: a potential mechanism of acquired resistance to EGFR inhibition in EGFR-mutant lung cancers that lack the second-site EGFR T790M mutation." *Cancer discovery*(2012): 922-33
30. Yu H, Arcila M, Rekhtman N, Sima C, Zakowski M, Pao W, Kris M, Miller V, Ladanyi M, Riely G "Analysis of tumor specimens at the time of acquired resistance to EGFR-TKI therapy in 155 patients with EGFR-mutant lung cancers." *Clinical cancer research : an official journal of the American Association for Cancer Research*(2013): 2240-7
31. Jänne P, Boss D, Camidge D, Britten C, Engelman J, Garon E, Guo F, Wong S, Liang J, Letrent S, Millham R, Taylor I, Eckhardt S, Schellens J "Phase I dose-escalation study of the pan-HER inhibitor, PF299804, in patients with advanced malignant solid tumors." *Clinical cancer research : an official journal of the American Association for Cancer Research*(2011): 1131-9
32. Jänne P, Yang J, Kim D, Planchard D, Ohe Y, Ramalingam S, Ahn M, Kim S, Su W, Horn L, Haggstrom D, Felip E, Kim J, Frewer P, Cantarini M, Brown K, Dickinson P, Ghiorghiu S, Ranson M "AZD9291 in EGFR inhibitor-resistant non-small-cell lung cancer." *The New England journal of medicine*(2015): 1689-99
33. Greig S "Osimertinib: First Global Approval." *Drugs*(2016): 263-73
34. Yang J, Ahn M, Kim D, Ramalingam S, Sequist L, Su W, Kim S, Kim J, Planchard D, Felip E, Blackhall F, Haggstrom D, Yoh K, Novello S, Gold K, Hirashima T, Lin C, Mann H, Cantarini M, Ghiorghiu S, Jänne P "Osimertinib in Pretreated T790M-Positive Advanced Non-Small-Cell Lung Cancer: AURA Study Phase II Extension Component." *Journal of clinical oncology : official journal of the American Society of Clinical Oncology*(2017): 1288-1296
35. Wang S, Cang S, Liu D "Third-generation inhibitors targeting EGFR T790M mutation in advanced non-small cell lung cancer." *Journal of hematology & oncology*(2016): 34
36. Watanabe S, Sone T, Matsui T, Yamamura K, Tani M, Okazaki A, Kurokawa K, Tambo Y, Takato H, Ohkura N, Waseda Y, Katayama N, Kasahara K "Transformation to small-cell lung cancer following treatment with EGFR tyrosine kinase inhibitors in a patient with lung adenocarcinoma." *Lung cancer* (Amsterdam, Netherlands)(2013): 370-2
37. Chang Y, Kim S, Choi Y, So K, Rho J, Kim W, Lee J, Chung J, Choi C "Neuroendocrine differentiation in acquired resistance to epidermal growth factor receptor tyrosine kinase inhibitor." *Tuberculosis and respiratory diseases*(2013): 95-103
38. Popat S, Wotherspoon A, Nutting C, Gonzalez D, Nicholson A, O'Brien M "Transformation to "high grade" neuroendocrine carcinoma as an acquired drug resistance mechanism in EGFR-mutant lung adenocarcinoma." *Lung cancer* (Amsterdam, Netherlands)(2013): 1-4
39. Sequist L, Waltman B, Dias-Santagata D, Digumarthy S, Turke A, Fidias P, Bergethon K, Shaw A, Gettinger S, Cosper A, Akhavanfar S, Heist R, Temel J, Christensen J, Wain J, Lynch T, Vernovsky K, Mark E, Lanuti M, Iafrate A, Mino-Kenudson M, Engelman J "Genotypic and histological evolution of lung cancers acquiring resistance to EGFR inhibitors." *Science translational medicine*(2011): 75ra26
40. Staal F, van der Luijt R, Baert M, van Drunen J, van Bakel H, Peters E, de Valk I, van Amstel H, Taphoorn M, Jansen G, van Veelen C, Burgering B, Staal G "A novel germline mutation of PTEN associated with brain tumours of multiple lineages." *British journal of cancer*(2002): 1586-91
41. Couto S, Cao M, Duarte P, Banach-Petrosky W, Wang S, Romanienko P, Wu H, Cardiff R, Abate-Shen C, Cunha G "Simultaneous haploinsufficiency of Pten and Trp53 tumor suppressor genes accelerates tumorigenesis in a mouse model of prostate cancer." *Differentiation; research in biological diversity*(2009): 103-11
42. Kwon C, Zhao D, Chen J, Alcantara S, Li Y, Burns D, Mason R, Lee E, Wu H, Parada L "Pten haploinsufficiency accelerates formation of high-grade astrocytomas." *Cancer research*(2008): 3286-94
43. Kwabi-Addo B, Giri D, Schmidt K, Podsypanina K, Parsons R, Greenberg N, Iltmann M "Haploinsufficiency of the Pten tumor suppressor gene promotes prostate cancer progression." *Proceedings of the National Academy of Sciences of the United States of America*(2001): 11563-8
44. Velickovic M, Delahunt B, McIver B, Grebe S "Intragenic PTEN/MMAC1 loss of heterozygosity in conventional (clear-cell) renal cell carcinoma is associated with poor patient prognosis." *Modern pathology : an official journal of the United States and Canadian Academy of Pathology, Inc*(2002): 479-85
45. Hollander M, Blumenthal G, Dennis P "PTEN loss in the continuum of common cancers, rare syndromes and mouse models." *Nature reviews. Cancer*(2011): 289-301
46. Rulle U, Tsourtzi Z, Casanova R, Deml K, Verbeken E, Thunnissen E, Warth A, Cheney R, Seida A, Speel E, Madsen L, Nonaka D, Navarro A, Sansano I, Marchetti A, Finn S, Monkhurst K, Kerr K, Haberecker M, Wu C, Zygoura P, Kammiller R, Geiger T, Gendreau S, Schulze K, Vrugt B, Wild P, Moch H, Weder W, Ciftlik A, Dafni U, Peters S, Bubendorf L, Stahel R, Soltermann A "Computer-Based Intensity Measurement Assists Pathologists in Scoring Phosphatase and Tensin Homolog Immunohistochemistry - Clinical Associations in NSCLC Patients of the European Thoracic Oncology Platform Lungscape Cohort." *Journal of thoracic oncology : official publication of the International Association for the Study of Lung Cancer*(2018): 1851-1863
47. Li X, Yang Y, Zhang H, Yue W, Zhang T, Lu B, Li J, Liu Z, Wang Q, Gao Y, Hu A, Zhang H, Shi H, Hu F, Li B "High levels of Phosphatase and Tensin Homolog Expression Predict Favorable Prognosis in Patients with Non-small Cell Lung Cancer." *Cell biochemistry and biophysics*(2015): 631-7

## References

48. Shin E, Choi C, Kim H, Jang S, Park Y "Immunohistochemical characterization of the mTOR pathway in stage-I non-small-cell lung carcinoma." *Lung cancer* (Amsterdam, Netherlands)(2015): 13-8
49. Trigka E, Levidou G, Saetta A, Chatziandreou I, Tomos P, Thalassinou N, Anastasiou N, Spartalis E, Kavantzis N, Patsouris E, Korkolopoulou P "A detailed immunohistochemical analysis of the PI3K/AKT/mTOR pathway in lung cancer: correlation with PIK3CA, AKT1, K-RAS or PTEN mutational status and clinicopathological features." *Oncology reports*(2013): 623-36
50. Ji Y, Zheng M, Ye S, Chen J, Chen Y "PTEN and Ki67 expression is associated with clinicopathologic features of non-small cell lung cancer." *Journal of biomedical research*(2014): 462-7
51. Yun F, Jia Y, Li X, Yuan L, Sun Q, Yu H, Shi L, Yuan H "Clinicopathological significance of PTEN and PI3K/AKT signal transduction pathway in non-small cell lung cancer." *International journal of clinical and experimental pathology*(2013): 2112-20
52. Malkoski S, Cleaver T, Thompson J, Sutton W, Haeger S, Rodriguez K, Lu S, Merrick D, Wang X "Role of PTEN in basal cell derived lung carcinogenesis." *Molecular carcinogenesis*(2014): 841-6
53. Xu C, Fillmore C, Koyama S, Wu H, Zhao Y, Chen Z, Herter-Sprie G, Akbay E, Tchaicha J, Altabef A, Reibel J, Walton Z, Ji H, Watanabe H, Jänne P, Castrillon D, Rustgi A, Bass A, Freeman G, Padera R, Dranoff G, Hammerman P, Kim C, Wong K "Loss of Lkb1 and Pten leads to lung squamous cell carcinoma with elevated PD-L1 expression." *Cancer cell*(2014): 590-604
54. Wu R, Hu T, Rehemtulla A, Fearon E, Cho K "Preclinical testing of PI3K/AKT/mTOR signaling inhibitors in a mouse model of ovarian endometrioid adenocarcinoma." *Clinical cancer research : an official journal of the American Association for Cancer Research*(2011): 7359-72
55. Dreyling M, Santoro A, Mollica L, et al. "COPANLISIB IN PATIENTS WITH RELAPSED OR REFRACTORY INDOLENT B-CELL LYMPHOMA (CHRONOS-1)" *J Clin Oncol* (2017): Abstract 108
56. André F, Ciruelos E, Rubovszky G, Campone M, Loibl S, Rugo H, Iwata H, Conte P, Mayer I, Kaufman B, Yamashita T, Lu Y, Inoue K, Takahashi M, Pápai Z, Longin A, Mills D, Wilke C, Hirawat S, Juric D "Alpelisib for PIK3CA-Mutated, Hormone Receptor-Positive Advanced Breast Cancer." *The New England journal of medicine*(2019): 1929-1940
57. Massaccesi C, di Tomaso E, Fretault N, Hirawat S "Challenges in the clinical development of PI3K inhibitors." *Annals of the New York Academy of Sciences*(2013): 19-23
58. Janku F, Tsimberidou A, Garrido-Laguna I, Wang X, Luthra R, Hong D, Naing A, Falchook G, Moroney J, Piha-Paul S, Wheeler J, Moulder S, Fu S, Kurzrock R "PIK3CA mutations in patients with advanced cancers treated with PI3K/AKT/mTOR axis inhibitors." *Molecular cancer therapeutics*(2011): 558-65
59. Grunt T, Mariani G "Novel approaches for molecular targeted therapy of breast cancer: interfering with PI3K/AKT/mTOR signaling." *Current cancer drug targets*(2013): 188-204
60. Turner N, Oliveira M, Howell S, et al. "Capivasertib and fulvestrant for patients with aromatase inhibitor-resistant hormone receptor-positive/human epidermal growth factor receptor 2-negative advanced breast cancer: results from the Phase III CAPitello-291 trial" *Cancer Research*(2023)
61. "Study of Induction SBRT and Olaparib Followed by Combination Pembrolizumab/Olaparib in Gastric and Gastroesophageal Junction (GEJ) Cancers" (2023)
62. "A Phase II Study of Olaparib in Patients With Advanced Biliary Tract Cancer With Aberrant DNA Repair Gene Mutations" (2023)
63. "Combination of HX008 And Niraparib in GErM-line-mutAted Metastatic Breast Cancer: a muLti-centEr Phase II Study" (2022)
64. "A Phase II Trial of the PARP Inhibitor, Niraparib, in BAP1 and Other DNA Damage Response (DDR) Pathway Deficient Neoplasms (UF-STO-ETI-001)" (2023)
65. "A Phase II Clinical Trial of the PARP Inhibitor Talazoparib in BRCA1 and BRCA2 Wild Type Patients With Advanced Triple Negative Breast Cancer and Homologous Recombination Deficiency or Advanced HER2 Negative Breast Cancer or Other Solid Tumors With a Mutation in Homologous Recombination Pathway Genes" (2023)
66. Mendes-Pereira A, Martin S, Brough R, McCarthy A, Taylor J, Kim J, Waldman T, Lord C, Ashworth A "Synthetic lethal targeting of PTEN mutant cells with PARP inhibitors." *EMBO molecular medicine*(2009): 315-22
67. Dedes K, Wetterskog D, Mendes-Pereira A, Natrajan R, Lambros M, Geyer F, Vatcheva R, Savage K, Mackay A, Lord C, Ashworth A, Reis-Filho J "PTEN deficiency in endometrioid endometrial adenocarcinomas predicts sensitivity to PARP inhibitors." *Science translational medicine*(2010): 53ra75
68. McEllin B, Camacho C, Mukherjee B, Hahm B, Tomimatsu N, Bachoo R, Burma S "PTEN loss compromises homologous recombination repair in astrocytes: implications for glioblastoma therapy with temozolomide or poly(ADP-ribose) polymerase inhibitors." *Cancer research*(2010): 5457-64
69. Sos M, Koker M, Weir B, Heynck S, Rabinovsky R, Zander T, Seeger J, Weiss J, Fischer F, Frommolt P, Michel K, Peifer M, Mermel C, Girard L, Peyton M, Gazdar A, Minna J, Garraway L, Kashkar H, Pao W, Meyerson M, Thomas R "PTEN loss contributes to erlotinib resistance in EGFR-mutant lung cancer by activation of Akt and EGFR." *Cancer research*(2009): 3256-61
70. Takeda H, Takigawa N, Ohashi K, Minami D, Kataoka I, Ichihara E, Ochi N, Tanimoto M, Kiura K "Vandetanib is effective in EGFR-mutant lung cancer cells with PTEN deficiency." *Experimental cell research*(2013): 417-23
71. Maeda M, Murakami Y, Watari K, Kuwano M, Izumi H, Ono M "CpG hypermethylation contributes to decreased expression of PTEN during acquired resistance to gefitinib in human lung cancer cell lines." *Lung cancer* (Amsterdam, Netherlands)(2015): 265-71
72. Kim H, Cho B, Shim H, Lim S, Kim S, Chang J, Kim D, Kim J "Prediction for response duration to epidermal growth factor receptor-tyrosine kinase inhibitors in EGFR mutated never smoker lung adenocarcinoma." *Lung cancer* (Amsterdam, Netherlands)(2014): 374-82
73. Thakur R, Mishra D "Pharmacological modulation of beta-catenin and its applications in cancer therapy." *Journal of cellular and molecular medicine*(2013): 449-56
74. Barr M, Gray S, Hoffmann A, Hilger R, Thomale J, O'Flaherty J, Fennell D, Richard D, O'Leary J, O'Byrne K "Generation and characterisation of cisplatin-resistant non-small cell lung cancer cell lines displaying a stem-like signature." *PLoS one*(2013): e54193
75. Miao Y, Wang L, Zhang X, Xu X, Jiang G, Fan C, Liu Y, Lin X, Yu J, Zhang Y, Wang E "Promoter methylation-mediated silencing of  $\beta$ -catenin enhances invasiveness of non-small cell lung cancer and predicts adverse prognosis." *PLoS one*(2014): e112258
76. Polakis P "The oncogenic activation of beta-catenin." *Current opinion in genetics & development*(1999): 15-21
77. Abraham S, Klimstra D, Wilentz R, Yeo C, Conlon K, Brennan M, Cameron J, Wu T, Hruban R "Solid-pseudopapillary tumors of the pancreas are genetically distinct from pancreatic ductal adenocarcinomas and almost always harbor beta-catenin mutations." *The American journal of pathology*(2002): 1361-9

## References

78. Tanaka Y, Kato K, Notohara K, Hojo H, Ijiri R, Miyake T, Nagahara N, Sasaki F, Kitagawa N, Nakatani Y, Kobayashi Y "Frequent beta-catenin mutation and cytoplasmic/nuclear accumulation in pancreatic solid-pseudopapillary neoplasm." *Cancer research*(2001): 8401-4
79. Takahashi Y, Hiraoka N, Onozato K, Shibata T, Kosuge T, Nimura Y, Kanai Y, Hirohashi S "Solid-pseudopapillary neoplasms of the pancreas in men and women: do they differ?" *Virchows Archiv : an international journal of pathology*(2006): 561-9
80. Notani D, Gottimukkala K, Jayani R, Limaye A, Damle M, Mehta S, Purbey P, Joseph J, Galande S "Global regulator SATB1 recruits beta-catenin and regulates T(H)2 differentiation in Wnt-dependent manner." *PLoS biology*(2010): e1000296
81. Lévy L, Wei Y, Labalette C, Wu Y, Renard C, Buendia M, Neuveut C "Acetylation of beta-catenin by p300 regulates beta-catenin-Tcf4 interaction." *Molecular and cellular biology*(2004): 3404-14
82. Braggio D, Costas C de Faria F, Koller D, Jin F, Zewdu A, Lopez G, Batte K, Casadei L, Welliver M, Horrigan S, Han R, Larson J, Strohecker A, Pollock R "Preclinical efficacy of the Wnt/ $\beta$ -catenin pathway inhibitor BC2059 for the treatment of desmoid tumors." *PloS one*(2022): e0276047
83. Savvidou I, Khong T, Whish S, Carmichael I, Sepehrizadeh T, Mithraprabhu S, Horrigan S, de Veer M, Spencer A "Combination of Histone Deacetylase Inhibitor Panobinostat (LBH589) with  $\beta$ -Catenin Inhibitor Tegavivint (BC2059) Exerts Significant Anti-Myeloma Activity Both In Vitro and In Vivo." *Cancers*(2022)
84. Nomura M, Rainusso N, Lee Y, Dawson B, Coarfa C, Han R, Larson J, Shuck R, Kurenbekova L, Yustein J "Tegavivint and the  $\beta$ -Catenin/ALDH Axis in Chemotherapy-Resistant and Metastatic Osteosarcoma." *Journal of the National Cancer Institute*(2019): 1216-1227
85. "A Phase 1/2 Study of FOG-001 in Participants With Locally Advanced or Metastatic Solid Tumors" (2023)
86. "A Phase 1/2 Study of Tegavivint (NSC#826393) in Children, Adolescents, and Young Adults With Recurrent or Refractory Solid Tumors, Including Lymphomas and Desmoid Tumors" (2023)
87. "A Phase 1/2 Exploratory Study of the TBL1 Inhibitor, Tegavivint (BC2059), in Patients With Advanced Hepatocellular Carcinoma" (2024)
88. Ruiz de Galarreta M, Bresnahan E, Molina-Sánchez P, Lindblad K, Maier B, Sia D, Puigvehi M, Miguela V, Casanova-Acebes M, Dhainaut M, Villacorta-Martin C, Singhi A, Moghe A, von Felden J, Tal Grinspan L, Wang S, Kamphorst A, Monga S, Brown B, Villanueva A, Llovet J, Merad M, Lujambio A " $\beta$ -Catenin Activation Promotes Immune Escape and Resistance to Anti-PD-1 Therapy in Hepatocellular Carcinoma." *Cancer discovery*(2019): 1124-1141
89. Harding J, Nandakumar S, Armenia J, Khalil D, Albano M, Ly M, Shia J, Hechtman J, Kundra R, El Dika I, Do R, Sun Y, Kingham T, D'Angelica M, Berger M, Hyman D, Jarnagin W, Klimstra D, Janjigian Y, Solit D, Schultz N, Abou-Alfa G "Prospective Genotyping of Hepatocellular Carcinoma: Clinical Implications of Next-Generation Sequencing for Matching Patients to Targeted and Immune Therapies." *Clinical cancer research : an official journal of the American Association for Cancer Research* (2019): 2116-2126
90. Matsuda A, Ishiguro K, Yan I, Patel T "Extracellular Vesicle-Based Therapeutic Targeting of  $\beta$ -Catenin to Modulate Anticancer Immune Responses in Hepatocellular Cancer." *Hepatology communications*(2019): 525-541
91. Ganesh S, Shui X, Craig K, Park J, Wang W, Brown B, Abrams M "RNAi-Mediated  $\beta$ -Catenin Inhibition Promotes T Cell Infiltration and Antitumor Activity in Combination with Immune Checkpoint Blockade." *Molecular therapy : the journal of the American Society of Gene Therapy*(2018): 2567-2579
92. Spranger S, Bao R, Gajewski T "Melanoma-intrinsic  $\beta$ -catenin signalling prevents anti-tumour immunity." *Nature*(2015): 231-5
93. Thorsson V, Gibbs D, Brown S, Wolf D, Bortone D, Ou Yang T, Porta-Pardo E, Gao G, Plaisier C, Eddy J, Ziv E, Culhane A, Paul E, Sivakumar I, Gentles A, Malhotra R, Farshidfar F, Colaprico A, Parker J, Mose L, Vo N, Liu J, Liu Y, Rader J, Dhankani V, Reynolds S, Bowlby R, Califano A, Cherniack A, Anastassiou D, Bedognetti D, Mokrab Y, Newman A, Rao A, Chen K, Krasnitz A, Hu H, Malta T, Noushmehr H, Pedamallu C, Bullman S, Ojesina A, Lamb A, Zhou W, Shen H, Choueiri T, Weinstein J, Guinney J, Saltz J, Holt R, Rabkin C, Lazar A, Serody J, Demicco E, Disis M, Vincent B, Shmulevich I "The Immune Landscape of Cancer." *Immunity*(2018): 812-830.e14
94. Tenbaum S, Ordóñez-Morán P, Puig I, Chicote I, Arqués O, Landolfi S, Fernández Y, Herance J, Gispert J, Mendizabal L, Aguilar S, Ramón y Cajal S, Schwartz S, Vivancos A, Espín E, Rojas S, Baselga J, Tabernero J, Muñoz A, Palmer H " $\beta$ -catenin confers resistance to PI3K and AKT inhibitors and subverts FOXO3a to promote metastasis in colon cancer." *Nature medicine*(2012): 892-901
95. Zhang H, Zhang Z, Che C, Mei Y, Shi Y "Array analysis for potential biomarker of gemcitabine identification in non-small cell lung cancer cell lines." *International journal of clinical and experimental pathology*(2013): 1734-46
96. Fang X, Gu P, Zhou C, Liang A, Ren S, Liu F, Zeng Y, Wu Y, Zhao Y, Huang B, Zhang Z, Yi X " $\beta$ -Catenin overexpression is associated with gefitinib resistance in non-small cell lung cancer cells." *Pulmonary pharmacology & therapeutics*(2014): 41-48
97. Farber L, Efrati E, Elkin H, Peerless Y, Sabo E, Ben-Izhak O, Herschkovitz D "Molecular morphometric analysis shows relative intra-tumoural homogeneity for KRAS mutations in colorectal cancer." *Virchows Archiv : an international journal of pathology*(2011): 487-93
98. Feldmann G, Beaty R, Hruban R, Maitra A "Molecular genetics of pancreatic intraepithelial neoplasia." *Journal of hepato-biliary-pancreatic surgery*(2007): 224-32
99. Han C, Ma J, Zhao J, Zhou Y, Jing W, Zou H "EGFR mutations, gene amplification, and protein expression and KRAS mutations in primary and metastatic tumors of non-small cell lung cancers and their clinical implications: a meta-analysis." *Cancer investigation*(2011): 626-34
100. Colombino M, Paliogiannis P, Cossu A, Santeufemia D, Sini M, Casula M, Palomba G, Manca A, Pisano M, Doneddu V, Palmieri G "EGFR, KRAS, BRAF, ALK, and cMET genetic alterations in 1440 Sardinian patients with lung adenocarcinoma." *BMC pulmonary medicine*(2019): 209
101. Li D, Ding L, Ran W, Huang Y, Li G, Wang C, Xiao Y, Wang X, Lin D, Xing X "Status of 10 targeted genes of non-small cell lung cancer in eastern China: A study of 884 patients based on NGS in a single institution." *Thoracic cancer*(2020): 2580-2589
102. Liu Y, Li H, Zhu J, Zhang Y, Liu X, Li R, Zhang Q, Cheng Y "The Prevalence and Concurrent Pathogenic Mutations of KRAS G12C in Northeast Chinese Non-small-cell Lung Cancer Patients." *Cancer management and research*(2021): 2447-2454
103. Liu S, Sun H, Zhou J, Jie G, Xie Z, Shao Y, Zhang X, Ye J, Chen C, Zhang X, Zhou Q, Yang J, Wu Y "Clinical characteristics and prognostic value of the KRAS G12C mutation in Chinese non-small cell lung cancer patients." *Biomarker research*(2020): 22
104. Cui W, Franchini F, Alexander M, Officer A, Wong H, IJzerman M, Desai J, Solomon B "Real world outcomes in KRAS G12C mutation positive non-small cell lung cancer." *Lung cancer (Amsterdam, Netherlands)*(2020): 310-317
105. Judd J, Abdel Karim N, Khan H, Naqash A, Baca Y, Xiu J, VanderWalde A, Mamdani H, Raez L, Nagasaka M, Pai S, Socinski M, Nieva J, Kim C, Wozniak A, Ikpeazu C, de Lima Lopes G, Spira A, Korn W, Kim E, Liu S, Borghaei H "Characterization of KRAS Mutation Subtypes in Non-small Cell Lung Cancer." *Molecular cancer therapeutics* (2021): 2577-2584

## References

106. Gainor J, Varghese A, Ou S, Kabraji S, Awad M, Katayama R, Pawlak A, Mino-Kenudson M, Yeap B, Riely G, Iafrate A, Arcila M, Ladanyi M, Engelman J, Dias-Santagata D, Shaw A "ALK rearrangements are mutually exclusive with mutations in EGFR or KRAS: an analysis of 1,683 patients with non-small cell lung cancer." *Clinical cancer research : an official journal of the American Association for Cancer Research*(2013): 4273-81
107. Lee B, Lee T, Lee S, Choi Y, Han J "Clinicopathologic characteristics of EGFR, KRAS, and ALK alterations in 6,595 lung cancers." *Oncotarget*(2016): 23874-84
108. Chatziandreu I, Tsioli P, Sakellariou S, Mourikioti I, Giannopoulou I, Levidou G, Korkolopoulou P, Patsouris E, Saetta A "Comprehensive Molecular Analysis of NSCLC; Clinicopathological Associations." *PloS one*(2015): e0133859
109. Paik P, Johnson M, D'Angelo S, Sima C, Ang D, Dogan S, Miller V, Ladanyi M, Kris M, Riely G "Driver mutations determine survival in smokers and never-smokers with stage IIIB/IV lung adenocarcinomas." *Cancer*(2012): 5840-7
110. Sholl L, Aisner D, Varela-Garcia M, Berry L, Dias-Santagata D, Wistuba I, Chen H, Fujimoto J, Kugler K, Franklin W, Iafrate A, Ladanyi M, Kris M, Johnson B, Bunn P, Minna J, Kwiatkowski D "Multi-institutional Oncogenic Driver Mutation Analysis in Lung Adenocarcinoma: The Lung Cancer Mutation Consortium Experience." *Journal of thoracic oncology : official publication of the International Association for the Study of Lung Cancer*(2015): 768-777
111. Cho J, Choi S, Lee J, Lee C, Lee S, Kim D, Yim J, Kim Y, Yoo C, Kim Y, Han S, Park Y "Proportion and clinical features of never-smokers with non-small cell lung cancer." *Chinese journal of cancer*(2017): 20
112. Zhao M, Zhan C, Li M, Yang X, Yang X, Zhang Y, Lin M, Xia Y, Feng M, Wang Q "Aberrant status and clinicopathologic characteristic associations of 11 target genes in 1,321 Chinese patients with lung adenocarcinoma." *Journal of thoracic disease*(2018): 398-407
113. Jin Y, Xue Q, Shen X, Zheng Q, Chen H, Zhou X, Li Y "PD-L1 Expression and Comprehensive Molecular Profiling Predict Survival in Nonsmall Cell Lung Cancer: A Real-World Study of a Large Chinese Cohort." *Clinical lung cancer*(2022): 43-51
114. Shirasawa M, Yoshida T, Shimoda Y, Takayanagi D, Shiraishi K, Kubo T, Mitani S, Matsumoto Y, Masuda K, Shinno Y, Okuma Y, Goto Y, Horinouchi H, Ichikawa H, Kohno T, Yamamoto N, Matsumoto S, Goto K, Watanabe S, Ohe Y, Motoi N "Differential Immune-Related Microenvironment Determines Programmed Cell Death Protein-1/Programmed Death-Ligand 1 Blockade Efficacy in Patients With Advanced NSCLC." *Journal of thoracic oncology : official publication of the International Association for the Study of Lung Cancer*(2021): 2078-2090
115. Lauko A, Kotecha R, Barnett A, Li H, Tatineni V, Ali A, Patil P, Mohammadi A, Chao S, Murphy E, Angelov L, Suh J, Barnett G, Pennell N, Ahluwalia M "Impact of KRAS mutation status on the efficacy of immunotherapy in lung cancer brain metastases." *Scientific reports*(2021): 18174
116. Kerr K, Thunnissen E, Dafni U, Finn S, Bubendorf L, Soltermann A, Verbeken E, Biernat W, Warth A, Marchetti A, Speel E, Pokharel S, Quinn A, Monkhurst K, Navarro A, Madsen L, Radonic T, Wilson J, De Luca G, Gray S, Cheney R, Savic S, Martorell M, Muley T, Baas P, Meldgaard P, Blackhall F, Dingemans A, Dziadziuszko R, Vansteenkiste J, Weder W, Polydoropoulou V, Geiger T, Kammler R, Peters S, Stahel R "A retrospective cohort study of PD-L1 prevalence, molecular associations and clinical outcomes in patients with NSCLC: Results from the European Thoracic Oncology Platform (ETOP) Lungscape Project." *Lung cancer (Amsterdam, Netherlands)* (2019): 95-103
117. Kim H, Kwon H, Park S, Park Y, Park E, Chung J "Clinicopathological analysis and prognostic significance of programmed cell death-ligand 1 protein and mRNA expression in non-small cell lung cancer." *PloS one*(2018): e0198634
118. Cui S, Dong L, Qian J, Ye L, Jiang L "Classifying Non-Small Cell Lung Cancer by Status of Programmed Cell Death Ligand 1 and Tumor-Infiltrating Lymphocytes on Tumor Cells." *Journal of Cancer*(2018): 129-134
119. Petrelli F, Maltese M, Tomasello G, Conti B, Borgonovo K, Cabiddu M, Ghilardi M, Ghidini M, Passalacqua R, Barni S, Brighenti M "Clinical and Molecular Predictors of PD-L1 Expression in Non-Small-Cell Lung Cancer: Systematic Review and Meta-analysis." *Clinical lung cancer*(2018): 315-322
120. Lan B, Ma C, Zhang C, Chai S, Wang P, Ding L, Wang K "Association between PD-L1 expression and driver gene status in non-small-cell lung cancer: a meta-analysis." *Oncotarget*(2018): 7684-7699
121. Albitar M, Sudarsanam S, Ma W, Jiang S, Chen W, Funari V, Blocker F, Agersborg S "Correlation of MET gene amplification and TP53 mutation with PD-L1 expression in non-small cell lung cancer." *Oncotarget*(2018): 13682-13693
122. Yang H, Chen H, Luo S, Li L, Zhou S, Shen R, Lin H, Xie X "The correlation between programmed death-ligand 1 expression and driver gene mutations in NSCLC." *Oncotarget*(2017): 23517-23528
123. Zhang M, Li G, Wang Y, Wang Y, Zhao S, Haihong P, Zhao H, Wang Y "PD-L1 expression in lung cancer and its correlation with driver mutations: a meta-analysis." *Scientific reports*(2017): 10255
124. Li D, Zhu X, Wang H, Li N "Association between PD-L1 expression and driven gene status in NSCLC: A meta-analysis." *European journal of surgical oncology : the journal of the European Society of Surgical Oncology and the British Association of Surgical Oncology*(2017): 1372-1379
125. Ameratunga M, Asadi K, Lin X, Walkiewicz M, Murone C, Knight S, Mitchell P, Boutros P, John T "PD-L1 and Tumor Infiltrating Lymphocytes as Prognostic Markers in Resected NSCLC." *PloS one*(2016): e0153954
126. Scheel A, Ansén S, Schultheis A, Scheffler M, Fischer R, Michels S, Hellmich M, George J, Zander T, Brockmann M, Stoelben E, Groen H, Timens W, Perner S, von Bergwelt-Baildon M, Büttner R, Wolf J "PD-L1 expression in non-small cell lung cancer: Correlations with genetic alterations." *Oncoimmunology*(2016): e1131379
127. Mansour M, Malmros K, Mager U, Ericson Lindquist K, Hejny K, Holmgren B, Seidal T, Dejmeck A, Dobra K, Planck M, Brunnström H "PD-L1 Expression in Non-Small Cell Lung Cancer Specimens: Association with Clinicopathological Factors and Molecular Alterations." *International journal of molecular sciences*(2022)
128. Yeh J, Routh E, Rubinas T, Peacock J, Martin T, Shen X, Sandler R, Kim H, Keku T, Der C "KRAS/BRAF mutation status and ERK1/2 activation as biomarkers for MEK1/2 inhibitor therapy in colorectal cancer." *Molecular cancer therapeutics*(2009): 834-43
129. Britten C "PI3K and MEK inhibitor combinations: examining the evidence in selected tumor types." *Cancer chemotherapy and pharmacology*(2013): 1395-409
130. Jänne P, van den Heuvel M, Barlesi F, Cobo M, Mazieres J, Crinò L, Orlov S, Blackhall F, Wolf J, Garrido P, Poltoratskiy A, Mariani G, Giorghiu D, Kilgour E, Smith P, Kohlmann A, Carille D, Lawrence D, Bowen K, Vansteenkiste J "Selumetinib Plus Docetaxel Compared With Docetaxel Alone and Progression-Free Survival in Patients With KRAS-Mutant Advanced Non-Small Cell Lung Cancer: The SELECT-1 Randomized Clinical Trial." *JAMA*(2017): 1844-1853
131. Puyol M, Martín A, Dubus P, Mulero F, Pizcueta P, Khan G, Guerra C, Santamaría D, Barbacid M "A synthetic lethal interaction between K-Ras oncogenes and Cdk4 unveils a therapeutic strategy for non-small cell lung carcinoma." *Cancer cell*(2010): 63-73



## References

132. Corcoran R, Cheng K, Hata A, Faber A, Ebi H, Coffee E, Greninger P, Brown R, Godfrey J, Cohoon T, Song Y, Lifshits E, Hung K, Shioda T, Dias-Santagata D, Singh A, Settleman J, Benes C, Mino-Kenudson M, Wong K, Engelman J "Synthetic lethal interaction of combined BCL-XL and MEK inhibition promotes tumor regressions in KRAS mutant cancer models." *Cancer cell*(2013): 121-8
133. Adjei A, Cohen R, Franklin W, Morris C, Wilson D, Molina J, Hanson L, Gore L, Chow L, Leong S, Maloney L, Gordon G, Simmons H, Marlow A, Litwiler K, Brown S, Poch G, Kane K, Haney J, Eckhardt S "Phase I pharmacokinetic and pharmacodynamic study of the oral, small-molecule mitogen-activated protein kinase kinase 1/2 inhibitor AZD6244 (ARRY-142886) in patients with advanced cancers." *Journal of clinical oncology : official journal of the American Society of Clinical Oncology*(2008): 2139-46
134. Manchado E, Weissmueller S, Morris J, Chen C, Wullenkord R, Lujambio A, de Stanchina E, Poirier J, Gainor J, Corcoran R, Engelman J, Rudin C, Rosen N, Lowe S "A combinatorial strategy for treating KRAS-mutant lung cancer." *Nature*(2016): 647-51
135. Infante J, Somer B, Park J, Li C, Scheulen M, Kasubhai S, Oh D, Liu Y, Redhu S, Steplewski K, Le N "A randomised, double-blind, placebo-controlled trial of trametinib, an oral MEK inhibitor, in combination with gemcitabine for patients with untreated metastatic adenocarcinoma of the pancreas." *European journal of cancer (Oxford, England : 1990)*(2014): 2072-81
136. Lito P, Saborowski A, Yue J, Solomon M, Joseph E, Gadad S, Saborowski M, Kastenhuber E, Fellmann C, Ohara K, Morikami K, Miura T, Lukacs C, Ishii N, Lowe S, Rosen N "Disruption of CRAF-mediated MEK activation is required for effective MEK inhibition in KRAS mutant tumors." *Cancer cell*(2014): 697-710
137. Hochster H, Uboha N, Messersmith W, Gold P, O'Neil B, Cohen D, Denlinger C, Cohen S, Leichman C, Leichman L, Lenz H "Phase II study of selumetinib (AZD6244, ARRY-142886) plus irinotecan as second-line therapy in patients with K-RAS mutated colorectal cancer." *Cancer chemotherapy and pharmacology*(2015): 17-23
138. Blumenschein G, Smit E, Planchard D, Kim D, Cadranel J, De Pas T, Dunphy F, Udud K, Ahn M, Hanna N, Kim J, Mazieres J, Kim S, Baas P, Rappold E, Redhu S, Puski A, Wu F, Jänne P "A randomized phase II study of the MEK1/MEK2 inhibitor trametinib (GSK1120212) compared with docetaxel in KRAS-mutant advanced non-small-cell lung cancer (NSCLC)." *Annals of oncology : official journal of the European Society for Medical Oncology*(2015): 894-901
139. Zhu Z, Aref A, Cohoon T, Barbie T, Imamura Y, Yang S, Moody S, Shen R, Schinzel A, Thai T, Reibel J, Tamayo J, Godfrey J, Qian Z, Page A, Maciag K, Chan E, Silkworth W, Labowsky M, Rozhansky L, Mesirov J, Gillanders W, Ogino S, Hacohen N, Gaudet S, Eck M, Engelman J, Corcoran R, Wong K, Hahn W, Barbie D "Inhibition of KRAS-driven tumorigenicity by interruption of an autocrine cytokine circuit." *Cancer discovery*(2014): 452-65
140. Gerber D, Camidge D, Morgensztern D, Cetnar J, Kelly R, Ramalingam S, Spigel D, Jeong W, Scagliioni P, Zhang S, Li M, Weaver D, Vaikus L, Keegan M, Horobin J, Burns T "Phase 2 study of the focal adhesion kinase inhibitor defactinib (VS-6063) in previously treated advanced KRAS mutant non-small cell lung cancer." *Lung cancer (Amsterdam, Netherlands)*(2020): 60-67
141. Mainardi S, Mulero-Sánchez A, Prahallad A, Germano G, Bosma A, Krimpenfort P, Liefink C, Steinberg J, de Wit N, Gonçalves-Ribeiro S, Nadal E, Bardelli A, Villanueva A, Bernards R "SHP2 is required for growth of KRAS-mutant non-small-cell lung cancer in vivo." *Nature medicine*(2018): 961-967
142. Ruess D, Heynen G, Ciecinski K, Ai J, Berninger A, Kabacaoglu D, Görgülü K, Dantes Z, Wörmann S, Diakopoulos K, Karpathaki A, Kowalska M, Kaya-Aksoy E, Song L, van der Laan E, López-Alberca M, Nazaré M, Reichert M, Saur D, Erkan M, Hopt U, Sainz B, Birchmeier W, Schmid R, Lesina M, Algül H "Mutant KRAS-driven cancers depend on PTPN11/SHP2 phosphatase." *Nature medicine*(2018): 954-960
143. Konstantinidou G, Ramadori G, Torti F, Kangasniemi K, Ramirez R, Cai Y, Behrens C, Dellinger M, Brekken R, Wistuba I, Heguy A, Teruya-Feldstein J, Scagliioni P "RHOA-FAK is a required signaling axis for the maintenance of KRAS-driven lung adenocarcinomas." *Cancer discovery*(2013): 444-57
144. Tang K, Constanzo J, Venkateswaran N, Melegari M, Ilcheva M, Morales J, Skoulidis F, Heymach J, Boothman D, Scagliioni P "Focal Adhesion Kinase Regulates the DNA Damage Response and Its Inhibition Radiosensitizes Mutant KRAS Lung Cancer." *Clinical cancer research : an official journal of the American Association for Cancer Research*(2016): 5851-5863
145. Chen Y, LaMarche M, Chan H, Fekkes P, Garcia-Fortanet J, Acker M, Antonakos B, Chen C, Chen Z, Cooke V, Dobson J, Deng Z, Fei F, Firestone B, Fodor M, Fridrich C, Gao H, Grunenfelder D, Hao H, Jacob J, Ho S, Hsiao K, Kang Z, Karki R, Kato M, Larrow J, La Bonte L, Lenoir F, Liu G, Liu S, Majumdar D, Meyer M, Palermo M, Perez L, Pu M, Price E, Quinn C, Shukya S, Shultz M, Slisz J, Venkatesan K, Wang P, Warmuth M, Williams S, Yang G, Yuan J, Zhang J, Zhu P, Ramsey T, Keen N, Sellers W, Stams T, Fortin P "Allosteric inhibition of SHP2 phosphatase inhibits cancers driven by receptor tyrosine kinases." *Nature*(2016): 148-52
146. Janes M, Zhang J, Li L, Hansen R, Peters U, Guo X, Chen Y, Babbar A, Firdaus S, Darjania L, Feng J, Chen J, Li S, Li S, Long Y, Thach C, Liu Y, Zariwala A, Ely T, Kucharski J, Kessler L, Wu T, Yu K, Wang Y, Yao Y, Deng X, Zarrinkar P, Brehmer D, Dhanak D, Lorenzi M, Hu-Lowe D, Patricelli M, Ren P, Liu Y "Targeting KRAS Mutant Cancers with a Covalent G12C-Specific Inhibitor." *Cell*(2018): 578-589.e17
147. Ostrem J, Peters U, Sos M, Wells J, Shokat K "K-Ras(G12C) inhibitors allosterically control GTP affinity and effector interactions." *Nature*(2013): 548-51
148. Patricelli M, Janes M, Li L, Hansen R, Peters U, Kessler L, Chen Y, Kucharski J, Feng J, Ely T, Chen J, Firdaus S, Babbar A, Ren P, Liu Y "Selective Inhibition of Oncogenic KRAS Output with Small Molecules Targeting the Inactive State." *Cancer discovery*(2016): 316-29
149. Nagasaka M, Potugari B, Nguyen A, Sukari A, Ou S "KRAS Inhibitors- yes but what next? Direct targeting of KRAS- vaccines, adoptive T cell therapy and beyond." *Cancer treatment reviews*(2021): 102309
150. "Sotorasib Edges Closer to Approval." *Cancer discovery*(2021): OF2
151. Li BT, Skoulidis F, Falchook G, et al. "CodeBreak 100: Registrational Phase 2 Trial of Sotorasib in KRAS p.G12C Mutated Non-small Cell Lung Cancer" *Journal of Thoracic Oncology*(2020): PS01.07
152. "A Phase 1/2, Open-label Study Evaluating the Safety, Tolerability, Pharmacokinetics, Pharmacodynamics, and Efficacy of AMG 510 Monotherapy in Subjects With Advanced Solid Tumors With KRAS p.G12C Mutation and AMG 510 Combination Therapy in Subjects With Advanced NSCLC With KRAS p.G12C Mutation (CodeBreak 100)" (2021)
153. Riely G, Ou S, Rybkin I, et al. "990\_PR - KRYSTAL-1: Activity and Preliminary Pharmacodynamic (PD) Analysis of Adagrasib (MRTX849) in Patients (Pts) With Advanced Non-Small-Cell Lung Cancer (NSCLC) Harboring KRASG12C Mutation" *Annals of Oncology*(2021)
154. Jänne P, Riely G, Gadgil S, Heist R, Ou S, Pacheco J, Johnson M, Sabari J, Leventakos K, Yau E, Bazhenova L, Negrao M, Pennell N, Zhang J, Anderes K, Der-Torossian H, Kheoh T, Velastegui K, Yan X, Christensen J, Chao R, Spira A "Adagrasib in Non-Small-Cell Lung Cancer Harboring a KRASG12C Mutation." *The New England journal of medicine*(2022): 120-131
155. Hong DS, Kuboki Y, Strickler JH, et al. "Sotorasib (Soto) plus panitumumab (Pmab) and FOLFIRI for previously treated KRAS G12C-mutated metastatic colorectal cancer (mCRC): CodeBreak 101 phase 1b safety and efficacy." *J Clin Oncol*(2023): 3513
156. Yaeger R, Weiss J, Pelster M, Spira A, Barve M, Ou S, Leal T, Bekaii-Saab T, Pawletz C, Heavey G, Christensen J, Velastegui K, Kheoh T, Der-Torossian H, Klempner S "Adagrasib with or without Cetuximab in Colorectal Cancer with Mutated KRAS G12C." *The New England journal of medicine*(2023): 44-54

## References

157. Klempner S, Weiss J, Pelster M, et al. "KRYSTAL-1: Updated efficacy and safety of adagrasib (MRTX849) with or without cetuximab in patients with advanced colorectal cancer (CRC) harboring a KRASG12C mutation" *Annals of Oncology*(2022)
158. Fakih M, Salvatore L, Esaki T, Modest D, Lopez-Bravo D, Taieb J, Karamouzis M, Ruiz-Garcia E, Kim T, Kuboki Y, Meriggi F, Cunningham D, Yeh K, Chan E, Chao J, Saportas Y, Tran Q, Cremolini C, Pietrantonio F "Sotorasib plus Panitumumab in Refractory Colorectal Cancer with Mutated KRAS G12C." *The New England journal of medicine*(2023): 2125-2139
159. "A Phase 1/2 Multiple Expansion Cohort Trial of MRTX849 in Patients With Advanced Solid Tumors With KRAS G12C Mutation KRYSTAL-1" (2024)
160. Landre T, Justeau G, Assié J, Chouahnia K, Davoine C, Taleb C, Chouaïd C, Duchemann B "Anti-PD-(L)1 for KRAS-mutant advanced non-small-cell lung cancers: a meta-analysis of randomized-controlled trials." *Cancer immunology, immunotherapy : CII*(2022): 719-726
161. Kartolo A, Feilottter H, Hopman W, Fung A, Robinson A "A single institution study evaluating outcomes of PD-L1 high KRAS-mutant advanced non-small cell lung cancer (NSCLC) patients treated with first line immune checkpoint inhibitors." *Cancer treatment and research communications*(2021): 100330
162. Xu Y, Wang Q, Xie J, Chen M, Liu H, Zhan P, Lv T, Song Y "The Predictive Value of Clinical and Molecular Characteristics or Immunotherapy in Non-Small Cell Lung Cancer: A Meta-Analysis of Randomized Controlled Trials." *Frontiers in oncology*(2021): 732214
163. Jeanson A, Tomasini P, Souquet-Bressand M, Brandone N, Boucekine M, Grangeon M, Chaleat S, Khobta N, Milia J, Mhanna L, Greillier L, Biemar J, Nanni I, Ouafik L, Garcia S, Mazières J, Barlesi F, Mascaux C "Efficacy of Immune Checkpoint Inhibitors in KRAS-Mutant Non-Small Cell Lung Cancer (NSCLC)." *Journal of thoracic oncology : official publication of the International Association for the Study of Lung Cancer*(2019): 1095-1101
164. Favazza L, Parseghian C, Kaya C, Nikiforova M, Roy S, Wald A, Landau M, Prokcell S, Dueker J, Johnston E, Brand R, Bahary N, Gorantla V, Rhee J, Pingpank J, Choudry H, Lee K, Paniccia A, Ongchin M, Zureikat A, Bartlett D, Singhi A "KRAS amplification in metastatic colon cancer is associated with a history of inflammatory bowel disease and may confer resistance to anti-EGFR therapy." *Modern pathology : an official journal of the United States and Canadian Academy of Pathology, Inc* (2020): 1832-1843
165. Eberhard D, Johnson B, Amler L, Goddard A, Heldens S, Herbst R, Ince W, Jänne P, Januario T, Johnson D, Klein P, Miller V, Ostland M, Ramies D, Sebisanoovic D, Stinson J, Zhang Y, Seshagiri S, Hillan K "Mutations in the epidermal growth factor receptor and in KRAS are predictive and prognostic indicators in patients with non-small-cell lung cancer treated with chemotherapy alone and in combination with erlotinib." *Journal of clinical oncology : official journal of the American Society of Clinical Oncology*(2005): 5900-9
166. Linardou H, Dahabreh I, Kanaloupiti D, Siannis F, Bafaloukos D, Kosmidis P, Papadimitriou C, Murray S "Assessment of somatic k-RAS mutations as a mechanism associated with resistance to EGFR-targeted agents: a systematic review and meta-analysis of studies in advanced non-small-cell lung cancer and metastatic colorectal cancer." *The Lancet. Oncology*(2008): 962-72
167. Ramos F, Macarulla T, Capdevila J, Elez E, Tabernero J "Understanding the predictive role of K-ras for epidermal growth factor receptor-targeted therapies in colorectal cancer." *Clinical colorectal cancer*(2008): S52-7
168. Campos-Parra A, Zuloaga C, Manríquez M, Avilés A, Borbolla-Escoboza J, Cardona A, Meneses A, Arrieta O "KRAS mutation as the biomarker of response to chemotherapy and EGFR-TKIs in patients with advanced non-small cell lung cancer: clues for its potential use in second-line therapy decision making." *American journal of clinical oncology*(2015): 33-40
169. De Roock W, De Vriendt V, Normanno N, Ciardiello F, Tejpar S "KRAS, BRAF, PIK3CA, and PTEN mutations: implications for targeted therapies in metastatic colorectal cancer." *The Lancet. Oncology*(2011): 594-603
170. Douillard J, Oliner K, Siena S, Tabernero J, Burkes R, Barugel M, Humblet Y, Bodoky G, Cunningham D, Jassem J, Rivera F, Kocákova I, Ruff P, Błasińska-Morawiec M, Šmakal M, Canon J, Rother M, Williams R, Rong A, Wiezorek J, Sidhu R, Patterson S "Panitumumab-FOLFOX4 treatment and RAS mutations in colorectal cancer." *The New England journal of medicine*(2013): 1023-34
171. Valtorta E, Misale S, Sartore-Bianchi A, Nagtegaal I, Paraf F, Lauricella C, Dimartino V, Hobor S, Jacobs B, Ercolani C, Lamba S, Scala E, Veronese S, Laurent-Puig P, Siena S, Tejpar S, Mottlesse M, Punt C, Gambacorta M, Bardelli A, Di Nicolantonio F "KRAS gene amplification in colorectal cancer and impact on response to EGFR-targeted therapy." *International journal of cancer*(2013): 1259-65
172. Li W, Shi Q, Wang W, Liu J, Ren J, Li Q, Hou F "KRAS status and resistance to epidermal growth factor receptor tyrosine-kinase inhibitor treatment in patients with metastatic colorectal cancer: a meta-analysis." *Colorectal disease : the official journal of the Association of Coloproctology of Great Britain and Ireland*(2014): O370-8
173. Mao C, Qiu L, Liao R, Du F, Ding H, Yang W, Li J, Chen Q "KRAS mutations and resistance to EGFR-TKIs treatment in patients with non-small cell lung cancer: a meta-analysis of 22 studies." *Lung cancer (Amsterdam, Netherlands)*(2010): 272-8
174. Ludovini V, Bianconi F, Pistola L, Chiari R, Minotti V, Colella R, Giuffrida D, Tofanetti F, Siggillino A, Flacco A, Baldelli E, Iacono D, Mameli M, Cavaliere A, Crinò L "Phosphoinositide-3-kinase catalytic alpha and KRAS mutations are important predictors of resistance to therapy with epidermal growth factor receptor tyrosine kinase inhibitors in patients with advanced non-small cell lung cancer." *Journal of thoracic oncology : official publication of the International Association for the Study of Lung Cancer*(2011): 707-15
175. Sun J, Hwang D, Ahn J, Ahn M, Park K "Prognostic and predictive value of KRAS mutations in advanced non-small cell lung cancer." *PLoS one*(2013): e64816
176. Pao W, Wang T, Riely G, Miller V, Pan Q, Ladanyi M, Zakowski M, Heelan R, Kris M, Varmus H "KRAS mutations and primary resistance of lung adenocarcinomas to gefitinib or erlotinib." *PLoS medicine*(2005): e17
177. Rulli E, Marabese M, Torri V, Farina G, Veronese S, Bettini A, Longo F, Moscetti L, Ganzinelli M, Lauricella C, Copreni E, Labianca R, Martelli O, Marsoni S, Brogginì M, Garassino M "Value of KRAS as prognostic or predictive marker in NSCLC: results from the TAILOR trial." *Annals of oncology : official journal of the European Society for Medical Oncology*(2015): 2079-84
178. Gomatou G, Syrigos N, Kotteas E "Osimertinib Resistance: Molecular Mechanisms and Emerging Treatment Options." *Cancers*(2023)
179. Leonetti A, Sharma S, Minari R, Perego P, Giovannetti E, Tiseo M "Resistance mechanisms to osimertinib in EGFR-mutated non-small cell lung cancer." *British journal of cancer*(2019): 725-737
180. Ernst S, Uzun S, Paats M, van Marion R, Atmodimedjo P, de Jonge E, van Schaik R, Aerts J, von der Thüsen J, Dubbink H, Dingemans A "Efficacy and Tolerability of Osimertinib and Sotorasib Combination Treatment for Osimertinib Resistance Caused by KRAS G12C Mutation: A Report of Two Cases." *JCO precision oncology*(2023): e2300451
181. Bahcall M, Awad M, Sholl L, Wilson F, Xu M, Wang S, Palakurthi S, Choi J, Ivanova E, Leonardi G, Ulrich B, Paweletz C, Kirschmeier P, Watanabe M, Baba H, Nishino M, Nagy R, Lanman R, Capelletti M, Chambers E, Redig A, VanderLaan P, Costa D, Imamura Y, Jänne P "Amplification of Wild-type KRAS Imparts Resistance to Crizotinib in MET Exon 14 Mutant Non-Small Cell Lung Cancer." *Clinical cancer research : an official journal of the American Association for Cancer Research*(2018): 5963-5976

## References

182. Doebele R, Pilling A, Aisner D, Kutateladze T, Le A, Weickhardt A, Kondo K, Linderman D, Heasley L, Franklin W, Varella-Garcia M, Camidge D "Mechanisms of resistance to crizotinib in patients with ALK gene rearranged non-small cell lung cancer." *Clinical cancer research : an official journal of the American Association for Cancer Research*(2012): 1472-82
183. Rossing H, Grauslund M, Urbanska E, Melchior L, Rask C, Costa J, Skov B, Sørensen J, Santoni-Rugiu E "Concomitant occurrence of EGFR (epidermal growth factor receptor) and KRAS (V-Ki-ras2 Kirsten rat sarcoma viral oncogene homolog) mutations in an ALK (anaplastic lymphoma kinase)-positive lung adenocarcinoma patient with acquired resistance to crizotinib: a case report." *BMC research notes*(2013): 489
184. Cargnelutti M, Corso S, Pergolizzi M, Mévellec L, Aisner D, Dziadziuszko R, Varella-Garcia M, Comoglio P, Doebele R, Vialard J, Giordano S "Activation of RAS family members confers resistance to ROS1 targeting drugs." *Oncotarget*(2015): 5182-94
185. Bordi P, Tiseo M, Rofi E, Petrini I, Restante G, Danesi R, Del Re M "Detection of ALK and KRAS Mutations in Circulating Tumor DNA of Patients With Advanced ALK-Positive NSCLC With Disease Progression During Crizotinib Treatment." *Clinical lung cancer*(2017): 692-697
186. Awad M, Liu S, Rybkin I, Arbour K, Dilly J, Zhu V, Johnson M, Heist R, Patil T, Riely G, Jacobson J, Yang X, Persky N, Root D, Lowder K, Feng H, Zhang S, Haigis K, Hung Y, Sholl L, Wolpin B, Wiese J, Christiansen J, Lee J, Schrock A, Lim L, Garg K, Li M, Engstrom L, Waters L, Lawson J, Olson P, Lito P, Ou S, Christensen J, Jänne P, Aguirre A "Acquired Resistance to KRASG12C Inhibition in Cancer." *The New England journal of medicine*(2021): 2382-2393
187. Tanaka N, Lin J, Li C, Ryan M, Zhang J, Kiedrowski L, Michel A, Syed M, Fella K, Sakhi M, Baiev I, Juric D, Gainor J, Klempner S, Lennerz J, Siravegna G, Bar-Peled L, Hata A, Heist R, Corcoran R "Clinical Acquired Resistance to KRASG12C Inhibition through a Novel KRAS Switch-II Pocket Mutation and Polyclonal Alterations Converging on RAS-MAPK Reactivation." *Cancer discovery*(2021): 1913-1922
188. Li BT, Velcheti V, Price TJ, et al. "Largest evaluation of acquired resistance to sotorasib in KRAS p.G12C-mutated non-small cell lung cancer (NSCLC) and colorectal cancer (CRC): Plasma biomarker analysis of CodeBreak100." *Journal of Clinical Oncology*(2022)
189. Di Federico A, Ricciotti I, Favorito V, Michelina S, Scaparoni P, Metro G, De Giglio A, Pecci F, Lamberti G, Ambrogio C, Ricciuti B "Resistance to KRAS G12C Inhibition in Non-small Cell Lung Cancer." *Current oncology reports*(2023): 1017-1029
190. Brown C, Lain S, Verma C, Fersht A, Lane D "Awakening guardian angels: drugging the p53 pathway." *Nature reviews. Cancer*(2009): 862-73
191. Malkin D, Li F, Strong L, Fraumeni J, Nelson C, Kim D, Kassel J, Gryka M, Bischoff F, Tainsky M "Germ line p53 mutations in a familial syndrome of breast cancer, sarcomas, and other neoplasms." *Science (New York, N.Y.)*(1990): 1233-8
192. Srivastava S, Zou Z, Pirollo K, Blattner W, Chang E "Germ-line transmission of a mutated p53 gene in a cancer-prone family with Li-Fraumeni syndrome." *Nature*(1991): 747-9
193. Santibáñez-Koref M, Birch J, Hartley A, Jones P, Craft A, Eden T, Crowther D, Kelsey A, Harris M "p53 germline mutations in Li-Fraumeni syndrome." *Lancet (London, England)*(1991): 1490-1
194. Wang Y, Lin R, Tan Y, Chen J, Chen C, Wang Y "Wild-type p53 overexpression and its correlation with MDM2 and p14ARF alterations: an alternative pathway to non-small-cell lung cancer." *Journal of clinical oncology : official journal of the American Society of Clinical Oncology*(2005): 154-64
195. Koga T, Hashimoto S, Sugio K, Yoshino I, Nakagawa K, Yonemitsu Y, Sugimachi K, Sueishi K "Heterogeneous distribution of P53 immunoreactivity in human lung adenocarcinoma correlates with MDM2 protein expression, rather than with P53 gene mutation." *International journal of cancer*(2001): 232-9
196. Kato S, Han S, Liu W, Otsuka K, Shibata H, Kanamaru R, Ishioka C "Understanding the function-structure and function-mutation relationships of p53 tumor suppressor protein by high-resolution missense mutation analysis." *Proceedings of the National Academy of Sciences of the United States of America*(2003): 8424-9
197. Houben R, Hesbacher S, Schmid C, Kauczok C, Flohr U, Haferkamp S, Müller C, Schrama D, Wischhusen J, Becker J "High-level expression of wild-type p53 in melanoma cells is frequently associated with inactivity in p53 reporter gene assays." *PloS one*(2011): e22096
198. Olivier M, Petitjean A, Marcel V, Pétrel A, Mounawar M, Plymoth A, de Fromental C, Hainaut P "Recent advances in p53 research: an interdisciplinary perspective." *Cancer gene therapy*(2009): 1-12
199. Chang Y, Wu C, Shih J, Lee Y "Comparison of p53 and epidermal growth factor receptor gene status between primary tumors and lymph node metastases in non-small cell lung cancers." *Annals of surgical oncology*(2011): 543-50
200. Jiang R, Zhang B, Teng X, Hu P, Xu S, Zheng Z, Liu R, Tang T, Ye F "Validating a targeted next-generation sequencing assay and profiling somatic variants in Chinese non-small cell lung cancer patients." *Scientific reports*(2020): 2070
201. Mattioni M, Soddu S, Prodosmo A, Visca P, Conti S, Alessandrini G, Facciolo F, Strigari L "Prognostic role of serum p53 antibodies in lung cancer." *BMC cancer*(2015): 148
202. Bircan A, Bircan S, Kapucuoglu N, Songur N, Ozturk O, Akkaya A "Maspin, VEGF and p53 expression in small biopsies of primary advanced lung cancer and relationship with clinicopathologic parameters." *Pathology oncology research : POR*(2010): 553-61
203. Kim Y, Hammerman P, Kim J, Yoon J, Lee Y, Sun J, Wilkerson M, Pedamallu C, Cibulskis K, Yoo Y, Lawrence M, Stojanov P, Carter S, McKenna A, Stewart C, Sivachenko A, Oh I, Kim H, Choi Y, Kim K, Shim Y, Kim K, Song S, Na K, Choi Y, Hayes D, Kim J, Cho S, Kim Y, Ahn J, Ahn M, Getz G, Meyerson M, Park K "Integrative and comparative genomic analysis of lung squamous cell carcinomas in East Asian patients." *Journal of clinical oncology : official journal of the American Society of Clinical Oncology*(2014): 121-8
204. Dong Z, Zhong W, Zhang X, Su J, Xie Z, Liu S, Tu H, Chen H, Sun Y, Zhou Q, Yang J, Yang X, Lin J, Yan H, Zhai H, Yan L, Liao R, Wu S, Wu Y "Potential Predictive Value of TP53 and KRAS Mutation Status for Response to PD-1 Blockade Immunotherapy in Lung Adenocarcinoma." *Clinical cancer research : an official journal of the American Association for Cancer Research*(2017): 3012-3024
205. Mansuet-Lupo A, Alifano M, Pécuchet N, Biton J, Becht E, Goc J, Germain C, Ouakrim H, Régnard J, Cremer I, Laurent-Puig P, Dieu-Nosjean M, Blons H, Damotte D "Intratumoral Immune Cell Densities Are Associated with Lung Adenocarcinoma Gene Alterations." *American journal of respiratory and critical care medicine*(2016): 1403-1412
206. Kadara H, Choi M, Zhang J, Parra E, Rodriguez-Canales J, Gaffney S, Zhao Z, Behrens C, Fujimoto J, Chow C, Yoo Y, Kalhor N, Moran C, Rimm D, Swisher S, Gibbons D, Heymach J, Kaftan E, Townsend J, Lynch T, Schllessinger J, Lee J, Lifton R, Wistuba I, Herbst R "Whole-exome sequencing and immune profiling of early-stage lung adenocarcinoma with fully annotated clinical follow-up." *Annals of oncology : official journal of the European Society for Medical Oncology*(2017): 75-82
207. Van Egeren D, Kohli K, Warner J, Bedard P, Riely G, Lepisto E, Schrag D, LeNoue-Newton M, Catalano P, Kehl K, Michor F "Genomic analysis of early-stage lung cancer reveals a role for TP53 mutations in distant metastasis." *Scientific reports*(2022): 19055

## References

208. Schuler P, Harasymczuk M, Visus C, Deleo A, Trivedi S, Lei Y, Argiris A, Gooding W, Butterfield L, Whiteside T, Ferris R "Phase I dendritic cell p53 peptide vaccine for head and neck cancer." *Clinical cancer research : an official journal of the American Association for Cancer Research*(2014): 2433-44
209. Vermeij R, Leffers N, van der Burg S, Melief C, Daemen T, Nijman H "Immunological and clinical effects of vaccines targeting p53-overexpressing malignancies." *Journal of biomedicine & biotechnology*(2011): 702146
210. Saito H, Ando S, Morishita N, Lee K, Dator D, Dy D, Shigemura K, Adhim Z, Nibu K, Fujisawa M, Shirakawa T "A combined lymphokine-activated killer (LAK) cell immunotherapy and adenovirus-p53 gene therapy for head and neck squamous cell carcinoma." *Anticancer research*(2014): 3365-70
211. Ma C, Janetka J, Piwnica-Worms H "Death by releasing the breaks: CHK1 inhibitors as cancer therapeutics." *Trends in molecular medicine*(2011): 88-96
212. Hirai H, Arai T, Okada M, Nishibata T, Kobayashi M, Sakai N, Imagaki K, Ohtani J, Sakai T, Yoshizumi T, Mizuarai S, Iwasawa Y, Kotani H "MK-1775, a small molecule Wee1 inhibitor, enhances anti-tumor efficacy of various DNA-damaging agents, including 5-fluorouracil." *Cancer biology & therapy*(2010): 514-22
213. Bridges K, Hirai H, Buser C, Brooks C, Liu H, Buchholz T, Molkentine J, Mason K, Meyn R "MK-1775, a novel Wee1 kinase inhibitor, radiosensitizes p53-defective human tumor cells." *Clinical cancer research : an official journal of the American Association for Cancer Research*(2011): 5638-48
214. Vilgelm A, Pawlikowski J, Liu Y, Hawkins O, Davis T, Smith J, Weller K, Horton L, McClain C, Ayers G, Turner D, Essaka D, Stewart C, Sosman J, Kelley M, Ecsedy J, Johnston J, Richmond A "Mdm2 and aurora kinase inhibitors synergize to block melanoma growth by driving apoptosis and immune clearance of tumor cells." *Cancer research*(2015): 181-93
215. Li Z, Sun Y, Chen X, Squires J, Nowroozizadeh B, Liang C, Huang J "p53 Mutation Directs AURKA Overexpression via miR-25 and FBXW7 in Prostatic Small Cell Neuroendocrine Carcinoma." *Molecular cancer research : MCR*(2015): 584-91
216. Katayama H, Sen S "Functional significance of Aurora kinase A regulatory interactions with p53-ERα complex in human breast cancer cells." *Hormones & cancer*(2011): 117-24
217. Tentler J, Ionkina A, Tan A, Newton T, Pitts T, Glogowska M, Kabos P, Sartorius C, Sullivan K, Espinosa J, Eckhardt S, Diamond J "p53 Family Members Regulate Phenotypic Response to Aurora Kinase A Inhibition in Triple-Negative Breast Cancer." *Molecular cancer therapeutics*(2015): 1117-29
218. Gully C, Velazquez-Torres G, Shin J, Fuentes-Mattei E, Wang E, Carlock C, Chen J, Rothenberg D, Adams H, Choi H, Guma S, Phan L, Chou P, Su C, Zhang F, Chen J, Yang T, Yeung S, Lee M "Aurora B kinase phosphorylates and instigates degradation of p53." *Proceedings of the National Academy of Sciences of the United States of America*(2012): E1513-22
219. Marxer M, Ma H, Man W, Poon R "p53 deficiency enhances mitotic arrest and slippage induced by pharmacological inhibition of Aurora kinases." *Oncogene*(2014): 3550-60
220. El-Deiry W "The role of p53 in chemosensitivity and radiosensitivity." *Oncogene*(2003): 7486-95
221. Miyasaka A, Oda K, Ikeda Y, Sone K, Fukuda T, Inaba K, Makii C, Enomoto A, Hosoya N, Tanikawa M, Uehara Y, Arimoto T, Kuramoto H, Wada-Hiraike O, Miyagawa K, Yano T, Kawana K, Osuga Y, Fujii T "PI3K/mTOR pathway inhibition overcomes radioresistance via suppression of the HIF1-α/VEGF pathway in endometrial cancer." *Gynecologic oncology*(2015): 174-80