

Guardant360 基因檢測服務報告

醫師姓名：廖映庭

受檢者姓名：KONG SANG

送檢編號：A1168951

檢測次數：第一次

提醒

基因數據乃屬個人隱私，切勿輕易向任何個人、團體或非您的授權者透漏本報告內容。若您有任何疑慮，歡迎來電洽詢，我們很樂意為您提供更詳細的諮詢服務。若因郵遞錯誤收此檔，請予銷毀，多謝合作。

詳細資訊



康誠生技股份有限公司 客戶服務中心

諮詢時間 | 週一～週五 9:00～17:00 (國定假日除外)


諮詢專線 | 02-55696099

客服信箱 | service.gb@healthconn.com

REPORTING	PHYSICIAN	 <div>Complete Tumor Response Map on page 2</div>
Report Date: OCT-09-2024	Ying Ting Liao	
Receipt Date: OCT-05-2024	Account: Genconn Biotech Co., LTD	
Collection Date: OCT-04-2024	Address: 5F., No. 54, Sec. 1, Jhongsiao E. Rd., Zhongzheng Dist., Beixin Rd, Xindian Dist, Taipei City, 100, Taiwan	
Specimen: Blood	Ph: +886 963 820 633 Fax: N/A	
Status: FINAL	Additional Recipient: N/A	

Summary of Detected Somatic Alterations, Immunotherapy Biomarkers & Associated Treatment Options

KEY  Approved in indication  Approved in other indication  Lack of response

Detected Alteration(s) / Biomarker(s)	Associated FDA-approved therapies	Clinical trial availability (see page 3)	% cfDNA or Amplification
EGFR E746_A750del (Exon 19 deletion)	 Afatinib, Amivantamab, Amivantamab+lazertinib, Dacomitinib, Erlotinib, Erlotinib+ramucirumab, Gefitinib, Osimertinib	Yes	0.7%
TP53 P151S	None	No	0.3%

Comments
Reported by: JV4

Additional Biomarkers

Biomarker	Additional Details
MSI-High	NOT DETECTED

We evaluated this sample for 74 genes, including the following guideline-recommended genes for NSCLC

EGFR(T790M and others)

ALK

ROS1

BRAF

MET

ERBB2(HER2)


RET

NTRK

KRAS

Guardant360 Tumor Response Map

The Guardant360 Tumor Response Map illustrates the variant allele fraction (% cfDNA) of observed somatic variants at each sample submission. Amplifications are not plotted, and only the first and last five test dates are plotted. Please see the Physician Portal (portal.guardanthealth.com) for the Tumor Response Map with all test dates.



Detected Alteration(s) / Biomarker(s)	% cfDNA or Amp
EGFR E746_A750del (Exon 19 deletion)	0.7%
TP53 P151S	0.3%

The table above annotates the variant allele fraction (% cfDNA) detected in this sample, listed in descending order.
§ See definitions section for more detail

Available Clinical Trials (within the same state as the ordering physician)

There may be additional trials not listed here. Visit: portal.guardanthealth.com or email clientservices@guardanthealth.com with A1168951 in the subject line of the email, for additional trials.

Alteration	Trial ID / Contact	Title	Phase	Site(s)
EGFR E746_A750del	NCT04077463 Study Contact,Participate-In-This-Study@its.jnj.com,844-434-4210	A Study of Lazertinib as Monotherapy or in Combination With Amivantamab in Participants With Advanced Non-small Cell Lung Cancer	Phase 1	Kaohsiung, Taiwan Taipei City, Taiwan Tainan, Taiwan Taichung, Taiwan
	NCT05215548 Jin-Shing Chen, M.D., Ph.D.,chenjs@ntu.edu.tw,886-2-2322-0322	Primary Tumor Resection With EGFR TKI for Stage IV NSCLC	Phase 2	Taipei, Taiwan (2)
	NCT05442060 Anna Hu,annahu@obipharma.com,886-2-27866589 x104	To Evaluate OBI-833/OBI-821 in Combination With First-Line Erlotinib in Patients With EGFR-Mutated, Globo H-Positive, Locally Advanced or Metastatic Non-Small Cell Lung Cancer	Phase 2	New Taipei City, Taiwan Taoyuan, Taiwan Taichung, Taiwan Taipei, Taiwan (4)
	NCT05801029 AstraZeneca Clinical Study Information Center,information.center@astrazeneca.com,1-877-240-9479	A Study to Investigate Safety and Efficacy of Osimertinib and Amivantamab in Participants With Non-small Cell Lung Cancer With Common Epidermal Growth Factor Receptor Mutations	Phase 2	Yunlin, Taiwan Taipei City, Taiwan Kaohsiung, Taiwan (2) Taipei, Taiwan (2) Additional trial sites available
	NCT06120140 Study Contact,Participate-In-This-Study@its.jnj.com,844-434-4210	Enhanced Dermatological Care to Reduce Rash and Paronychia in Epidermal Growth Factor Receptor (EGFR)-Mutated Non-Small Cell Lung Cancer (NSCLC) Treated First-line With Amivantamab Plus Lazertinib	Phase 2	Kaohsiung City, Taiwan Taipei, Taiwan Taoyuan City, Taiwan Taichung City, Taiwan Additional trial sites available

Visit portal.guardanthealth.com for trials not within the same state as the physician's office

More clinical trial options available at portal.guardanthealth.com

Definitions

Deletion (Del): The following alteration was detected in this patient: *EGFR* E746_A750del. Guardant360 detects short deletions in exons of certain genes (see Table 1), including potential splice site-disrupting events.

Interpretation

Somatic alterations were detected in the circulating cell-free DNA isolated from this patient's blood specimen. These genomic alterations are cancer-associated somatic variants, some of which have been associated with either increased or reduced clinical response to specific treatments. The percentage of altered cell-free DNA circulating (% cfDNA) in blood is related to the unique tumor biology of each patient. Factors that may affect the % cfDNA of detected somatic alterations include tumor growth, turn over, size, heterogeneity, vascularization, disease progression, and treatment.

Method and Limitations

Guardant360 sequences 74 cancer-associated genes to identify somatic alterations. Cell-free DNA (cfDNA) is extracted from plasma, enriched for targeted regions, and sequenced using the Illumina platform and hg19 as the reference genome. All exons are sequenced in some genes; only clinically significant exons are sequenced in other genes. The types of genomic alterations detected by Guardant360 include single nucleotide variants, gene amplifications, fusions, short insertions/deletions (longest detected, 70 base pairs), and splice site disrupting events (see Table 1). Microsatellite Instability (MSI) is assessed for all cancer types by evaluating somatic changes in the length of repetitive sequences on the Guardant360 panel. A "Not Detected" result in samples where the highest % cfDNA is < 0.2% is an inconclusive result because it does not preclude MSI-High status in tissue. MSI status is currently not reported for earlier panel versions. This version of the Guardant360 test is not validated for the detection of other types of genomic alterations, such as complex rearrangements or gene deletions. Certain sample or variant characteristics, such as low cfDNA concentration, may result in reduced analytic sensitivity. Guardant360 cannot discern the source of circulating cfDNA, and for some variants in the range of ~40 to 60% cfDNA, the test cannot easily distinguish germline variants from somatic alterations. Guardant360 is not validated for the detection of germline or de novo variants that are associated with hereditary cancer risk. Tissue genotyping should be considered when plasma genotyping is negative, if clinically appropriate.

Table 1: Genes on the Guardant360 Panel

Guardant360 reports single nucleotide variants, splice site mutations, and insertion and deletion variants (indels) in all clinically relevant exons in 74 genes and reports other variant types in select genes as indicated below.

AKT1	ALK #	APC	AR †	ARAF	ARID1A	ATM	BRAF †	BRCA1
BRCA2	CCND1 †	CCND2 †	CCNE1 †	CDH1	CDK12	CDK4 †	CDK6 †	CDKN2A
CTNNB1	DDR2	EGFR †	ERBB2 †	ESR1	EZH2	FBXW7	FGFR1 †	FGFR2 † #
FGFR3 #	GATA3	GNA11	GNAQ	GNAS	HNF1A	HRAS	IDH1	IDH2
JAK2	JAK3	KIT †	KRAS †	MAP2K1	MAP2K2	MAPK1	MAPK3	MET †
MLH1	MPL	MTOR	MYC †	NF1	NFE2L2	NOTCH1	NPM1	NRAS
NTRK1 #	NTRK3	PDGFRA †	PIK3CA †	PTEN	PTPN11	RAF1 †	RB1	RET #
RHEB	RHOA	RIT1	ROS1 #	SMAD4	SMO	STK11	TERT ‡	TP53
TSC1	VHL							

‡ Guardant360 reports alterations in the promoter region of this gene.
Guardant360 reports fusion events involving this gene.
† Guardant360 reports amplifications of this gene.

About the Test

The Guardant360 assay was developed and its performance characteristics were determined by Guardant Health, Inc. This test has not been cleared or approved by the U.S. Food and Drug Administration (FDA). The FDA has determined that such clearance or approval is not necessary. This test may be used for clinical purposes and should not be regarded as investigational or for research only. Guardant Health's clinical reference laboratory is certified under the Clinical Laboratory Improvement Amendments of 1988 (CLIA) as qualified to perform high complexity clinical laboratory testing. The laboratory report should be interpreted in the context of other clinical information and laboratory, pathology, and imaging studies by a qualified medical professional prior to initiating or changing a patient's treatment plan. The selection of any, all, or none of the drugs associated with potential clinical benefit (or potential lack of clinical benefit) is entirely at the discretion of the treating medical professional. Drug and trial information are based on the diagnosis written on the submitted test request form; this information is not based on any supplemental information provided by the requesting medical professional, including pathology reports or other molecular studies. Some drugs listed in this report may not be approved or cleared by the FDA for the indicated use. Guardant Health makes no endorsement, express or implied, of any product, physician, or procedure contained in this report. This report makes no promises or guarantees that a particular medication will affect (or not affect) the clinical outcome of any patient.

Testing Performed at: Guardant Health

Laboratory Director: Martina Lefterova, MD PhD | CLIA ID: 05D2070300 | CAP #: 8765297 | 505 Penobscot Drive, Redwood City, CA, 94063, USA

Additional information is available

Any therapeutic annotations are based on publicly available information. This information is described in the "Detailed Therapy Results" and "Relevance of Detected Alterations" sections.

Visit portal.guardanthealth.com or email clientservices@guardanthealth.com with A1168951 in the subject line of the email for:

- Additional clinical trials
- Detailed Therapy Results
- Relevance of Detected Alterations
- References

If you would like to receive this additional information with every Guardant360 report, please call client services at [855.698.8887](tel:855.698.8887) to opt-in.

Additional information begins on the next page.

List of Available Clinical Trials

Alteration	Trial ID / Contact	Title	Phase	Site (number in parenthesis is count of trial sites)
EGFR E746_A750del	NCT03574402 Yi-Long Wu, Professor, syylwu@live.cn, 862083827812	Phase II Umbrella Study Directed by Next Generation Sequencing	Phase 2	China
	NCT04077463 Study Contact, Participate-In-This-Study@its.jnj.com, 844-434-4210	A Study of Lazertinib as Monotherapy or in Combination With Amivantamab in Participants With Advanced Non-small Cell Lung Cancer	Phase 1	Seattle, WA; Detroit, MI; Saint Louis, MO; Philadelphia, PA; Portland, OR; Salt Lake City, UT; Tampa, FL; Fairfax, VA; Boston, MA (2); New York, NY (2); CA (5); Puerto Rico; Japan (7); China (9); Taiwan (4); Korea, Republic of (4); Italy (5); France (7); Germany (8); Spain (8)
	NCT04841811 Yi-Long Wu, doctor, syylwu@live.cn, 86-13544561166	MRD Guiding Treatment After Almonertinib Induction Therapy for EGFR+ Stage III NSCLC in the MDT Diagnostic Model.	Phase 3	China
	NCT04922138 Baohui Han, doctor, 18930858216@163.com, 18930858216	Aumolertinib Adjuvant Therapy of Resectable Stage I EGFR+ NSCLC With High-grade Patterns	Phase 2	China (2)
	NCT05215548 Jin-Shing Chen, M.D., Ph.D., chenjs@ntu.edu.tw, 886-2-2322-0322	Primary Tumor Resection With EGFR TKI for Stage IV NSCLC	Phase 2	Taiwan (2)
	NCT05442060 Anna Hu, annahu@obipharma.com, 886-2-27866589 x104	To Evaluate OBI-833/OBI-821 in Combination With First-Line Erlotinib in Patients With EGFR-Mutated, Globo H-Positive, Locally Advanced or Metastatic Non-Small Cell Lung Cancer	Phase 2	Taiwan (7)
	NCT05801029 AstraZeneca Clinical Study Information Center, information.center@astrazeneca.com, 1-877-240-9479	A Study to Investigate Safety and Efficacy of Osimertinib and Amivantamab in Participants With Non-small Cell Lung Cancer With Common Epidermal Growth Factor Receptor Mutations	Phase 2	Glendale, CA; New York, NY; Washington, DC; Sacramento, CA; Beverly Hills, CA; Fairfax, VA; Canada (2); Singapore (3); Hong Kong (4); Taiwan (9); Korea, Republic of (7); Malaysia (6); Thailand (5)
	NCT06041776 Shun Lu, M.D., shunlu@sjtu.edu.cn, +86 21 2220 0000	Adjuvant Befotertinib in Stage IB-IIIB Non-small Cell Lung Cancer With Positive EGFR Sensitive Mutations	Phase 3	China (3)
	NCT06043973 Degan Lu, Professor, deganlu@126.com, 18753157623	Almonertinib Combined With Anlotinib as First-line Treatment for Advanced Non-small Cell Lung Cancer	Phase 3	China
	NCT06120140 Study Contact, Participate-In-This-Study@its.jnj.com, 844-434-4210	Enhanced Dermatological Care to Reduce Rash and Paronychia in Epidermal Growth Factor Receptor (EGFR)-Mutated Non-Small Cell Lung Cancer (NSCLC) Treated First-line With Amivantamab Plus Lazertinib	Phase 2	Renton, WA; Hinsdale, IL; Westbury, NY; Cleveland, OH; W. Salem, WI; Reno, NV; Springfield, MO; Fairfax, VA; Flemington, NJ; Wilson, NC; CA (15); Argentina (5); Turkey (8); China (11); Taiwan (5); Korea, Republic of (3); Brazil (11); Malaysia (4); France (3); Germany (6); Spain (9)

Detailed Therapy Results

Alteration	Drug	Trade Name	Target	Current Status
EGFR E746_A750del (Exon 19 deletion)	ABT-101		Egfr/Her2 inhibitor.	Phase 1 (Non-small cell lung carcinoma (NSCLC)) Phase 1 (Head and neck squamous cell carcinoma (HNSCC))
	Afatinib	Gilotrif	Irreversible pan-ErbB kinase inhibitor.	Phase 3 (Non-small cell lung carcinoma (NSCLC)) FDA Approved in other indications (Lung squamous cell carcinoma, EGFR-mutant NSCLC)
	Amivantamab	Rybrevant	Bispecific anti-Met/Egfr antibody.	Phase 3 (Non-small cell lung carcinoma (NSCLC)) FDA Approved in other indications (NSCLC with EGFR exon 20 insertion, NSCLC with EGFR exon 19 del /L858R)
	Amivantamab+lazertinib	Rybrevant+Lazcluze	Bispecific anti-Met/Egfr antibody+Egfr tyrosine kinase inhibitor.	Phase 3 (Non-small cell lung carcinoma (NSCLC)) FDA Approved in other indications (NSCLC with EGFR exon 19 del/L858R)
	Aumolertinib		Egfr T790M inhibitor.	Phase 3 (Non-small cell lung carcinoma (NSCLC))
	Avitinib		Irreversible mutation-specific Egfr kinase inhibitor.	Phase 2 (Non-small cell lung carcinoma (NSCLC)) Phase 1 (Non-Hodgkin lymphoma (NHL))
	BAY2927088		Egfr/Her2 kinase inhibitor.	Phase 1 (Non-small cell lung carcinoma (NSCLC))
	BBP-398		Shp-2 inhibitor.	Phase 1 (Non-small cell lung carcinoma (NSCLC)) Phase 1 (Solid Tumor)
	BDTX-1535		Irreversible brain-penetrant fourth generation Egfr inhibitor.	Phase 1 (Non-small cell lung carcinoma (NSCLC)) Phase 1 (Glioma)
	Befotertinib		Third generation mutation-specific (T790M, L858R, exon 19 deletion) Egfr tyrosine kinase inhibitor.	Phase 3 (Non-small cell lung carcinoma (NSCLC))
	BLU-945		Fourth generation Egfr inhibitor targeting T790M and T790M/C797S mutations.	Phase 2 (Non-small cell lung carcinoma (NSCLC))
	BPI-361175		Fourth generation Egfr inhibitor targeting T790M and T790M/C797S mutations.	Phase 2 (Non-small cell lung carcinoma (NSCLC))
	CLN-081		Covalent mutation-specific (L858R, T790M, exon 19 deletion, exon 20 insertion) Egfr tyrosine kinase inhibitor.	Phase 2 (Non-small cell lung carcinoma (NSCLC))
	CM93		Third generation mutation-specific (T790M, L858R, exon 19 deletion) Egfr tyrosine kinase inhibitor.	Phase 1 (Glioblastoma)
	Dacomitinib	Vizimpro	Pan-ErbB family tyrosine kinase inhibitor.	Phase 3 (Non-small cell lung carcinoma (NSCLC)) FDA Approved in other indications (EGFR-mutant lung carcinoid, EGFR-mutant lung large cell neuroendocrine carcinoma, EGFR-mutant NSCLC)

Detailed Therapy Results

Alteration	Drug	Trade Name	Target	Current Status
	ERAS-601		Shp-2 inhibitor.	Phase 2 (Solid Tumor)
	Erlotinib	Tarceva	Egfr tyrosine kinase inhibitor.	Phase 3 (Non-small cell lung carcinoma (NSCLC)) FDA Approved in other indications (EGFR-mutant lung carcinoid, EGFR-mutant lung large cell neuroendocrine carcinoma, NSCLC with EGFR exon 19 del/L858R, Pancreatic carcinoma, EGFR-mutant NSCLC)
	Erlotinib+bevacizumab	Tarceva+Avastin	Egfr tyrosine kinase inhibitor + anti-VEGF-A monoclonal antibody combination.	Phase 3 (Non-small cell lung carcinoma (NSCLC)) Phase 3 (Lung cancer)
	Erlotinib+ramucirumab	Tarceva+Cyramza	Egfr tyrosine kinase inhibitor + anti-VEGFR-2 monoclonal antibody combination.	Phase 3 (Non-small cell lung carcinoma (NSCLC)) FDA Approved in other indications (NSCLC with EGFR exon 19 del/L858R)
	ET0038		Shp-2 inhibitor.	Phase 1 (Solid Tumor)
	Furmonertinib		Third generation mutation-specific (T790M, L858R, exon 19 deletion) Egfr tyrosine kinase inhibitor.	Phase 2 (Non-small cell lung carcinoma (NSCLC))
	FWD1509		Egfr/Her2 kinase inhibitor.	Phase 2 (Non-small cell lung carcinoma (NSCLC))
	Gefitinib	Iressa	Egfr tyrosine kinase inhibitor.	Phase 3 (Non-small cell lung carcinoma (NSCLC)) FDA Approved in other indications (EGFR-mutant lung carcinoid, EGFR-mutant lung large cell neuroendocrine carcinoma, EGFR-mutant NSCLC)
	H002		Fourth generation Egfr inhibitor targeting exon 19del /L858R, T790M, and C797S mutations.	Phase 2 (Non-small cell lung carcinoma (NSCLC))
	HBI-2376		Shp-2 inhibitor.	Phase 1 (Solid Tumor)
	Hemay022		Egfr tyrosine kinase inhibitor.	Phase 1 (Breast carcinoma (HER2+))
	Icotinib	Conmana	Egfr inhibitor.	Phase 3 (Non-small cell lung carcinoma (NSCLC)) Phase 2 (Esophageal carcinoma)
	JIN-A02		Fourth generation Egfr inhibitor targeting T790M and T790M/C797S mutations.	Phase 2 (Non-small cell lung carcinoma (NSCLC))
	Lazertinib	Lazcluze	Third generation mutation-specific Egfr tyrosine kinase inhibitor.	Phase 2 (Non-small cell lung carcinoma (NSCLC))
	Lifirafenib		Dual Braf/Egfr inhibitor.	Phase 1 (Non-small cell lung carcinoma (NSCLC)) Phase 1 (Solid Tumor, Brain and Central Nervous System Tumors)
	MCLA-129		Anti-EGFR/c-Met bispecific antibody.	Phase 2 (Non-small cell lung carcinoma (NSCLC)) Phase 2 (Gastric carcinoma, Head and neck squamous cell carcinoma (HNSCC), Esophageal squamous cell carcinoma)

Detailed Therapy Results

Alteration	Drug	Trade Name	Target	Current Status
	Mobocertinib	Exkivity	Mutation-specific Egfr/Her2 inhibitor.	Phase 3 (Non-small cell lung carcinoma (NSCLC)) Phase 3 (NSCLC with EGFR exon 20 insertion, Lung cancer)
	Modotuximab		Anti-EGFR antibody.	Phase 1 (Gastric carcinoma, Colorectal carcinoma (CRC))
	Nazartinib		Third generation EGFR mutant-specific (T790M, L858R, exon 19 deletion) tyrosine kinase inhibitor.	Phase 2 (Non-small cell lung carcinoma (NSCLC))
	Neratinib	Nerlynx	Egfr/Her2/ErbB4 inhibitor.	Phase 2 (Non-small cell lung carcinoma (NSCLC)) FDA Approved in other indications (Breast carcinoma (HER2+))
	NX-019		Egfr inhibitor.	Phase 1 (Solid Tumor)
	Olafertinib		Third generation mutation-specific (T790M, L858R, exon 19 deletion) Egfr tyrosine kinase inhibitor.	Phase 2 (Non-small cell lung carcinoma (NSCLC)) Phase 1 (Solid Tumor)
	Osimertinib	Tagrisso	Egfr T790M inhibitor.	Phase 3 (Non-small cell lung carcinoma (NSCLC)) FDA Approved in other indications (EGFR-mutant lung carcinoid, EGFR-mutant lung large cell neuroendocrine carcinoma, EGFR-mutant NSCLC)
	Pirotinib		ErbB family inhibitor.	Phase 1 (Solid Tumor)
	Pozotinib		Egfr/Her2/ErbB4 kinase inhibitor.	Phase 2 (Non-small cell lung carcinoma (NSCLC)) Phase 2 (Gastric carcinoma, Head and neck squamous cell carcinoma (HNSCC), Breast carcinoma, Esophageal squamous cell carcinoma, Colorectal carcinoma (CRC))
	Pyrotinib		Egfr/Her2 kinase inhibitor.	Phase 2 (Non-small cell lung carcinoma (NSCLC)) Phase 3 (Breast carcinoma)
	Rezivertinib		Egfr T790M inhibitor.	Phase 3 (Non-small cell lung carcinoma (NSCLC))
	SKLB1028		Egfr/Flt3/c-Abl inhibitor.	Phase 2 (Acute myeloid leukemia (AML))
	Sunvozertinib		Bispecific anti-Egfr/Her2 monoclonal antibody.	Phase 2 (Non-small cell lung carcinoma (NSCLC)) Phase 2 (Non-Hodgkin lymphoma (NHL))
	TAS2940		Egfr/Her2 kinase inhibitor.	Phase 1 (Solid Tumor)
	TAS3351		Fourth generation Egfr inhibitor targeting T790M and T790M/C797S mutations.	Phase 2 (Non-small cell lung carcinoma (NSCLC))
	TAVO412		Anti-c-Met/anti-EGFR/anti-VEGF trispecific antibody.	Phase 1 (Solid Tumor)
	Varlitinib		Egfr/Her2 kinase inhibitor.	Phase 2 (Gastric carcinoma, Hepatocellular carcinoma (HCC), Pancreatic carcinoma, Cholangiocarcinoma)
	WSD0922-FU		Blood-brain barrier penetrable EGFR/EGFRvIII	Phase 1 (Non-small cell lung carcinoma (NSCLC)) Phase 1 (Glioblastoma,

Detailed Therapy Results

Alteration	Drug	Trade Name	Target	Current Status
			inhibitor.	Anaplastic astrocytoma)
	ZN-e4		Egfr T790M inhibitor.	Phase 1 (Non-small cell lung carcinoma (NSCLC))
	Zorifertinib		Egfr tyrosine kinase inhibitor.	Phase 2 (Non-small cell lung carcinoma (NSCLC))

Relevance of Detected Alterations

Alteration	Role in Disease	Effect on Drug Sensitivity	Effect on Drug Resistance
EGFR E746_A750del	The presence of an EGFR abnormality (mutation, amplification, or overexpression) can result in an overabundance or overactivity of Egfr protein, which can lead to excessive proliferation. ⁽¹⁾ .	The presence of a sensitizing EGFR mutation in a tumor is the strongest biological predictor of sensitivity to an Egfr tyrosine kinase inhibitor (TKI). Compared with conventional chemotherapy, Egfr TKIs have been shown to improve progression-free survival in non-small cell lung cancer patients whose tumors harbor EGFR mutations. ⁽²⁻⁵⁾ . The Egfr TKIs erlotinib, afatinib, gefitinib, osimertinib, and dacomitinib, as well as the combination of amivantamab plus lazertinib, have been approved by the FDA for the treatment of non-small cell lung cancer (NSCLC) with exon 19 deletion or L858R EGFR mutations; osimertinib has additionally been approved for the treatment of NSCLC with EGFR T790M. ^(2,5-14) . Afatinib has additionally been FDA-approved for the treatment of NSCLC with S768I, L861Q, and/or G719X mutations. ⁽¹⁵⁾ . The combination of erlotinib and ramucirumab as well as osimertinib plus platinum-based chemotherapy have been FDA-approved for the treatment of metastatic NSCLC patients with tumors harboring an EGFR exon 19 deletion or the exon 21 L858R mutation. ⁽¹⁶⁻¹⁸⁾ . Amivantamab in combination with carboplatin and pemetrexed has been FDA-approved for the treatment of adult patients with locally advanced or metastatic NSCLC harboring EGFR Exon 19 deletions or Exon 21 L858R substitution mutations whose disease has progressed on or after treatment with an EGFR TKI. ^(12,19) . Amivantamab has also been approved by the FDA for NSCLC patients with EGFR exon 20 insertions, whose disease has progressed on or after platinum-based chemotherapy and as frontline therapy in combination with carboplatin and pemetrexed. The accelerated FDA approval of mobocertinib for NSCLC patients with EGFR exon 20 insertions has been withdrawn due to lack of progression-free survival benefit in the confirmatory Phase 3 trial. ⁽²⁰⁻²³⁾ . Studies have reported non-squamous NSCLC patients with metastatic disease and tumors harboring an EGFR exon 19 deletion or L858R mutation to be sensitive to osimertinib, erlotinib, afatinib, gefitinib, dacomitinib, and the combination of erlotinib plus ramucirumab. ^(2,5-7,11,16,24) . Less	Some patients with EGFR-mutant NSCLC exhibit resistance to Egfr inhibition; resistance has been associated with insertions in EGFR exon 20, the T790M mutation in EGFR, and amplification of either MET or ERBB2. ⁽²⁶⁻³⁰⁾ . Third generation irreversible Egfr TKIs that target the EGFR T790M mutation have shown efficacy in T790M-mutant NSCLC, including osimertinib, which has received approval by the FDA, EMA, and PMDA for the treatment of EGFR T790M-mutant metastatic NSCLC. ⁽³¹⁻³⁵⁾ . Several studies have reported that resistance to Egfr TKIs in NSCLC is mediated by the transformation of NSCLC cell types to those of SCLC with neuroendocrine features. ⁽³⁶⁻³⁹⁾ .

Relevance of Detected Alterations

Alteration	Role in Disease	Effect on Drug Sensitivity	Effect on Drug Resistance
		common activating EGFR mutations have variable sensitivity to EGFR tyrosine kinase inhibitors. ⁽²⁵⁾	
TP53 P151S	Loss of tumor suppressor p53, which is encoded by the TP53 gene, is common in aggressive advanced cancers. ⁽⁴⁰⁾ . Carriers of a germline mutation in TP53 have Li-Fraumeni Syndrome, an inherited cancer syndrome resulting in multiple tumors in early adulthood, including breast cancer, brain tumors, and leukemias. ⁽⁴¹⁻⁴³⁾ . Expression of p53 in normal cells is low; however, TP53 alterations, including those that result in loss of p53 tumor suppressor function, may lead to stabilization and increased expression of p53, particularly in the nucleus, and several studies have shown that it may have oncogenic gain-of-function effects. ⁽⁴⁴⁻⁴⁸⁾ . TP53 alterations are believed to be early events in NSCLC, preceding lymph node metastasis. ⁽⁴⁹⁾ . TP53 mutation and expression of p53 have been correlated with the lung squamous cell carcinoma subtype, and p53 expression in lung squamous cell carcinoma has also been associated with disease stage and higher grade tumors. ⁽⁵⁰⁻⁵³⁾ . TP53 mutation has been associated with PD-L1 expression and T-cell infiltration in lung adenocarcinoma samples. ⁽⁵⁴⁻⁵⁸⁾ . TP53 mutations have been significantly associated with the development of distant metastases after diagnosis in early-stage NSCLC in a cohort of 759 patients. ⁽⁵⁹⁾ .	At present, there are no approved therapies targeting TP53 alterations, despite their high prevalence in cancer. Therapeutic approaches under investigation include gene therapy for TP53 and (dendritic cell-based) TP53 vaccines. ⁽⁶⁰⁻⁶²⁾ . Inhibition of components of the DNA damage checkpoint, including Wee1, has been reported to enhance the activity of DNA-damaging agents in preclinical cancer models with deficiency of p53 function. ⁽⁶³⁻⁶⁵⁾ . Clinical trials of the Wee1 inhibitor adavosertib (MK-1775) are currently underway for patients with solid tumors and hematologic malignancies. Studies have reported Aurora kinase A to be activated in cells harboring TP53 mutation, and Aurora kinase A and B inhibitors have been reported to activate wild-type p53 in cellular assays; thus, tumors retaining a wild-type TP53 allele may benefit from Aurora kinase inhibitors. ⁽⁶⁶⁻⁷¹⁾ .	Mutations in TP53 may increase resistance to ionizing radiation therapy. ^(72,73) .

References

1. Ciardiello F, Tortora G "EGFR antagonists in cancer treatment." *The New England journal of medicine*(2008): 1160-74
2. Mok T, Wu Y, Thongprasert S, Yang C, Chu D, Saijo N, Sunpaweravong P, Han B, Margono B, Ichinose Y, Nishiwaki Y, Ohe Y, Yang J, Chewaskulyong B, Jiang H, Duffield E, Watkins C, Armour A, Fukuoka M "Gefitinib or carboplatin-paclitaxel in pulmonary adenocarcinoma." *The New England journal of medicine*(2009): 947-57
3. Rosell R, Moran T, Queralt C, Porta R, Cardenal F, Camps C, Majem M, Lopez-Vivanco G, Isla D, Provencio M, Insa A, Massuti B, Gonzalez-Larriba J, Paz-Ares L, Bover I, Garcia-Campelo R, Moreno M, Catot S, Rolfo C, Reguart N, Palmero R, Sánchez J, Bastus R, Mayo C, Bertran-Alamillo J, Molina M, Sanchez J, Taron M "Screening for epidermal growth factor receptor mutations in lung cancer." *The New England journal of medicine*(2009): 958-67
4. Tsao M, Sakurada A, Cutz J, Zhu C, Kamel-Reid S, Squire J, Lorimer I, Zhang T, Liu N, Daneshmand M, Marrano P, da Cunha Santos G, Lagarde A, Richardson F, Seymour L, Whitehead M, Ding K, Pater J, Shepherd F "Erlotinib in lung cancer - molecular and clinical predictors of outcome." *The New England journal of medicine* (2005): 133-44
5. Rosell R, Carcereny E, Gervais R, Vergnenegre A, Massuti B, Felip E, Palmero R, Garcia-Gomez R, Pallares C, Sanchez J, Porta R, Cobo M, Garrido P, Longo F, Moran T, Insa A, De Marinis F, Corre R, Bover I, Illiano A, Dansin E, de Castro J, Milella M, Reguart N, Altavilla G, Jimenez U, Provencio M, Moreno M, Terrasa J, Muñoz-Langa J, Valdivia J, Isla D, Domine M, Molinier O, Mazieres J, Baize N, Garcia-Campelo R, Robinet G, Rodriguez-Abreu D, Lopez-Vivanco G, Gebbia V, Ferrera-Delgado L, Bombardieri P, Bernabe R, Bearz A, Artañ A, Cortesi E, Rolfo C, Sanchez-Ronco M, Drozdowski A, Queralt C, de Aguirre I, Ramirez J, Sanchez J, Molina M, Taron M, Paz-Ares L "Erlotinib versus standard chemotherapy as first-line treatment for European patients with advanced EGFR mutation-positive non-small-cell lung cancer (EORTC): a multicentre, open-label, randomised phase 3 trial." *The Lancet. Oncology*(2012): 239-46
6. Soria J, Ohe Y, Vansteenkiste J, Reungwetwattana T, Chewaskulyong B, Lee K, Dechaphunkul A, Imamura F, Nogami N, Kurata T, Okamoto I, Zhou C, Cho B, Cheng Y, Cho E, Voon P, Planchard D, Su W, Gray J, Lee S, Hodge R, Marotti M, Rukazekov Y, Ramalingam S "Osimertinib in Untreated EGFR-Mutated Advanced Non-Small-Cell Lung Cancer." *The New England journal of medicine*(2018): 113-125
7. Wu Y, Cheng Y, Zhou X, Lee K, Nakagawa K, Niho S, Tsui F, Linke R, Rosell R, Corral J, Migliorino M, Pluzanski A, Sbar E, Wang T, White J, Nadanaciva S, Sandin R, Mok T "Dacomitinib versus gefitinib as first-line treatment for patients with EGFR-mutation-positive non-small-cell lung cancer (ARCHER 1050): a randomised, open-label, phase 3 trial." *The Lancet. Oncology*(2017): 1454-1466
8. Mok T, Cheng Y, Zhou X, Lee K, Nakagawa K, Niho S, Lee M, Linke R, Rosell R, Corral J, Migliorino M, Pluzanski A, Sbar E, Wang T, White J, Wu Y "Improvement in Overall Survival in a Randomized Study That Compared Dacomitinib With Gefitinib in Patients With Advanced Non-Small-Cell Lung Cancer and EGFR-Activating Mutations." *Journal of clinical oncology : official journal of the American Society of Clinical Oncology*(2018): 2244-2250
9. Shepherd F, Rodrigues Pereira J, Ciuleanu T, Tan E, Hirsh V, Thongprasert S, Campos D, Maoleekoonpiroj S, Smylie M, Martins R, van Kooten M, Dediu M, Findlay B, Tu D, Johnston D, Bezjak A, Clark G, Santabárbara P, Seymour L "Erlotinib in previously treated non-small-cell lung cancer." *The New England journal of medicine*(2005): 123-32
10. Sequist L, Yang J, Yamamoto N, O'Byrne K, Hirsh V, Mok T, Geater S, Orlov S, Tsai C, Boyer M, Su W, Bannouna J, Kato T, Gorbunova V, Lee K, Shah R, Massey D, Zazulina V, Shahidi M, Schuler M "Phase III study of afatinib or cisplatin plus pemetrexed in patients with metastatic lung adenocarcinoma with EGFR mutations." *Journal of clinical oncology : official journal of the American Society of Clinical Oncology*(2013): 3327-34
11. Douillard J, Ostoros G, Cobo M, Ciuleanu T, McCormack R, Webster A, Milenkova T "First-line gefitinib in Caucasian EGFR mutation-positive NSCLC patients: a phase-IV, open-label, single-arm study." *British journal of cancer*(2014): 55-62
12. Passaro A, Wang J, Wang Y, Lee S, Melosky B, Shih J, Wang J, Azuma K, Juan-Vidal O, Cobo M, Felip E, Girard N, Cortot A, Califano R, Cappuzzo F, Owen S, Popat S, Tan J, Salinas J, Tomasini P, Gentzler R, William W, Reckamp K, Takahashi T, Ganguly S, Kowalski D, Bearz A, MacKean M, Barala P, Bourla A, Girvin A, Greger J, Millington D, Withelder M, Xie J, Sun T, Shah S, Diorio B, Knoblauch R, Baum J, Campelo R, Cho B "Amivantamab plus chemotherapy with and without lazertinib in EGFR-mutant advanced NSCLC after disease progression on osimertinib: primary results from the phase III MARIPOSA-2 study." *Annals of oncology : official journal of the European Society for Medical Oncology*(2024): 77-90
13. "A Phase III, Randomized, Double-blind, Placebo-controlled, Multicenter, International Study of Osimertinib as Maintenance Therapy in Patients With Locally Advanced, Unresectable EGFR Mutation-positive Non-Small Cell Lung Cancer (Stage III) Whose Disease Has Not Progressed Following Definitive Platinum-based Chemoradiation Therapy (LAURA)." (2024)
14. Lu S, Kato T, Dong X, Ahn M, Quang L, Soparattanapisarn N, Inoue T, Wang C, Huang M, Yang J, Cobo M, Özgüroğlu M, Casarini I, Khiem D, Sriuranpong V, Cronemberger E, Takahashi T, Runglodvatana Y, Chen M, Huang X, Grainger E, Ghiorgiu D, van der Gronde T, Ramalingam S "Osimertinib after Chemoradiotherapy in Stage III EGFR-Mutated NSCLC." *The New England journal of medicine*(2024): 585-597
15. Yang J, Sequist L, Geater S, Tsai C, Mok T, Schuler M, Yamamoto N, Yu C, Ou S, Zhou C, Massey D, Zazulina V, Wu Y "Clinical activity of afatinib in patients with advanced non-small-cell lung cancer harbouring uncommon EGFR mutations: a combined post-hoc analysis of LUX-Lung 2, LUX-Lung 3, and LUX-Lung 6." *The Lancet. Oncology*(2015): 830-8
16. Nakagawa K, Garon E, Seto T, Nishio M, Ponce Aix S, Paz-Ares L, Chiu C, Park K, Novello S, Nadal E, Imamura F, Yoh K, Shih J, Au K, Moro-Sibilot D, Enatsu S, Zimmermann A, Frimodt-Moller B, Visseren-Grul C, Reck M "Ramucicirumab plus erlotinib in patients with untreated, EGFR-mutated, advanced non-small-cell lung cancer (RELAY): a randomised, double-blind, placebo-controlled, phase 3 trial." *The Lancet. Oncology*(2019): 1655-1669
17. Planchard D, Jänne P, Cheng Y, Yang J, Yanagitani N, Kim S, Sugawara S, Yu Y, Fan Y, Geater S, Laktionov K, Lee C, Valdiviezo N, Ahmed S, Maurel J, Andrasina I, Goldman J, Ghiorgiu D, Rukazekov Y, Todd A, Kobayashi K "Osimertinib with or without Chemotherapy in EGFR-Mutated Advanced NSCLC." *The New England journal of medicine*(2023): 1935-1948
18. "A Phase III, Open-label, Randomized Study of Osimertinib With or Without Platinum Plus Pemetrexed Chemo, as First-line Treatment in Patients With Epidermal Growth Factor Receptor (EGFR) Mutation Positive, Locally Advanced or Metastatic Non-small Cell Lung Cancer (FLAURA2)." (2024)
19. "A Phase 3, Open-Label, Randomized Study of Amivantamab and Lazertinib in Combination With Platinum-Based Chemotherapy Compared With Platinum-Based Chemotherapy in Patients With EGFR-Mutated Locally Advanced or Metastatic Non-Small Cell Lung Cancer After Osimertinib Failure" (2024)
20. Cho B, Lee K, Cho E, et al. "Amivantamab (JNJ-61186372), an EGFR-MET bispecific antibody, in combination with lazertinib, a 3rd-generation tyrosine kinase inhibitor (TKI), in advanced EGFR NSCLC" *Annals of Oncology*(2020): Abstract 1258O
21. Park K, John T, Kim S, et al. "Amivantamab (JNJ-61186372), an anti-EGFR-MET bispecific antibody, in patients with EGFR exon 20 insertion (exon20ins)-mutated non-small cell lung cancer (NSCLC)." *Journal of Clinical Oncology*(2020)

References

22. Riely G, Neal J, Camidge D, Spira A, Piotrowska Z, Costa D, Tsao A, Patel J, Gadgeel S, Bazhenova L, Zhu V, West H, Mekhail T, Gentzler R, Nguyen D, Vincent S, Zhang S, Lin J, Bunn V, Jin S, Li S, Jänne P "Activity and Safety of Mobocertinib (TAK-788) in Previously Treated Non-Small Cell Lung Cancer with EGFR Exon 20 Insertion Mutations from a Phase I/II Trial." *Cancer discovery*(2021): 1688-1699
23. Zhou C, Ramalingam S, Li B, et al. "Mobocertinib in NSCLC With EGFR Exon 20 Insertions: Results From EXCLAIM and Pooled Platinum-Pretreated Patient Populations" *Journal of Thoracic Oncology*(2021)
24. Yang J, Wu Y, Schuler M, Sebastian M, Popat S, Yamamoto N, Zhou C, Hu C, O'Byrne K, Feng J, Lu S, Huang Y, Geater S, Lee K, Tsai C, Gorbunova V, Hirsh V, Bennouna J, Orlov S, Mok T, Boyer M, Su W, Lee K, Kato T, Massey D, Shahidi M, Zazulina V, Sequist L "Afatinib versus cisplatin-based chemotherapy for EGFR mutation-positive lung adenocarcinoma (LUX-Lung 3 and LUX-Lung 6): analysis of overall survival data from two randomised, phase 3 trials." *The Lancet. Oncology* (2015): 141-51
25. O'Kane G, Bradbury P, Feld R, Leigh N, Liu G, Pisters K, Kamel-Reid S, Tsao M, Shepherd F "Uncommon EGFR mutations in advanced non-small cell lung cancer." *Lung cancer* (Amsterdam, Netherlands)(2017): 137-144
26. Engelman J, Zejnullahu K, Mitsudomi T, Song Y, Hyland C, Park J, Lindeman N, Gale C, Zhao X, Christensen J, Kosaka T, Holmes A, Rogers A, Cappuzzo F, Mok T, Lee C, Johnson B, Cantley L, Jänne P "MET amplification leads to gefitinib resistance in lung cancer by activating ERBB3 signaling." *Science* (New York, N.Y.)(2007): 1039-43
27. Greulich H, Chen T, Feng W, Jänne P, Alvarez J, Zappaterra M, Bulmer S, Frank D, Hahn W, Sellers W, Meyerson M "Oncogenic transformation by inhibitor-sensitive and -resistant EGFR mutants." *PLoS medicine*(2005): e313
28. Kwak E, Sordella R, Bell D, Godin-Heymann N, Okimoto R, Brannigan B, Harris P, Driscoll D, Fidias P, Lynch T, Rabindran S, McGinnis J, Wissner A, Sharma S, Isselbacher K, Settleman J, Haber D "Irreversible inhibitors of the EGF receptor may circumvent acquired resistance to gefitinib." *Proceedings of the National Academy of Sciences of the United States of America*(2005): 7665-70
29. Takezawa K, Pirazzoli V, Arcila M, Nebhan C, Song X, de Stanchina E, Ohashi K, Janjigian Y, Spitzler P, Melnick M, Riely G, Kris M, Miller V, Ladanyi M, Politi K, Pao W "HER2 amplification: a potential mechanism of acquired resistance to EGFR inhibition in EGFR-mutant lung cancers that lack the second-site EGFR T790M mutation." *Cancer discovery*(2012): 922-33
30. Yu H, Arcila M, Rekhtman N, Sima C, Zakowski M, Pao W, Kris M, Miller V, Ladanyi M, Riely G "Analysis of tumor specimens at the time of acquired resistance to EGFR-TKI therapy in 155 patients with EGFR-mutant lung cancers." *Clinical cancer research : an official journal of the American Association for Cancer Research*(2013): 2240-7
31. Jänne P, Boss D, Camidge D, Britten C, Engelman J, Garon E, Guo F, Wong S, Liang J, Letrent S, Millham R, Taylor I, Eckhardt S, Schellens J "Phase I dose-escalation study of the pan-HER inhibitor, PF299804, in patients with advanced malignant solid tumors." *Clinical cancer research : an official journal of the American Association for Cancer Research*(2011): 1131-9
32. Jänne P, Yang J, Kim D, Planchard D, Ohe Y, Ramalingam S, Ahn M, Kim S, Su W, Horn L, Haggstrom D, Felip E, Kim J, Frewer P, Cantarini M, Brown K, Dickinson P, Ghiorghiu S, Ranson M "AZD9291 in EGFR inhibitor-resistant non-small-cell lung cancer." *The New England journal of medicine*(2015): 1689-99
33. Greig S "Osimertinib: First Global Approval." *Drugs*(2016): 263-73
34. Yang J, Ahn M, Kim D, Ramalingam S, Sequist L, Su W, Kim S, Kim J, Planchard D, Felip E, Blackhall F, Haggstrom D, Yoh K, Novello S, Gold K, Hirashima T, Lin C, Mann H, Cantarini M, Ghiorghiu S, Jänne P "Osimertinib in Pretreated T790M-Positive Advanced Non-Small-Cell Lung Cancer: AURA Study Phase II Extension Component." *Journal of clinical oncology : official journal of the American Society of Clinical Oncology*(2017): 1288-1296
35. Wang S, Cang S, Liu D "Third-generation inhibitors targeting EGFR T790M mutation in advanced non-small cell lung cancer." *Journal of hematology & oncology*(2016): 34
36. Watanabe S, Sone T, Matsui T, Yamamura K, Tani M, Okazaki A, Kurokawa K, Tambo Y, Takato H, Ohkura N, Waseda Y, Katayama N, Kasahara K "Transformation to small-cell lung cancer following treatment with EGFR tyrosine kinase inhibitors in a patient with lung adenocarcinoma." *Lung cancer* (Amsterdam, Netherlands)(2013): 370-2
37. Chang Y, Kim S, Choi Y, So K, Rho J, Kim W, Lee J, Chung J, Choi C "Neuroendocrine differentiation in acquired resistance to epidermal growth factor receptor tyrosine kinase inhibitor." *Tuberculosis and respiratory diseases*(2013): 95-103
38. Popat S, Wotherspoon A, Nutting C, Gonzalez D, Nicholson A, O'Brien M "Transformation to "high grade" neuroendocrine carcinoma as an acquired drug resistance mechanism in EGFR-mutant lung adenocarcinoma." *Lung cancer* (Amsterdam, Netherlands)(2013): 1-4
39. Sequist L, Waltman B, Dias-Santagata D, Digumarthy S, Turke A, Fidias P, Bergethon K, Shaw A, Gettinger S, Cosper A, Akhavanfar S, Heist R, Temel J, Christensen J, Wain J, Lynch T, Vernovsky K, Mark E, Lanuti M, Iafrate A, Mino-Kenudson M, Engelman J "Genotypic and histological evolution of lung cancers acquiring resistance to EGFR inhibitors." *Science translational medicine*(2011): 75ra26
40. Brown C, Lain S, Verma C, Fersht A, Lane D "Awakening guardian angels: drugging the p53 pathway." *Nature reviews. Cancer*(2009): 862-73
41. Malkin D, Li F, Strong L, Fraumeni J, Nelson C, Kim D, Kassel J, Gryka M, Bischoff F, Tainsky M "Germ line p53 mutations in a familial syndrome of breast cancer, sarcomas, and other neoplasms." *Science* (New York, N.Y.)(1990): 1233-8
42. Srivastava S, Zou Z, Pirollo K, Blattner W, Chang E "Germ-line transmission of a mutated p53 gene in a cancer-prone family with Li-Fraumeni syndrome." *Nature*(1991): 747-9
43. Santibáñez-Koref M, Birch J, Hartley A, Jones P, Craft A, Eden T, Crowther D, Kelsey A, Harris M "p53 germline mutations in Li-Fraumeni syndrome." *Lancet* (London, England)(1991): 1490-1
44. Wang Y, Lin R, Tan Y, Chen J, Chen C, Wang Y "Wild-type p53 overexpression and its correlation with MDM2 and p14ARF alterations: an alternative pathway to non-small-cell lung cancer." *Journal of clinical oncology : official journal of the American Society of Clinical Oncology*(2005): 154-64
45. Koga T, Hashimoto S, Sugio K, Yoshino I, Nakagawa K, Yonemitsu Y, Sugimachi K, Sueishi K "Heterogeneous distribution of P53 immunoreactivity in human lung adenocarcinoma correlates with MDM2 protein expression, rather than with P53 gene mutation." *International journal of cancer*(2001): 232-9
46. Kato S, Han S, Liu W, Otsuka K, Shibata H, Kanamaru R, Ishioka C "Understanding the function-structure and function-mutation relationships of p53 tumor suppressor protein by high-resolution missense mutation analysis." *Proceedings of the National Academy of Sciences of the United States of America*(2003): 8424-9
47. Houben R, Hesbacher S, Schmid C, Kauczok C, Flohr U, Haferkamp S, Müller C, Schrama D, Wischhusen J, Becker J "High-level expression of wild-type p53 in melanoma cells is frequently associated with inactivity in p53 reporter gene assays." *PLoS one*(2011): e22096
48. Olivier M, Petitjean A, Marcel V, Pétrel A, Mounawar M, Plymoth A, de Fromental C, Hainaut P "Recent advances in p53 research: an interdisciplinary perspective." *Cancer gene therapy*(2009): 1-12

References

49. Chang Y, Wu C, Shih J, Lee Y "Comparison of p53 and epidermal growth factor receptor gene status between primary tumors and lymph node metastases in non-small cell lung cancers." *Annals of surgical oncology*(2011): 543-50
50. Jiang R, Zhang B, Teng X, Hu P, Xu S, Zheng Z, Liu R, Tang T, Ye F "Validating a targeted next-generation sequencing assay and profiling somatic variants in Chinese non-small cell lung cancer patients." *Scientific reports*(2020): 2070
51. Mattioni M, Soddu S, Prodosmo A, Visca P, Conti S, Alessandrini G, Facciolo F, Strigari L "Prognostic role of serum p53 antibodies in lung cancer." *BMC cancer*(2015): 148
52. Bircan A, Bircan S, Kapucuoglu N, Songur N, Ozturk O, Akkaya A "Maspin, VEGF and p53 expression in small biopsies of primary advanced lung cancer and relationship with clinicopathologic parameters." *Pathology oncology research : POR*(2010): 553-61
53. Kim Y, Hammerman P, Kim J, Yoon J, Lee Y, Sun J, Wilkerson M, Pedamallu C, Cibulskis K, Yoo Y, Lawrence M, Stojanov P, Carter S, McKenna A, Stewart C, Sivachenko A, Oh I, Kim H, Choi Y, Kim K, Shim Y, Kim K, Song S, Na K, Choi Y, Hayes D, Kim J, Cho S, Kim Y, Ahn J, Ahn M, Getz G, Meyerson M, Park K "Integrative and comparative genomic analysis of lung squamous cell carcinomas in East Asian patients." *Journal of clinical oncology : official journal of the American Society of Clinical Oncology*(2014): 121-8
54. Dong Z, Zhong W, Zhang X, Su J, Xie Z, Liu S, Tu H, Chen H, Sun Y, Zhou Q, Yang J, Yang X, Lin J, Yan H, Zhai H, Yan L, Liao R, Wu S, Wu Y "Potential Predictive Value of TP53 and KRAS Mutation Status for Response to PD-1 Blockade Immunotherapy in Lung Adenocarcinoma." *Clinical cancer research : an official journal of the American Association for Cancer Research*(2017): 3012-3024
55. Scheel A, Ansén S, Schultheis A, Scheffler M, Fischer R, Michels S, Hellmich M, George J, Zander T, Brockmann M, Stoelben E, Groen H, Timens W, Perner S, von Bergwelt-Baildon M, Büttner R, Wolf J "PD-L1 expression in non-small cell lung cancer: Correlations with genetic alterations." *Oncoimmunology*(2016): e1131379
56. Albitar M, Sudarsanam S, Ma W, Jiang S, Chen W, Funari V, Blocker F, Agersborg S "Correlation of MET gene amplification and TP53 mutation with PD-L1 expression in non-small cell lung cancer." *Oncotarget*(2018): 13682-13693
57. Mansuet-Lupo A, Alifano M, Pécuchet N, Biton J, Becht E, Goc J, Germain C, Ouakrim H, Régnard J, Cremer I, Laurent-Puig P, Dieu-Nosjean M, Blons H, Damotte D "Intratumoral Immune Cell Densities Are Associated with Lung Adenocarcinoma Gene Alterations." *American journal of respiratory and critical care medicine*(2016): 1403-1412
58. Kadara H, Choi M, Zhang J, Parra E, Rodriguez-Canales J, Gaffney S, Zhao Z, Behrens C, Fujimoto J, Chow C, Yoo Y, Kalhor N, Moran C, Rimm D, Swisher S, Gibbons D, Heymach J, Kaftan E, Townsend J, Lynch T, Schlessinger J, Lee J, Lifton R, Wistuba I, Herbst R "Whole-exome sequencing and immune profiling of early-stage lung adenocarcinoma with fully annotated clinical follow-up." *Annals of oncology : official journal of the European Society for Medical Oncology*(2017): 75-82
59. Van Egeren D, Kohli K, Warner J, Bedard P, Riely G, Lepisto E, Schrag D, LeNoue-Newton M, Catalano P, Kehl K, Michor F "Genomic analysis of early-stage lung cancer reveals a role for TP53 mutations in distant metastasis." *Scientific reports*(2022): 19055
60. Schuler P, Harasymczuk M, Visus C, Deleo A, Trivedi S, Lei Y, Argiris A, Gooding W, Butterfield L, Whiteside T, Ferris R "Phase I dendritic cell p53 peptide vaccine for head and neck cancer." *Clinical cancer research : an official journal of the American Association for Cancer Research*(2014): 2433-44
61. Vermeij R, Leffers N, van der Burg S, Melief C, Daemen T, Nijman H "Immunological and clinical effects of vaccines targeting p53-overexpressing malignancies." *Journal of biomedicine & biotechnology*(2011): 702146
62. Saito H, Ando S, Morishita N, Lee K, Dator D, Dy D, Shigemura K, Adhim Z, Nibu K, Fujisawa M, Shirakawa T "A combined lymphokine-activated killer (LAK) cell immunotherapy and adenovirus-p53 gene therapy for head and neck squamous cell carcinoma." *Anticancer research*(2014): 3365-70
63. Ma C, Janetka J, Piwnicka-Worms H "Death by releasing the breaks: CHK1 inhibitors as cancer therapeutics." *Trends in molecular medicine*(2011): 88-96
64. Hirai H, Arai T, Okada M, Nishibata T, Kobayashi M, Sakai N, Imagaki K, Ohtani J, Sakai T, Yoshizumi T, Mizuarai S, Iwasawa Y, Kotani H "MK-1775, a small molecule Wee1 inhibitor, enhances anti-tumor efficacy of various DNA-damaging agents, including 5-fluorouracil." *Cancer biology & therapy*(2010): 514-22
65. Bridges K, Hirai H, Buser C, Brooks C, Liu H, Buchholz T, Molkenhine J, Mason K, Meyn R "MK-1775, a novel Wee1 kinase inhibitor, radiosensitizes p53-defective human tumor cells." *Clinical cancer research : an official journal of the American Association for Cancer Research*(2011): 5638-48
66. Vilgelm A, Pawlikowski J, Liu Y, Hawkins O, Davis T, Smith J, Weller K, Horton L, McClain C, Ayers G, Turner D, Essaka D, Stewart C, Sosman J, Kelley M, Ecsedy J, Johnston J, Richmond A "Mdm2 and aurora kinase inhibitors synergize to block melanoma growth by driving apoptosis and immune clearance of tumor cells." *Cancer research*(2015): 181-93
67. Li Z, Sun Y, Chen X, Squires J, Nowroozizadeh B, Liang C, Huang J "p53 Mutation Directs AURKA Overexpression via miR-25 and FBXW7 in Prostatic Small Cell Neuroendocrine Carcinoma." *Molecular cancer research : MCR*(2015): 584-91
68. Katayama H, Sen S "Functional significance of Aurora kinase A regulatory interactions with p53-ERα complex in human breast cancer cells." *Hormones & cancer*(2011): 117-24
69. Tentler J, Ionkina A, Tan A, Newton T, Pitts T, Glogowska M, Kabos P, Sartorius C, Sullivan K, Espinosa J, Eckhardt S, Diamond J "p53 Family Members Regulate Phenotypic Response to Aurora Kinase A Inhibition in Triple-Negative Breast Cancer." *Molecular cancer therapeutics*(2015): 1117-29
70. Gully C, Velazquez-Torres G, Shin J, Fuentes-Mattei E, Wang E, Carlock C, Chen J, Rothenberg D, Adams H, Choi H, Guma S, Phan L, Chou P, Su C, Zhang F, Chen J, Yang T, Yeung S, Lee M "Aurora B kinase phosphorylates and instigates degradation of p53." *Proceedings of the National Academy of Sciences of the United States of America*(2012): E1513-22
71. Marxer M, Ma H, Man W, Poon R "p53 deficiency enhances mitotic arrest and slippage induced by pharmacological inhibition of Aurora kinases." *Oncogene*(2014): 3550-60
72. El-Deiry W "The role of p53 in chemosensitivity and radiosensitivity." *Oncogene*(2003): 7486-95
73. Miyasaka A, Oda K, Ikeda Y, Sone K, Fukuda T, Inaba K, Makii C, Enomoto A, Hosoya N, Tanikawa M, Uehara Y, Arimoto T, Kuramoto H, Wada-Hiraike O, Miyagawa K, Yano T, Kawana K, Osuga Y, Fujii T "PI3K/mTOR pathway inhibition overcomes radioresistance via suppression of the HIF1-α/VEGF pathway in endometrial cancer." *Gynecologic oncology*(2015): 174-80