



Sample Information

Patient Name: 向駿  
Gender: Male  
ID No.: F103921860  
History No.: 32259522  
Age: 70

Ordering Doctor: DOC4205A 柯博伸  
Ordering REQ.: H49EPA5  
Signing in Date: 2024/02/23

Path No.: M113-00054  
MP No.: MY24007  
Assay: Oncomine Myeloid Assay  
Sample Type: Bone Marrow  
Bone Marrow Aspirating Date: 2024/02/22

Reporting Doctor: DOC5444B 楊靜芬 (Phone: 8#5444)

Note:

Sample Cancer Type: Acute Myeloid Leukemia

Table of Contents	Page	Report Highlights
Variants (Exclude variant in Taiwan BioBank with >1% allele frequency)	2	2 Relevant Biomarkers
Biomarker Descriptions	2	11 Therapies Available
Relevant Therapy Summary	5	0 Clinical Trials
Relevant Therapy Details	6	
Diagnostic Details	13	

Relevant Biomarkers

Tier	Genomic Alteration	Relevant Therapies (In this cancer type)	Relevant Therapies (In other cancer type)	Clinical Trials
IA	IDH2 p.(R140Q) c.419G>A isocitrate dehydrogenase (NADP(+)) 2 Allele Frequency: 37.80%	enasidenib <sup>1</sup> azacitidine decitabine enasidenib + chemotherapy venetoclax + chemotherapy	None	0

Public data sources included in relevant therapies: FDA<sup>1</sup>, NCCN, EMA<sup>2</sup>, ESMO  
Public data sources included in diagnostic significance: NCCN, ESMO

Relevant Biomarkers (continued)

Tier	Genomic Alteration	Relevant Therapies (In this cancer type)	Relevant Therapies (In other cancer type)	Clinical Trials
IA	<i>NPM1</i> p.(W288Cfs*12) <i>c.863_864insTCTG</i> nucleophosmin 1 Allele Frequency: 40.84%	allogeneic stem cells cytarabine cytarabine + daunorubicin cytarabine + idarubicin cytarabine + mitoxantrone gemtuzumab ozogamicin + chemotherapy	None	0
Diagnostic significance: Acute Myeloid Leukemia				

Public data sources included in relevant therapies: FDA1, NCCN, EMA2, ESMO  
Public data sources included in diagnostic significance: NCCN, ESMO

Prevalent cancer biomarkers without relevant evidence based on included data sources  
*DNMT3A* p.(M682Cfs\*23) *c.2043delC*, *NRAS* p.(G12D) *c.35G>A*, *PTPN11* p.(G503R) *c.1507G>A*, *PTPN11* p.(Q510H) *c.1530G>C*, *TET2* p.(E1405\*) *c.4213G>T*

Variants (Exclude variant in Taiwan BioBank with >1% allele frequency)

DNA Sequence Variants								
Gene	Amino Acid Change	Coding	Variant ID	Locus	Allele Frequency	Transcript	Variant Effect	Coverage
NRAS	p.(G12D)	c.35G>A	COSM564	chr1:115258747	5.81%	NM_002524.5	missense	1995
DNMT3A	p.(M682Cfs*23)	c.2043delC	.	chr2:25464469	44.61%	NM_022552.4	frameshift Deletion	1975
TET2	p.(E1405*)	c.4213G>T	.	chr4:106193751	48.77%	NM_001127208.2	nonsense	1999
NPM1	p.(W288Cfs*12)	c.863_864insTCTG	COSM17559	chr5:170837547	40.84%	NM_002520.6	frameshift Insertion	1993
PTPN11	p.(G503R)	c.1507G>A	COSM14259	chr12:112926887	12.20%	NM_002834.5	missense	2000
PTPN11	p.(Q510H)	c.1530G>C	COSM1318058	chr12:112926910	28.71%	NM_002834.5	missense	1999
IDH2	p.(R140Q)	c.419G>A	COSM41590	chr15:90631934	37.80%	NM_002168.4	missense	2000
ZRSR2	p.(N382K)	c.1146C>A	.	chrX:15841062	99.55%	NM_005089.3	missense	1998

Biomarker Descriptions

PTPN11 p.(G503R) c.1507G>A, PTPN11 p.(Q510H) c.1530G>C

protein tyrosine phosphatase non-receptor type 11

Background: The PTPN11 gene encodes a tyrosine phosphatase non-receptor type 11 protein, and is also known as Src homology region 2 domain-containing phosphatase-2 (SHP-2)<sup>1</sup>. PTPN11 is a member of the protein tyrosine phosphatase (PTP) family that is ubiquitously expressed and regulates cellular growth, differentiation, mitotic cycle, and oncogenic transformation. PTPN11 contains two tandem N-terminal Src homology-2 domains (N-SH2 and C-SH2), a PTP catalytic domain, and uncharacterized C-terminal domain<sup>2</sup>. PTPN11 regulates various signaling processes including the RAS/RAF/MEK/ERK, PI3K/AKT/MTOR, and JAK/STAT pathways<sup>3,4</sup>. Germline mutations in PTPN11 are associated with LEOPARD syndrome and Noonan syndrome with a predisposition to juvenile myelomonocytic leukemia (JMML) or myeloproliferative neoplasms (MPN)<sup>5,6</sup>. Somatic mutations in PTPN11 are associated with JMML<sup>7,8</sup> and solid tumors such as lung, colon, and thyroid<sup>2,9</sup>

## Biomarker Descriptions (continued)

**Alterations and prevalence:** Somatic alterations in PTPN11 include mutations and amplification<sup>5,10</sup>. PTPN11 mutations occur in 6% of uterine carcinoma and 5% of acute myeloid leukemia (AML) cases<sup>11</sup>. Mutations including E76K and D61Y result in PTPN11 activation and are associated with 30% of JMML<sup>4</sup>.

**Potential relevance:** Currently, no therapies are approved for PTPN11 aberrations. Somatic mutations in PTPN11 confer drug resistance to venetoclax and azacitidine in AML<sup>12,13</sup>.

### DNMT3A p.(M682Cfs\*23) c.2043delC

*DNA methyltransferase 3 alpha*

**Background:** The DNMT3A gene encodes the DNA methyltransferase 3 alpha which functions as a de novo methyltransferase (DNMT) with equal methylation efficiency for unmethylated and hemimethylated DNA<sup>14</sup>. Methylation of DNA occurs at CpG islands, a region of DNA consisting of sequential cytosine/guanine dinucleotide pairs. CpG island methylation plays an important role in development as well as stem cell regulation. Alterations to global DNA methylation patterns are dependent on DNMTs, which are associated with cancer initiation and progression<sup>15,16</sup>.

**Alterations and prevalence:** DNMT3A mutations are observed in approximately 25% of all acute myeloid leukemia (AML) including 29-34% of AML with normal karyotype (NK-AML)<sup>17,18,19,20,21,22,23</sup>. Mutations in DNMT3A are also reported in 12-18% of myelodysplastic syndromes (MDS) as well as 4-6% of melanoma, lung adenocarcinoma, and uterine cancer<sup>6,22</sup>. The majority of mutations in DNMT3A are missense however, frameshift, nonsense, and splice site mutations have also been reported<sup>17,22</sup>. Missense mutations at R882 are most prevalent and are observed to coexist with NPM1 and FLT3 mutations<sup>24,25</sup>. The R882 mutations occur at the dimer/tetramer interface within the catalytic domain, which leads to disruption of DNMT3A tetramerization and loss of CpG methylation<sup>26,27</sup>. However, DNMT3A mutations observed in AML at positions other than R882 also contribute to pathogenesis by mechanisms that do not involve methyltransferase activity<sup>28</sup>.

**Potential relevance:** DNMT3A mutations confer shorter overall survival (OS) in patients with AML including those with NK-AML<sup>17,20,21,25</sup>. DNMT3A mutations are a useful in the diagnosis of angioimmunoblastic T-cell lymphoma (AITCL) when trying to differentiate from other peripheral T-cell lymphomas (PTCL)<sup>29</sup>.

### NPM1 p.(W288Cfs\*12) c.863\_864insTCTG

*nucleophosmin 1*

**Background:** The NPM1 gene encodes the nucleophosmin protein, a histone chaperone of the nucleophosmin/nucleoplasmin family, which also includes NPM2 and NPM3<sup>30</sup>. NPM1 functions as an oncogene and tumor suppressor, and is important in maintaining genomic stability, DNA repair, and apoptosis<sup>30,31</sup>. NPM1 has a highly conserved N-terminal region which constitutes the core domain responsible for oligomerization, an acidic domain, a nuclear localization signal, and a disorganized C-terminal region which is required for nucleolar localization<sup>30</sup>. Oligomerization of NPM1 localizes the protein in the nucleus of proliferating cells where it binds to Akt in response to growth factor stimulation and escapes proteolytic degradation by caspase activity, thereby promoting cell survival<sup>30,31</sup>. NPM1 is one of the most frequently altered genes in hematological cancers<sup>32</sup>. Most NPM1 mutations occur in the C-terminus, impacting protein folding or the nucleolar localization signal, and result in the localization of NPM1 to the cytoplasm (NPMc) instead of to the nucleus<sup>30</sup>.

**Alterations and prevalence:** NPM1 mutations are observed in 45-60% of AML with a normal karyotype (NK-AML), 28-35% of de novo acute myeloid leukemia (AML) and are frequently co-mutated with DNMT3A and/or FLT3-ITD<sup>23,33,34</sup>. NPM1 fusions are associated with distinct partner genes in acute promyelocytic leukemia (APL), anaplastic large-cell lymphoma (ALCL), AML, and myelodysplasia<sup>32</sup>. Specifically, NPM1::ALK fusion is found in 30% of all ALCL and this specific fusion is observed in 85% of ALK-positive ALCL<sup>30</sup>. The t(5;17)(q35;q21) translocation that results in NPM1::RARA fusion is observed in APL<sup>35</sup>.

**Potential relevance:** Mutation of NPM1 is recognized as a diagnostic entity for AML with NPM1 mutation by the World Health Organization (WHO)<sup>36</sup>. NPM1 mutations are associated with better outcomes, increased complete remission, improved overall survival, and favorable risk in AML<sup>23,34,37</sup>. Concurrent expression of FLT-ITD with mutant or wild-type NPM1 (when lacking adverse risk genetic lesions) confers intermediate risk in AML<sup>23,37</sup>. The NPM1 frameshift mutation W288fs\*12 is associated with poor prognosis in myelodysplastic syndrome (MDS)<sup>6</sup>. The ALK::NPM1 fusion and translocation t(2;5)(p23;q35) which leads to an ALK::NPM1 fusion is diagnostic of ALK-positive anaplastic large cell lymphoma<sup>29,38</sup>.

### TET2 p.(E1405\*) c.4213G>T

*tet methylcytosine dioxygenase 2*

**Background:** TET2 encodes the tet methylcytosine dioxygenase 2 protein and belongs to a family of ten-eleven translocation (TET) proteins that also includes TET1 and TET3<sup>39</sup>. TET2 is involved in DNA methylation, specifically in the conversion of 5-methylcytosine to

## Biomarker Descriptions (continued)

5-hydroxymethylcytosine<sup>40,41</sup>. The TET proteins contain a C-terminal core catalytic domain that contains a cysteine-rich domain and a double stranded  $\beta$ -helix domain (DSBH)<sup>42</sup>. TET2 is a tumor suppressor gene. Loss of function mutations in TET2 are associated with loss of catalytic activity and transformation to hematological malignancies<sup>39,40,41</sup>

**Alterations and prevalence:** Somatic TET2 mutations, including nonsense, frameshift, splice site, and missense, are observed in 20-25% of myelodysplastic syndrome (MDS) associated diseases, including 40%-60% chronic myelomonocytic leukemia (CMML)<sup>6</sup>. TET2 mutations at H1881 and R1896 are frequently observed in myeloid malignancies<sup>40,43</sup>. TET2 mutations are also observed in 9% of uterine, 8% of melanoma and acute myeloid leukemia (AML), as well as 6% of diffuse large B-cell lymphoma (DLBCL).

**Potential relevance:** The presence of TET2 mutations may be used as one of the major diagnostic criteria in pre-primary myelofibrosis (pre-PMF) and overt PMF in the absence of JAK2/CALR/MPL mutations<sup>44</sup>. TET2 mutations are associated with poor prognosis in PMF and increased rate of transformation to leukemia<sup>44,45</sup>

### IDH2 p.(R140Q) c.419G>A

*isocitrate dehydrogenase (NADP(+)) 2*

**Background:** The IDH1 and IDH2 genes encode homologous isocitrate dehydrogenase enzymes that catalyze the conversion of isocitrate to  $\alpha$ -ketoglutarate ( $\alpha$ -KG)<sup>46</sup>. The IDH1 gene encodes the NADP+ dependent cytoplasmic isocitrate dehydrogenase enzyme; IDH2 encodes the mitochondrial isoform.

**Alterations and prevalence:** Recurrent somatic mutations in IDH1 and IDH2 are mutually exclusive and observed in several malignancies including glioma, chondrosarcoma, intrahepatic cholangiocarcinoma, acute myeloid leukemia (AML), and myelodysplastic syndrome (MDS)<sup>47</sup>. Recurrent IDH2 variants include predominately R140Q and R172K plus other substitutions at lower frequencies. These gain of function variants confer neomorphic enzyme activity<sup>48</sup>. Although wild-type enzymatic activity is ablated, recurrent IDH2 variants catalyze the conversion of  $\alpha$ -KG to D-2-hydroxyglutarate, an oncometabolite with diverse effects on cellular metabolism, epigenetic regulation, redox states, and DNA repair<sup>46,49</sup>. Recurrent IDH2 mutations are present in 10-20% of patients with AML and 5% of patients with MDS<sup>50,51,52</sup>.

**Potential relevance:** Enasidenib<sup>53</sup> is FDA approved (2017) for the treatment of AML patients with IDH2 R140G/L/Q/W and R172G/K/M/S/W mutations. The FDA also granted fast track designation (2023) to the small-molecule IDH1 and IDH2 selective inhibitor, vorasidenib, for IDH-mutant (Grade 2) gliomas<sup>54</sup>. In AML, acquired resistance to enasidenib has been associated with the emergence of Q316E or I319M mutations<sup>55</sup>. IDH2 R172 and R140Q variants are associated with poor prognosis in MDS but have been shown to confer improved prognosis in lower grade gliomas<sup>6,56,57</sup>. Additionally, IDH2 mutations are associated with inferior overall survival in polycythemia vera (PV) and essential thrombocythemia (ET) as well as inferior leukemia-free survival in primary myelofibrosis (PMF)<sup>44</sup>. Mutations in IDH2 are diagnostic of astrocytoma IDH-mutant and oligodendroglioma IDH-mutant and 1p/19q-codeleted subtypes of central nervous system (CNS) tumors<sup>58</sup>

### NRAS p.(G12D) c.35G>A

*NRAS proto-oncogene, GTPase*

**Background:** The NRAS proto-oncogene encodes a GTPase that functions in signal transduction and is a member of the RAS superfamily which also includes KRAS and HRAS. RAS proteins mediate the transmission of growth signals from the cell surface to the nucleus via the PI3K/AKT/MTOR and RAS/RAF/MEK/ERK pathways, which regulate cell division, differentiation, and survival<sup>59,60,61</sup>.

**Alterations and prevalence:** Recurrent mutations in RAS oncogenes cause constitutive activation and are found in 20-30% of cancers. NRAS mutations are particularly common in melanomas (up to 25%) and are observed at frequencies of 5-10% in acute myeloid leukemia, colorectal, and thyroid cancers<sup>22,62</sup>. The majority of NRAS mutations consist of point mutations at G12, G13, and Q61<sup>22,63</sup>. Mutations at A59, K117, and A146 have also been observed but are less frequent<sup>11,64</sup>.

**Potential relevance:** Currently, no therapies are approved for NRAS aberrations. The EGFR antagonists, cetuximab<sup>65</sup> and panitumumab<sup>66</sup>, are contraindicated for treatment of colorectal cancer patients with NRAS mutations in exon 2 (codons 12 and 13), exon 3 (codons 59 and 61), and exon 4 (codons 117 and 146)<sup>64</sup>. In 2022, the FDA has granted fast track designation to the pan-RAF inhibitor, KIN-2787<sup>67</sup>, for the treatment of NRAS-mutant metastatic or unresectable melanoma. In 2023, the FDA has granted fast track designation to the pan-RAF inhibitor, naporafenib, in combination with trametinib<sup>68</sup> for NRAS-mutated unresectable or metastatic melanoma. NRAS mutations are associated with poor prognosis in patients with low-risk myelodysplastic syndrome<sup>6</sup> as well as melanoma<sup>69</sup>. In a phase III clinical trial in patients with advanced NRAS-mutant melanoma, binimetinib improved progression free survival (PFS) relative to dacarbazine with median PFS of 2.8 and 1.5 months, respectively<sup>70</sup>.

## Relevant Therapy Summary

☒ In this cancer type    
 ☐ In other cancer type    
 ☒ In this cancer type and other cancer types    
 ✕ No evidence

### IDH2 p.(R140Q) c.419G>A

Relevant Therapy	FDA	NCCN	EMA	ESMO	Clinical Trials*
enasidenib	●	●	✕	●	✕
azacitidine	✕	●	✕	✕	✕
decitabine	✕	●	✕	✕	✕
enasidenib + azacitidine	✕	●	✕	✕	✕
venetoclax + azacitidine	✕	●	✕	✕	✕
venetoclax + cytarabine	✕	●	✕	✕	✕
venetoclax + decitabine	✕	●	✕	✕	✕

### NPM1 p.(W288Cfs\*12) c.863\_864insTCTG

Relevant Therapy	FDA	NCCN	EMA	ESMO	Clinical Trials*
Allogeneic hematopoietic stem cell transplantation	✕	●	✕	✕	✕
cytarabine	✕	●	✕	✕	✕
cytarabine + daunorubicin	✕	●	✕	✕	✕
cytarabine + idarubicin	✕	●	✕	✕	✕
cytarabine + mitoxantrone	✕	●	✕	✕	✕
gemtuzumab ozogamicin + cytarabine	✕	●	✕	✕	✕
gemtuzumab ozogamicin + cytarabine + daunorubicin	✕	●	✕	✕	✕
gemtuzumab ozogamicin + cytarabine + fludarabine + idarubicin + filgrastim	✕	●	✕	✕	✕

## Relevant Therapy Details

### Current FDA Information

☒ In this cancer type    ☐ In other cancer type    ☐ In this cancer type and other cancer types

FDA information is current as of 2024-01-17. For the most up-to-date information, search [www.fda.gov](https://www.fda.gov).

#### IDH2 p.(R140Q) c.419G>A

##### ☒ enasidenib

**Cancer type:** Acute Myeloid Leukemia

**Label as of:** 2023-12-21

**Variant class:** IDH2 R140Q mutation

**Indications and usage:**

IDHIFA® is an isocitrate dehydrogenase-2 inhibitor indicated for the treatment of adult patients with relapsed or refractory acute myeloid leukemia (AML) with an isocitrate dehydrogenase-2 (IDH2) mutation as detected by an FDA-approved test.

**Reference:**

[https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2023/209606s006lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2023/209606s006lbl.pdf)

## Current NCCN Information

- ☒ In this cancer type    ☐ In other cancer type    ☒ In this cancer type and other cancer types

NCCN information is current as of 2024-01-02. To view the most recent and complete version of the guideline, go online to NCCN.org.

For NCCN International Adaptations & Translations, search [www.nccn.org/global/what-we-do/international-adaptations](http://www.nccn.org/global/what-we-do/international-adaptations).

Some variant specific evidence in this report may be associated with a broader set of alterations from the NCCN Guidelines. Specific variants listed in this report were sourced from approved therapies or scientific literature. These therapeutic options are appropriate for certain population segments with cancer. Refer to the NCCN Guidelines® for full recommendation.

All guidelines cited below are referenced with permission from the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®) National Comprehensive Cancer Network, Inc. 2023. All rights reserved. NCCN makes no warranties regarding their content.

### IDH2 p.(R140Q) c.419G>A

#### ● venetoclax + azacitidine

Cancer type: Acute Myeloid Leukemia

Variant class: IDH2 R140 mutation

NCCN Recommendation category: 1

Population segment (Line of therapy):

- (Induction therapy); Preferred intervention

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 6.2023]

#### ● enasidenib

Cancer type: Acute Myeloid Leukemia

Variant class: IDH2 R140 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- Relapsed, Refractory (Line of therapy not specified)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 6.2023]

#### ● venetoclax + decitabine

Cancer type: Acute Myeloid Leukemia

Variant class: IDH2 R140 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Induction therapy); Preferred intervention

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 6.2023]

## IDH2 p.(R140Q) c.419G>A (continued)

### ● azacitidine

Cancer type: Acute Myeloid Leukemia

Variant class: IDH2 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Induction therapy); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 6.2023]

### ● decitabine

Cancer type: Acute Myeloid Leukemia

Variant class: IDH2 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Induction therapy); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 6.2023]

### ● enasidenib

Cancer type: Acute Myeloid Leukemia

Variant class: IDH2 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Induction therapy); Preferred intervention

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 6.2023]

### ● venetoclax + cytarabine

Cancer type: Acute Myeloid Leukemia

Variant class: IDH2 mutation

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Induction therapy); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 6.2023]

### ● enasidenib + azacitidine

Cancer type: Acute Myeloid Leukemia

Variant class: IDH2 mutation

NCCN Recommendation category: 2B

Population segment (Line of therapy):

- (Induction therapy); Useful in certain circumstances

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 6.2023]



**NPM1 p.(W288Cfs\*12) c.863\_864insTCTG****● Allogeneic hematopoietic stem cell transplantation**

Cancer type: Acute Myeloid Leukemia

Variant class: NPM1 mutation

Other criteria: FLT3 ITD negative

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Consolidation therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 6.2023]

**● cytarabine**

Cancer type: Acute Myeloid Leukemia

Variant class: NPM1 mutation

Other criteria: FLT3 ITD negative

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Consolidation therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 6.2023]

**● cytarabine + daunorubicin**

Cancer type: Acute Myeloid Leukemia

Variant class: NPM1 mutation

Other criteria: FLT3 ITD negative

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Consolidation therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 6.2023]

**● cytarabine + daunorubicin**

Cancer type: Acute Myeloid Leukemia

Variant class: NPM1 mutation

Other criteria: FLT3 ITD negative

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Induction therapy); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 6.2023]

**NPM1 p.(W288Cfs\*12) c.863\_864insTCTG (continued)****● cytarabine + idarubicin**

Cancer type: Acute Myeloid Leukemia

Variant class: NPM1 mutation

Other criteria: FLT3 ITD negative

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Consolidation therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 6.2023]

**● cytarabine + idarubicin**

Cancer type: Acute Myeloid Leukemia

Variant class: NPM1 mutation

Other criteria: FLT3 ITD negative

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Induction therapy); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 6.2023]

**● cytarabine + mitoxantrone**

Cancer type: Acute Myeloid Leukemia

Variant class: NPM1 mutation

Other criteria: FLT3 ITD negative

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Consolidation therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 6.2023]

**● cytarabine + mitoxantrone**

Cancer type: Acute Myeloid Leukemia

Variant class: NPM1 mutation

Other criteria: FLT3 ITD negative

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Induction therapy); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 6.2023]

**NPM1 p.(W288Cfs\*12) c.863\_864insTCTG (continued)****● gemtuzumab ozogamicin + cytarabine**

Cancer type: Acute Myeloid Leukemia

Variant class: NPM1 mutation

Other criteria: CD33 positive, FLT3 ITD negative

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Consolidation therapy)

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 6.2023]

**● gemtuzumab ozogamicin + cytarabine + daunorubicin**

Cancer type: Acute Myeloid Leukemia

Variant class: NPM1 mutation

Other criteria: CD33 positive, FLT3 ITD negative

NCCN Recommendation category: 2A

Population segment (Line of therapy):

- (Consolidation therapy)
- (Induction therapy); Preferred intervention

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 6.2023]

**● gemtuzumab ozogamicin + cytarabine + fludarabine + idarubicin + filgrastim**

Cancer type: Acute Myeloid Leukemia

Variant class: NPM1 mutation

Other criteria: FLT3 ITD negative

NCCN Recommendation category: 2B

Population segment (Line of therapy):

- (Induction therapy); Other recommended intervention

Reference: NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 6.2023]

## Current ESMO Information

☒ In this cancer type    ☐ In other cancer type    ☒ In this cancer type and other cancer types

ESMO information is current as of 2024-01-02. For the most up-to-date information, search [www.esmo.org](http://www.esmo.org).

### IDH2 p.(R140Q) c.419G>A

#### ☒ enasidenib

Cancer type: Acute Myeloid Leukemia

Variant class: IDH2 mutation

ESMO Level of Evidence/Grade of Recommendation: IV / B

Population segment (Line of therapy):

- Relapsed, Refractory (Second-line therapy)

Reference: ESMO Clinical Practice Guidelines - ESMO-Acute Myeloblastic Leukaemia in Adult Patients [Ann Oncol (2020); 31(6): 697-712.]

## Diagnostic Details

### Current ESMO Information

ESMO information is current as of 2024-01-02. For the most up-to-date information, search [www.esmo.org](http://www.esmo.org).

**NPM1 p.(W288Cfs\*12) c.863\_864insTCTG**

**Diagnostic significance: Acute Myeloid Leukemia**

**Variant class:** NPM1 mutation

**Diagnostic notes:**

- AML with recurrent genetic abnormalities; WHO classification of AML

**Reference:** ESMO Clinical Practice Guidelines - ESMO-Acute Myeloblastic Leukaemia in Adult Patients [Ann Oncol (2020); 31(6): 697-712.]

## References

1. Tartaglia et al. Somatic PTPN11 mutations in childhood acute myeloid leukaemia. *Br. J. Haematol.* 2005 May;129(3):333-9. PMID: 15842656
2. Chan et al. The tyrosine phosphatase Shp2 (PTPN11) in cancer. *Cancer Metastasis Rev.* 2008 Jun;27(2):179-92. PMID: 18286234
3. Steelman et al. Roles of the Raf/MEK/ERK and PI3K/PTEN/Akt/mTOR pathways in controlling growth and sensitivity to therapy-implications for cancer and aging. *Aging (Albany NY)*. 2011 Mar;3(3):192-222. PMID: 21422497
4. Liu et al. Inhibition of the Gab2/PI3K/mTOR signaling ameliorates myeloid malignancy caused by Ptpn11 (Shp2) gain-of-function mutations. *Leukemia*. 2017 Jun;31(6):1415-1422. PMID: 27840422
5. Tartaglia et al. Somatic mutations in PTPN11 in juvenile myelomonocytic leukemia, myelodysplastic syndromes and acute myeloid leukemia. *Nat. Genet.* 2003 Jun;34(2):148-50. PMID: 12717436
6. NCCN Guidelines® - NCCN-Myelodysplastic Syndromes [Version 3.2023]
7. Tartaglia et al. Mutations in PTPN11, encoding the protein tyrosine phosphatase SHP-2, cause Noonan syndrome. *Nat. Genet.* 2001 Dec;29(4):465-8. PMID: 11704759
8. Tartaglia et al. Diversity and functional consequences of germline and somatic PTPN11 mutations in human disease. *Am. J. Hum. Genet.* 2006 Feb;78(2):279-90. PMID: 16358218
9. Jongmans et al. Cancer risk in patients with Noonan syndrome carrying a PTPN11 mutation. *Eur. J. Hum. Genet.* 2011 Aug;19(8):870-4. PMID: 21407260
10. Liu et al. Gain-of-function mutations of Ptpn11 (Shp2) cause aberrant mitosis and increase susceptibility to DNA damage-induced malignancies. *Proc. Natl. Acad. Sci. U.S.A.* 2016 Jan 26;113(4):984-9. PMID: 26755576
11. Cerami et al. The cBio cancer genomics portal: an open platform for exploring multidimensional cancer genomics data. *Cancer Discov.* 2012 May;2(5):401-4. PMID: 22588877
12. Chyla et al. Genetic Biomarkers Of Sensitivity and Resistance to Venetoclax Monotherapy in Patients With Relapsed Acute Myeloid Leukemia. *Am. J. Hematol.* 2018 May 17. PMID: 29770480
13. Zhang et al. Biomarkers Predicting Venetoclax Sensitivity and Strategies for Venetoclax Combination Treatment. *Blood* 2018 132:175; doi: <https://doi.org/10.1182/blood-2018-175.132:175>
14. Okano et al. Cloning and characterization of a family of novel mammalian DNA (cytosine-5) methyltransferases. *Nat Genet.* 1998 Jul;19(3):219-20. PMID: 9662389
15. Fernandez et al. A DNA methylation fingerprint of 1628 human samples. *Genome Res.* 2012 Feb;22(2):407-19. PMID: 21613409
16. Jones et al. The epigenomics of cancer. *Cell.* 2007 Feb 23;128(4):683-92. PMID: 17320506
17. Ley et al. DNMT3A mutations in acute myeloid leukemia. *N. Engl. J. Med.* 2010 Dec 16;363(25):2424-33. PMID: 21067377
18. Marková et al. Prognostic impact of DNMT3A mutations in patients with intermediate cytogenetic risk profile acute myeloid leukemia. *Eur. J. Haematol.* 2012 Feb;88(2):128-35. PMID: 21967546
19. Yang et al. DNMT3A in haematological malignancies. *Nat. Rev. Cancer.* 2015 Mar;15(3):152-65. PMID: 25693834
20. Renneville et al. Prognostic significance of DNA methyltransferase 3A mutations in cytogenetically normal acute myeloid leukemia: a study by the Acute Leukemia French Association. *Leukemia*. 2012 Jun;26(6):1247-54. PMID: 22289988
21. Marcucci et al. Age-related prognostic impact of different types of DNMT3A mutations in adults with primary cytogenetically normal acute myeloid leukemia. *J. Clin. Oncol.* 2012 Mar 1;30(7):742-50. PMID: 22291079
22. Weinstein et al. The Cancer Genome Atlas Pan-Cancer analysis project. *Nat. Genet.* 2013 Oct;45(10):1113-20. PMID: 24071849
23. NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 6.2023]
24. Kumar et al. DNMT3A (R882) mutation features and prognostic effect in acute myeloid leukemia in Coexistent with NPM1 and FLT3 mutations. *Hematol Oncol Stem Cell Ther.* 2018 Jun;11(2):82-89. PMID: 29079128
25. Thol et al. Incidence and prognostic influence of DNMT3A mutations in acute myeloid leukemia. *J. Clin. Oncol.* 2011 Jul 20;29(21):2889-96. PMID: 21670448
26. Sandoval et al. Mutations in the DNMT3A DNA methyltransferase in acute myeloid leukemia patients cause both loss and gain of function and differential regulation by protein partners. *J. Biol. Chem.* 2019 Mar 29;294(13):4898-4910. PMID: 30705090
27. Holz-Schietinger et al. Mutations in DNA methyltransferase (DNMT3A) observed in acute myeloid leukemia patients disrupt processive methylation. *J. Biol. Chem.* 2012 Sep 7;287(37):30941-51. PMID: 22722925
28. Russler-Germain et al. The R882H DNMT3A mutation associated with AML dominantly inhibits wild-type DNMT3A by blocking its ability to form active tetramers. *Cancer Cell.* 2014 Apr 14;25(4):442-54. PMID: 24656771
29. NCCN Guidelines® - NCCN-T-Cell Lymphomas [Version 1.2024]
30. Box et al. Nucleophosmin: from structure and function to disease development. 2016 Aug 24;17(1):19. PMID: 27553022

## References (continued)

31. Koike et al. Recruitment of phosphorylated NPM1 to sites of DNA damage through RNF8-dependent ubiquitin conjugates. *Cancer Res.* 2010 Sep 1;70(17):6746-56. PMID: 20713529
32. Sportoletti. How does the NPM1 mutant induce leukemia?. *Pediatr Rep.* 2011 Jun 22;3 Suppl 2:e6. PMID: 22053282
33. Hourigan et al. Accurate Medicine: Indirect Targeting of NPM1-Mutated AML. 2016 Oct;6(10):1087-1089. PMID: 27698101
34. Jain et al. Mutated NPM1 in patients with acute myeloid leukemia in remission and relapse. *Leuk. Lymphoma.* 2014 Jun;55(6):1337-44. PMID: 24004182
35. Redner et al. The t(5;17) acute promyelocytic leukemia fusion protein NPM-RAR interacts with co-repressor and co-activator proteins and exhibits both positive and negative transcriptional properties. *Blood.* 2000 Apr 15;95(8):2683-90. PMID: 10753851
36. Khoury et al. The 5th edition of the World Health Organization Classification of Haematolymphoid Tumours: Myeloid and Histiocytic/Dendritic Neoplasms. *Leukemia.* 2022 Jul;36(7):1703-1719. PMID: 35732831
37. Döhner et al. Diagnosis and management of AML in adults: 2022 recommendations from an international expert panel on behalf of the ELN. *Blood.* 2022 Sep 22;140(12):1345-1377. PMID: 35797463
38. NCCN Guidelines® - NCCN-Primary Cutaneous Lymphomas [Version 1.2024]
39. Pan et al. The TET2 interactors and their links to hematological malignancies. *IUBMB Life.* 2015 Jun;67(6):438-45. PMID: 26099018
40. Ko et al. Impaired hydroxylation of 5-methylcytosine in myeloid cancers with mutant TET2. *Nature.* 2010 Dec 9;468(7325):839-43. PMID: 21057493
41. Solary et al. The Ten-Eleven Translocation-2 (TET2) gene in hematopoiesis and hematopoietic diseases. *Leukemia.* 2014 Mar;28(3):485-96. PMID: 24220273
42. An et al. TET family dioxygenases and DNA demethylation in stem cells and cancers. *Exp. Mol. Med.* 2017 Apr 28;49(4):e323. PMID: 28450733
43. Kosmider et al. TET2 mutation is an independent favorable prognostic factor in myelodysplastic syndromes (MDSs). *Blood.* 2009 Oct 8;114(15):3285-91. PMID: 19666869
44. NCCN Guidelines® - NCCN-Myeloproliferative Neoplasms [Version 1.2024]
45. Lundberg et al. Clonal evolution and clinical correlates of somatic mutations in myeloproliferative neoplasms. *Blood.* 2014 Apr 3;123(14):2220-8. PMID: 24478400
46. Molenaar et al. Wild-type and mutated IDH1/2 enzymes and therapy responses. *Oncogene.* 2018 Apr;37(15):1949-1960. PMID: 29367755
47. Yan et al. IDH1 and IDH2 mutations in gliomas. *N. Engl. J. Med.* 2009 Feb 19;360(8):765-73. PMID: 19228619
48. Dang et al. Cancer-associated IDH1 mutations produce 2-hydroxyglutarate. *Nature.* 2009 Dec 10;462(7274):739-44. PMID: 19935646
49. Ward et al. The common feature of leukemia-associated IDH1 and IDH2 mutations is a neomorphic enzyme activity converting alpha-ketoglutarate to 2-hydroxyglutarate. *Cancer Cell.* 2010 Mar 16;17(3):225-34. PMID: 20171147
50. Paschka et al. IDH1 and IDH2 mutations are frequent genetic alterations in acute myeloid leukemia and confer adverse prognosis in cytogenetically normal acute myeloid leukemia with NPM1 mutation without FLT3 internal tandem duplication. *J. Clin. Oncol.* 2010 Aug 1;28(22):3636-43. PMID: 20567020
51. Chou et al. The prognostic impact and stability of Isocitrate dehydrogenase 2 mutation in adult patients with acute myeloid leukemia. *Leukemia.* 2011 Feb;25(2):246-53. PMID: 21079611
52. Marcucci et al. IDH1 and IDH2 gene mutations identify novel molecular subsets within de novo cytogenetically normal acute myeloid leukemia: a Cancer and Leukemia Group B study. *J. Clin. Oncol.* 2010 May 10;28(14):2348-55. PMID: 20368543
53. [https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2023/209606s006lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2023/209606s006lbl.pdf)
54. <https://www.biospace.com/article/releases/new-analyses-from-pivotal-phase-3-indigo-study-reinforce-vorasidenib-s-potential-to-change-the-treatment-paradigm-for-idh-mutant-diffuse-glioma/?s=86>
55. Intlekofer et al. Acquired resistance to IDH inhibition through trans or cis dimer-interface mutations. *Nature.* 2018 Jul;559(7712):125-129. PMID: 29950729
56. Cancer Genome Atlas Research Network. Comprehensive, Integrative Genomic Analysis of Diffuse Lower-Grade Gliomas. *N Engl J Med.* 2015 Jun 25;372(26):2481-98. doi: 10.1056/NEJMoa1402121. Epub 2015 Jun 10. PMID: 26061751
57. Houillier et al. IDH1 or IDH2 mutations predict longer survival and response to temozolomide in low-grade gliomas. *Neurology.* 2010 Oct 26;75(17):1560-6. PMID: 20975057
58. Louis et al. The 2021 WHO Classification of Tumors of the Central Nervous System: a summary. *Neuro Oncol.* 2021 Aug 2;23(8):1231-1251. PMID: 34185076

## References (continued)

59. Pylayeva-Gupta et al. RAS oncogenes: weaving a tumorigenic web. *Nat. Rev. Cancer*. 2011 Oct 13;11(11):761-74. PMID: 21993244
60. Karnoub et al. Ras oncogenes: split personalities. *Nat. Rev. Mol. Cell Biol.* 2008 Jul;9(7):517-31. PMID: 18568040
61. Scott et al. Therapeutic Approaches to RAS Mutation. *Cancer J.* 2016 May-Jun;22(3):165-74. doi: 10.1097/PPO.000000000000187. PMID: 27341593
62. Janku et al. PIK3CA mutations frequently coexist with RAS and BRAF mutations in patients with advanced cancers. *PLoS ONE*. 2011;6(7):e22769. PMID: 21829508
63. Ohashi et al. Characteristics of lung cancers harboring NRAS mutations. *Clin. Cancer Res.* 2013 May 1;19(9):2584-91. PMID: 23515407
64. Allegra et al. Extended RAS Gene Mutation Testing in Metastatic Colorectal Carcinoma to Predict Response to Anti-Epidermal Growth Factor Receptor Monoclonal Antibody Therapy: American Society of Clinical Oncology Provisional Clinical Opinion Update 2015. *J. Clin. Oncol.* 2016 Jan 10;34(2):179-85. PMID: 26438111
65. [https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2021/125084s279lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2021/125084s279lbl.pdf)
66. [https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2021/125147s210lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2021/125147s210lbl.pdf)
67. <https://investors.kinnate.com/news-releases/news-release-details/kinnate-biopharma-inc-receives-fast-track-designation-us-food>
68. <https://investors.erasca.com/node/7891/pdf>
69. Johnson et al. Treatment of NRAS-Mutant Melanoma. *Curr Treat Options Oncol.* 2015 Apr;16(4):15. doi: 10.1007/s11864-015-0330-z. PMID: 25796376
70. Dummer et al. Binimetinib versus dacarbazine in patients with advanced NRAS-mutant melanoma (NEMO): a multicentre, open-label, randomised, phase 3 trial. *Lancet Oncol.* 2017 Apr;18(4):435-445. PMID: 28284557