

Searching 21-cm Absorption Systems in Chinese Radio Telescopes

Hao-Ran Yu^{1,2}, Ue-Li Pen^{2,3,4,5}, Tong-Jie Zhang⁶, Di Li^{7,8}, Xuelei Chen⁹

¹ Kavli Institute for Astronomy & Astrophysics, Peking University, Beijing 100871, China;

² Canadian Institute for Theoretical Astrophysics, University of Toronto, Toronto, M5S 3H8, ON, Canada;

³ Dunlap Institute for Astronomy and Astrophysics, University of Toronto, Toronto, M5S 3H4, ON, Canada;

⁴ Canadian Institute for Advanced Research, Program in Cosmology and Gravitation;

⁵ Perimeter Institute for Theoretical Physics, Waterloo, ON, N2L 2Y5, Canada;

⁶ Department of Astronomy, Beijing Normal University, Beijing, 100875, China;

⁷ National Astronomical Observatories, Chinese Academy of Sciences, Beijing, China;

⁸ Key Laboratory of Radio Astronomy, Chinese Academy of Sciences, Beijing, China;

⁹ Key Laboratory for Computational Astrophysics, National Astronomical Observatories, Chinese Academy of Sciences, Beijing, 100012, China.

Abstract

Key words:

1 INTRODUCTION

Acknowledgements HRY acknowledges General Financial Grant No.2015M570884 and Special Financial Grant No. 2016T90009 from the China Postdoctoral Science Foundation. HRY and ULP acknowledge the support of the National Science and Engineering Research Council of Canada.