

Evaluating Magnetic Reconnection Through the Nanoflare Heating Model: Evidence from Modern Solar Observations

Jun He*

Millbrook School, Millbrook, NY 12545, USA

**Corresponding author: Jun He.*

Abstract

The solar corona reaches temperatures exceeding one million Kelvin, far higher than the approximately 5,800 K temperature of the photosphere, giving rise to the long-standing coronal heating problem. This paper reviews modern observational evidence related to the nanoflare heating model and discusses the extent to which it supports magnetic reconnection as a mechanism for coronal heating. A qualitative literature review was conducted using three representative studies that provide complementary perspectives: theoretical predictions of nanoflare heating, observational evidence from solar active regions, and the thermal evolution of individual coronal loops. The studies were compared in terms of observed heating signatures, consistency with theoretical predictions, and their main limitations. The reviewed evidence shows several behaviors consistent with impulsive nanoflare heating, including temperature-dependent emission sequences, heating and cooling patterns in coronal loops, and signatures associated with small-scale episodic energy release. These characteristics are observed across different spatial scales and generally agree with predictions of nanoflare-based models. However, the evidence remains indirect because individual nanoflares and their associated magnetic reconnection events cannot yet be directly resolved by current instruments. In addition, observational uncertainties and differences between modeled and observed loop behavior limit the strength of the conclusions. Overall, modern solar observations support nanoflares as an important potential contributor to coronal heating, but they do not establish magnetic reconnection through nanoflares as the sole primary heating mechanism. Further improvements in observational resolution and theoretical modeling are required to better determine the role of nanoflares in coronal heating.

Keywords

coronal heating, nanoflares, magnetic reconnection, solar corona, coronal loops

1. Introduction

The Sun, the closest star to Earth, serves as the primary laboratory for studying stellar atmospheres and plasma physics. Like other stars, it includes several distinct layers: the core, radiative zone, convective zone, photosphere, chromosphere, and corona. Energy generated by nuclear fusion in the core is transported outward via radiation and convection before escaping from the photosphere as electromagnetic radiation. Based on this energy transport, one would expect temperatures to decrease with increasing distance from the core. However, observations reveal a striking contradiction: while the photosphere has an average temperature of

approximately 5,800 K, the overlying corona reaches temperatures exceeding one million kelvin [1]. This unexpected increase, known as the coronal heating problem, remains one of the longest-standing unresolved questions in solar physics [1,2]. Understanding coronal heating is essential not only for explaining the Sun's atmospheric structure but also for advancing our knowledge of stellar magnetic activity, plasma physics, and space weather. The corona serves as the source of the solar wind and numerous eruptive events that can affect satellites, communication systems, and power infrastructure on Earth. Hence, identifying the mechanism responsible for coronal heating has become a major goal in both theoretical and observational solar physics.

Over the past several decades, several mechanisms have been proposed to transport energy into the corona and convert it into heat. Among these, two broad classes of models have received particular attention. The first is wave heating, in which magnetohydrodynamic (MHD) waves generated below the solar surface propagate into the corona and dissipate their energy through various damping processes [1-4]. The second is magnetic reconnection, in which stressed magnetic field lines rapidly change their connectivity, converting stored magnetic energy into thermal energy, plasma motion, and energetic particles [5]. Both mechanisms enjoy substantial theoretical and observational support, yet neither has been established as a complete explanation for all observed coronal properties [1,2]. Rather than attempting a comprehensive comparison of all proposed mechanisms, this review focuses on the nanoflare model—a reconnection-based explanation for coronal heating.

The nanoflare heating model, first proposed by Parker, suggests that the solar corona may be heated by a bunch of extremely small magnetic energy release events. Individually, these nanoflares are too small to be distinguished directly by current observational instruments. However, if they occur frequently throughout the corona, their cumulative energy release could account for the observed million-kelvin temperatures [6]. Since Parker's original proposal, theoretical models have identified several observable signatures that should accompany impulsive nanoflare heating, including intensity fluctuations, temperature-dependent time lags, characteristic emission measure distributions, and the presence of hot plasma [7]. Detecting these signatures has thus become a primary method for evaluating the nanoflare hypothesis.

Recent advances in solar observations have greatly improved the ability to investigate these predictions. The Solar Dynamics Observatory (SDO) and its Atmospheric Imaging Assembly (AIA) provide high-cadence, multiwavelength observations of the solar atmosphere [8,9]. Hinode has also supplied extensive observations of coronal plasma and magnetic structures [10], while Parker Solar Probe [11] and Solar Orbiter [12] have expanded the observational basis for investigating solar magnetic activity and coronal processes. These missions and instruments have enabled researchers to compare theoretical predictions with observations across a wide range of spatial and temporal scales. This review examines observational evidence for the nanoflare heating model and evaluates what this evidence implies about magnetic reconnection by analyzing three representative studies. Together, these studies approach the problem from complementary perspectives: theoretical predictions, active region observations, and the thermal evolution of individual coronal loops. By comparing their methodologies, findings, and limitations, this review aims to assess the current strength of observational support for magnetic reconnection as a mechanism for coronal heating.

Specifically, this paper addresses the following research question: To what extent do modern solar observations support magnetic reconnection—through the nanoflare heating model—as the primary explanation for coronal heating?

2. Literature Review

2.1 The Coronal Heating Problem

The solar corona, the Sun's outermost atmospheric layer, reaches temperatures exceeding one million kelvin, while the underlying photosphere is only about 5,800 K [1]. This temperature inversion is unexpected because energy from nuclear fusion in the core is transported outward through the radiative and convective zones before escaping from the photosphere; under normal conditions, temperature would decrease with distance from the energy source. The fact that the corona is nearly two orders of magnitude hotter than the photosphere thus defines the coronal heating problem. Despite decades of spectroscopic and space-based observations confirming the corona's extreme heat, the physical mechanism that sustains it remains uncertain. The corona continuously loses energy via radiation, thermal conduction, and the solar wind, so any viable

heating mechanism must supply a continuous energy input while also reproducing the observed temperature structure and variability [1,2]. Moreover, the corona is a highly magnetized plasma environment, where energy storage and release are governed by complex magnetic processes rather than by simple thermal transport [1,5]. Consequently, explaining how energy is transferred from the solar interior into the corona and converted into heat remains a central challenge in solar physics. Over the past decades, various competing mechanisms have been proposed, each aiming to account for this energy conversion.

2.2 Wave Heating

The wave heating model is one of the leading explanations for coronal heating. It proposes that energy generated by convective motions beneath the Sun's surface is transported into the corona via magneto hydrodynamic (MHD) waves. These waves, including Alfvén waves, travel along magnetic field lines and dissipate their energy through phase mixing, resonant absorption, and turbulent damping [1,3,4]. Observations have confirmed the widespread presence of MHD waves throughout the solar atmosphere, suggesting a potential contribution to coronal heating [3]. Despite this observational support, the ability of wave heating alone to account for the corona's temperature remains uncertain. Although MHD waves can carry substantial energy, the locations and efficiency of their dissipation in the corona still present theoretical and observational challenges [1-3]. Consequently, wave heating remains an active research area rather than a universally accepted solution to the coronal heating problem. Since no single mechanism has yet been shown to fully explain all observed coronal properties, alternative models continue to be explored. Among these, magnetic reconnection has become one of the most extensively studied explanations and is the focus of this review.

2.3 Magnetic Reconnection and the Nanoflare Hypothesis

Magnetic reconnection is a leading mechanism for coronal heating, as it converts magnetic energy into thermal energy. Convective motions beneath the Sun's surface continuously twist, stretch, and braid the magnetic field lines that extend into the corona. As magnetic stresses accumulate, oppositely directed field lines may reconnect, releasing stored magnetic energy as heat, plasma motion, and energetic particles [5]. Building on this idea, Parker proposed that the corona could be heated not only by large reconnection events but also by countless small-scale events termed nanoflares. Individual nanoflares release only modest amounts of energy, insufficient to sustain the corona alone. However, if such impulsive events occur frequently throughout the corona, their cumulative energy could account for the required heating [6]. A key challenge for the nanoflare hypothesis is that individual events are expected to occur at spatial and temporal scales below the resolution of current instruments, precluding direct detection. Instead, the hypothesis is tested by searching for observable signatures in coronal plasma that are predicted to result from impulsive heating [7]. These indirect diagnostics provide the basis for assessing whether nanoflare reconnection contributes significantly to heating the solar corona. In addition to observational diagnostics, numerical hydrodynamic models have become essential for interpreting coronal loop evolution. Models such as the Enthalpy-based Thermal Evolution of Loops (EBTEL) allow researchers to simulate the plasma response to impulsive energy inputs. These simulations predict specific cooling times and emission measure distributions, providing quantitative benchmarks against which observed loop brightening and fading can be compared. However, these models often rely on one-dimensional approximations along the magnetic field, which may not fully capture the transverse structure or broadening observed in actual loops [13].

2.4 Observational Signatures of Nanoflare Heating

Since individual nanoflares lie below the spatial and temporal resolution of current instruments, the nanoflare heating model must be tested indirectly. Klimchuk outlined several observable signatures that should arise if impulsive nanoflare heating is significant [7]. These signatures provide a base for comparing theoretical predictions with observations. First, intensity variability: if heating is impulsive rather than steady, coronal emission should fluctuate as plasma cycles through heating and cooling. Second, time delays between temperature-sensitive channels: after an impulsive event, plasma cools across different temperature ranges, producing a predictable brightening sequence in extreme-ultraviolet (EUV) channels. Third, the emission measure distribution—the amount of plasma at each temperature—should exhibit specific shapes, reflecting a mix of rapidly heated and cooling material. Finally, the presence of very hot plasma, which is above approximately 5 MK, is also predicted, as impulsive energy release can generate temperatures exceeding those

from steady heating models. While these signatures offer key tests of the nanoflare hypothesis, they remain indirect evidence of reconnection rather than direct observations of individual nanoflares. As summarized in Table 1, the four main observational signatures provide complementary indicators for evaluating whether coronal plasma behavior is consistent with impulsive nanoflare heating. Hence, the central question is not merely whether such signatures exist, but whether recent observations consistently match nanoflare model predictions.

Table 1: Observational signatures used to evaluate the nanoflare heating model

Observational signature	Expected behavior under impulsive heating	Relevance to the nanoflare model
Intensity variability	Coronal emission varies as plasma is repeatedly heated and cooled.	Tests whether heating is impulsive rather than purely steady.
Time lags depend on temperature	Emission peaks occur sequentially as plasma cools through different temperature ranges.	Tests for the predicted heating–cooling cycle.
Emission measure distribution	Plasma is distributed across temperatures in a way influenced by impulsive heating and cooling.	Tests whether observed thermal structure is consistent with nanoflare models.
Very hot plasma (>5 MK)	Impulsive heating can produce transient plasma with high temperature.	Tests whether sufficiently energetic impulsive heating occurs.

Source: Adapted/synthesized from Klimchuk [7].

Early observations from TRACE and SOHO/EIT revealed that coronal loops often exhibit persistent but variable intensities, challenging steady-state heating models [14]. This prompted the development of the ‘nanoflare storm’ concept, where overlapping impulsive events produce continuous emission. However, instrument response functions often cover up individual heating events. As a result, subsequent studies have gradually shifted towards two complementary strategies: statistical analysis of entire active regions and detailed tracking of isolated individual loops [13,15].

3. Methodology

This study employs a qualitative literature review to evaluate the extent to which modern solar observations support magnetic reconnection through the nanoflare heating model as an explanation for coronal heating. Rather than collecting or analyzing new observational data, this research synthesizes evidence from previously published studies. Three papers were selected for detailed analysis based on their relevance to the research question and their complementary perspectives on nanoflare heating. Together, the selected studies represent three levels of investigation: a theoretical synthesis of existing evidence [7], observations of heating signatures across solar active regions [15], and detailed observations of individual coronal loops [13]. As shown in Table 2, these studies differ in their type of evidence, spatial scale, and primary diagnostic, allowing them to serve complementary roles in this review. Examining studies at different spatial scales provides a broader basis for assessing whether evidence for nanoflare heating is consistent across observational approaches.

Table 2: Selected studies and analytical roles

Study	Type of evidence	Spatial scale	Primary diagnostic	Role in this review
Klimchuk (2017)	Theoretical synthesis	Multiple scales	Four observational signatures	Establishes predictions
Viall & Klimchuk (2011)	Active region observations	Active region	EUV time lag analysis	Tests cooling signatures
Li et al. (2015)	Coronal loop observations	Individual loops	Heating/cooling evolution	Tests thermal evolution

To ensure a systematic comparison, each study was evaluated using the same analytical framework based on three criteria. The first criterion, observed signatures, examines which predicted indicators of nanoflare heating are reported, including those outlined in the literature review [7]. The second criterion, consistency with theoretical predictions, evaluates how closely the reported observations correspond to the expected behavior of impulsive heating models. The third criterion, study limitations and uncertainties, considers factors that may affect interpretation, including observational constraints, instrumental resolution, and assumptions within numerical models.

Applying the same comparison framework to each study allows similarities and differences to be identified despite variations in observational methods and spatial scale. This approach provides a consistent basis for evaluating the evidence without treating the three studies as independent measurements of the same phenomenon.

4. Results

This section summarizes the findings of the three selected studies using the comparison framework established in the methodology. Each study is presented according to three criteria: the reported observational evidence, its agreement with theoretical predictions of the nanoflare heating model, and the limitations acknowledged by the authors.

4.1 Theoretical Framework and Observational Signatures of Nanoflare Heating

The theoretical synthesis draws on observational and theoretical research, including data from Hinode and the Solar Dynamics Observatory (SDO), as well as hydrodynamic simulations such as the Enthalpy-based Thermal Evolution of Loops (EBTEL) model [7]. The review identifies four observational signatures predicted by the nanoflare model: intensity fluctuations, time delays between temperature-sensitive channels, characteristic emission measure distributions, and plasma hotter than approximately 5 MK.

The review reports that existing observations are generally consistent with these predictions. Various techniques detect one or more of the proposed signatures, though no single study identifies all four simultaneously [7]. The review thus presents these observations as a collection of evidence supporting impulsive heating behavior.

The review also acknowledges important limitations. Individual nanoflares remain below the spatial and temporal resolution of current instruments, making direct observation unfeasible. Consequently, the evidence is indirect, derived from signatures rather than from direct detection of reconnection events.

4.2 Active Region Evidence from EUV Time Lag Analysis

Viall and Klimchuk investigated an active solar region using six extreme-ultraviolet (EUV) channels of the Atmospheric Imaging Assembly (AIA) aboard SDO [15]. They analyzed light curves from coronal loops and diffuse emission, comparing the timing of emission peaks across temperature-sensitive channels with predictions from nanoflare storm models.

The observations showed cooling patterns consistent with nanoflare storm heating. The ordering of peak intensities across AIA channels generally matched the predicted behavior of plasma cooling after impulsive heating [15]. Broader statistical patterns in active region emission were also consistent with impulsive heating.

Several limitations complicated interpretation. Multiple coronal structures overlap along the line of sight, hindering the isolation of individual structures. Additionally, AIA channels have broad temperature responses rather than single-temperature sensitivity, which complicates time-lag interpretation. The authors therefore noted that some observed patterns were less straightforward to interpret than others.

4.3 Thermal Evolution of Individual Coronal Loops

Li et al. [13] examined the thermal evolution of individual coronal loops in active region AR 11850 using SDO/AIA and HMI observations. The study tracked the spatial and temporal evolution of loops through multiple temperature-sensitive channels. During the heating phase, the loops showed transient brightening in Fe XVIII emission (corresponding to approximately 7.2 MK plasma) while simultaneously dimming at lower temperatures.

After the initial heating, the loops brightened sequentially in progressively cooler channels, indicating cooling over roughly one hour. The observed heating-cooling sequence was compared with a one-dimensional hydrodynamic loop model, showing agreement with the nanoflare process in most aspects of the evolution.

However, a discrepancy emerged: the observed loops broadened as they brightened, a behavior that the one-dimensional model could not reproduce. This limitation suggests that the model did not capture every aspect of the observed loop evolution.

5. Discussion

5.1 Complementary Evidence Across Different Scales

The three studies provide complementary rather than fully independent evidence for nanoflare heating. The theoretical synthesis outlines the expected observational signatures of impulsive heating [7], while the two observational studies examine these signatures at different spatial scales. Viall and Klimchuk analyze emission patterns across an entire active region [15], whereas Li et al. track the thermal evolution of individual coronal loops [13]. Together, these approaches show that behavior consistent with impulsive heating can be identified in both large scale active region emission and individual coronal structures. This combination strengthens the overall case by demonstrating that the relevant signatures are not restricted to a single observational scale. However, the studies also share theoretical assumptions and observational frameworks, meaning that their agreement should not be interpreted as three completely independent confirmations. The evidence is therefore complementary rather than statistically independent.

5.2 The Inference Problem

The most important limitation is that none of the three studies directly observes magnetic reconnection or an individual nanoflare. Instead, reconnection is inferred from the thermal and radiative behavior predicted by impulsive energy release [7,13,15]. Time-dependent brightening, cooling sequences, emission measure distributions, and very hot plasma can therefore provide evidence consistent with nanoflare heating, but they do not uniquely prove that magnetic reconnection produced the observed energy. This distinction is critical because individual nanoflares are expected to occur on spatial and temporal scales smaller than those currently accessible to solar instruments. The observations consequently test the consequences predicted by the nanoflare model rather than the reconnection events themselves. The results therefore support the model at the level of indirect observational consistency, but they cannot establish a direct causal connection between the observed plasma behavior and individual reconnection events.

5.3 Limitations of Current Models

The comparison also reveals that agreement between observations and nanoflare models is not complete. Li et al., for example, found that while a one-dimensional hydrodynamic model generally reproduced the observed thermal evolution of a coronal loop, it did not reproduce the observed broadening of the loop [13]. This discrepancy is important because it indicates that impulsive heating alone may not capture every aspect of coronal loop evolution. The broadening could reflect additional physical processes or limitations in the simplified one dimensional description, though the observations alone cannot determine which explanation is responsible. More broadly, the limitations identified across the studies show that matching temperature evolution does not necessarily mean that current models reproduce the full physical behavior of the corona. Nanoflare heating may therefore operate together with other processes rather than functioning as an isolated mechanism.

5.4 Dependence and Scope of the Evidence

A further limitation concerns the independence and scope of the selected literature. Two of the three core studies share James Klimchuk as an author, and the four observational signatures used as the primary evaluation framework are also drawn from Klimchuk's theoretical synthesis [7,15]. This creates a degree of conceptual and author overlap within the evidence base. The three studies should therefore not be treated as completely independent tests derived from unrelated research programs. At the same time, their different observational scales and analytical approaches provide useful complementary evidence. Within the scope of this review, the observations consistently exhibit behavior predicted by impulsive heating models, but the evidence remains indirect and subject to model and instrumental limitations. Thus, modern observations

support magnetic reconnection through nanoflares as a plausible and important contributor to coronal heating, but the available evidence is not sufficient to establish it as the primary mechanism.

6. Conclusion

The coronal heating problem originates from the unexpectedly high temperature of the solar corona, which exceeds 1 million K while the photosphere is only about 5,800 K. This review assessed the extent to which modern solar observations support magnetic reconnection via the nanoflare heating model as an explanation for this temperature disparity. By comparing three studies—covering theoretical predictions, active region observations, and individual coronal loop analyses—this evaluation was based on three criteria: observed heating signatures, consistency with predictions, and study limitations.

Overall, the evidence provides substantial and consistent support for nanoflare heating as a significant process in the corona. The examined studies report multiple behaviors predicted by impulsive heating models, including emission sequences that depend on temperature and other signatures associated with episodic energy release. These patterns appear across observational scales, from individual loops to entire active regions. The consistency of these observations with nanoflare models indicates that impulsive heating plays an important role in coronal energy release.

However, the evidence does not establish magnetic reconnection through nanoflares as the definitively primary mechanism for all coronal heating. The central limitation is that none of the studies directly observes an individual nanoflare or the associated reconnection process. Instead, reconnection is inferred from plasma behavior that matches the predictions of nanoflare models. Since other physical processes may produce similar observational signatures, consistency with these predictions cannot uniquely identify the causal mechanism. Furthermore, current one dimensional models fail to reproduce all observed loop properties, indicating that nanoflare heating models do not yet capture every aspect of coronal evolution. Therefore, in answer to the research question, modern solar observations substantially support magnetic reconnection through nanoflares as an important contributor to coronal heating, but they do not yet establish it as the sole primary mechanism. Future progress will depend on observations with finer spatial and temporal resolution, improved magnetic field measurements, and more comprehensive models. Such advances may eventually allow individual small scale energy release events to be resolved directly, providing stronger evidence for whether nanoflares themselves—and the associated magnetic reconnection—are responsible for heating the corona.

References

- [1] Klimchuk, J. A. (2006). On solving the coronal heating problem. *Solar Physics*, 234(1), 41–77. <https://doi.org/10.1007/s11207-006-0055-z>
- [2] Parnell, C. E., & De Moortel, I. (2012). A contemporary view of coronal heating. *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences*, 370(1970), 3217–3240. <https://doi.org/10.1098/rsta.2012.0113>
- [3] De Moortel, I., & Nakariakov, V. M. (2012). Magnetohydrodynamic waves and coronal heating. *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences*, 370(1970), 3193–3216. <https://doi.org/10.1098/rsta.2011.0640>
- [4] Priest, E. R. (2014). *Magnetohydrodynamics of the Sun*. Cambridge University Press. <https://doi.org/10.1017/CBO9781139020732>
- [5] Priest, E. R., & Forbes, T. G. (2000). *Magnetic Reconnection: MHD Theory and Applications*. Cambridge University Press. <https://doi.org/10.1017/CBO9780511525087>
- [6] Parker, E. N. (1988). Nanoflares and the solar X-ray corona. *The Astrophysical Journal*, 330, 474–479. <https://doi.org/10.1086/166485>
- [7] Klimchuk, J. A. (2017). Nanoflare heating: Observations and theory. arXiv preprint arXiv:1709.07320.
- [8] Pesnell, W. D., Thompson, B. J., & Chamberlin, P. C. (2012). The Solar Dynamics Observatory (SDO). *Solar Physics*, 275, 3–15. <https://doi.org/10.1007/s11207-011-9841-3>

- [9] Lemen, J. R., Title, A. M., Akin, D. J., Boerner, P. F., Chou, C., Drake, J. F., Duncan, D. W., Edwards, C. G., Friedlaender, F. M., Heyman, G. F., Hurlburt, N. E., Katz, N. L., Kushner, G. D., Levay, M., Lindgren, R. W., Mathur, D. P., McFeaters, E. L., Mitchell, S., Rehse, R. A., Schrijver, C. J., Springer, L. A., Stern, R. A., Tarbell, T. D., Wülser, J.-P., Wolfson, C. J., Yanari, C., Bookbinder, J. A., Cheimets, P. N., Caldwell, D., DeLuca, E. E., Gates, R., Golub, L., Park, S., Podgorski, W. A., Bush, R. I., Scherrer, P. H., Gumm, M. A., Smith, P., Auken, G., Jerram, P., Pool, P., Soufli, R., Windt, D. L., Beardsley, S., Clapp, M., Lang, J., & Waltham, N. (2012). The Atmospheric Imaging Assembly (AIA) on the Solar Dynamics Observatory (SDO). *Solar Physics*, 275, 17–40. <https://doi.org/10.1007/s11207-011-9776-8>
- [10] Kosugi, T., Matsuzaki, K., Sakao, T., Shimizu, T., Sone, Y., Tachikawa, S., Hashimoto, T., Minesugi, K., Ohnishi, A., Yamada, T., Tsuneta, S., Hara, H., Ichimoto, K., Suematsu, Y., Shimojo, M., Watanabe, T., Shimada, S., Davis, J. M., Hill, L. D., Owens, J. K., Title, A. M., Culhane, J. L., Harra, L. K., Doschek, G. A., & Golub, L. (2007). The Hinode (Solar-B) Mission: An Overview. *Solar Physics*, 243, 3–17. <https://doi.org/10.1007/s11207-007-9014-6>
- [11] Fox, N. J., Velli, M. C., Bale, S. D., Decker, R., Driesman, A., Howard, R. A., Kasper, J. C., Kinnison, J., Kusterer, M., Lario, D., Lockwood, M. K., McComas, D. J., Raouafi, N. E., & Szabo, A. (2016). The Solar Probe Plus Mission: Humanity's First Visit to Our Star. *Space Science Reviews*, 204, 7–48. <https://doi.org/10.1007/s11214-015-0211-6>
- [12] Müller, D., St. Cyr, O. C., Zouganelis, I., Gilbert, H. R., Marsden, R., Nieves-Chinchilla, T., Antonucci, E., Auchère, F., Berghmans, D., Horbury, T. S., Howard, R. A., Krucker, S., Maksimovic, M., Owen, C. J., Rochus, P., Rodriguez-Pacheco, J., Romoli, M., Solanki, S. K., Bruno, R., Carlsson, M., Fludra, A., Harra, L., Hassler, D. M., Livi, S., Louarn, P., Peter, H., Schühle, U., Teriaca, L., del Toro Iniesta, J. C., Wimmer-Schweingruber, R. F., Marsch, E., Velli, M., De Groof, A., Walsh, R. W., & Williams, D. (2020). The Solar Orbiter mission: Science overview. *Astronomy & Astrophysics*, 642, A1. <https://doi.org/10.1051/0004-6361/202038467>
- [13] Li, L. P., Peter, H., Chen, F., & Zhang, J. (2015). Heating and cooling of coronal loops observed by SDO. *Astronomy & Astrophysics*, 583, A109. <https://doi.org/10.1051/0004-6361/201526912>
- [14] Reale, F. (2014). Coronal loops: Observations and modeling of confined plasma. *Living Reviews in Solar Physics*, 11, 4. <https://doi.org/10.12942/lrsp-2014-4>
- [15] Viall, N. M., & Klimchuk, J. A. (2011). Patterns of nanoflare storm heating exhibited by an active region observed with Solar Dynamics Observatory/Atmospheric Imaging Assembly. *The Astrophysical Journal*, 738(1), 24. <https://doi.org/10.1088/0004-637X/738/1/24>

Funding

This research received no external funding.

Conflicts of Interest

The authors declare no conflict of interest.

Acknowledgment

This paper is an output of the science project.

Copyrights

Copyright for this article is retained by the author (s), with first publication rights granted to the journal. This is an open-access article distributed under the terms and conditions of the Creative Commons Attribution license (<http://creativecommons.org/licenses/by/4.0/>).