

### 3 Sequences

#### 3.1 Sequences

##### Definition 3.1

A **sequence** is a function  $f$  defined on all integers  $n \geq n_0$  (usually  $n_0 = 0$  or  $1$ ).

We write  $\{a_n\}_{n=n_0}^\infty$  where  $f(n) = a_n$ , for  $n \geq n_0$ .

Often, we just write  $\{a_n\}$ , if the  $n_0$  is clear.

Note that  $n$  in  $a_n$  is an **index**.

The indices of  $\{a_n\}_{n=3}^\infty$  are  $3, 4, 5, \dots$ .

##### Example 3.2

$\{(-1)^n\}_{n=1}^\infty: -1, 1, -1, 1, \dots$

$\{1 - \frac{1}{n}\}_{n=1}^\infty: 0, \frac{1}{2}, \frac{2}{3}, \dots$

$\{e^n\}_{n=1}^\infty$

##### Definition 3.3

A sequence  $\{a_n\}_{n=1}^\infty$  is **recursive** or **inductive** if  $a_1$  is given, and  $a_{n+1} = f(a_n)$  for  $n \geq 1$ .

##### Example 3.4

Suppose we have the sequence  $a_1 = \sqrt{2}$ ,  $a_2 = \sqrt{2 + \sqrt{2}}$ ,  $\sqrt{2 + \sqrt{2 + \sqrt{2}}}, \dots$

Here we can see that,  $a_2 = \sqrt{2 + a_1}$ , and  $a_3 = \sqrt{2 + a_2}$

So we can define the sequence as,  $a_{n+1} = \sqrt{2 + a_n}$  for  $n \geq 1$ .

##### Definition 3.5

$\{a_n\}_{n=n_0}^\infty$  **converges** to **number**  $L$  if for each  $\epsilon > 0$ , there is  $N^*$  or  $N_\epsilon$  so that  $n \geq N^* \implies |a_n - L| < \epsilon$ . Then, we write  $\lim_{n \rightarrow \infty} a_n = L$ , and often we will write  $a_n \rightarrow L$ .

Otherwise, if no such  $L$  exists, then  $\{a_n\}_{n=n_0}^\infty$  **diverges**.

Note that there are two types of divergence.

$\{(-1)^n\}_{n=1}^\infty$  "wobbles".

$\{n^2\}_{n=1}^\infty$ . Then,  $\lim_{n \rightarrow \infty} n^2 = \infty$ .

##### Note 3.6

1.  $\lim_{n \rightarrow \infty} a_n = a$  if and only if  $\lim_{n \rightarrow \infty} (a_n - a) = 0$

2. Prop 2.6:  $\lim_{n \rightarrow \infty} \frac{1}{n} = 0$

*Proof.* Let  $\epsilon > 0$  be arbitrary. By the Archimedean Property, there is  $n_\epsilon > 0$  such that  $0 < \frac{1}{n_\epsilon} < \epsilon$ .

Then,  $n \geq n_\epsilon \implies 0 < \frac{1}{n} < \frac{1}{n_\epsilon} < \epsilon$ . Thus,  $\lim_{n \rightarrow \infty} \frac{1}{n} = 0$ .  $\square$

3.  $\lim_{n \rightarrow \infty} a_n = \infty$  if for any number  $M > 0$ , there is an  $n_M$  so that  $n \geq n_M \implies a_n > M$ .

#### 3.2 Properties of convergence

##### Property 3.7 (Comparison Property - Lemma 2.9)

Assume that  $a_n \rightarrow a$ , and  $\{b_n\}_{n=1}^\infty$  is a sequence, and  $b$  is a number.

If  $c \geq 0$  is a number so that  $|b_n - b| \leq c|a_n - a|$ , for all  $n \geq n^*$ , then  $b_n \rightarrow b$ .

*Proof.* Note  $n \geq n^* \implies |b_n - b| \leq c|a_n - a| \xrightarrow{n} 0$

$\square$

**Example 3.8**

Show that  $\lim_{n \rightarrow \infty} \frac{1}{n^2+n} = 0$ .

Solution: for  $n \geq 1$ ,  $0 < \frac{1}{n^2+n} \leq \frac{1}{n} \xrightarrow{n} 0$  by Prop 2.6. So by the Comparison Property,  $\frac{1}{n^2+n} \xrightarrow{n} 0$

**Definition 3.9**

$\{a_n\}_{n=n_0}^{\infty}$  is **bounded** if there is some number  $M$  so that  $|a_n| \leq M$  for all  $n$ .

**Theorem 3.10** (Thm 2.18 \*\*)

If  $\{a_n\}_{n=n_0}^{\infty}$  converges, then it is bounded.

*Proof.* Let  $\epsilon > 0$  be arbitrary. If  $a_n \rightarrow a$ , then there is an  $N^*$  so that  $n \geq N^* \implies |a_n - a| < \epsilon$ .  
Let  $M = \max\{|a_1|, |a_2|, \dots, |a_{N^*}|, |a| + \epsilon\}$ . Then we have that  $|a_n| \leq M$  for all  $n \geq 0$ .  $\square$

**Theorem 3.11** (Sum Rule)

If  $a_n \rightarrow a$ ,  $b_n \rightarrow b$ , then  $a_n + b_n \rightarrow a + b$ .

*Proof.*  $|a_n + b_n - (a + b)| \leq |a_n - a| + |b_n - b| \rightarrow 0$ , because  $|a_n - a| \rightarrow 0$  and  $|b_n - b| \rightarrow 0$ .  $\square$

**Theorem 3.12** (Product Rule)

If  $a_n \rightarrow a$ ,  $b_n \rightarrow b$ , then  $a_n b_n \rightarrow ab$ .

*Proof.*  $\lim_{n \rightarrow \infty} b_n = b$  implies, by Thm 2.18,  $|b_n| \leq M$  for all  $n \geq n_0$ .  
Then,  $|a_n b_n - ab| \leq |a_n b_n - ab_n| + |ab_n - ab| = |a_n - a||b_n| + |a||b_n - b| \leq M|a_n - a| + |a||b_n - b| \rightarrow 0$ , so  $a_n b_n \rightarrow ab$ , where the first inequality is created by the triangle inequality and adding an extra  $(ab_n - ab_n)$ .  $\square$

**Theorem 3.13** (Quotient Rule)

If  $a_n \rightarrow a$ ,  $b_n \rightarrow b$ , then if  $b \neq 0$ , then  $\frac{a_n}{b_n} \rightarrow \frac{a}{b}$ .

**Note 3.14**

Suppose we want to prove  $\lim_{n \rightarrow \infty} (\sqrt{n^2+n} - \sqrt{n^2})$  exists and find it.  
We would multiply by the conjugate as follows:

$$\begin{aligned} \lim_{n \rightarrow \infty} (\sqrt{n^2+n} - \sqrt{n^2}) &= \lim_{n \rightarrow \infty} (\sqrt{n^2+n} - \sqrt{n^2}) \left( \frac{\sqrt{n^2+n} + \sqrt{n^2}}{\sqrt{n^2+n} + \sqrt{n^2}} \right) \\ &= \lim_{n \rightarrow \infty} \frac{(n^2+n) - n^2}{\sqrt{n^2+n} + \sqrt{n^2}} \\ &= \lim_{n \rightarrow \infty} \frac{n}{\sqrt{n^2+n} + \sqrt{n^2}} \\ &= \lim_{n \rightarrow \infty} \frac{n}{n+n} \\ &= \frac{1}{2} \end{aligned}$$