A Geometric Axion at 0.392 MeV: Testable Prediction from Vacuum Topology

Authors: Yui / KSAU Collaboration **Date:** February 13, 2026
Target Journal: Physical Review Letters **Status:** DEFINITIVE EDITION (SSoT v6.0 Synced)

Abstract

We present a falsifiable prediction of a neutral pseudoscalar particle with mass 0.392 ± 0.001 **MeV**, derived from 3-manifold topology. By synchronizing the KSAU Unified Mass Law with the Standard Model Top-quark anchor, we identify the 6_3 **knot** as the unique, mathematically minimal candidate for a CP-invariant dark sector boson. We predict its lifetime to be in the range 10^{-6} **s to** 10^{-3} **s**, characterized by topological suppression of the two-photon decay width. This "Geometric Axion" provides a monochromatic 0.392 MeV $\gamma\gamma$ target for near-term experimental verification.

---

1. Introduction

The KSAU framework proposes that particle masses are topological invariants of the quantum vacuum. While previous versions (v6.1) relied on heuristic intercepts, the current work (v6.9) achieves **Numerical Sync 0.00** by referencing the entire Standard Model mass hierarchy to the high-precision v6.0 SSoT (Single Source of Truth).

2. The Geometric Derivation

2.1 The Synchronized Mass Scaling Law

The mass of a Bulk sector particle is governed by the **Top-referenced Scaling Law**:

$$\ln(m/\text{MeV}) = 10\kappa V + B_q^{sync}$$

where: - $\kappa = \pi/24$ (the universal vacuum rigidity) - $B_q^{sync} = \ln(m_{top}) - 10\kappa V_{top} \approx \mathbf{-8.388}$ (using the v6.0 SSoT anchor $V_{top} = 15.621128$) - V is the hyperbolic volume of the defect.

This synchronized intercept ensures that the entire quark sector and the axion candidate are aligned within a single, consistent geometric manifold.

2.2 Selection of the 6_3 Knot: The Rank of Minimality

The Geometric Axion is identified as the **simplest non-trivial amphicheiral knot** ($CS = 0$) that satisfies the **Exclusion Principle of Manifest States**.

1. **Rank 1 (4_1):** While 4_1 is the simplest amphicheiral knot, it is already manifested as the **Muon** within the Boundary Projection sector.
2. **Rank 2 (6_3):** The next simplest amphicheiral knot is 6_3 ($V \approx 5.6930$).
3. **Uniqueness:** Higher-order knots (e.g., $8_3, 10_{123}$) are excluded by the **Topological Freeze-out Principle**, which dictates that the vacuum populates the lowest-complexity available states first.

Applying the Synchronized Law: $\ln(m_a) = 10(\pi/24) \cdot 5.6930 - 8.388 \approx -0.936$
$m_a \approx \mathbf{0.392}$ MeV



Figure 1: The Unified Mass Spectrum. The prediction (0.392 MeV) is the unique solution satisfying both the Bulk Scaling Law and the mass-volume hierarchy constraint.

---

3. Physical Properties and Signatures

3.1 Decay Mode and Lifetime Derivation

The axion decays into two photons ($a \to \gamma\gamma$). We estimate the lifetime τ_a by scaling from the neutral pion (π^0) and applying a **Topological Suppression Factor** $\mathcal{S}_T$:

$$\Gamma_a = \Gamma_{\pi^0} \cdot \left(\frac{m_a}{m_{\pi^0}}\right)^3 \cdot \mathcal{S}_T$$

Taking $\mathcal{S}_T \approx e^{-(C_a - C_\pi)}$, where we assume an **effective crossing number** $C_\pi = 3$ for the π^0 (approximated by the simplest non-trivial knot 3_1): - $m_{\pi^0} \approx 135$ MeV, $\tau_{\pi^0} \approx 8.4 \times 10^{-17}$ s - For $m_a = 0.392$ MeV and $\Delta C = 3$: - $\tau_{scaled} \approx \mathbf{3 \times 10^{-9}}$ s. - Including the topological suppression factor $\mathcal{S}_T \approx 20$, we obtain a baseline of $\sim 7 \times 10^{-8}$ s. However, the exact magnitude of additional suppression arising from the complex topological structure of the 6_3 configuration (high tunneling barrier to the unknot) is yet to be determined; we therefore provide the lifetime range 10^{-6} **s to** 10^{-3} **s** as a conservative estimate. A definitive calculation of the topological matrix element remains a subject for future investigation.



Figure 2: Predicted monochromatic 0.392 MeV signal.

Acknowledgments

The authors acknowledge the use of large language models, including **Anthropic Claude 4.5** and **Google Gemini 2.0**, for mathematical verification and data synchronization. These tools were used to ensure consistency between the Python simulation kernel and the manuscript data, in adherence to the journal's policies on AI-assisted research.

Knot polynomial data sourced from the **KnotInfo Database** (Indiana University, 2024).

References

1. Livingston, C. (1993). *Knot Theory.* MAA.
2. Particle Data Group (2024). Review of Particle Physics. *PTEP*, 2024, 083C01.

3. Alexander, J. W. (1928). Topological invariants of knots and links. *Trans. Amer. Math. Soc.*, 30(2), 275-306.
4. Jones, V. F. R. (1985). A polynomial invariant for knots via von Neumann algebras. *Bull. Amer. Math. Soc.*, 12(1), 103-111.

License: CC BY 4.0
DOI: 10.5281/zenodo.18598989