

This file is to collect various notes on our project on the higher symmetry and Seiberg duality.

1 Duality map

The duality map among $Spin$, SO_+ and SO_- was first found in [1]

$$\begin{aligned} Spin(N_c) &\leftrightarrow SO_-(N'_c), \\ SO_+(N_c) &\leftrightarrow SO_+(N'_c), \\ SO_-(N_c) &\leftrightarrow Spin(N'_c) \end{aligned} \tag{1.1}$$

where $N'_c = N_f - N_c + 4$.

Razamat-Willett [2] performed a rather extensive check of this mapping by means of localization on the lens space times S^1 .

Note that there is a natural action of $SL(2, \mathbb{Z})$ on the theories with \mathbb{Z}_2 1-form symmetry. Requiring that this is compatible with the Seiberg duality, one finds that the mapping should in fact be

$$\begin{aligned} Spin(N_c) &\leftrightarrow T(SO_-(N'_c)), \\ SO_+(N_c) &\leftrightarrow T(SO_+(N'_c)), \\ SO_-(N_c) &\leftrightarrow T(Spin(N'_c)) \end{aligned} \tag{1.2}$$

as discussed in [3, Sec. 6] and [4].

2 References on fermionic zero modes on monopoles

Index theorem on the monopole background: Callias [5] Bott and Seeley [6]¹

General reviews: Harvey [7, Lecture 4].

3 Explicit configurations detecting anomalies

Here we describe geometries detecting $\int_{M_5} B\beta E$, $\int_{M_4} B\beta w_2$, etc. All cohomologies in this section is \mathbb{Z}_2 -valued.

This is to confirm that these expressions are not secretly trivial.

Klein bottle: We start from the Klein bottle K as a nontrivial S^1 bundle over S^1 . Let us denote by a the Poincaré dual to the fiber S^1 , and t the Poincaré dual to the base S^1 .

We have $\beta a = ta$, since $\int_K \beta a = \int_K w_1 a$.

¹These are on CMP. Papers on CMP are not open access via Springer (which is reachable by doi) but is open access at Project Euclid. I'd like a way to include links in the references appropriately.

T^4 bundle over S^1 : We now consider a T^4 bundle over S^1 . We denote four directions of T^4 as 1, 2, 3 and 4, and we let the directions 1 and 3 to flip the orientation when we go around S^1 . We let $a_{1,2,3,4} \in H^1(T^4)$ be the dual basis to the S^1 along four directions.

We now take $B = a_1 a_2$ and $E = a_3 a_4$. Then $\beta B = tB$ and $\beta E = tE$, and $\int B \beta E = 1$.

Realizing as $SO(3)$ bundles We now look for $SO(3)$ bundles realizing these B and E as w_2 in this T^4 bundle over S^1 .

We note that an $SO(3)$ bundle over T^2 with two commuting holonomies around two directions

$$R_x = \text{diag}(+1, -1, -1), \quad R_y = \text{diag}(+1, -1, -1)$$

has a nontrivial w_2 , since their lift to $SU(2)$ is given by $i\sigma_x$ and $i\sigma_y$ which anticommute.

Luckily, these R_x and R_y are of order two, so we can put it over our T^4 bundle. Done.

4 Anomaly of trifundamental

Lee-kun's computation says that

$$(D\Omega^{\text{spin}})^6(B[SU(2)^3/\mathbb{Z}_2^2]) = \mathbb{Z}_2$$

generated by

$$\int w_2 \beta w'_2.$$

We would like to know if a trifundamental fermion has this anomaly.

A Computation of $\Omega_5^{\text{spin}}(B(\frac{SO(4) \times SU(2)}{\mathbb{Z}_2}))$

Let us consider the simplest case of the Seiberg duality and examine the anomaly consequences. For $SO(4)$ gauge theory with $N_f = 2$ flavors, fermions are charged under $\frac{SO(4) \times SU(2)}{\mathbb{Z}_2}$.

A.1 Leray-Serre SS (preparatory)

For the various input of cohomology groups, see Appendix A of our WZW paper. For the fibration

$$BSO(3) \rightarrow B(SO(3) \times SO(3)) = B\left(\frac{SO(4)}{\mathbb{Z}_2}\right) \rightarrow BSO(3) \quad (\text{A.1})$$

one has

$$E_2^{p,q} = H^p(BSO(3); H^q(BSO(3); \mathbb{Z})) \quad H^{p+q}(B\left(\frac{SO(4)}{\mathbb{Z}_2}\right); \mathbb{Z}) \quad (A.2)$$

6	\mathbb{Z}_2						
5							
4	\mathbb{Z}						
3	\mathbb{Z}_2			\mathbb{Z}_2	\mathbb{Z}_2		
2							
1							
0	\mathbb{Z}			\mathbb{Z}_2	\mathbb{Z}		\mathbb{Z}_2
		0	1	2	3	4	5

\longrightarrow

6	$\mathbb{Z}_2^{\oplus 3}$						
5							
4	$\mathbb{Z}^{\oplus 2}$						
3	$\mathbb{Z}_2^{\oplus 2}$						
2							
1							
0	\mathbb{Z}						

Here we expect non-trivial differentials to be absent (for the region of interest) from the explicit consideration of generators (since there are W_3 and W'_3 , there should be $(W_3)^2$, $(W'_3)^2$, and $W_3 W'_3$) or by requiring proper reproduction of the \mathbb{Z}_2 cohomology (which we expect to be generated by w_2 , w'_2 , w_3 , and w'_3). Then, for the fibration

$$BSU(2) \rightarrow B\left(\frac{SO(4) \times SU(2)}{\mathbb{Z}_2}\right) \rightarrow B\left(\frac{SO(4)}{\mathbb{Z}_2}\right) \quad (A.3)$$

we can further plug it into

$$E_2^{p,q} = H^p(B\left(\frac{SO(4)}{\mathbb{Z}_2}\right); H^q(BSU(2); \mathbb{Z})) \quad H^{p+q}(B\left(\frac{SO(4) \times SU(2)}{\mathbb{Z}_2}\right); \mathbb{Z}) \quad (A.4)$$

6							
5							
4	\mathbb{Z}						
3							
2							
1							
0	\mathbb{Z}			$\mathbb{Z}_2^{\oplus 2}$	$\mathbb{Z}_2^{\oplus 2}$	\mathbb{Z}_2	$\mathbb{Z}_2^{\oplus 3}$
		0	1	2	3	4	5

\longrightarrow

6	$\mathbb{Z}_2^{\oplus 3}$						
5							
4	$\mathbb{Z}^{\oplus 3}$						
3	$\mathbb{Z}_2^{\oplus 2}$						
2							
1							
0	\mathbb{Z}						

Taking the normalization of instanton number into account (see Ohmori-san's e-mail on 2020-08-19), the differential $d_2 : E_{0,4} \rightarrow E_{5,0}$ seems to be non-trivial.

So we believe the integral cohomology structure to be

d		0	1	2	3	4	5	6	\dots
$H^d(B\left(\frac{SO(4) \times SU(2)}{\mathbb{Z}_2}\right); \mathbb{Z})$	\mathbb{Z}	0	0	$\mathbb{Z}_2^{\oplus 2}$	$\mathbb{Z}^{\oplus 3}$	0	$\mathbb{Z}_2^{\oplus 3}$	\dots	
generator	1	—	—	W_3	p_1	—	$(W_3)^2$	\dots	
				W'_3	p'_1		$(W'_3)^2$		
					$2c_2$		$W_3 W'_3$		

where the reduction to \mathbb{Z}_2 cohomology are

$$\begin{aligned} W_3 &\rightarrow w_3 \\ p_1 &\rightarrow (w_2)^2 \end{aligned} \quad (\text{A.6})$$

A.2 Atiyah-Hirzebruch SS

Having obtained (co)homology groups, one can fill in the E^2 -page of the AHSS:

$$E_{p,q}^2 = H_p\left(B\left(\frac{SO(4) \times SU(2)}{\mathbb{Z}_2}\right); \Omega_q^{\text{spin}}\right) \quad \tilde{\Omega}_{p+q}^{\text{spin}}\left(B\left(\frac{SO(4) \times SU(2)}{\mathbb{Z}_2}\right)\right) \quad (\text{A.7})$$

Based on our belief, $d^2 : E_{4,0}^2 \rightarrow E_{2,1}^2$ and $d^2 : E_{4,1}^2 \rightarrow E_{2,2}^2$ should be a dual of

$$\begin{aligned} Sq^2 w_2 &= (w_2)^2 \\ Sq^2 w'_2 &= (w'_2)^2 \end{aligned} \quad (\text{A.8})$$

and also $d^2 : E_{5,0}^2 \rightarrow E_{3,1}^2$ and $d^2 : E_{5,1}^2 \rightarrow E_{3,2}^2$ should be a dual of

$$\begin{aligned} Sq^2 w_3 &= w_2 w_3 \\ Sq^2 w'_3 &= w'_2 w'_3 \end{aligned} \quad (\text{A.9})$$

and finally $d^2 : E_{6,0}^2 \rightarrow E_{4,1}^2$ should be a dual of

$$Sq^2(w_2 w'_2) = w_3 w'_3 + (w_2)^2 w'_2 + w_2 (w'_2)^2 \quad (\text{A.10})$$

then the would-be- E_3 -page is given by

A.3 Adams SS

According to our naive guess, the module $\tilde{H}^*(B(\frac{SO(4) \times SU(2)}{\mathbb{Z}_2}); \mathbb{Z}_2)_{\leq 5}$ consists of

(A.12)


To be consistent with the AHSS computation, it seems that $w_3w'_2 + w_2w'_3$ should be modded out (is it an obvious consequence of the transgression in LSSS...?) and the remaining part (*) turns out to be

(A.13)

and therefore one concludes

$$\tilde{H}^*(B(\frac{SO(4) \times SU(2)}{\mathbb{Z}_2}); \mathbb{Z}_2)_{\leq 5} = J[2] \oplus J[2] \oplus \mathcal{A}_1 // \mathcal{E}_0[4] \oplus J[5]. \quad (\text{A.14})$$

This leads to the following Adams chart:



and it indeed seems to be compatible with the AHSS computation. If the above argument (and beliefs) is correct, then the anomaly should be captured by

$$w_2 w'_3 (= w_3 w'_2). \quad (\text{A.15})$$

A.4 Seiberg dual

The dual theory is $SO(2)$ gauge theory with $N_f = 2$ flavors, and the fermions are charged under

$$\frac{SO(2) \times SU(2)}{\mathbb{Z}_2} = U(2) \quad (\text{A.16})$$

Its spin bordism is known, and the relevant group turns out to be trivial:

$$\Omega_5^{\text{spin}}(BU(2)) = 0. \quad (\text{A.17})$$

This means there is no anomaly for fermions on the dual side, and thus the $B\beta E$ anomaly cannot be canceled by fermions.

References

- [1] O. Aharony, N. Seiberg, and Y. Tachikawa, *Reading Between the Lines of Four-Dimensional Gauge Theories*, *JHEP* **08** (2013) 115, [arXiv:1305.0318 \[hep-th\]](#).
- [2] S. S. Razamat and B. Willett, *Global Properties of Supersymmetric Theories and the Lens Space*, *Commun. Math. Phys.* **334** (2015) 661–696, [arXiv:1307.4381 \[hep-th\]](#).
- [3] D. Gaiotto, A. Kapustin, N. Seiberg, and B. Willett, *Generalized Global Symmetries*, *JHEP* **02** (2015) 172, [arXiv:1412.5148 \[hep-th\]](#).
- [4] L. Bhardwaj, Y. Lee, and Y. Tachikawa, *$SL(2, \mathbb{Z})$ action on QFTs with \mathbb{Z}_2 symmetry and the Brown-Kervaire invariants*, *JHEP* **11** (2020) 141, [arXiv:2009.10099 \[hep-th\]](#).
- [5] C. Callias, *Index Theorems on Open Spaces*, *Commun. Math. Phys.* **62** (1978) 213–234.
- [6] R. Bott and R. Seeley, *Some Remarks on the Paper of Callias*, *Commun. Math. Phys.* **62** (1978) 235–245.
- [7] J. A. Harvey, *Magnetic Monopoles, Duality and Supersymmetry*, in *ICTP Summer School in High-Energy Physics and Cosmology*. 3, 1996. [arXiv:hep-th/9603086](#).