


Small ball probabilities for Gaussian processes



Alma mater:
St Petersburg State
University, Russia



Yulia Petrova

PUC-Rio, Pontifical Catholic University of Rio de Janeiro
Department of Mathematics

<https://yulia-petrova.github.io/>



Spring School “Multiplicative chaos and cascades”
TU Darmstadt, 19 February 2024



TECHNISCHE
UNIVERSITÄT
DARMSTADT

Surveys on the topic:

- W.V. Li and Q.Shao, 2001. Gaussian processes: inequalities, small ball probabilities and applications
- A. Nazarov and Y. Petrova, 2023. L_2 -small ball asymptotics for Gaussian random functions: A survey
- Bibliography (by M.A. Lifshits): <https://airtable.com/shrMG0nNxI9SiGxII/tbl7Xj1mZW2VuYurm>

Small ball probabilities (small deviations)

Let $(\mathcal{X}, \|\cdot\|)$ be a Banach space (f.e. $C[0, 1]$ or $L^2[0, 1]$).

Small ball probabilities (small deviations)

Let $(\mathcal{X}, \|\cdot\|)$ be a Banach space (f.e. $C[0, 1]$ or $L^2[0, 1]$).

Definition

An \mathcal{X} -valued random vector X is a measurable mapping

$$X : (\Omega, \mathbb{P}) \rightarrow \mathcal{X} \tag{1}$$

Small ball probabilities (small deviations)

Let $(\mathcal{X}, \|\cdot\|)$ be a Banach space (f.e. $C[0, 1]$ or $L^2[0, 1]$).

Definition

An \mathcal{X} -valued random vector X is a measurable mapping

$$X : (\Omega, \mathbb{P}) \rightarrow \mathcal{X} \tag{1}$$

We will consider centered random vector, that is $\mathbb{E}X = 0$.

Small ball probabilities (small deviations)

Let $(\mathcal{X}, \|\cdot\|)$ be a Banach space (f.e. $C[0, 1]$ or $L^2[0, 1]$).

Definition

An \mathcal{X} -valued random vector X is a measurable mapping

$$X : (\Omega, \mathbb{P}) \rightarrow \mathcal{X} \quad (1)$$

We will consider centered random vector, that is $\mathbb{E}X = 0$.

Definition

Small ball probability problem consists in finding the asymptotics

$$\mathbb{P}(\|X\| < \varepsilon) \quad \text{as } \varepsilon \rightarrow 0 \quad (2)$$

Small ball probabilities (small deviations)

Let $(\mathcal{X}, \|\cdot\|)$ be a Banach space (f.e. $C[0, 1]$ or $L^2[0, 1]$).

Definition

An \mathcal{X} -valued random vector X is a measurable mapping

$$X : (\Omega, \mathbb{P}) \rightarrow \mathcal{X} \quad (1)$$

We will consider centered random vector, that is $\mathbb{E}X = 0$.

Definition

Small ball probability problem consists in finding the asymptotics

$$\mathbb{P}(\|X\| < \varepsilon) \quad \text{as } \varepsilon \rightarrow 0 \quad (2)$$

Actually, it can be formulated as a problem in measure theory.

Small ball probabilities (small deviations)

Let $(\mathcal{X}, \|\cdot\|)$ be a Banach space (f.e. $C[0, 1]$ or $L^2[0, 1]$).

Definition

An \mathcal{X} -valued random vector X is a measurable mapping

$$X : (\Omega, \mathbb{P}) \rightarrow \mathcal{X} \quad (1)$$

We will consider centered random vector, that is $\mathbb{E}X = 0$.

Definition

Small ball probability problem consists in finding the asymptotics

$$\mathbb{P}(\|X\| < \varepsilon) \quad \text{as } \varepsilon \rightarrow 0 \quad (2)$$

Actually, it can be formulated as a problem in measure theory. Let:

- P — distribution of X — a measure in \mathcal{X} , given by $P(A) = \mathbb{P}(X \in A)$
- $U := \{x \in \mathcal{X} : \|x\| \leq 1\}$ — unit ball in \mathcal{X}

We want to study the measure of the small balls:

$$P(\varepsilon U), \quad \text{as } \varepsilon \rightarrow 0$$

Gaussian random vectors

Gaussian random vector extends the notion of a normally distributed random variable.

Definition

We call a random vector X , taking value in a linear topological space \mathcal{X} , Gaussian, if for every continuous linear functional $g \in \mathcal{X}^$ the random variable $g(X)$ has a normal distribution.*

The distribution of a Gaussian vector is uniquely determined by:

- means of $\{g(X) : g \in \mathcal{X}^*\}$;
- covariances of $\{g(X) : g \in \mathcal{X}^*\}$.

Main example

Wiener process $W(t)$ — a random element in $C[0, 1]$ or in $L^2[0, 1]$:

- $\mathbb{E}W(t) \equiv 0$;
- $\text{cov}(W(s), W(t)) = \min(s, t)$.


Example

Typical answer:

$$\mathbb{P}(\|X\| < \varepsilon) \sim D \cdot \varepsilon^C \cdot \exp(-B\varepsilon^{-A}), \quad \varepsilon \rightarrow 0$$

A, B — *logarithmic* asymptotics; A, B, C, D — *exact* asymptotics

Example: $\mathcal{X} = C[0, 1]$, $X = W(t)$ — Wiener process



$$\mathbb{P}\left(\sup_{0 \leq t \leq 1} |W(t)| < \varepsilon\right) \sim \frac{4}{\pi} \exp\left(-\frac{\pi^2}{8} \varepsilon^{-2}\right)$$

Methods

“...there is no royal road to small ball probabilities...” M.A. Lifshits

Methods

“...there is no royal road to small ball probabilities...” M.A. Lifshits

① via metric entropy:


- works for general classes of processes (e.g. in separable Banach spaces)
- allows to get only logarithmic asymptotics
- J. Kuelbs, W. Li, W. Linde, T. Dunker, F. Gao, M. Lifshits, F. Aurzada, T. Kuhn, E. Belinsky, R. Blei, W. Salkeld etc

Methods

“...there is no royal road to small ball probabilities...” M.A. Lifshits

① via metric entropy:

- works for general classes of processes (e.g. in separable Banach spaces)
- allows to get only logarithmic asymptotics
- J. Kuelbs, W. Li, W. Linde, T. Dunker, F. Gao, M. Lifshits, F. Aurzada, T. Kuhn, E. Belinsky, R. Blei, W. Salkeld etc




Methods

“...there is no royal road to small ball probabilities...” M.A. Lifshits

① via metric entropy:

- works for general classes of processes (e.g. in separable Banach spaces)
- allows to get only logarithmic asymptotics
- J. Kuelbs, W. Li, W. Linde, T. Dunker, F. Gao, M. Lifshits, F. Aurzada, T. Kuhn, E. Belinsky, R. Blei, W. Salkeld etc



Metric entropy: $\ln N_L(\varepsilon)$, where $N_L(\varepsilon)$ — covering numbers:


$$N_L(\varepsilon) = \inf \left\{ n : \exists \{x_j\}_{j \leq n}, \{Lh : \|h\|_H \leq 1\} \subset \bigcup_{j=1}^n B_\varepsilon(x_j) \right\}$$

Methods

“...there is no royal road to small ball probabilities...” M.A. Lifshits

① via metric entropy:

- works for general classes of processes (e.g. in separable Banach spaces)
- allows to get only logarithmic asymptotics
- J. Kuelbs, W. Li, W. Linde, T. Dunker, F. Gao, M. Lifshits, F. Aurzada, T. Kuhn, E. Belinsky, R. Blei, W. Salkeld etc



Metric entropy: $\ln N_L(\varepsilon)$, where $N_L(\varepsilon)$ — covering numbers:

$$N_L(\varepsilon) = \inf \left\{ n : \exists \{x_j\}_{j \leq n}, \{Lh : \|h\|_H \leq 1\} \subset \bigcup_{j=1}^n B_\varepsilon(x_j) \right\}$$

X — Gaussian vector in \mathcal{X} \leftrightarrow L — associated linear compact operator

Example: $\ln \mathbb{P}(\|X\| < \varepsilon) \approx -\varepsilon^{-\frac{2\beta}{2-\beta}}$ \iff $\ln N_L(\varepsilon) \approx \varepsilon^{-\beta}$, as $\varepsilon \rightarrow 0$

Methods

“...there is no royal road to small ball probabilities...” M.A. Lifshits

② via spectral theory:

- works for \mathcal{X} being a Hilbert space
- allows to get exact asymptotics
- St Petersburg school:
started by I. Ibragimov, M. Lifshits, Ya. Nikitin, A. Nazarov, and
followed by R. Pusev, A. Karol, N. Rastegaev, Yu. Petrova, etc

Methods

“...there is no royal road to small ball probabilities...” M.A. Lifshits

② via spectral theory:

- works for \mathcal{X} being a Hilbert space
- allows to get exact asymptotics
- St Petersburg school:
started by I. Ibragimov, M. Lifshits, Ya. Nikitin, A. Nazarov, and
followed by R. Pusev, A. Karol, N. Rastegaev, Yu. Petrova, etc

Karhunen-Loeve expansion (KL-expansion):

(K. Karhunen'1947, M. Loève'1948) Let $\mathcal{X} = L^2[0, 1]$. Then

$$X(t) \stackrel{d}{=} \sum_{k \in \mathbb{N}} u_k(t) \sqrt{\mu_k} \xi_k$$

- ξ_k , $k \in \mathbb{N}$, — iid standard normal rv
- $u_k(t)$, μ_k — orthonormal eigenfunc., eigenval. of covariance operator \mathbb{G}

Methods

“...there is no royal road to small ball probabilities...” M.A. Lifshits

② via spectral theory:

- works for \mathcal{X} being a Hilbert space
- allows to get exact asymptotics
- St Petersburg school:
started by I. Ibragimov, M. Lifshits, Ya. Nikitin, A. Nazarov, and
followed by R. Pusev, A. Karol, N. Rastegaev, Yu. Petrova, etc

Karhunen-Loeve expansion (KL-expansion):

(K. Karhunen'1947, M. Loève'1948) Let $\mathcal{X} = L^2[0, 1]$. Then

$$X(t) \stackrel{d}{=} \sum_{k \in \mathbb{N}} u_k(t) \sqrt{\mu_k} \xi_k$$

- $\xi_k, k \in \mathbb{N}$, — iid standard normal rv
- $u_k(t), \mu_k$ — orthonormal eigenfunc., eigenval. of covariance operator \mathbb{G}

$$\mathbb{P}(\|X\|_2 < \varepsilon) = \mathbb{P}\left(\sum_{k=1}^{\infty} \mu_k \xi_k^2 < \varepsilon^2\right)$$

Hilbert structure \implies spectral problem

Stochastic processes

 $X(t), t \in (0, 1), -$

- Gaussian process
- $\mathbb{E}X(t) \equiv 0$
- $G(s, t) = \mathbb{E}X(s)X(t).$

Small ball asymptotics

$$\mathbb{P}(\|X\|_2 < \varepsilon) = \mathbb{P}\left(\sum \mu_k \xi_k^2 < \varepsilon^2\right)$$

Spectral theory

 $\mathbb{G} : L_2[0, 1] \rightarrow \text{Im}(\mathbb{G})$

- integral operator of trace class

$$(\mathbb{G}u)(s) = \int_0^1 G(s, t)u(t) dt$$

- eigenvalues: $\sum \mu_k < \infty$

Asymptotics of eigenvalues μ_k Find “good” approximation to μ_k

Comments:

- the whole sequence of eigenvalues μ_k is important (in contrast to large deviations where only the first eigenvalue is sufficient to know)
- If \mathbb{G}^{-1} is a differential operator, then spectral theory for ODEs helps — my research interests

Example of a general theorem (Nazarov, Petrova' 2016)

Let the eigenvalues μ_k have the asymptotics

$$\mu_k = (\vartheta(k + \delta + F(k) + O(k^{-1})))^{-d},$$

where $F(k)$ is a slowly varying function (SVF) at ∞ .

Then for the small deviation probabilities



$$\mathbb{P}(\|X\|_2 < \varepsilon) \sim D \cdot \exp\left(\frac{1}{2}F_{-1}(\varepsilon^{-2})\right) \cdot \varepsilon^C \exp(B\varepsilon^A), \quad \varepsilon \rightarrow 0,$$

where $A = A(d)$, $B = B(d, \vartheta)$, $C = C(d, \vartheta, \delta)$, $D = D(\{\mu_k\})$:

$$A = -\frac{2}{d-1}, \quad B = -\frac{d-1}{2} \left(\frac{\pi/d}{\vartheta \sin(\pi/d)} \right)^{\frac{d}{d-1}},$$

$$C = \frac{2-d-2\delta d}{2(d-1)}, \quad F_{-1}(t) = \int_1^t \frac{F(x)}{x} dx \quad \text{also SVF}$$

Vielen Dank für eure Aufmerksamkeit!



Questions? Comments?

Contact: <https://yulia-petrova.github.io/>
yu.pe.petrova@gmail.com

Literature

Surveys / books :

- Site with all bibliography around small ball probabilities (collected by M. Lifshits): <https://airtable.com/shrMG0nNxI9SiGxII/tbl7Xj1mZW2VuYurm>
- Li, W.V. and Shao, Q.M., 2001. Gaussian processes: inequalities, small ball probabilities and applications. Handbook of Statistics, 19, pp.533-597.
- Lifshits, M., 2012. Lectures on Gaussian processes. Springer Berlin Heidelberg.
- Nazarov, A. and Petrova, Y., 2023. L_2 -small ball asymptotics for Gaussian random functions: A survey. Probability Surveys, 20, pp.608-663.