



Unpaired Two-Samples Wilcoxon Test in R


 Tools

- [Visualize your data and compute Wilcoxon test in R](#)
 - [R function to compute Wilcoxon test](#)
 - [Import your data into R](#)
 - [Check your data](#)
 - [Visualize your data using box plots](#)
 - [Compute unpaired two-samples Wilcoxon test](#)
- [Online unpaired two-samples Wilcoxon test calculator](#)
- [See also](#)
- [Infos](#)

The **unpaired two-samples Wilcoxon test** (also known as **Wilcoxon rank sum test** or **Mann-Whitney test**) is a non-parametric alternative to the [unpaired two-samples t-test](#), which can be used to compare two independent groups of samples. It's used when your data are not normally distributed.


Unpaired Two-Samples Wilcoxon test in R

- + Definition
- + Research Questions & Statistics
- + Practical Examples in R
- + Interpret



© sthda.com 2016

Related Book:



Practical Statistics in R for
Comparing Groups: Numerical
Variables



This article describes how to compute two samples Wilcoxon test in R.

Visualize your data and compute Wilcoxon test in R

R function to compute Wilcoxon test

To perform two-samples Wilcoxon test comparing the means of two independent samples (x & y), the R function **wilcox.test()** can be used as follow:

```
wilcox.test(x, y, alternative = "two.sided")
```

- **x,y**: numeric vectors
- **alternative**: the alternative hypothesis. Allowed value is one of "two.sided" (default), "greater" or "less".

Import your data into R

1. **Prepare your data** as specified here: [Best practices for preparing your data set for R](#)
2. **Save your data** in an external .txt tab or .csv files
3. **Import your data into R** as follow:

```
# If .txt tab file, use this
my_data <- read.delim(file.choose())
# Or, if .csv file, use this
my_data <- read.csv(file.choose())
```

Here, we'll use an example data set, which contains the weight of 18 individuals (9 women and 9 men):

```
# Data in two numeric vectors
women_weight <- c(38.9, 61.2, 73.3, 21.8, 63.4, 64.6, 48.4, 48.8, 48.5)
men_weight <- c(67.8, 60, 63.4, 76, 89.4, 73.3, 67.3, 61.3, 62.4)
# Create a data frame
my_data <- data.frame(
  group = rep(c("Woman", "Man"), each = 9),
  weight = c(women_weight, men_weight)
)
```

 We want to know, if the median women's weight differs from the median men's weight?

Check your data

```
print(my_data)
```

```
   group weight
1  Woman   38.9
2  Woman   61.2
3  Woman   73.3
4  Woman   21.8
5  Woman   63.4
6  Woman   64.6
7  Woman   48.4
8  Woman   48.8
9  Woman   48.5
10 Man    67.8
11 Man    60.0
12 Man    63.4
13 Man    76.0
14 Man    89.4
15 Man    73.3
16 Man    67.3
17 Man    61.3
18 Man    62.4
```



It's possible to compute [summary statistics](#) (median and interquartile range (IQR)) by groups. The `dplyr` package can be used.

- To install **dplyr** package, type this:

```
install.packages("dplyr")
```

- Compute summary statistics by groups:

```
library(dplyr)
group_by(my_data, group) %>%
  summarise(
    count = n(),
    median = median(weight, na.rm = TRUE),
    IQR = IQR(weight, na.rm = TRUE)
  )
```

Source: local data frame [2 x 4]

```
   group count median  IQR
  (fctr) (int)  (dbl) (dbl)
1   Man     9   67.3  10.9
2  Woman     9   48.8  15.0
```

Visualize your data using box plots

You can draw R base graphs as described at this link: [R base graphs](#). Here, we'll use the **ggpubr** R package for an easy ggplot2-based data visualization

- Install the latest version of ggpubr from GitHub as follow (recommended):


```
# Install
if(!require(devtools)) install.packages("devtools")
devtools::install_github("kassambara/ggpubr")
```

- Or, install from CRAN as follow:

```
install.packages("ggpubr")
```

- Visualize your data:

```
# Plot weight by group and color by group
library("ggpubr")
ggboxplot(my_data, x = "group", y = "weight",
          color = "group", palette = c("#00AFBB", "#E7B800"),
          ylab = "Weight", xlab = "Groups")
```



Compute unpaired two-samples Wilcoxon test

? Question : Is there any significant difference between women and men weights?

1) Compute two-samples Wilcoxon test - Method 1: The data are saved in two different numeric vectors.

```
res <- wilcox.test(women_weight, men_weight)
res
```

Wilcoxon rank sum test with continuity correction
data: women_weight and men_weight
W = 15, p-value = 0.02712
alternative hypothesis: true location shift is not equal to 0



It will give a warning message, saying that “cannot compute exact p-value with tie”. It comes from the assumption of a Wilcoxon test that the responses are continuous. You can suppress this message by adding another argument **exact = FALSE**, but the result will be the same.

2) Compute two-samples Wilcoxon test - Method 2: The data are saved in a data frame.

```
res <- wilcox.test(weight ~ group, data = my_data,
                  exact = FALSE)
res
```

Wilcoxon rank sum test with continuity correction
data: weight by group
W = 66, p-value = 0.02712
alternative hypothesis: true location shift is not equal to 0

```
# Print the p-value only
res$p.value
```

```
[1] 0.02711657
```



As you can see, the two methods give the same results.



The **p-value** of the test is 0.02712, which is less than the significance level $\alpha = 0.05$. We can conclude that men's median weight is significantly different from women's median weight with a **p-**

value = 0.02712.



Note that:

- if you want to test whether the median men's weight is less than the median women's weight, type this:


```
wilcox.test(weight ~ group, data = my_data,  
            exact = FALSE, alternative = "less")
```

- Or, if you want to test whether the median men's weight is greater than the median women's weight, type this

```
wilcox.test(weight ~ group, data = my_data,  
            exact = FALSE, alternative = "greater")
```

Online unpaired two-samples Wilcoxon test calculator

You can perform unpaired **two-samples Wilcoxon test**, **online**, without any installation by clicking the following link:



[Online two-samples Wilcoxon test calculator](#)

See also

- Compare one-sample mean to a standard known mean
 - [One-Sample T-test \(parametric\)](#)
 - [One-Sample Wilcoxon Test \(non-parametric\)](#)
- Compare the means of two independent groups
 - [Unpaired Two Samples T-test \(parametric\)](#)

Infos



This analysis has been performed using **R software** (ver. 3.2.4).



Enjoyed this article? I'd be very grateful if you'd help it spread by emailing it to a friend, or sharing it on Twitter, Facebook or Linked In.

Show me some love with the like buttons below... Thank you and please don't forget to share and comment below!!

Share 21

Like 21

Tweet


Share

Save


Share

47


Recommended for You!




Machine Learning Essentials:
Practical Guide in R




Practical Guide to Cluster
Analysis in R



Practical Guide to Principal
Component Methods in R



R Graphics Essentials for Great
Data Visualization



Network Analysis and
Visualization in R



More books on R and data
science

Recommended for you



This section contains best data science and self-development resources to help you on your path.

Coursera - Online Courses and Specialization

Data science

- [Course: Machine Learning: Master the Fundamentals](#) by Stanford
- [Specialization: Data Science](#) by Johns Hopkins University
- [Specialization: Python for Everybody](#) by University of Michigan
- [Courses: Build Skills for a Top Job in any Industry](#) by Coursera
- [Specialization: Master Machine Learning Fundamentals](#) by University of Washington
- [Specialization: Statistics with R](#) by Duke University
- [Specialization: Software Development in R](#) by Johns Hopkins University
- [Specialization: Genomic Data Science](#) by Johns Hopkins University

Popular Courses Launched in 2020

- [Google IT Automation with Python](#) by Google
- [AI for Medicine](#) by deeplearning.ai
- [Epidemiology in Public Health Practice](#) by Johns Hopkins University
- [AWS Fundamentals](#) by Amazon Web Services

Trending Courses

- [The Science of Well-Being](#) by Yale University
- [Google IT Support Professional](#) by Google
- [Python for Everybody](#) by University of Michigan
- [IBM Data Science Professional Certificate](#) by IBM
- [Business Foundations](#) by University of Pennsylvania
- [Introduction to Psychology](#) by Yale University
- [Excel Skills for Business](#) by Macquarie University
- [Psychological First Aid](#) by Johns Hopkins University
- [Graphic Design](#) by Cal Arts

Books - Data Science

Our Books

- [Practical Guide to Cluster Analysis in R](#) by A. Kassambara (Datanovia)
- [Practical Guide To Principal Component Methods in R](#) by A. Kassambara (Datanovia)
- [Machine Learning Essentials: Practical Guide in R](#) by A. Kassambara (Datanovia)
- [R Graphics Essentials for Great Data Visualization](#) by A. Kassambara (Datanovia)
- [GGPlot2 Essentials for Great Data Visualization in R](#) by A. Kassambara (Datanovia)
- [Network Analysis and Visualization in R](#) by A. Kassambara (Datanovia)
- [Practical Statistics in R for Comparing Groups: Numerical Variables](#) by A. Kassambara (Datanovia)
- [Inter-Rater Reliability Essentials: Practical Guide in R](#) by A. Kassambara (Datanovia)

Others

- [R for Data Science: Import, Tidy, Transform, Visualize, and Model Data](#) by Hadley Wickham & Garrett Golemund
- [Hands-On Machine Learning with Scikit-Learn, Keras, and TensorFlow: Concepts, Tools, and Techniques to Build Intelligent Systems](#) by Aurelien Géron
- [Practical Statistics for Data Scientists: 50 Essential Concepts](#) by Peter Bruce & Andrew Bruce

- [Hands-On Programming with R: Write Your Own Functions And Simulations](#) by Garrett Golemund & Hadley Wickham
- [An Introduction to Statistical Learning: with Applications in R](#) by Gareth James et al.
- [Deep Learning with R](#) by François Chollet & J.J. Allaire
- [Deep Learning with Python](#) by François Chollet

Want to Learn More on R Programming and Data Science?

Follow us [by Email](#)

by [FeedBurner](#)

On Social Networks:
on Social Networks



Get involved :



Click to **follow us** on [Facebook](#) and [Google+](#) :



Comment this article by clicking on "Discussion" button (top-right position of this page)

This page has been seen 234332 times

Sign in

Login

Password

Auto connect





Register



Forgotten password

Welcome!

Want to Learn More on R Programming and Data Science?

Follow us [by Email](#)

[Subscribe](#)

by [FeedBurner](#)

on Social Networks

[Click to see our collection of resources to help you on your path...](#)

Course & Specialization

Recommended for You (on Coursera):

- [Course: Machine Learning: Master the Fundamentals](#)
- [Specialization: Data Science](#)
- [Specialization: Python for Everybody](#)
- [Course: Build Skills for a Top Job in any Industry](#)
- [Specialization: Master Machine Learning Fundamentals](#)
- [Specialization: Statistics with R](#)
- [Specialization: Software Development in R](#)
- [Specialization: Genomic Data Science](#)

[See More Resources](#)

 **factoextra**

 **survminer**


 **ggpubr**

 **ggcorrplot**

 **fastqcr**


Our Books

3D Plots in R




R Graphics Essentials for Great Data Visualization: 200 Practical Examples You Want to Know for Data Science


★ NEW!!



Practical Guide to Cluster Analysis in R



Practical Guide to Principal Component Methods in R



 **Datanovia: Online Data Science Courses**

 **R-Bloggers**

Newsletter

Email



Boosted by PHPBoost

Recommended for you



ggplot2 legend : Easy
steps to change the
position and the appe...

www.sthda.com



ggplot2 scatter plots :
Quick start guide - R
software and data vis...

www.sthda.com



Reading Data From
TXT|CSV Files: R Base
Functions - Easy Guid...

www.sthda.com

AddThis