

ON A THEOREM OF MINGFENG CHEN, AND THE PROOF OF THE RIEMANN HYPOTHESIS

MINGFENG CHEN, JIAQI HOU, AND YU LUO

Dedicated to Yiyu Wang on his 25th birthday

ABSTRACT. We give a quick application for Chen's structure theorem of abelian groups, to attack the building block of mathematics.

CONTENTS

1. Introduction	1
2. p -adic number revisite	2
3. Proof of main theorem	2
4. Proof of Riemann hypothesis	2
References	2

1. INTRODUCTION

In his celebrated homework [1], Chen proved the following result:

Theorem 1.1. [1] *Given finite abelian groups G and H , we have $G \simeq H$ if and only if $|G| = |H|$.*¹

In this paper, we will give a quick application for this theorem:

Theorem 1.2. *Given any prime number p and q , one has $p = q$. In particular, one has $2 = 3$.*

Corollary 1.3. *Riemann zeta function has trivial analytic continuation, and Riemann hypothesis holds.*

1.4. Acknowledgements. The second author wants to thanks for the video games American Truck Simulator and Sid Meier's Civilization VI which give him happiness for a long time. The third author wants to thank McDonald's for supporting his daily meals.

The work of authors are funded by Mingfeng Chen's sub for Math 340 discussions and lunch. They also would like to thank Mingfeng for sharing his unpublished homework.

Date: February 11, 2024.

¹Zero points for Q1. - Simon Marshall Sep 20 2023

2. p -ADIC NUMBER REVISIT

Theorem 2.1. \mathbb{Z}_p is pure of characteristic p .

Proof. Recall that $\mathbb{Z}_p = \varprojlim \mathbb{Z}/p^n$. By Theorem 1.1, one has $\mathbb{Z}/p^n \simeq (\mathbb{Z}/p)^n$, hence $p\mathbb{Z}_p = 0$. \square

3. PROOF OF MAIN THEOREM

An immediate consequence of Theorem 2.1 is the following:

Corollary 3.1. \mathbb{Q}_p is pure of characteristic p .

Corollary 3.2. \mathbb{C} is pure of characteristic p .

Proof. By the classical identification $\mathbb{C} \simeq \overline{\mathbb{Q}_p}$. \square

Corollary 3.3. For any prime numbers p and q , one has $p = q$.

Proof. One has $\overline{\mathbb{Q}_p} \simeq \overline{\mathbb{Q}_q}$, hence $p = q$ by comparing the characteristic. \square

4. PROOF OF RIEMANN HYPOTHESIS

By Corollary 3.2, the Riemann zeta function is equal to the local zeta function for any prime p , in particular, one has

$$(4.0.1) \quad \zeta(s) = \frac{1}{1 - 2^{-s}}.$$

Hence it has trivial analytic continuation, and there is no zeros in the plane. As a result, all nontrivial zeros lie in $\text{Re}(s) = \frac{1}{2}$. \square

REFERENCES

- [1] Mingfeng, Chen, Math 749 Analytic number theory Homework 1, unpublished note. 1

UNIVERSITY OF WISCONSIN-MADISON, DEPARTMENT OF MATHEMATICS, 480 LINCOLN DRIVE, MADISON, WI 53706, USA

Email address: `mchen454@wisc.edu`

UNIVERSITY OF WISCONSIN-MADISON, DEPARTMENT OF MATHEMATICS, 480 LINCOLN DRIVE, MADISON, WI 53706, USA

Email address: `jhou39@wisc.edu`

UNIVERSITY OF WISCONSIN-MADISON, DEPARTMENT OF MATHEMATICS, 480 LINCOLN DRIVE, MADISON, WI 53706, USA

Email address: `yluo237@wisc.edu`