

Classifying Space and Bar Construction

Taught by Yuri Berest

Notes by Yun Liu
yl2649@cornell.edu

Cornell University
Fall, 2022

Last updated April 25, 2024.

Contents

1	Introduction	5
1.1	Outline	5
2	Category Theory	6
2.1	Small Object Argument	6
2.2	Quillen functors and Quillen Equivalences	8
3	Model Categories Generated by Small Categories	10
3.1	Motivation	10
3.2	Universal Homotopy Categories	11
3.3	Bousfield-Kan Formula for Homotopy Colimits	13
3.3.1	Classical Nerves	13
3.3.2	Categorical Realization	14
3.3.3	Homotopy Inverse of \mathcal{N}_*	14
3.4	BK formula	15
3.4.1	Homotopy Inverse of Nerves	15
4	Cyclic Objects	18
4.1	Cyclic Category	18
4.1.1	Crossed Group Categories	18
4.1.2	Dwyer-Hopkins-Kan presentation	18
4.2	Cyclic Homology	20
4.2.1	Quillen's Ayclisim Formalism	22
4.2.2	Cyclic (Co)homology and Algebra Extensions	23
4.2.3	Chern-Simons Formalism for Algebra Cochains	24
4.3	Functor Homology	24
4.3.1	Hochschild Homology	25
4.3.2	Cyclic Homology	26
4.4	Topological Interpretation of Cyclic Homology	29
5	Homotopy Colimits	37
5.1	Homotopy Kan Extension	37
5.2	Grothendieck Construction	39
5.2.1	Universal Mapping Property	40
5.2.2	Fibered/cofibered Categories	41
5.2.3	Fibered/Cofibered Functors	41
5.3	Thomason's Formula	43
5.4	Homotopy Coends	45
5.4.1	Twisted Arrow Category / Factorization Category	45
6	Simplicial Bar Construction	48

6.1	Tensor and Cotensor	48
6.2	Functor Tensor Products	49
6.3	Two-sided Simplicial Bar Construction	50
6.3.1	Examples	51
6.4	Cyclic Bar Construction of a Discrete Group	52
6.4.1	Simplicial Models of Circles	55
6.5	Nerves	56
6.5.1	Twisted Nerve of a Group	56
6.5.2	Crossed Modules	57
6.6	Cyclic Nerve	63
6.6.1	Burghelea Decomposition	65
6.6.2	Cyclic Homology of Group Algebras	69
6.7	Homotopy Simplicial Groups	70
6.7.1	Cyclic Homology	72
7	Realizable (Voedvodsky) Homotopy Colimits	73
7.1	Relative Categories	73
7.2	2-Category	77
8	Stable Homotopy Theory	82
8.1	Motivation Results	82
8.2	Γ -Spaces and Homotopy-theoretic generalization of Quillen homology	84

Contents by Lecture

Lecture 01 on 22 August 2022	5
Lecture 02 on 24 August 2022	5
Lecture 03 on 29 August 2022	5
Lecture 04 on 31 August 2022	6
Lecture 05 on 07 September 2022	9
Lecture 06 on Bousfield-Kan Formula for Homotopy Colimits	13
Lecture 07 on 14 September 2022	14
Lecture 08 on 19 September 2022	18
Lecture 09 on 21 September 2022	20
Lecture 10 on 26 Sep 2022	24
Lecture 11 on 28 September 2022	29
Lecture 12 on 03 October 2022	32
Lecture 13 on 05 October 2022	36
Lecture 14 on 12 October 2022	39
Lecture 15 on 17 October 2022	44
Lecture 16 on 19 October 2022	47
Lecture 17 on 24 October 2022	51
Lecture 18 on 26 October 2022	53
Lecture 19 on 31 October 2022	56
Lecture 20 on 2 November 2022	59
Lecture 21 on 7 November 2022	62
Lecture 22 on 14 November 2022	64
Lecture 23 on 16 November 2022	69

Lecture 24 on 21 November 2022	71
Lecture 25 on 28 November 2022	73
Lecture 26 on 30 November 2022	77
Lecture 27 on 05 December 2022	80
Lecture 28 on 07 December 2022	82
Lecture 29 on 09 December 2022	85

1 Introduction

1.1 Outline

1. Nerves of discrete groups.
2. Classifying space of compact Lie groups and finite loop spaces.
3. Nerves of categories.

2 Category Theory

2.1 Small Object Argument

Correction Let \mathcal{M} be a model category and $I \subset \text{Mor}(\mathcal{M})$ a set of maps.

Definition 2.1. I-fibration = the class of maps in \mathcal{M} which has RLP with respect to I .

I-cofibrations = LLP with respect to I-fib.

Definition 2.2 (Hir, 11.1.2). A model category \mathcal{M} is cofibrantly generated if

1. there is a set of generating cofibrations morphisms $I \subset \text{Mor}(\mathcal{M})$
 - (a) I permits SOA.
 - (b) $\text{RLP}(I) = \text{Fib}(\mathcal{M}) \cap \mathcal{W}$.
2. there is a set of generating acyclic cofibrations J such that
 - (a) J permits SOA.
 - (b) $\text{RLP}(J) = \text{Fib}(\mathcal{M})$.

Theorem 2.3 (Recognition Theorem). Assume $(\mathcal{M}, \mathcal{W})$ satisfies (MC1), (MC2), (MC3). Assume I, J

1. SOA.
2. $J - \text{cof} \subseteq I - \text{cof} \cap \mathcal{W}$
3. $I - \text{Fib} \subseteq J - \text{Fib} \cap \mathcal{W}$
4. either (2) or (3) is equality.

Then $(\mathcal{M}, I, J, \mathcal{W})$ is a cofibrantly generated model category with

- weak equivalences \mathcal{W} ,
- fibrations = $\text{RLP}(J)$,
- cofibrations = $\text{LLP}(\text{Fib} \cap \mathcal{W})$.

Theorem 2.4 (Promoting Model Structure). Let $(\mathcal{M}, I, J, \mathcal{W})$ be a cofibrantly generated model category. Let \mathcal{N} be a bicomplete category such that there is a pair of adjoint functors

$$F : \mathcal{M} \rightleftarrows \mathcal{N} : U$$

Define $F(I) = \{F(f) | f \in I\}$ and $F(J) = \{F(f) | f \in J\}$. Assume

1. $F(I)$ and $F(J)$ permits soa.
2. U maps relative $F(J)$ -cell complexes to weak equivalences in \mathcal{M} .

then $(\mathcal{N}, F(I), F(J), \mathcal{W}_{\mathcal{N}})$ where $\mathcal{W}_{\mathcal{N}} = \{f \in \text{Mor}(\mathcal{N}) | Uf \in \mathcal{W}_{\mathcal{M}}\}$ is cofibrantly generated model category. Moreover (F, U) is a Quillen pair.

Notation Let \mathcal{M} be a model category and $I \subset \text{Mor}(\mathcal{M})$, \mathcal{C} a small category.
 $\mathcal{M}^{\mathcal{C}} = \text{Fun}(\mathcal{C}, \mathcal{M})$.

$$F_I^{\mathcal{C}} = \{\varphi \in \text{Mor}(\mathcal{M}^{\mathcal{C}}) \mid \varphi_c \in I, \forall c \in \text{Ob}(\mathcal{C})\}.$$

Theorem 2.5 (Hir, 11.6.17). *Let (\mathcal{M}, I, J, W) be a cofibrantly generated model category, and let \mathcal{C} be any small category, then $(\mathcal{M}^{\mathcal{C}}, F_I^{\mathcal{C}}, F_J^{\mathcal{C}}, F_W^{\mathcal{C}})$ is a cofibrantly generated model category with objectwise weak equivalences and objectwise fibrations.*

Sketch of proof. Based on Theorem 1 and 2.

Step 1. Let $i : \mathcal{C}^{\delta} \rightarrow \mathcal{C}$ be the discrete subcategory of \mathcal{C} , with

- objects are objects in \mathcal{C}
- morphisms are $\text{Hom}_{\mathcal{C}^{\delta}}(c, c) = \{\text{Id}_c\}, \text{Hom}_{\mathcal{C}^{\delta}}(c, d) = \emptyset, c \neq d$.

then $\mathcal{M}^{\mathcal{C}^{\delta}} = \prod_{c \in \text{Ob}(\mathcal{C})} \mathcal{M}$ is cofibrantly generated model category.

Step 2. The forgetful functor $U = i^* : \mathcal{M}^{\mathcal{C}} \rightarrow \mathcal{M}^{\mathcal{C}^{\delta}}$ has a left adjoint, the free diagram functor

$$F : \mathcal{M}^{\mathcal{C}^{\delta}} \longrightarrow \mathcal{M}^{\mathcal{C}}$$

$$F(X) := \coprod_{c \in \text{Ob}(\mathcal{C})} X(c) \otimes F_*^c : \mathcal{C} \rightarrow \mathcal{M} \text{ so that for any } d \in \text{Ob}(\mathcal{C}),$$

$$F(X)(d) = \coprod_{c \in \text{Ob}(\mathcal{C})} \coprod_{\text{Hom}(c, d)} X(c)$$

where for $S : \mathcal{C} \rightarrow \mathbf{Set}$,

$$S \otimes \mathcal{M} : \mathcal{C} \longrightarrow \mathcal{M}$$

$$c \longmapsto S(c) \otimes A = \coprod_{c \in \text{Ob}(\mathcal{C})} A$$

here

$$S = F_*^c : \mathcal{C} \longrightarrow \mathbf{Set}$$

$$d \longmapsto \text{hom}_{\mathcal{C}}(c, d)$$

Step 3. Check conditions of Theorem 2 for (F, U) . □

2.2 Quillen functors and Quillen Equivalences

Lemma 2.6. *A Quillen pair*

$$F : \mathcal{M} \rightleftarrows \mathcal{N} : U$$

is an adjunction such that one of the following equivalent conditions hold.

1. *F preserves cofibrations and U preserves fibrations.*
2. *F preserves cofibrations and acyclic cofibrations.*
3. *G preserves fibrations and acyclic fibrations.*

Definition 2.7. A **Quillen equivalence** is a Quillen pair

$$F : \mathcal{M} \rightleftarrows \mathcal{N} : U$$

such that for any $X \in \text{Ob}(\mathcal{M}^{\text{cof}})$ and $A \in \text{Ob}(\mathcal{M}^{\text{fib}})$, $f : F(X) \rightarrow A$ is a weak equivalence in \mathcal{N} if and only if $f^\# : X \rightarrow U(A)$ is a weak equivalence in \mathcal{M} , where

$$\text{hom}_{\mathcal{N}}(FX, A) \xrightarrow{\cong} \text{hom}_{\mathcal{M}}(X, UA)$$

$$f \longmapsto f^\#$$

Remark. A map $\mathcal{M} \rightarrow \mathcal{N}$ is a left Quillen functor.

Definition 2.8 (Dugger). A Quillen homotopy between (left) Quillen functors $F_1, F_2 : \mathcal{M} \rightarrow \mathcal{N}$, is a natural transformation $\varphi : F_1 \Rightarrow F_2$ such that for any $X \in \text{Ob}(\mathcal{M}^{\text{cof}})$, $\varphi_X : F_1(X) \rightarrow F_2(X)$ are weak equivalences.

Two functors $F, F' : \mathcal{M} \rightarrow \mathcal{N}$ are **Quillen homotopic** if there is a chain of Quillen homotopies

$$F = F_0 \Leftarrow F_1 \Rightarrow \cdots \Leftarrow F_n = F'$$

Derived Functors $F : \mathcal{M} \rightarrow \mathcal{N}$ a functor between model categories.

Total left derived functors are right Kan extensions

Alternative Characterization Left derived functor

$LF : \mathcal{M} \rightarrow \text{Ho}(\mathcal{N})$ is a universal homotopical functor if there is a natural transformation $\varepsilon : LF \Rightarrow \gamma_{\mathcal{M}} \circ F$ which is terminal among all homotopical functors to $\gamma_{\mathcal{M}} \circ F$

$$LF = \mathbb{F} \circ \gamma_{\mathcal{M}}.$$

Theorem 2.9 (Quillen Adjunction Theorem). 1. Any Quillen pair

$$F : \mathcal{M} \rightleftarrows \mathcal{N} : U$$

induces an adjunction of homotopy categories

$$LF : \mathrm{Ho}(\mathcal{M}) \rightleftarrows \mathrm{Ho}(\mathcal{N}) : RU$$

where $LF(\bar{X}) = F(QX)$ where $QX \xrightarrow{\sim} X$ is a cofibrant replacement and $RG(\bar{A}) = G(RA)$ where $A \xrightarrow{\sim} RA$ is a fibrant replacement.

2. If (F, U) is a Quillen equivalence, then

$$LF : \mathcal{M} \rightleftarrows \mathcal{N} : RU$$

is an equivalence of categories.

Next time.

1. Universal model categories.
2. Model approximation.

We say a map $L : \mathcal{M} \rightarrow \mathcal{N}$ between model categories to be a Quillen pair

$$L : \mathcal{M} \rightleftarrows \mathcal{N} : R.$$

3 Model Categories Generated by Small Categories

3.1 Motivation

Let \mathcal{C} be a small category and let $\text{Pre}(\mathcal{C}) = \text{Fun}(\mathcal{C}^{\text{op}}, \mathbf{Set})$ be the category of presheaves on \mathcal{C} .

Remark. When X is a topological space, $\mathcal{C} = \text{Open}(X)$ the poset of open sets of X , then any functor $F : \mathcal{C}^{\text{op}} \rightarrow \mathbf{Set}$ is equivalent to a presheaf on X .

The category $\text{Pre}(\mathcal{C})$ has the UMP of being “free” cocomplete categories generated by \mathcal{C} .

Proposition 3.1. 1. *There is a natural fully faithful functor*

$$h : \mathcal{C} \hookrightarrow \text{Pre}(\mathcal{C})$$

$$X \mapsto (h_X : Y \mapsto \text{Hom}_{\mathcal{C}}(Y, X))$$

satisfying the following universal property: given any cocomplete category \mathcal{D} with

$$\begin{array}{ccc} \mathcal{C} & \xrightarrow{\gamma} & \mathcal{D} \\ \downarrow & \nearrow \text{Re} & \\ \text{Pre}(\mathcal{C}) & & \end{array}$$

there is a functor $\text{Re} : \text{Pre}(\mathcal{C}) \rightarrow \mathcal{D}$ such that $\text{Re} \circ h \xrightarrow{\sim} \gamma$ which is unique up to isomorphism.

2. $\text{Re} : \text{Pre}(\mathcal{C}) \rightarrow \mathcal{D}$ has a right adjoint Sing .

Proof. Note every presheaf on \mathcal{C} can be canonically expressed as a colimit of representable presheaves in $\text{Pre}(\mathcal{C})$.

Recall the Kan extension construction

$$\begin{array}{ccc} \mathcal{C} & \xrightarrow{\gamma} & \mathcal{D} \\ \downarrow G & \nearrow & \\ \mathcal{E} & & \end{array}$$

when \mathcal{D} is cocomplete,

$$\text{Lan}_G(\gamma)(e) = \text{colim}_{G \downarrow e} (G \downarrow e \xrightarrow{\pi} \mathcal{C} \xrightarrow{\gamma} \mathcal{D})$$

where $G \downarrow e$ is the slice category. In particular when G is fully faithful, $\text{Lan}_G(\gamma) \circ G \cong \gamma$.

For any presehaf $\mathcal{F} \in \text{Pre}(\mathcal{C})$,

$$\mathcal{F} \cong \text{colim}_{\mathcal{C} \downarrow \mathcal{F}} (\mathcal{C} \downarrow \mathcal{F} \xrightarrow{\pi} \mathcal{C} \xrightarrow{h} \text{Pre}(\mathcal{C})).$$

Similarly,

$$\text{Re}(\mathcal{F}) = \text{colim}_{\mathcal{C} \downarrow \mathcal{F}} (\pi^* \gamma).$$

This can be viewed as a realization of formal colimits in \mathcal{D} .

The right adjoint of Re is given by

$$\begin{aligned} \text{Sing} : \mathcal{D} &\longrightarrow \text{Pre}(\mathcal{C}) \\ d &\longmapsto (\text{Sing}(d) : X \mapsto \text{Hom}_{\mathcal{D}}(\gamma X, d)) \end{aligned}$$

□

Example 3.2. 1. geometric realization

2. nerve of small categories. categorization

Remark. This construction extends to an equivalence of categories

$$\mathcal{D}^{\mathcal{C}} \simeq \text{Adj}(\text{Pre}(\mathcal{C}), \mathcal{D}).$$

Goal: a generalization (homotopical completion of \mathcal{C}) by adding all homotopy colimits.

3.2 Universal Homotopy Categories

Given a small category \mathcal{C} and two model categories \mathcal{M} and \mathcal{N} together with two functors

$$\begin{array}{ccc} \mathcal{C} & \xrightarrow{\gamma} & \mathcal{M} \\ \downarrow r & & \\ \mathcal{N} & & \end{array}$$

we define the factorization of γ through r to be a triple (L, R, η) where

1. $L : \mathcal{N} \rightleftarrows \mathcal{M} : R$ is a Quillen pair.
2. $\eta : L \circ r \Rightarrow \gamma$ is a natural weak equivalence.

The category of factorizations $\text{Fact}_{\mathcal{N}}(\gamma)$ is defined by

- objects are triples (L, R, η) , and

- morphisms are natural transformations $\alpha : L \rightarrow L'$ such that

$$\begin{array}{ccc} L \circ r & \xrightarrow{\quad} & L' \circ r \\ & \searrow \quad \swarrow & \\ & \gamma & \end{array}$$

Definition 3.3. Let $\mathcal{UC} = \mathbf{sPre}(\mathcal{C}) = \mathbf{Fun}(\mathcal{C}^{\mathrm{op}}, \mathbf{sSet})$ be the category of simplicial presheaves equipped with projective model structure.

There is a natural functor

$$r : \mathcal{C} \xhookrightarrow{h} \mathbf{Pre}(\mathcal{C}) \longrightarrow \mathbf{sPre}(\mathcal{C})$$

$$X \longmapsto h_X \longmapsto h_X : [n] \mapsto h_X$$

Theorem 3.4 (Dugger). $(\mathcal{UC}, r : \mathcal{C} \rightarrow \mathcal{UC})$ is the universal model category generated by \mathcal{C} in the sense

1. any functor $\gamma : \mathcal{C} \rightarrow \mathcal{M}$ from \mathcal{C} to a model category \mathcal{M} factors

$$\begin{array}{ccc} \mathcal{C} & \xrightarrow{\gamma} & \mathcal{M} \\ \downarrow r & \nearrow & \\ \mathcal{UC} & & \end{array}$$

up to natural weak equivalences $\eta : \mathrm{Re} \circ r \xrightarrow{\mathrm{sim}} \gamma$.

2. Moreover, then factorization is homotopically unique in the sense that $\mathbf{Fact}_{\mathcal{UC}}(\mathcal{UC})$ is contractible, i.e. $\mathbf{BFact}_{\mathcal{UC}}(\mathcal{UC}) \simeq \mathbf{pt}$.

Remark. Notice that a category being contractible is a natural generalization of the property of being unique up to unique isomorphism.

Remark. Given a factorization is equivalent to give a cosimplicial resolution of diagrams $\mathcal{C} \xrightarrow{\gamma} \mathcal{M}$.

Next step is to construct a set of maps S in \mathcal{UC} such that

$$\mathrm{Re} : \mathcal{UC}/S = \mathcal{L}_S(\mathcal{UC}) \xrightleftharpoons{\quad} \mathcal{M} : \mathbf{Sing}$$

is a Quillen equivalence.

Example 3.5 (Homotopy Simplicial Groups). Let $\mathcal{G} \subset \mathbf{Gr}$ be the full subcategory of finite generated free groups, then $\mathbf{Gr} \simeq \mathbf{Pre}^{\otimes}(\mathcal{G})$ product-preserving presheaves. Then $\mathbf{sGr} \simeq \mathbf{sPre}^{\otimes}(\mathcal{G})$.

Definition 3.6. A homotopy simplicial group $\Gamma : \mathcal{G}^{\text{op}} \rightarrow \mathbf{sSet}$ is one such that

$$\Gamma(\langle n \rangle) \simeq \prod_n \Gamma(\langle 1 \rangle)$$

Theorem 3.7 (Badzioch). *There is a Quillen equivalence $\mathbf{sGr}^h \mathbf{sGr}$.*

3.3 Bousfield-Kan Formula for Homotopy Colimits

Recall given a small category \mathcal{C} together with $\text{Pre}(\mathcal{C})$ the universal category generated by \mathcal{C}

$$\begin{array}{ccc} \mathcal{C} & \xrightarrow{\quad} & \mathcal{D} \\ \downarrow h & \nearrow \text{Re} & \nearrow \text{Sing} \\ \text{Pre}(\mathcal{C}) & & \end{array}$$

Key fact: denote $\mathcal{C} \downarrow \mathcal{F} = h \downarrow \mathcal{F}$, then

$$\pi^* h : \mathcal{C} \downarrow \mathcal{F} \xrightarrow{\pi} \mathcal{C} \xrightarrow{h} \text{Pre}(\mathcal{C})$$

we have $\mathcal{F} \cong \text{colim}_{\mathcal{C} \downarrow \mathcal{F}} (\pi^* h)$.

Consider

$$\begin{array}{ccc} \mathcal{C} & \xrightarrow{\gamma} & \mathcal{M} \\ \downarrow h & \nearrow \text{Re} & \nearrow \text{Sing} \\ \mathcal{UC} & & \end{array}$$

Remark. In general, we cannot ask for η to be a natural isomorphism. If γ takes X to $\gamma(X)$ not cofibrant, then the above diagram cannot commute. Note $r(X)$ is cofibrant in \mathcal{UC} , thus $\text{Re}(\gamma(X))$ is cofibrant.

We want a universal expression of simplicial presheaves in terms of representable presheaves rX . The answer is BK formula for homotopy colimits.

3.3.1 Classical Nerves

The simplex category Δ
 nonstrictly order preserving maps
 generators: coface codegeneracy
 Simplicial relations

Simplicial Sets presentation X_* with face and degeneracy maps.

Example 3.8 (Standard n -simplex). $\Delta[n]_*$ corepresents a functor

$$(-)_n : \mathbf{sSet} \longrightarrow \mathbf{Set}$$

$$X_* \longmapsto X_n$$

3.3.2 Categorical Realization

$$\Delta \hookrightarrow \mathbf{Cat}$$

$$[n] \mapsto \vec{n}$$

$$c : \mathbf{sSet} \rightleftarrows \mathbf{Cat} : \mathcal{N}$$

description of \mathcal{N}_* :

description of c : the free category generated by the graph at level ≤ 2

Exercise 3.9. c is a left inverse, i.e. \mathcal{N}_* is fully faithful.

3.3.3 Homotopy Inverse of \mathcal{N}_*

Construction Category of elements (BK construction) Given \mathcal{C} a small category and $X : \mathcal{C} \rightarrow \mathbf{Set}$ $\mathcal{C}_{\mathcal{F}} = \mathcal{C} \int \mathcal{F}$

Definition 3.10. $\mathrm{hocolim}_{\mathcal{C}}^{\mathbf{sSet}}(\mathcal{F}) := \mathcal{N}_*(\mathcal{C}_{\mathcal{F}})$

Definition 3.11. For $\mathcal{C} \in \mathrm{Ob}(\mathbf{Cat})$, $B\mathcal{C} := |\mathcal{N}_*\mathcal{C}|$.

Definition 3.12. $\mathrm{hocolim}_{\mathcal{C}}^{\mathbf{Top}}(\mathcal{F}) := B(\mathcal{C}_{\mathcal{F}})$.

Exercise 3.13. $X_* : \Delta^{\mathrm{op}} \rightarrow \mathbf{sSet}$. There is a weak homotopy equivalence $|X| \simeq \mathrm{hocolim}_{\Delta^{\mathrm{op}}} (X_*) =: \|X\|$.

next time: $\Delta C, \Delta G$,
Borel fiber sequence

$$\|X\| \longrightarrow \|X\|^{\mathrm{cy}} \simeq ES^1 \times_{S^1} |X| \longrightarrow BS^1$$

Example 3.14.

$$\mathrm{Ad} : \underline{G} \longrightarrow \mathbf{Set}$$

$$* \longrightarrow G$$

$$g \in G \longmapsto \mathrm{Ad}_g : G \rightarrow G$$

$\mathrm{hocolim}_G^{\mathbf{Top}}(\mathrm{Ad}) \simeq \mathcal{L}(BG)$
cyclic nerve of G

3.4 BK formula

There is a Quillen adjunction

$$|-| : \mathbf{sSet} \rightleftarrows \mathbf{Top} : \text{Sing}$$

There is an equivalence of homotopy categories

$$\text{Ho}(\mathbf{sSet}) \simeq \text{Ho}(\mathbf{Top})$$

Example 3.15 (Classifying Space, or simplicial nerves). Consider the composite

$$B : \mathbf{Cat} \xrightarrow{\mathcal{N}_*} \mathbf{sSet} \xrightarrow{|-|} \mathbf{Top}$$

$$\mathcal{C} \longmapsto \mathcal{N}_* \mathcal{C} \longmapsto B\mathcal{C}$$

There are two generalizations

1. Is there a similar construction when we replace **Top** by $G - \mathbf{Top}$? What is such ΔG ?

There is one example for $G = S^1$ with ΔC the cyclic categories. And this leads to the concept of crossed simplicial groups.

endieck, Cisinski (07'), Maltsionitis How to characterize small categories \mathcal{A} that behave like Δ such that $\text{Ho}(\text{Pre}(\mathcal{A})) \simeq \text{Ho}(\mathbf{Top})$?

Apart from Δ , there is also cubical category \square . (See Voevodsky's lectures in Cambridge about cubical sets, IN Institute.)

3.4.1 Homotopy Inverse of Nerves

Recall the category of elements \mathcal{C}_F in $F : \mathcal{C} \rightarrow \mathbf{Set}$.

Example 3.16. Let $X : \Delta^{\text{op}} \rightarrow \mathbf{Set}$ be a simplicial set. Then Δ_X^{op} is the category with

- objects are $([n], x)$ where $[n] \in \text{Ob}(\Delta^{\text{op}})$ and $x \in X_n$, and
- morphisms $f : ([m], y) \rightarrow ([n], x)$ are $f \in \text{Mor}_{\Delta^{\text{op}}}([m], [n])$ such that $X(f)(y) = x$.

There is an alternative construction. Consider

$$h : \Delta \hookrightarrow \mathbf{sSet}$$

$$[n] \mapsto \Delta[n]_*$$

Define $\Delta X := h \downarrow X = \Delta \downarrow X$ the category of simplices of X , then

- objects are $([n], \chi)$ where $[n] \in \text{Ob}(\Delta)$ and $\chi : \Delta[n]_* \rightarrow X$, or equivalently, objects are elements in X_n , $\forall n \geq 0$.
- morphisms $\varphi : ([n], \chi) \rightarrow ([m], \psi)$ are $\varphi \in \text{Mor}_\Delta([n], [m])$ such that $\psi \circ \Delta\varphi = \chi$.

By Yoneda lemma, we can see that $\Delta_X^{\text{op}} \cong (\Delta X)^{\text{op}}$.

Exercise 3.17. Consider a small category \mathcal{C} and take $\Delta\mathcal{C} := \Delta(\mathcal{N}_*\mathcal{C})$, consider $\Delta : \Delta \rightarrow \mathbf{Cat}$ then

$$\Delta\mathcal{C} \cong \Delta \downarrow \mathcal{C}.$$

Define

$$W_\infty := \{F : \mathcal{C} \rightarrow \mathcal{D} \mid B\mathcal{C} \simeq B\mathcal{D}\}$$

(the most fundamental localizers)

then there are equivalences of homotopy categories

$$\overline{\mathcal{N}} : \text{Ho}(\mathbf{Cat}) \simeq \text{Ho}(\mathbf{sSet}) \simeq \text{Ho}(\mathbf{Top})$$

Theorem 3.18 (Latch, 1987). *The functor*

$$i_\Delta : \mathbf{sSet} \rightarrow \mathbf{Cat}$$

$$X \longmapsto \Delta X$$

is a homotopy inverse of $\overline{\mathcal{N}}$ (but not adjoint to \mathcal{N}_*).

Idea Look at analogue of i_Δ for Δ replaced by \mathcal{A} .

Notation: Let \mathcal{A} be a small category and write $\mathcal{A} - \mathbf{Set} = \text{Pre}(\mathcal{A})$.

$$i_{\mathcal{A}} : \mathcal{A} - \mathbf{Set} \longrightarrow \mathbf{Cat}$$

$$X \longmapsto i_{\mathcal{A}}(X) := h^{\mathcal{A}} \downarrow X$$

The functor $i_{\mathcal{A}}$ has a right adjoint

$$i_{\mathcal{A}} : \mathcal{A} - \mathbf{Set} \rightleftarrows \mathbf{Cat} : i_{\mathcal{A}}^*$$

Let $\varepsilon_{\mathcal{A}} : i_{\mathcal{A}}^* i_{\mathcal{A}} \Rightarrow \text{Id}_{\mathbf{Cat}}$ be the counit and $\varepsilon_{\mathcal{C}} : i_{\mathcal{A}}^* i_{\mathcal{A}} \rightarrow \mathcal{C}$.

Definition 3.19 (Grothendieck). \mathcal{A} is called a **test category** if

1. $B\mathcal{A} \simeq \mathbf{pt}$ is contractible.
2. For any $\mathcal{C} \in \text{Ob}(\mathbf{Cat})$ and for any $c \in \text{Ob}(\mathcal{C})$, $B(\varepsilon_{\mathcal{C}} \downarrow \mathcal{C}) \simeq \mathbf{pt}$.

Theorem 3.20 (Cisinski). *For any test category \mathcal{A} , the category $\mathcal{A} - \mathbf{Set}$ has a closed model category structure with*

- $\mathbf{Cof} =$ injective maps of $\mathcal{A}^{\mathbf{op}}$ -diagrams of sets.
- $\mathbf{We} = \{f : X \rightarrow Y \text{ in } \mathcal{A} - \mathbf{Set} \text{ such that } B(i_{\mathcal{A}}f) : B(i_{\mathcal{A}}X) \xrightarrow{\sim} B(i_{\mathcal{A}}Y) \text{ in } \mathbf{Top}\}.$

There is a Quillen equivalence

$$i_{\mathcal{A}} : \mathcal{A} - \mathbf{Set} \rightleftarrows \mathbf{Cat} : i_{\mathcal{A}}^*$$

See [1].

Definition 3.21. The fundamental localizers $W = \{F : \mathcal{C} \rightarrow \mathcal{D}\}$ (the classes of weak equivalences) such that

1. 2-of-3 property
2. retraction property
3. $\{\mathcal{C} \rightarrow \mathbf{pt}\} \in W$ whenever \mathcal{C} has a terminal objects.

Conjecture 3.22. $W_{\infty} = \bigcap \{\text{all fundamental localizers } W\}$

4 Cyclic Objects

4.1 Cyclic Category

Definition 4.1. ΔC has the same objects as Δ

1. objects $[n], n \geq 0$.
 2. there is an embedding $\Delta \hookrightarrow \Delta C$ such that Δ is a subcategory of ΔC .
 3. $\text{Aut}_{\Delta C}([n]) = C_{n+1} = \mathbb{Z}/(n+1)\mathbb{Z}$
 4. unique factorization (crossed relations)
- crossed group category, Berger and M... 2014

Exercise 4.2. $\Delta C \cong \Delta^{\text{op}}$. Connes duality

Cyclic Homology $\text{HC}_*(k[\Gamma]) \cong H_*(\text{ES}^1 \times_{S^1} \mathcal{L}(B\Gamma), k)$.

4.1.1 Crossed Group Categories

Recall we have the following equivalence of homotopy categories

$$\text{Ho}(\mathbf{sSet}) \simeq \text{Ho}(\mathbf{Top})$$

and we 'd like to generalize to see if there is any equivalence of homotopy categories

$$\text{Ho}(\mathbf{Set}^{\Delta G}) \simeq \text{Ho}(G - \mathbf{Top})$$

4.1.2 Dwyer-Hopkins-Kan presentation

Definition 4.3 (Duplicial Category). The category K^{op} is an amalgamation of Δ^{op} and Δ by identifying (in each dimension) the degeneracy operators $s_i \in \Delta^{\text{op}}$ with all but one coface operator $d^i \in \Delta$ and the codegeneracy operators s^j with all but one face operator $d_j \in \Delta^{\text{op}}$. Explicitly, K^{op} is defined by

- objects are the same as Δ , i.e. finite natural numbers $[n], n \geq 0$,
- morphism are generated by d_i, s_j subject to the relations

$$\begin{aligned} d_i d_j &= d_{j-1} d_i, i \leq j \\ s_j s_i &= s_i s_{j-1}, i \leq j \\ d_i s_j &= s_{j-1} d_i : [n], 0 < j-1 \leq n, \\ &= \text{Id} - 1 \leq j-i \leq 0, \\ s_j d_{i-1} &, j-i < -1. \end{aligned}$$

Note $d_0 s_{n+1} \neq s_n d_0$.

Observe that there are two natural functors

$$j : \Delta^{\text{op}} \hookrightarrow \mathcal{K}^{\text{op}}$$

and an isomorphism

$$m : \mathcal{K}^{\text{op}} \rightarrow \mathcal{K}$$

$$d_i \mapsto s_{n-i}$$

$$s_i \mapsto d_{n-i}$$

Proposition 4.4. 1. *There is a curious endomorphism of $\text{Id} : \mathcal{K}^{\text{op}} \rightarrow \mathcal{K}^{\text{op}}$*

$$v = \{v_n = d_0 s_{n+1}^{n+1} : [n] \rightarrow [n]\}$$

2. *DHK presentation: $\Delta C^{\text{op}} \cong \mathcal{K}^{\text{op}} / (v_n, n \geq 0)$ with a functor*

$$\mathcal{K}^{\text{op}} \longrightarrow \Delta C^{\text{op}}$$

$$d_0 s_{n+1} \mapsto t_{n+1} = \tau_n^{\text{op}}$$

$$d_i \longmapsto d_i$$

$$s_j \longmapsto s_j$$

“Analogy” Δ or $\Delta^{\text{op}}, \mathcal{K} \longleftrightarrow$ “differential operators” on varieties \longleftrightarrow Differential operators.

$d_i \longleftrightarrow$ “coordinates” on varieties $q_i = x_i$, Elliptic Hall Algebras

$s_j \longleftrightarrow$ “derivations” in $p^i = \frac{\partial}{x_i}$

Dold-Kan correspondence \longleftrightarrow Cannings-Holland Correspondence, Derived Hall Algebra

Crossed simplicial groups \longleftrightarrow Double affine Hecke algebras

Theorem 4.5. ΔC is generated by

1. $\text{Aut}_{\Delta C}([n]) =: C_n \cong \mathbb{Z}/(n+1), n \geq 0$
2. *any morphism $f : [n] \rightarrow [n]$ factors uniquely as $f = \varphi \circ \gamma$ where $\gamma \in C_n$ and $\varphi \in \text{Mor}(\Delta)$.*

The assignment $[n] \mapsto C_n$ extends to a functor

$$C_* : \Delta C^{\text{op}} \rightarrow \mathbf{Set}$$

$$[n] \longmapsto C_n$$

$$a \longmapsto a_*$$

where $a_* : C_n \rightarrow C_m$

Example 4.6 (Cocyclic Space). $\tau_n : (x_0, x_1, \dots, x_n) \mapsto (x_1, \dots, x_n, x_0)$.

Theorem 4.7. 1. Let $X : \Delta C^{\text{op}} \rightarrow \mathbf{Top}$ be a cyclic space, then $|X|$ the geometric realization of X as a simplicial space carries a natural S^1 -action.

2. It extends to a functor

$$\mathbf{Top}^{\Delta C^{\text{op}}} \longrightarrow S^1 - \mathbf{Top}$$

3. $HC_*(k[X]) \cong H_*^{S^1}(|X|, k)$.

4.2 Cyclic Homology

Let ΔC be Connes' category.

Let k be a commutative ring.

Consider the following functor

$$C_* : \Delta C^{\text{op}} \rightarrow \text{Mod}_k$$

$$[n] \longmapsto C_n$$

be a cyclic module. Define $\tilde{t}_n := (-1)^n t_n$, it comes with relations

$$\tilde{t}_n^{n+1} = \text{Id}$$

$$d_i \tilde{t}_n = -\tilde{t}_{n-1} d_{i-1}, 1 \leq i \leq n$$

$$s_i \tilde{t}_n = -\tilde{t}_n s_{i-1}, 1 \leq i \leq n$$

$$d_0 \tilde{t}_n = (-1)^n d_n$$

$$s_0 \tilde{t}_n = (-1)^n t_{n+1} s_n$$

This is Chern-Simons formalism.

Definition 4.8 (Tsygan, Loday-Quillen). The **cyclic bicomplex** $CC_*(C)$ associated to C_* is defined by

$$\begin{array}{ccccccc}
 & \dots & & \dots & & \dots & & \dots \\
 & \downarrow & & \downarrow & & \downarrow & & \downarrow \\
 \dots & \longrightarrow & C_2 & \xrightarrow{1-\tilde{t}} & C_2 & \xrightarrow{1-\tilde{t}} & C_2 & \xrightarrow{1-\tilde{t}} & C_2 \\
 & & \downarrow b & & \downarrow -b' & & \downarrow b & & \downarrow -b' \\
 \dots & \longrightarrow & C_2 & \xrightarrow{1-\tilde{t}} & C_2 & \xrightarrow{1-\tilde{t}} & C_2 & \xrightarrow{1-\tilde{t}} & C_2 \\
 & & \downarrow b & & \downarrow -b' & & \downarrow b & & \downarrow -b' \\
 \dots & \longrightarrow & C_2 & \xrightarrow{1-\tilde{t}} & C_2 & \xrightarrow{1-\tilde{t}} & C_2 & \xrightarrow{1-\tilde{t}} & C_2 \\
 & & \downarrow & & \downarrow & & \downarrow & & \downarrow \\
 & & 0 & & 0 & & 0 & & 0
 \end{array}$$

Define the total complex of CC_* as

$$\text{Tot}(CC_*) := \left(\bigoplus_{p+q=n} C_{p,q}, d = d^h + d^v \right)$$

and cyclic homology is defined as

$$HC_*(C) := H_*(\text{Tot}_{\text{ast}}(CC_*)).$$

Let $CC_*^{\{2\}}$ be the subcomplex consisting of the first two columns, then Hochschild homology is

$$HH_*(C) := H_*(\text{Tot}(CC_*^{\{2\}}))$$

Remark. If (C_*, b') is acyclic, then the natural projection

$$CC_*^{\{2\}} \xrightarrow[\text{q-sim}]{\sim} (C_*, b)$$

induces an isomorphism

$$HH_*(C) \cong H_*(C_*, b)$$

Theorem 4.9 (Connes' Periodicity Exact Sequence). *For any cyclic module, there is a long exact sequence ("ISB"):*

$$\cdots \longrightarrow HH_n(C) \xrightarrow{I} HC_n(C) \xrightarrow{S} HC_{n-2}(C) \xrightarrow{B} HH_{n-1}(C) \xrightarrow{I} \cdots$$

Proof. There is the following short exact sequence of bicomplexes

$$0 \longrightarrow CC_*^{\{2\}} \hookrightarrow CC_* \longrightarrow CC_*[2,0] \longrightarrow 0$$

where $CC_*[2,0]_{p,q} := CC_{p-2,q}$ is a bidegree shift. This induces an exact sequence in total complexes

$$0 \longrightarrow \text{Tot}(CC_*^{\{2\}}) \hookrightarrow \text{Tot}(CC_*) \longrightarrow CC_*[2,0] \longrightarrow 0$$

□

Remark. The relation $b(1 - \tilde{t}_n) = (1 - \tilde{t}_{n-1})b'$ implies that

$$C_*^\lambda(C) := (C_*/(1 - \tilde{t}_*, b))$$

is a well-defined cyclic (Connes') complex. If $k \supseteq \mathbb{Q}$, then

$$CC_* \longrightarrow C_* \twoheadrightarrow C_*^\lambda$$

gives

$$HC_*(C) \cong H_*(C_*^\lambda, b).$$

In general, there is a spectral sequence with E^1 page

$$E_{p,q}^1 := H_p^{EM}(\mathbb{Z}/(n+1), C_q) \Rightarrow HC_{p+q}(C)$$

Quillen is trying to connect this to Bott periodicity.

G. Segal (2011)

Example 4.10. Let A be an associative algebra with 1. Define

$$\begin{aligned} C_*(A) : \Delta C^{\text{op}} &\longrightarrow \text{Mod}_k \\ [n] &\longmapsto A^{\otimes(n+1)} \end{aligned}$$

with

$$\begin{aligned} d_i(a_0, \dots, a_n) &= \begin{cases} (a_0, \dots, a_i a_{i+1}, \dots, a_n), & 0 \leq i < n, \\ (a_n a_0, a_1, \dots, a_{n-1}), & i = n. \end{cases} \\ s_i(a_0, \dots, a_n) &= (a_0, \dots, a_i, 1, a_{i+1}, \dots, a_n) \\ t_n(a_0, \dots, a_n) &= (a_n, a_0, \dots, a_{n-1}) \end{aligned}$$

then $HH_*(C_*(A)) = HH_*(A)$.

If A is unital, $(C_*(A), b')$ is an acyclic complex.

Consider

$$B : C_n(A) \longrightarrow C_{n+1}(A)$$

where

$$B(a_0, \dots, a_n) = \sum_{i=0}^n (-1)^{ni} (1, a_i, \dots, a_n, a_0, \dots, a_{i-1}) = (-1)^{ni} (a_i, 1, a_{i+1}, \dots, a_n, a_0, \dots, a_{i-1})$$

$B^2 = b^2 = 0$ and $Bb + bB = 0$. Then

$$HC_*(A) := HC(C_*(A)).$$

4.2.1 Quillen's Ayclisim Formalism

Theorem 4.11 (Loday-Quillen-Tsygan Theorem). *Let $k \supseteq \mathbb{Q}$ be a field, and let A be a unital associative algebra. Define*

$$gl_{\infty}(A) := \lim_{n \rightarrow \infty} gl_n(A)$$

where $gl_n(A) = (\mathbb{M}_n(A), [-, -])$. Then

$$H_*^{CE}(gl_{\infty}(A); k) \cong \lim_{n \rightarrow \infty} H_*^{CE}(gl_n(A); k) \cong \Lambda_k[HC_{*-1}(A)]$$

where Λ_k is graded symmetric algebra. This is analogue of Quillen's Q -construction.

Note $H_*^{\text{CE}}(\mathfrak{gl}_\infty(A); k)$ is a graded Hopf algebra.

Compare this with $GL_\infty(A)$ and higher K-theory construction

$$K_i(A) := \pi_i(\text{BGL}_\infty(A)^+)$$

Let $K_i(A)_\mathbb{Q} = K_i(A) \otimes_{\mathbb{Z}} \mathbb{Q}$, then

$$H_*(GL_\infty(A), \mathbb{Q}) \cong \bigwedge_{\mathbb{Q}} (K_*(A)_\mathbb{Q}).$$

This can be thought as an infinitesimal K-theory.

4.2.2 Cyclic (Co)homology and Algebra Extensions

Remark. This is the true Grothendieck understanding.

Let A be an associative k -algebra.

Consider the category of all algebra extensions $\mathbf{Alg}_k \downarrow A$ with

- objects are algebra extensions $\{f : R \twoheadrightarrow A\}$
- morphisms are $f : R \rightarrow R'$ such that

$$\begin{array}{ccc} R & \xrightarrow{\varphi} & R \\ & \searrow f & \swarrow f' \\ & A & \end{array}$$

Higher Trace Functor

$$\mathbf{Alg}_k \downarrow A \longrightarrow \text{Mod}_k$$

$$(R \twoheadrightarrow A) \longmapsto R / (I^{n+1} + [R, R])$$

Theorem 4.12. For $n \geq 0$,

$$HC_n(A) \cong \begin{cases} \lim_{\mathbf{Alg}_k \downarrow A} R / (I^{n+1} + [R, R]), & n = 2k, \\ \lim_{\mathbf{Alg}_k \downarrow A} (I^{k+1} / (I^k, I)), & n = 2k + 1 \end{cases}$$

even trace

Question 4.13. *odd trace?*

4.2.3 Chern-Simons Formalism for Algebra Cochains

Reduced Bar Construction Let A be a nonunital algebra. Define

$$\overline{B}(A) := \bigoplus_{n=1}^{\infty} \overline{B}(A)_n$$

where $\overline{B}(A)_n = A^{\oplus n}$ with

$$b'(a_1, \dots, a_n) = \sum_{i=1}^{n-1} (-1)^{i-1} (a_1, \dots, a_i a_{i+1}, \dots, a_n)$$

and

$$\Delta : \overline{B} \longrightarrow \overline{B} \otimes \overline{B}$$

$$(a_1, \dots, a_n) \longmapsto \sum_{i=1}^{n-1} (a_1, \dots, a_i) \otimes (a_{i+1}, \dots, a_n)$$

Claim 4.14. $(\overline{B}(A), b', \Delta)$ is a DG coalgebra with $d(ab) = d(a)b + (-1)^{|a|}ad(b)$.

On algebras, there are two natural constructions

- universal traces

$$\mathrm{Tr}^{\mathrm{un}} : A \longrightarrow A_{\#} = A / [A, A]$$

Universal cotrace on DG coalgebras

$$C^{\#} := \ker \left(C \xrightarrow{\Delta - \tau \circ \Delta} C \otimes C \right)$$

we have

$$(\overline{B}A)^{\#}[1] \cong C^{\lambda}(A)$$

Exercise 4.15. What is $CC_*(A)$?

Koszul dual construction
noncommutative de Rham complex

4.3 Functor Homology

Let \mathcal{C} be a small category, and let k be a commutative ring (or a field). We may consider

$$\mathrm{Mod}_k(\mathcal{C}) := \mathrm{Fun}(\mathcal{C}, \mathrm{Mod}_k)$$

as left \mathcal{C} -modules and

$$\mathrm{Mod}_k(\mathcal{C}^{\mathrm{op}}) := \mathrm{Fun}(\mathcal{C}^{\mathrm{op}}, \mathrm{Mod}_k)$$

as right \mathcal{C} -modules, which are abelian categories with enough projective and injective objects.

Define functor tensor product

$$\begin{aligned} - \otimes_{\mathcal{C},k} - : \text{Mod}_k(\mathcal{C}^{\text{op}}) \times \text{Mod}_k(\mathcal{C}) &\longrightarrow \text{Mod}_k \\ (\mathcal{N}, \mathcal{M}) &\longmapsto \mathcal{N} \otimes_{\mathcal{C},k} \mathcal{M} \end{aligned}$$

where

$$\mathcal{N} \otimes_{\mathcal{C},k} \mathcal{M} := \bigoplus_{c \in \text{Ob}(\mathcal{C})} \mathcal{N}(c) \otimes \mathcal{M}(c)$$

which is right exact (in each argument) and (left) balanced, which induces derived functors

$$- \otimes_{\mathcal{C},k}^{\mathbb{L}} - : \mathcal{D}(\text{Mod}_k(\mathcal{C}^{\text{op}})) \times \mathcal{D}(\text{Mod}_k(\mathcal{C})) \longrightarrow \mathcal{D}(k) = \mathcal{D}(\text{Mod}_k)$$

and we can define

Definition 4.16. $\text{Tor}_*^{\mathcal{C}}(\mathcal{N}, \mathcal{M}) := H_*[\mathcal{C} \otimes_{\mathcal{C},k}^{\mathbb{L}} \mathcal{M}]$

Example 4.17. Consider k -linearized Yoneda functor: fix $A \in \text{Ob}(\mathcal{C})$,

$$\begin{aligned} k[h^A] : \mathcal{C} &\xrightarrow{h^A} \text{Mod}_k \\ X &\longmapsto \text{Hom}_{\mathcal{C}}(A, X) \mapsto k[\text{Hom}_{\mathcal{C}}(A, X)] \\ k[h_A] : \mathcal{C} &\xrightarrow{h_A} \text{Mod}_k \\ X &\longmapsto \text{Hom}_{\mathcal{C}}(X, A) \mapsto k[\text{Hom}_{\mathcal{C}}(X, A)] \end{aligned}$$

There is a (linearized) Yoneda lemma: for any $A \in \text{Ob}(\mathcal{C})$, $\mathcal{N} \in \text{Mod}_k(\mathcal{C})$, $\mathcal{M} \in \text{Mod}_k(\mathcal{C})$,

$$\begin{aligned} \mathcal{N} \otimes_{\mathcal{C},k} k[h^A] &\cong \mathcal{N}(A) \\ k[h^A] \otimes_{\mathcal{C},k} \mathcal{M} &\cong \mathcal{M}(A) \end{aligned}$$

4.3.1 Hochschild Homology

Given a simplicial module

$$\begin{aligned} E_* : \Delta^{\text{op}} &\rightarrow \text{Mod}_k \\ [n] &\longmapsto E_n \end{aligned}$$

can define

$$H_*(E) := H_*(C_*(E), d)$$

where $d = \sum_{i=0}^n (-1)^i d_i$.

Example 4.18. Let A be an associative k -algebra. The cyclic module

$$C_*(A) : \Delta^{op} \rightarrow \text{Mod}_k$$

$$[n] \longmapsto A^{n+1}$$

can be restricted to a simplicial module.

Given an A -bimodule M , we can introduce coefficients

$$C_*(A, M) : \Delta^{op} \rightarrow \text{Mod}_k$$

$$[n] \longmapsto M \otimes_k A^n$$

where

$$d_i(m, a_1, \dots, a_n) := \begin{cases} (ma_1, a_2, \dots, a_n), & i = 0, \\ (m, a_1, \dots, a_i a_{i+1}, \dots, a_n), & 0 < i < n, \\ (a_n m, a_1, \dots, a_{n-1}), & i = n. \end{cases}$$

then classical Hochschild homology is defined as

$$HH_*(A, M) := H_*(C_*(A, M)).$$

4.3.2 Cyclic Homology

Question 4.19. What are coefficients for cyclic homology?

Answer. One possible answer is by Kaledin.

Example 4.20. Suppose $X : \Delta^{op} \rightarrow \mathbf{Set}$ is a simplicial set, we can linearize it to a functor then

$$H_*(k[X]) \cong H_*(|X|, k).$$

Proposition 4.21. For any simplicial module $E : \Delta^{op} \rightarrow \text{Mod}_k$

$$\text{Tor}_*^{\Delta^{op}}(k, E) \cong H_*(E)$$

In particular, if $E = C_*(A, M)$, then

$$\text{Tor}_*^{\Delta^{op}}(k, C_*(A, M)) \cong HH_*(A, M)$$

where

$$k : \Delta \rightarrow \text{Mod}_k$$

$$[n] \longmapsto k$$

is a trivial cosimplicial module.

Proof. There is a natural isomorphism

$$\mathrm{Tor}_{*}^{\mathcal{C}}(\mathcal{N}, \mathcal{M}) \cong \mathrm{Tor}_{*}^{\mathcal{C}^{\mathrm{op}}}(\mathcal{M}, \mathcal{N})$$

we need to prove

$$\mathrm{Tor}_{*}^{\Delta}(E, k) \cong H_{*}(E)$$

We construct for $k : \Delta \rightarrow \mathrm{Mod}_k$ the projective resolution as follows.

Let

$$\begin{aligned} \Delta_n = h^{[n]} : \Delta &\longrightarrow \mathbf{Set} \\ [m] &\longmapsto \mathrm{Hom}_{\Delta}([n], [m]) \end{aligned}$$

and we may linearize Δ to be a functor

$$\begin{aligned} K_n = k[\Delta_n] : \Delta &\longrightarrow \mathrm{Mod}_k \\ [m] &\longmapsto k[\mathrm{Hom}_{\Delta}([n], [m])] \end{aligned}$$

which composes together to form a cosimplicial complex

$$\begin{aligned} K_* : \Delta \rightarrow \mathrm{Mod}_k^{\Delta^{\mathrm{op}}} &\xrightarrow{C_*} \mathrm{Ch}_k \\ [m] &\longmapsto K_*[m] : [n] \mapsto k[\mathrm{Hom}_{\Delta}([n], [m])] \end{aligned}$$

and

$$\begin{aligned} k : \Delta \rightarrow \mathrm{Mod}_k &\hookrightarrow \mathrm{Ch}_k \\ [m] &\longmapsto k \longmapsto [0 \leftarrow k \leftarrow 0] \end{aligned}$$

with augmentation

$$\varepsilon_{[m]} : K_0([m]) \cong k[\{0, \dots, m\}] \rightarrow k \ni \delta_{i0}$$

□

Lemma 4.22. $K_*(\bullet) \rightarrow k$ is a projective resolution of k as a trivial Δ -module.

Proof. For each $m \geq 0$, we have

$$H_i(K_*([m])) \cong \begin{cases} k, & i = 0 \\ 0, & i > 0 \end{cases}$$

where we have isomorphisms

$$H_i(K_*([m]))$$

Each K_n is a projective object in Mod_k^Δ .

$$\text{Tor}_*^{\Delta^{\text{op}}}(k, E) \cong \text{Tor}_*^\Delta(E, k) \cong H_*(E \bigotimes_{\Delta} K_*) \cong H_*(E).$$

where the second last isomorphism follows from the lemma and the last isomorphism is from one previous example. \square

Theorem 4.23 (Connes). *For any cyclic module, $C : \Delta C^{\text{op}} \rightarrow \text{Mod}_k$, there is a natural isomorphism*

$$\text{Tor}_*^{\Delta C^{\text{op}}}(k, C) \cong \text{HC}_*(C).$$

Sketch. For $(p, q) \in \mathbb{Z}_{\geq 0} \times \mathbb{Z}_{\geq 0}$,

$$\begin{aligned} (K(p, q))_* : \Delta &\longrightarrow \text{Mod}_k \\ [n] &\longmapsto k[\text{Hom}_\Delta([q], [n])] \end{aligned}$$

\square

extend this to ΔC .

Given $g \in \text{Aut}_{\Delta C}([n])$, let g act by \circ ,

$$\begin{aligned} (K(p, q))_* : \Delta &\longrightarrow \text{Mod}_k \\ g : ([n] \xrightarrow{f} [n]) &\mapsto ([q] \xrightarrow{f} [n] \xrightarrow{g} [n]) \end{aligned}$$

For each $n \geq 0$, (with p fixed)

$$\begin{aligned} (K(p, \bullet))_n : \Delta^{\text{op}} &\longrightarrow \text{Mod}_k \\ [q] &\longmapsto (K(p, q))_n \end{aligned}$$

which gives rise to

$$\begin{aligned} (K(p, \bullet))_n : \Delta^{\text{op}} &\longrightarrow \text{Mod}_k^{\Delta C} \\ [q] &\longmapsto (K(p, q))_n \end{aligned}$$

which gives us a complex of ΔC -module, then we can get a biresolution of

ΔC -module.

$$\begin{array}{ccccc}
 (K(0,2))_* & \xleftarrow{N_2} & (K(1,2))_* & \xleftarrow{1-\bar{t}_2} & (K(2,2))_* \\
 & & \downarrow & & \downarrow \\
 (K(0,1))_* & \xleftarrow{N_1} & (K(1,1))_* & \xleftarrow{1-\bar{t}_1} & (K(1,1))_* \\
 & & \downarrow & & \downarrow \\
 (K(0,0))_* & \xleftarrow{N_0} & (K(1,0))_* & \xleftarrow{1-\bar{t}_0} & (K(0,0))_* \\
 & & \downarrow & & \downarrow \\
 0 & & 0 & & 0
 \end{array}$$

Next time:

$$|X| \longrightarrow ES^1 \times_{S^1} |X| \longrightarrow BS^1$$

Gysin sequence and Connes' Periodicity Sequence

4.4 Topological Interpretation of Cyclic Homology

Cyclic Spaces A cyclic space is a functor

$$\begin{aligned}
 X : \Delta C^{\text{op}} &\rightarrow \mathbf{Top} \\
 [n] &\longmapsto X_n
 \end{aligned}$$

The underlying simplicial space is given by precomposing with $\Delta^{\text{op}} \hookrightarrow \Delta C^{\text{op}}$.

We assume that X is a “good simplicial space” (in the sense of G. Segal “Categories and Cohomology Theories”, Appendix A), so that geometric realization

$$|X| = \left(\coprod_{n \geq 0} X_n \times \Delta^n \right) / \sim$$

where the equivalence relation is defined as

$$\begin{aligned}
 (d_i^X(x), (t_0, \dots, t_n)) &\sim (x, d_{\Delta}^i(t_0, \dots, t_n)) \\
 (s_j^X(x), (t_0, \dots, t_n)) &\sim (x, s_{\Delta}^j(t_0, \dots, t_n))
 \end{aligned}$$

The pathology is, it may happen that $f : X_* \rightarrow Y_*$ such that $f_n : X_n \rightarrow Y_n$ for all $n \geq 0$, but $|f| : |X_*| \not\cong |Y_*|$.

Definition 4.24 (Segal). A fat realization

$$\|X\| := \coprod_{n \geq 0} (X_n \times \Delta^n) / \sim$$

where \sim is only generated by the face relations.

There is a canonical comparison map

$$f_X : \|X\| \longrightarrow |X|$$

$X : \Delta^{\text{op}} \rightarrow \mathbf{Top}$ is **good** if f_X is homotopy equivalence.

Theorem 4.25 (Segal, Proposition A1). *Let $X : \Delta^{\text{op}} \rightarrow \mathbf{Top}$*

1. *If all X_n are of homotopy type of CW complex, so is $\|X\|$.*
2. *If $f : X_* \rightarrow Y_*$ such that $f_n : X_n \xrightarrow{\sim} Y_n$ for all $n \geq 0$, then $\|X\| \xrightarrow{\|f\|} \|Y\|$.*
3. *$\|X \times Y\| \simeq \|X\| \times \|Y\|$.*

Example 4.26. For a bisimplicial set $X_{*,*} : \Delta^{\text{op}} \rightarrow \mathbf{sSet}$, $|X_{*,*}| : \Delta^{\text{op}} \rightarrow \mathbf{sSet} \xrightarrow{|\cdot|} \mathbf{Top}$ is always a good simplicial space.

Remark (C. Berger-K. Ratlcovic, “Gabriel=Morita theorem”, 2019). [Wendli 2018, P. May, Ajay and Yuri] Let $\Delta_{\text{inj}} \subseteq \Delta$ be the subcategory of injectives generated by d^i 's.

$$i : \Delta_{\text{inj}}^{\text{op}} \hookrightarrow \Delta^{\text{op}}$$

induces an adjunction

$$i_! : \Delta_{\text{inj}}^{\Delta^{\text{op}}} \rightleftarrows \mathbf{Set}^{\Delta^{\text{op}}} : i^*$$

which gives $\Delta_{\text{inj}}^{\Delta^{\text{op}}}$ induced model structure with weak equivalences maps $f : X \rightarrow Y$ such that $i_! f : i_! X \xrightarrow{\sim} i_! Y$ is a weak equivalence in \mathbf{sSet} .

Curious Fact There is no model structure on $\Delta_{\text{inj}}^{\Delta^{\text{op}}}$, thus Δ_{inj} is not a test category. Thus the above adjunction can never be a Quillen adjunction, but a right model approximation.

Note we always have $\|X\| \sim |i_!(X)|$, but $|i_! i^*(X)| \simeq |X|$ provided X is fibrant, i.e. a Kan complex.

Claim 4.27. $(i_!, i^*)$ is a right model approximation of $\mathbf{Set}^{\Delta_{\text{inj}}^{\text{op}}}$ such that

$$\text{Ho}(\mathbf{Set}^{\Delta_{\text{inj}}}) \subseteq \text{Ho}(\mathbf{sSet}).$$

Theorem 4.28. *Let $X : \Delta^{\text{op}} \rightarrow \mathbf{Top}$ be a (good) cyclic space. Then*

1. *$|X|$ carries a canonical S^1 -action.*
2. *$X \mapsto |X|$ extends to a functor*

$$|\cdot| : \text{GCS} \longrightarrow S^1\text{-Sp}$$

How can we define S^1 -action?

Recall

$$C_* : \Delta C^{\text{op}} \rightarrow \mathbf{Set}$$

$$[n] \longmapsto C_n = \text{Aut}_{\Delta^{\text{op}}}([n]) \cong \mathbb{Z}/(n+1)$$

there is another way to define $C_* \cong S_*^1 = \Delta[1]_* / \partial\Delta[1]_*$ with $|C_1| \cong S^1$.

Naive idea is to define a map

$$\text{ev} : C_n \times X_n \rightarrow X_n$$

$$(g, x) \longmapsto X(g)x$$

Exercise 4.29. This is not a simplicial map, so it does not induce a map on geometric realization unless X is a trivial cyclic space.

The modification is to introduce a new “twisted” simplicial set $F(X)$ with two properties

1. $|F(X)| \sim |C_* \times X_*| \cong S^1 \times |X|$.
2. there is a canonical map $F(X) \rightarrow X$ inducing the action map $S^1 \times |X| \rightarrow |X|$.

Lemma 4.30. *The forgetful functor*

$$F : \mathbf{Top}^{\Delta^{\text{op}}} \rightleftarrows \mathbf{Top}^{\Delta C^{\text{op}}} : U$$

with three properties

1. $F(\mathbf{pt}) = C_*$.
2. the map $F(X) \rightarrow X$ comes from the counit $\epsilon_X : FU(X) \rightarrow X$.
3. for any simplicial space X , there is canonical homeomorphism

$$|F(X)| \cong |C_*| \times |X| \cong S^1 \times |X|.$$

Proof. Recall in ΔC^{op} we defined the two operations

1. for all $g \in C_m$,

$$g^* : \text{Hom}_{\Delta^{\text{op}}}([m], [n]) \rightarrow \text{Hom}_{\Delta^{\text{op}}}([m], [n])$$

$$a \longmapsto g^*(a)$$

2. for all $\alpha \in \text{Hom}_{\Delta C^{\text{op}}}([m], [n])$,

$$\alpha_* : C_m \rightarrow C_n$$

$$g \longmapsto \alpha_*(g)$$

which are defined by the unique factorization property $\alpha \circ g = \alpha_*(g) \circ g^*(\alpha)$ in ΔC^{op} :

$$\begin{array}{ccc} [n] & \xleftarrow{\alpha} & [m] \\ \alpha_* g \uparrow & & \uparrow g \\ [n] & \xleftarrow{g^* \alpha} & [m] \end{array}$$

□

Explicitly,

$$F : \mathbf{Top}^{\Delta C^{\text{op}}} \longrightarrow \mathbf{Top}^{\Delta C^{\text{op}}}$$

$$Y \longmapsto F(Y)_* : \Delta C^{\text{op}} \rightarrow \mathbf{Top}$$

where $F(Y)_n = C_n \times Y_n$.

For any $f \in \text{Hom}_{\Delta C^{\text{op}}}([m], [n])$, write $f = h \circ \alpha$ where $\alpha \in \text{Hom}_{\Delta C^{\text{op}}}([m], [n])$ and $h \in \text{Aut}_{\Delta C^{\text{op}}}([n]) = C_n$. Then

$$F(Y)_f : F(Y)_m \longrightarrow F(Y)_n$$

$$(g, y) \longmapsto (h \cdot \alpha_*(g), Y(g^*(\alpha))(y))$$

Define for any cyclic space $X : \Delta C^{\text{op}} \rightarrow \mathbf{Top}$

$$ev : F(UX) = F(X)_* \longrightarrow X_*$$

$$ev_m : C_m \times X_m \longrightarrow X_m$$

$$(g, x) \longmapsto X(g)(x) =: g_*(x)$$

Need to check

$$\begin{array}{ccc} F(X)_m & \xrightarrow{ev_m} & X_m \\ F(X)_\alpha \downarrow & & \downarrow \alpha_* \\ F(X)_n & \xrightarrow{ev_n} & X_n \end{array}$$

for all $\alpha \in \text{Hom}_{\Delta C^{\text{op}}}([m], [n])$, i.e. whether the following diagram

$$\begin{array}{ccc} (g, x) & \longmapsto & g_*(x) \\ \downarrow & & \downarrow \\ (\alpha_*(g), (g^*(\alpha))_*(x)) & \xrightarrow{?} & \alpha_* g_*(x) \end{array}$$

Check

$$\begin{aligned}
 \text{ev}_n(F(X)_a(g, x)) &= \text{ev}_n(a_*(g), (g^*(a))_*(x)) \\
 &= (a_*(g)) (g^*(a))_*(x) \\
 &= (a_*(g) \circ g^*(a))_*(x) \\
 &= (a \circ g)_*(x) \\
 &= a_*(g_*(x)).
 \end{aligned}$$

Now we can define

$$\begin{aligned}
 \Phi : \text{Hom}_{\mathbf{Top}^{\Delta\text{cop}}}(F(Y), X) &\longrightarrow \text{Hom}_{\mathbf{Top}^{\Delta\text{op}}}(Y, UX) \\
 (F(Y) \xrightarrow{\alpha} X) &\longmapsto (Y \xrightarrow{i} F(Y) \xrightarrow{\alpha} X) \\
 \Psi : \text{Hom}_{\mathbf{Top}^{\Delta\text{op}}}(Y, UX) &\longrightarrow \text{Hom}_{\mathbf{Top}^{\Delta\text{cop}}}(F(Y), X) \\
 (\beta : Y \rightarrow UX) &\longmapsto (\text{ev} \circ F : F(Y) \rightarrow F(UX) = F(X) \xrightarrow{\text{ev}} X)
 \end{aligned}$$

Check

$$\Phi \circ \Psi = \text{Id}, \quad \Psi \circ \Phi = \text{Id}.$$

It remains to prove for any simplicial topological space X ,

$$|F(X)| \xrightarrow{\cong} |C| \times |X|$$

We will define

$$p_1 : |F(X)| \twoheadrightarrow |C|$$

$$p_2 : |F(X)| \twoheadrightarrow |X|$$

and show that

$$(p_1, p_2) : |F(X)| \xrightarrow{\cong} |C| \times |X|$$

1. p_1 is induced by the map of simplicial space

$$p_{1,*} : F(X)_* \rightarrow C_*$$

$$(g, x) \longmapsto g$$

2. p_2 is defined as follows.

Recall $\Delta^* : \Delta \rightarrow \mathbf{Top}$ can be extended to $\Delta C \rightarrow \mathbf{Top}$ where $\tau_n : [n] \rightarrow [n]$ acts by

$$e_i \mapsto \begin{cases} e_{i-1}, & 1 \leq i \leq n \\ e_n, & i = 0 \end{cases}$$

so it acts on Δ^n by

$$\tau_n(x_0, \dots, x_n) = (x_n, x_0, \dots, x_{n-1}).$$

Now we can define

$$p_2(g, x, u) = (x, g^*(u))$$

where g^* is the standard action of $g \in \text{Aut}([n])$ on Δ^* . We need to check p_2 is well-defined. For any $f \in \text{Hom}_{\Delta^{\text{op}}}([m], [n])$,

$$\begin{aligned} p_2(f(g, x); u) &= p_2(f_*(g), g^*(f)_*(x); u) \\ &= (g^*(f)_*(x); (f_*(g))^*(u)) \\ &= (x, g^*f^*(f_*(g))^*(u)) \\ &= (x, (g^*(f) \circ f_*(g))^*(u)) \\ &= (x, (g \circ f)^*(u)) \\ &= (x, g^*(f^*(u))) \\ &= p_2(g, x; f^*(u)) \end{aligned}$$

3. $(p_1, p_2) : |F(X)| \rightarrow |C| \times |X|$ is well-defined, and we can define a natural map

$$h_X : |F(X)| \xrightarrow{\sim} |C \times X|$$

$$(g, x; u) \mapsto (g^{-1}, x; g^*(u))$$

h_X is a homeomorphism.

4. To prove (2) we factor (p_1, p_2) as follows:

$$(p_1, p_2) : |F(X)| \xrightarrow[\cong]{h_X} |C \times X| \xrightarrow[\cong]{(|p_1|, |p_2|)} |C| \times |X| \xrightarrow[\cong]{h_{\mathbf{pt}} \times \text{Id}} |C| \times |X|$$

If we identify $|C| \cong S^1$ via $C_* \cong S^1_*$, use evaluation map we define the action map

$$\xi_X : |C| \times |X| \xrightarrow{(|p_1|, |p_2|)} |F(X)| \xrightarrow{\text{ev}} |X|.$$

Lemma 4.31. $\xi_{|C|} : |C| \times |C| \rightarrow |C|$ is the standard group structure on S^1 .

Proof. The diagram commutes

$$|C| \times |C| \xleftarrow{(|p_1|, |p_2|)} |F(C)| \xrightarrow{\text{ev}} |C|$$

Consider $F(C) = \{C_n \times C_n\}_{n \geq 0}$.

- there are three nondegenerate simplices in dim 1: $(1, t_1), (t_1, 1), (t_1, t_1)$.

- there are two nondegenerate simplices in dim 2: $(t_2, t_2), (t_2^2, t_2^2)$.

The triangulation for $S^1 \times S^1$ is as follows

•

For $(u, v) \in [0, 1]^2$,

$$(u, v) \in \begin{cases} \{(t_2^2, t_2^2), \Delta^2\} & u + v \leq 1 \\ \{(t_2, t_2), \Delta^2\} & u + v \geq 1 \end{cases}$$

Check

$$ev : F(C)_* \longrightarrow C_*$$

$$(t_2, t_2) \longmapsto t_2^2 = s_0(t_1)$$

$$(t_2^2, t_2^2) \longmapsto t_2^4 = t_2 = s_1(t_1)$$

so

$$|ev|(u, v) = \begin{cases} u + v, & \in \{t_1\} \times \Delta^2 \\ u + v - 1, & \in \{t_1\} \times \Delta^2 \end{cases}$$

identify this with multiplication of S^1 .

it remains to check that $\xi \circ (1 \times \xi) = \xi(\xi \times 1)$ and ξ is well-defined.

Given any $X \in \mathbf{Top}^{\Delta C^{op}}$ define

$$\mu : F(F(X)) \longrightarrow F(X)$$

$$(g, h, x) \mapsto (gh, X)$$

and

$$p'_1 : |F(F(X))| \twoheadrightarrow |F(C)|$$

$$(g, h, x; u) \mapsto (gh; u)$$

together with

$$p'_2 : |F(F(X))| \longrightarrow |X|$$

$$(g, h, x; u) \mapsto (x, (gh)_* u)$$

Lemma 4.32. p'_1, p'_2 are well-defined and

$$(p'_1, p'_2) : |F(F(X))| \xrightarrow{\cong} |F(C)| \times |X|$$

is a homeomorphism.

Same argument as in Lemma 1 for p_1, p_2 .

Now consider the following commutative diagram

$$\begin{array}{ccccc}
 |C| \times |C| \times |X| & \xleftarrow[\cong]{(p_1, p_2) \times} & |F(C)| \times |X| & \xrightarrow{|ev| \times 1} & |C|C \times |X| \\
 \uparrow \cong \scriptstyle 1 \times (p_1, p_2) & & \uparrow \cong \scriptstyle (p'_1, p'_2) & & \uparrow \cong \scriptstyle (p_1, p_2) \\
 |C| \times |F(X)| & \xleftarrow[\cong]{(p_1, p_2)} & |F(F(X))| & \xrightarrow{|\mu|} & |F(X)| \\
 \downarrow \scriptstyle 1 \times |ev_X| & & \downarrow \scriptstyle F(ev) & & \downarrow \scriptstyle |ev| \\
 |C| \times |X| & \xleftarrow[\cong]{(p_1, p_2)} & |F(X)| & \xrightarrow{|ev|} & |X|
 \end{array}$$

where the top, left, bottom and right composites are what we want. □

Hodge decomposition of string topology

5 Homotopy Colimits

AFTER Cisinski 2006 global Mike Shulman, Homotopy colimits 2009.

5.1 Homotopy Kan Extension

Notation Let \mathcal{M} be a cofibrantly generated model category with all (small) limits and colimits.

1. Let $\mathcal{A}, \mathcal{B}, \mathcal{C}, \mathcal{J}$ be small categories.
2. $\mathcal{M}^{\mathcal{A}}$ \mathcal{A} -diagrams with projective (Bousfield-Kan) model structure.

Now consider a functor $f : \mathcal{A} \rightarrow \mathcal{B}$ in **Cat**, there is an adjunction

$$\begin{aligned} f_! : \mathcal{M}^{\mathcal{A}} &\rightleftarrows \mathcal{M}^{\mathcal{B}} : f^* \\ \left(\mathcal{A} \xrightarrow{f} \mathcal{B} \xrightarrow{X} \mathcal{M} \right) &\longleftarrow \left(\mathcal{B} \xrightarrow{X} \mathcal{M} \right) \\ (X : \mathcal{A} \rightarrow \mathcal{M}) &\longmapsto (\mathrm{Lan}_f(X) : \mathcal{B} \rightarrow \mathcal{M}) \end{aligned}$$

Lemma 5.1. (5.1) is a Quillen pair.

Proof. f^* preserves both weak equivalences and fibrations. □

Thus we have a derived adjunction

$$\mathbb{L}f_! : \mathrm{Ho}(\mathcal{M}^{\mathcal{A}}) \rightleftarrows \mathrm{Ho}(\mathcal{M}^{\mathcal{B}}) : f^*$$

Definition 5.2. $\mathbb{L}f_!$ is called homotopy left Kan extension. And

$$\mathbb{L} \mathrm{colim}_{\mathcal{A}} := \mathbb{L}(\mathcal{A} \rightarrow *)$$

is the homotopy colimit functor.

Theorem 5.3 (Axiomatics after Grothendieck and Cisinski). *Fix \mathcal{M} as above.*

1. 2-functoriality: the pullback functors f^* fits together to give a strict 2-functor

$$\mathrm{Ho}(\overline{\mathcal{M}}) : \mathbf{Cat}^{\mathrm{op}} \longrightarrow \mathbf{CAT}$$

$$\mathcal{A} \longmapsto \mathrm{Ho}(\mathcal{M}^{\mathcal{A}})$$

In particular, this imply canonical isomorphism

$$\mathbb{L}(f \circ g)_! \cong \mathbb{L}(f)_! \circ \mathbb{L}(g)_!$$

for composable $f, g \in \mathbf{Cat}$.

2. $\mathrm{Ho}(\mathcal{M}^{\mathcal{A}})$ is weakly product preserving in the sense that for any (possibly empty) set $\{\mathcal{A}_s\}_{s \in S}$ in $\mathrm{Ob}(\mathbf{Cat})$,

$$\mathcal{A} = \coprod_{s \in S} \mathcal{A}_s$$

then $i_s : \mathcal{A}_s \rightarrow \mathcal{A}$ induces $i_s^* : \mathcal{M}^{\mathcal{A}} \rightarrow \mathcal{M}^{\mathcal{A}_s}$ thus a map

$$i^* : \mathrm{Ho}(\mathcal{M}^{\mathcal{A}}) \xrightarrow{\cong} \prod_{s \in S} \mathrm{Ho}(\mathcal{M}^{\mathcal{A}_s})$$

which is an equivalence of categories.

3. (Reflexity) For any small category \mathcal{A} , the canonical inclusion $i : \mathcal{A}^\delta \rightarrow \mathcal{A}$ of discrete subcategory gives

$$i^* : \mathrm{Ho}(\mathcal{M}^{\mathcal{A}}) \rightarrow \mathrm{Ho}(\mathcal{M}^{\mathcal{A}^\delta})$$

is conservative, i.e. for any $\alpha : X \rightarrow X'$, if $i^* \alpha : i^* X \rightarrow i^* X'$ is a weak equivalence in $\mathcal{M}^{\mathcal{A}^\delta}$ then $\alpha \in \mathrm{We}(\mathcal{M}^{\mathcal{A}})$.

4. (Base change) Recall for $f : \mathcal{A} \rightarrow \mathcal{B}$, and any $b \in \mathrm{Ob}(\mathcal{B})$, we have a sliced category $f \downarrow b$ such that we have the following diagram

$$\begin{array}{ccc} f \downarrow b & \xrightarrow{\pi} & \mathcal{A} \\ \downarrow p & & \downarrow f \\ * & \xrightarrow{b} & \mathcal{B} \end{array}$$

which gives a natural weak equivalence

$$\mathbb{L}p_! \circ \pi^* \xrightarrow{\sim} b^* \circ \mathbb{L}f_!$$

and for any $X : \mathcal{A} \rightarrow \mathcal{M}$,

$$\mathbb{L} \operatorname{colim}_{f \downarrow b} (\pi^*(X)) \simeq \mathbb{L}f_!(X)(b)$$

i.e.

$$f_!(X)(b) \cong \operatorname{colim}_{f \downarrow b} (X \rightarrow \pi)$$

Remark. Properties (2)-(4) can be facily summarized by saying that

$$\mathrm{Ho}(\mathcal{M}^-) : \mathbf{Cat}^{\mathrm{op}} \rightarrow \mathbf{CAT}$$

$$\mathcal{A} \longmapsto \mathrm{Ho}(\mathcal{M}^{\mathcal{A}})$$

is a (weak left) Grothendieck derivator associated to cofibrantly generated model category \mathcal{M} .

Theorem 5.4. *This object can be associated to an arbitrary model category.*

Groth, Handbook of Homotopy

A deeper theorem is 5.16.

Definition 5.5 (Quillen, HAK, Chapter 1). $f : \mathcal{A} \rightarrow \mathcal{B}$ is called **right homotopy cofinal** if for any object $b \in \text{Ob}(\mathcal{B})$, $b \downarrow f$ is contractible, i.e. $B(b \downarrow f) \simeq \mathbf{pt}$.

Lemma 5.6. *If $f : \mathcal{A} \rightarrow \mathcal{B}$ has a left adjoint, then it is right homotopy cofinal.*

Proof.

$$g : \mathcal{B} \rightleftarrows \mathcal{A} : f$$

gives the initial object

$$(b, \eta_b : gf(b) \rightarrow b)$$

in $b \downarrow f$, thus $B(b \downarrow f) \simeq \mathbf{pt}$. □

5.2 Grothendieck Construction

ref:

1. Grothendieck SGG1 Expose IV
2. Quillen HAK Chapter 1
3. R. Thomason, Homotopy Colimits in Cat, op-lax functors

Given a strict diagram of small categories

$$F : \mathcal{C} \longrightarrow \mathbf{Cat}$$

its Grothendieck Construction $\mathcal{C} \int F$ is defined to be the following category

1. objects are (c, x) where $c \in \text{Ob}(\mathcal{C})$ and $x \in \text{Ob}(F(c))$, and
2. morphisms are given by

$$\text{Hom}_{\mathcal{C} \int F}((c, x), (d, y)) = \{(\varphi, f) : \varphi \in \text{Hom}_{\mathcal{C}}(c, d), f \in \text{Hom}_{F(c)}(F\varphi(x), y)\}$$

3. composition is given as follows: for $(\varphi : c \rightarrow c', f : F(\varphi)x \rightarrow x'), (\varphi' : c' \rightarrow c'', f' : F(\varphi')x' \rightarrow x'')$, their composition is given by

$$(\varphi' \circ \varphi : c \rightarrow c'', f' \circ F(\varphi')(f) : F(\varphi' \circ \varphi)x \rightarrow F(\varphi')x' \rightarrow x'')$$

Notation . For $\mathcal{R} \subset \mathbf{Cat}$ a subcategory, $i_{\mathcal{R}} : \mathcal{R} \rightarrow \mathbf{Cat}$, and for $F : \mathcal{C} \rightarrow \mathcal{R}$ we write $\mathcal{C} \int F := \mathcal{C} \int i_{\mathcal{R}} \circ F$.

Example 5.7. For $\mathcal{R} = \mathbf{Gr}$, $i_{\mathbf{Gr}} : \mathbf{Gr} \hookrightarrow \mathbf{Cat}$ where we view groups as one object categories. Given $G, N \in \text{Ob}(\mathbf{Gr})$ and a group action $F : G \rightarrow \text{Aut}(N)$, it gives a functor

$$\begin{aligned} F_N : \underline{G} &\longrightarrow \mathbf{Gr} \\ * &\longmapsto N \\ g \in G &\mapsto F(g) : N \rightarrow N \end{aligned}$$

In this case, $G \int F_N = G \ltimes N$.

Grothendieck's construction is a generalization of semi-direct product of groups.

Question 5.8. What is a “deformed” Grothendieck construction?

Example 5.9. Let $\mathcal{R} = \mathbf{Set}$ with $i_{\mathbf{Set}} : \mathbf{Set} \hookrightarrow \mathbf{Cat}$. For $F : \mathcal{C} \rightarrow \mathbf{Set}$, $\mathcal{C} \int F = \mathcal{C}_F$ is Bousfield-Kan's category of elements of F .

5.2.1 Universal Mapping Property

Let \mathcal{C}, \mathcal{D} be two fixed categories, and let $F : \mathcal{C} \rightarrow \mathbf{Cat}$ be a fixed diagram.

Proposition 5.10. There is a natural bijection between the set of all functors $G : \mathcal{C} \int F \rightarrow \mathcal{D}$ and the following data

1. for each $c \in \text{Ob}(\mathcal{C})$ a functor $G(c) : F(c) \rightarrow \mathcal{D}$.
2. for every $\varphi : c \rightarrow c'$ in \mathcal{C} , a natural transformation of functors $F(c) \rightarrow \mathcal{D}$

$$G(\varphi) : G(c) \Longrightarrow G(c') \circ F(\varphi)$$

such that

- (a) $G(\text{Id})_{\mathcal{C}} = \text{Id}$.
- (b) for any composite $(c \xrightarrow{\varphi} c' \xrightarrow{\varphi'} c'')$, $G(\varphi' \circ \varphi) = G(\varphi') \circ G(\varphi)$ in \mathcal{C} .

Proof. Given the data above, we define

$$G : \mathcal{C} \int F \rightarrow \mathcal{D}$$

by

- $G(c, x) = G(c)x$ on objects.
- $G(\varphi : c \rightarrow c', F(c)x \xrightarrow{f} x')$

On the other hand, given a functor $G : \mathcal{C} \int F \rightarrow \mathcal{D}$

□

5.2.2 Fibered/cofibered Categories

Note there is a projection forgetful functor $p : \mathcal{C} \int F \rightarrow \mathcal{C}$ which simply forgets the second argument.

Fix $\mathcal{C} \in \mathbf{Cat}$,

$$\mathcal{C} \int (-) : \mathbf{Fun}(\mathcal{C}, \mathbf{Cat}) \rightarrow \mathbf{Cat} \downarrow \mathcal{C}$$

$$F \longmapsto (p : \mathcal{C} \int F \rightarrow \mathcal{C})$$

$$(F \Rightarrow F') \longmapsto (\mathcal{C} \int \alpha : \mathcal{C} \int F \rightarrow \mathcal{C} \int F')$$

$$\mathcal{C} \int \alpha : \mathcal{C} \int F \longrightarrow \mathcal{C} \int F'$$

where $(c, x) \longmapsto (c, \alpha_c(x))$

$$(c \xrightarrow{\varphi} c', F(\varphi)x \xrightarrow{f} x') \longmapsto (c \xrightarrow{\varphi} c', \alpha_c)$$

Theorem 5.11 (Grothendieck). $\mathcal{C} \int (-)$ is fully faithful with its image being small categories, cofibered over \mathcal{C} .

5.2.3 Fibered/Cofibered Functors

Given $\pi : \mathcal{A} \rightarrow \mathcal{B}$, for any $b \in \mathbf{Ob}(\mathcal{B})$, there are three different categories over b :

- fiber category at b : $\pi^{-1}(b)$. Objects are $a \in \mathbf{Ob}(\mathcal{A})$ such that $\pi(a) = b$. Morphisms are $f : a \rightarrow a' \in \mathbf{Mor}(\mathcal{A})$ such that $\pi(f) = \text{Id}_b$.
- slice category $b \downarrow \pi$. Objects are pairs (a, f) , $a \in \mathbf{Ob}(\mathcal{A})$, $f : b \rightarrow \pi(a)$.
- slice category $\pi \downarrow b$. Objects are pairs (a, g) , $a \in \mathbf{Ob}(\mathcal{A})$, $g : \pi(a) \rightarrow b$.

Theorem 5.12 (Grothendieck). $\mathcal{C} \int (-)$ fully faithful with its images being small categories cofibered over \mathcal{C} .

Consider the following two functors

$$i^* : \pi^{-1}(b) \hookrightarrow b \downarrow \pi$$

$$a \longmapsto (a, \text{Id}_b)$$

$$j_* : \pi^{-1}(b) \hookrightarrow \pi \downarrow b$$

$$a \longmapsto (a, \text{Id}_b)$$

Definition 5.13. $\pi : \mathcal{A} \rightarrow \mathcal{B}$ is **prefibered** if for any $b \in \text{Ob}(\mathcal{B})$, i_* has right adjoint $i^!$.

$\pi : \mathcal{A} \rightarrow \mathcal{B}$ is **precofibered** if for any $b \in \text{Ob}(\mathcal{B})$, j_* has left adjoint j^* .

Suppose $\pi : \mathcal{A} \rightarrow \mathcal{B}$ is prefibered, we can define a base change functor

$$f^* : \pi^{-1}(b') \rightarrow \pi^{-1}(b)$$

for every $f : b \rightarrow b'$ by the following composite

$$f^* : \pi^{-1}(b') \xrightarrow{i'_*} b' \downarrow \pi \xrightarrow{f \downarrow \pi} b \downarrow \pi \xrightarrow{i^!} \pi^{-1}(b)$$

such that for any composite $b \xrightarrow{f} b' \xrightarrow{g} b''$,

$$(g \circ f)^* = f^* \circ g^* : \pi^{-1}(b'') \rightarrow \pi^{-1}(b).$$

These two are related by the following natural transformation

$$\begin{array}{ccccc}
 \pi^{-1}(b'') & \xrightarrow{i''_*} & b'' \downarrow \pi & \xrightarrow{g \downarrow \pi} & b' \downarrow \pi & \xrightarrow{(i')^!} & \pi^{-1}(b') \\
 & & & & \searrow \text{Id} & \downarrow i'_* & \\
 & & & & & b' \downarrow \pi & \\
 & & & & & \downarrow f \downarrow \pi & \\
 & & & & & b \downarrow \pi & \\
 & & & & & \downarrow i^! & \\
 & & & & & \pi^{-1}(b) & \\
 & \searrow (gf)^* & & & & &
 \end{array}$$

together with natural transformation $f^* g^* \Rightarrow (gf)^*$.

Definition 5.14. $\pi : \mathcal{A} \rightarrow \mathcal{B}$ is called **fibered** if

1. it is prefibered, and
 2. $\alpha_{f,g}$ is an isomorphism for all composable morphisms.
- Dually, we can define cofibered functors and similar 1-1 correspondences.

5.3 Thomason's Formula

for homotopy colimits in **Cat**. (Chacholski-Sschrez, Homotopy theory of diagrams, Chapter 26)

Base Change Axiom For $f : \mathcal{A} \rightarrow \mathcal{B}$,

$$\begin{array}{ccc} f \downarrow b & \xrightarrow{\pi} & \mathcal{A} \\ \text{be} \downarrow p & & \downarrow f \\ * & \longrightarrow & \mathcal{B} \end{array}$$

$$\mathbb{L}p_! \pi^* \xrightarrow{\sim} b^* \mathbb{L}f_!$$

For any $X : \mathcal{A} \rightarrow \mathcal{M}$, \mathcal{M} a model category.

$$\mathbb{L}f_!(X)(b) = \text{hocolim}_{f \downarrow b} (\pi^* X)$$

Lemma 5.15. *If $f : \mathcal{A} \rightarrow \mathcal{B}$ is precofibered.*

Theorem 5.16 (Cofinality Theorem). *If $f : \mathcal{A} \rightarrow \mathcal{B}$ is right homotopy cofinal, i.e. $f \downarrow b$ is contractible, then $f^* : \text{hocolim}_{\mathcal{A}}(f^*(X)) \xrightarrow{\sim} \text{hocolim}_{\mathcal{B}}(X)$ for any $X : \mathcal{B} \rightarrow \mathcal{M}$.*

{thm:cofinal}

Example 5.17. For a compact conencted Lie group G , Fix a prime p , let \mathcal{B}_p be the poset of all p -subgroups and \mathcal{A}_p be the poset of all nontrivial elementary abelian subgroups. Then $f : \mathcal{A}_p \hookrightarrow \mathcal{B}_p$ is right homotopy cofinal.

Lemma 5.18. *If $f : \mathcal{A} \rightarrow \mathcal{B}$ is precofibered, then $\mathbb{L}f_! X(b) \simeq \text{hocolim}_{f^{-1}(b)} (i^{\text{ast}} X)$.*

This implies the following theorem

Theorem 5.19 (Generalized Thomason Formula, Chacholski-Schmer, 2002). *Let \mathcal{M} be a cofibrantly generated model category and $X : \mathcal{C} \int F \rightarrow \mathcal{M}$. There is a natural weak equivalence*

$$\text{hocolim}_{\mathcal{C} \int F} (X) \cong \text{hocolim}_{\mathcal{C} \in \mathcal{C}} (\text{hocolim}_{F(c)} X(c)),$$

where $X(c) := i_c^*(X)$ and

$$i_c : F(c) \rightarrow \mathcal{C} \int F$$

$$x \longmapsto (c, x)$$

$$(x \xrightarrow{f} x') \mapsto (\text{Id}_c, f)$$

When we take $\mathcal{M} = \mathbf{sSet}$, we have the classical Thomason's formula.

Corollary 5.20 (Thomason's Formula). $\mathrm{hocolim}_{\mathcal{C}}^{\mathbf{sSet}}(\mathcal{N}_*F) \simeq \mathcal{N}_*(\mathcal{C} \int F)$ is a weak equivalence in \mathbf{sSet} .

Corollary 5.21 (BK construction). Let $\mathcal{M} = \mathbf{sSet}$ and consider $F : \mathcal{C} \rightarrow \mathbf{sSet}$, then

$$\mathrm{hocolim}_{\mathcal{C}} F \cong \mathcal{N}_*(\mathcal{C}_F).$$

Example 5.22. $\mathrm{hocolim}_{\Delta \mathcal{C}^{\mathrm{op}}}(\mathcal{B}^{\mathrm{cyc}}(\Gamma)) \simeq \mathrm{ES}^1 \times_{\mathcal{S}^1} \mathcal{L}(\mathcal{B}\Gamma).$

Sketch. Recall $p : \mathcal{C} \int F \rightarrow \mathcal{C}$ is canonically cofibered over \mathcal{C} .

$p : \mathcal{A} \rightarrow \mathcal{B}$ is precofiber if $j : p^{-1}(b) \hookrightarrow p \downarrow b$ has left adjoint.
cofibered means

1. p is precofibered, and
2. $p^{-1} : \mathcal{B} \rightarrow \mathbf{Cat}$, $b \mapsto p^{-1}(b)$ is a functor, i.e. composable morphisms maps to composable functors between categories.

Consider the following diagram

$$\begin{array}{ccc} p^{-1}(b) & \xhookrightarrow{i} & \mathcal{A} \xrightarrow{X} \mathcal{M} \\ \downarrow j & \nearrow \pi & \\ p \downarrow b & & \end{array}$$

Lemma 5.23. If $p : \mathcal{A} \rightarrow \mathcal{B}$ is precofibered, then for any $X : \mathcal{A} \rightarrow \mathcal{M}$, we have

$$\mathbb{L}p_!(X)(b) \simeq \mathrm{hocolim}_{p^{-1}(b)}(i^*(X)) = \mathbb{L}(p^{-1}(b) \rightarrow *) (i^*(X))$$

where $i : p^{-1}(b) \hookrightarrow \mathcal{A}$.

Proof. Observe $p : \mathcal{A} \rightarrow \mathcal{B}$ is cofibered implies $j : p^{-1}(b) \rightarrow p \downarrow b$ is right adjoint, thus right homotopy cofinal, or equivalently, $\mathcal{B}(b \downarrow p) \sim \mathbf{pt}$.

$$\begin{aligned} \mathbb{L}p_!(X)(b) &\simeq \mathrm{hocolim}_{\pi \downarrow b}(\pi^*(X)) \\ &\simeq \mathrm{hocolim}_{p^{-1}(b)}(j^*\pi^*X) \\ &\simeq \mathrm{hocolim}_{p^{-1}(b)}(i^*X) \\ &= \mathrm{hocolim}_{p^{-1}(b)}(j^*(X)). \end{aligned}$$

□

Since $p : \mathcal{C} \int F \rightarrow \mathcal{C}$ is precofibered and $p^{-1}(c) = F(c)$, so by lemma

$$\mathrm{hocolim}_{\mathcal{C} \int F}(X) \simeq \mathrm{hocolim}_{\mathcal{C} \in \mathcal{C}}(\mathrm{hocolim}_{F(c)} X(c)).$$

□

5.4 Homotopy Coends

Define \mathcal{C} -bifunctor to be a functor

$$F : \mathcal{C}^{\text{op}} \times \mathcal{C} \rightarrow \mathcal{M}$$

5.4.1 Twisted Arrow Category/Factorization Category

See Quillen HAK.

Consider

$$\text{Hom} : \mathcal{C}^{\text{op}} \times \mathcal{C} \longrightarrow \mathbf{Set}$$

$$(c, d) \longmapsto \text{Hom}_{\mathcal{C}}(c, d)$$

Let

$$\mathcal{F}(\mathcal{C}) = (\mathcal{C}^{\text{op}} \times \mathcal{C}) \int \text{Hom} =: (\mathcal{C}^{\text{op}} \times \mathcal{C})_{\text{Hom}}$$

Note $\mathcal{F}(\mathcal{C})^{\text{op}} \neq \mathcal{F}(\mathcal{C}^{\text{op}})$.

The category $\mathcal{F}(\mathcal{C})$ is given by

1. Objects: $\text{Ob}(\mathcal{F}(\mathcal{C})) = \{(i \xrightarrow{f} j) : f \in \text{Mor}(\mathcal{C})\}$.
2. Morphisms: $\text{Hom}_{\mathcal{F}(\mathcal{C})}(i \xleftarrow{f} j, i' \xleftarrow{f'} j') := \{(i \xrightarrow{\beta} i', j \xleftarrow{\alpha} j') : g = \beta \circ f \circ \alpha\}$.

$$\begin{array}{ccc} i & \xrightarrow{\beta} & i' \\ f \uparrow & & \uparrow g \\ j & \xleftarrow{\alpha} & j' \end{array}$$

In opposite category $\mathcal{F}(\mathcal{C})^{\text{op}}$ we reverse all arrows.

Define $p = s \times t : \mathcal{F}(\mathcal{C}) \rightarrow \mathcal{C}^{\text{op}} \times \mathcal{C}$, then by definition p is precofibrated.

$$s^{\text{op}} : \mathcal{F}(\mathcal{C})^{\text{op}} \longrightarrow \mathcal{C}^{\text{op}}$$

$$(c \xrightarrow{f} d) \longmapsto c$$

(5.1) $\{\{\text{eq:s-t-op}\}\}$

$$t^{\text{op}} : \mathcal{F}(\mathcal{C})^{\text{op}} \longrightarrow \mathcal{C}$$

$$(c \xrightarrow{f} d) \longmapsto d$$

Lemma 5.24 (Quillen). *Both functors 5.1 are right homotopy cofinal.*

Proof. 1. $s \times t : \mathcal{F}(\mathcal{C}) \rightarrow \mathcal{C}^{\text{op}} \times \mathcal{C}$ is cofibered implies $s : \mathcal{F}(\mathcal{C}) \rightarrow \mathcal{C}^{\text{op}}$ and $t : \mathcal{F}(\mathcal{C}) \rightarrow \mathcal{C}$ are cofibered, so for any $c, d \in \text{Ob}(\mathcal{C})$

$$s^{-1}(c) \hookrightarrow s \downarrow c$$

$$t^{-1}(d) \hookrightarrow t \downarrow d$$

are homotopy cofinal, i.e $B(s^{-1}(c)) \simeq B(s \downarrow c)$ and $B(t^{-1}(d)) \simeq B(t \downarrow d)$, so

$$\begin{aligned} B(s \downarrow c) &\simeq B(c \downarrow \mathcal{C}) \simeq \mathbf{pt} \\ B(t \downarrow d) &\simeq B((\mathcal{C} \downarrow d)^{\text{op}}) \simeq \mathbf{pt}. \end{aligned}$$

Note $B(d \downarrow \mathcal{C}^{\text{op}}) \simeq \mathbf{pt}$.

□

Corollary 5.25. For any $X : \mathcal{C} \rightarrow \mathcal{M}$ and $Y : \mathcal{C}^{\text{op}} \rightarrow \mathcal{M}$,

$$(s^{\text{op}})^* : \text{hocolim}_{\mathcal{F}(\mathcal{C})^{\text{op}}} ((s^{\text{op}})^* X) \xrightarrow{\sim} \text{hocolim}_{\mathcal{C}}(X)$$

$$(t^{\text{op}})^* : \text{hocolim}_{\mathcal{F}(\mathcal{C})^{\text{op}}} ((t^{\text{op}})^* Y) \xrightarrow{\sim} \text{hocolim}_{\mathcal{C}^{\text{op}}}(Y)$$

Definition 5.26. For any bifunctor $D : \mathcal{C}^{\text{op}} \times \mathcal{C} \rightarrow \mathcal{M}$, define homotopy coend

$$\int_{\mathbb{L}}^{c \in \mathcal{C}} D(c, c) := \text{hocolim}_{\mathcal{F}(\mathcal{C})^{\text{op}}} (\pi^* D)$$

where

$$\pi^{\text{op}} := t^{\text{op}} \times s^{\text{op}} : \mathcal{F}(\mathcal{C})^{\text{op}} \longrightarrow \mathcal{C}^{\text{op}} \times \mathcal{C} \xrightarrow{D} \mathcal{M}$$

and $\pi^* := (\pi^{\text{op}})^*$. Consider

$$\mathcal{C}^{\text{op}} \times \mathcal{C} \xrightarrow{\pi_1} \mathcal{C}^{\text{op}} \xrightarrow{Y} \mathcal{M}$$

$$\mathcal{C}^{\text{op}} \times \mathcal{C} \xrightarrow{\pi_2} \mathcal{C} \xrightarrow{X} \mathcal{M}$$

then

$$\begin{aligned} \text{hocolim}_{\mathcal{C}}(X) &\simeq \int_{\mathbb{L}}^{c \in \mathcal{C}} (\pi_2)^*(X) \\ \text{hocolim}_{\mathcal{C}}(Y) &\simeq \int_{\mathbb{L}}^{c \in \mathcal{C}} (\pi_1)^*(Y) \end{aligned}$$

Recall for a bifunctor $X : I^{\text{op}} \times I \rightarrow \mathcal{C}$, the coend

$$\text{coend}(X) := \int^{i \in I} X(i, i)$$

is the universal (initial) cocone of X in \mathcal{C} .

Lemma 5.27. $\text{coend}(X) \cong \text{colim}_{\mathcal{F}(\mathcal{C})^{\text{op}}}(\pi^* X)$.

Remark. The reason to work with a bigger category is because you may have more “free” categories connected to $\mathcal{F}(\mathcal{C})^{\text{op}}$, i.e. functor $\mathcal{F}(\mathcal{C})^{\text{op}} \rightarrow \mathcal{D}$ which is right homotopy cofinal, so you can transfer computation to \mathcal{D} -diagrams.

Properties of Homotopy coend:

1. (interchange) The flip functor $\tau : \mathcal{C}^{\text{op}} \times \mathcal{C} \rightarrow \mathcal{C} \times \mathcal{C}^{\text{op}}$ gives

$$\int_{\mathbb{L}}^{\mathcal{C} \in \mathcal{C}} \tau^*(X)(c, c) \simeq \int_{\mathbb{L}}^{\mathcal{C} \in \mathcal{C}^{\text{op}}} X(c, c)$$

2. (Fubini Theorem) For $X : \mathcal{C} \times \mathcal{D} \rightarrow \mathcal{M}$,

$$\int_{\mathbb{L}}^{(c, d) \in \mathcal{C} \times \mathcal{D}} X(c, d; c, d) \simeq \int_{\mathbb{L}}^{\mathcal{D}} \int_{\mathbb{L}}^{\mathcal{C}} X(c, d; c, d)$$

3. (Diagrams of diagrams) For $X : \mathcal{C} \times \mathcal{C}^{\text{op}} \rightarrow \mathcal{M}^{\mathcal{J}}$

$$\left(\int_{\mathbb{L}}^{\mathcal{C} \in \mathcal{C}} X(c, c) \right)_j = \int_{\mathbb{L}}^{\mathcal{C} \in \mathcal{C}} X(c, c)_j.$$

4. (Commutativity with left Quillen derived functors) Given a left Quillen functor $F : \mathcal{M} \rightarrow \mathcal{N}$,

Proposition 5.28 (Shapiro Lemma for Model Categories). *Let \mathcal{M} be a simplicial model category and \mathcal{C} be a small category*

6 Simplicial Bar Construction

6.1 Tensor and Cotensor

Notation/Terminology Let \mathcal{M} be a simplicial (i.e. enriched over **sSet**) category. For any $X, Y \in \text{Ob}(\mathcal{M})$, morphisms $\mathbf{Hom}(X, Y) \in \text{Ob}(\mathbf{sSet})$ and the composition

$$\mathbf{Hom}_{\mathcal{M}}(Y, Z) \times \mathbf{Hom}_{\mathcal{M}}(X, Y) \xrightarrow{\circ} \mathbf{Hom}_{\mathcal{M}}(X, Z)$$

is a morphism in **sSet** which is natural in X, Y, Z .

Remark. The notation comes from the observation that $\mathbf{Hom}_{\mathcal{M}}(X, Y) = \text{Hom}_{\mathcal{M}}(X, Y)$ and $\mathbf{Hom}_{\mathcal{M}}(X, Y)$ is the simplicial homotopies between maps.

Definition 6.1. \mathcal{M} is **tensor**ed over **sSet** if there is a bifunctor

$$\boxtimes : \mathbf{sSet} \times \mathcal{M} \longrightarrow \mathcal{M}$$

$$(K, X) \longmapsto K \boxtimes X$$

such that there is a natural isomorphism

$$\mathbf{Hom}_{\mathcal{M}}(K \boxtimes X, Y) \cong \mathbf{Hom}_{\mathbf{sSet}}(K, \mathbf{Hom}_{\mathcal{M}}(X, Y)).$$

Definition 6.2. \mathcal{M} is **cotensor**ed over **sSet** if there is a bifunctor

$$\{-, -\} : \mathbf{sSet}^{\text{op}} \times \mathcal{M} \longrightarrow \mathcal{M}$$

$$(K, X) \longmapsto \{K, X\}$$

such that there is a natural isomorphism

$$\mathbf{Hom}_{\mathcal{M}}(X, \{K, Y\}) \cong \mathbf{Hom}_{\mathbf{sSet}}(K, \mathbf{Hom}_{\mathcal{M}}(X, Y)).$$

Example 6.3. $\mathcal{M} = \mathbf{sC}$ where \mathcal{C} is bicomplete is a category tensor and cotensor over **sSet** with

- tensor $K \boxtimes X := \coprod_{K_n} X_n$.
- $(K \times L) \boxtimes X = K \boxtimes (L \boxtimes X)$ and $\Delta[0]_* \boxtimes X \cong X$.
- $\mathbf{Hom}_{\mathbf{sC}}(X, Y) := \{\mathbf{Hom}_{\mathcal{M}}(\Delta[n]_* \boxtimes X, Y)\}_{n \geq 0}$.
- cotensor is defined by

$$\mathbf{Hom}_{\mathcal{M}}(K \boxtimes X, Y) \cong \mathbf{Hom}_{\mathcal{M}}(X, \{K, Y\}).$$

Example 6.4. 1. For $\mathcal{C} = \mathbf{Set}$, $\mathcal{M} = \mathbf{sSet}$,

- $K \boxtimes X = K \times X = \{K_n \times X_n\}_{n \geq 0}$.
- $\{K, X\} = \mathbf{Hom}_{\mathbf{sSet}}(K, X) := \{\mathbf{Hom}_{\mathbf{sSet}}(\Delta[n]_* \times K, X)\}$.

2. For $\mathcal{M} = \mathbf{Mod}_R$ where R is a ring and \mathbf{Mod}_R is the category of left R -modules, $\mathcal{M} = \mathbf{sMod}_R$

- $K \boxtimes M := \{R[K_n] \otimes_R M_n\}_{n \geq 0} \cong \{\bigoplus_{K_n} M_n\}_{n \geq 0}$.
- $\mathbf{Hom}_{\mathbf{sMod}_R}(M, N) = \{\mathbf{Hom}_{\mathbf{sMod}_R}(R[\Delta[n]_*] \otimes_R M, N)\}_{n \geq 0}$.
- $\{K, M\} = \mathbf{Hom}_{\mathbf{sSet}}(K, M)$ with pointwise simplicial R -module structure from M .

3. $\mathcal{C} = \mathbf{Comm}_k$ where k is a commutative ring, $\mathcal{M} = \mathbf{sComm}_k$.

- $K \boxtimes X = \{K_n \otimes X_n\}_{n \geq 0}$
- $\{K, X\} = \{\mathbf{Hom}_{\mathbf{sSet}}(\Delta[n]_*, X)\}$

4. $\mathcal{C} = \mathbf{Top} = \mathbf{CGWH}$.

- $K \boxtimes X = |K| \times X$.
- $\{K, X\} = \mathbf{Hom}_{\mathbf{Top}}(|K|, X)$.

6.2 Functor Tensor Products

Let \mathcal{C} be a small category. Let \mathcal{M} be a simplicial category tensored and cotensored over \mathbf{sSet} .

Give $G : \mathcal{C}^{\text{op}} \rightarrow \mathbf{sSet}$ and $F : \mathcal{C} \rightarrow \mathcal{M}$, define

$$G \boxtimes_{\mathcal{C}} F := \int^{c \in \text{Ob}(\mathcal{C})} G(c) \boxtimes F(c) := \text{coeq} \left\{ \coprod_{c \rightarrow c'} G(c') \boxtimes F(c) \xrightarrow[1 \times f_*]{f^* \times 1} \coprod_{c \in \text{Ob}(\mathcal{C})} G(c) \boxtimes F(c) \right\}$$

Example 6.5. If $G : \mathcal{C}^{\text{op}} \rightarrow \mathbf{sSet}$, $c \mapsto \Delta[0]_* = \{*\}$, then

$$* \boxtimes_{\mathcal{C}} F = \text{colim}_{\mathcal{C}} F.$$

Intuition $G \boxtimes_{\mathcal{C}} F$ is the weighted colimit of F with respect to weights G .

Example 6.6 ((Left) Kan extension). For \mathcal{D} cocomplete

$$\begin{array}{ccc} \mathcal{C} & \xrightarrow{F} & \mathcal{D} \\ \downarrow H & \nearrow \text{Lan}_H(F) & \\ \mathcal{E} & & \end{array}$$

For any $e \in \text{Ob}(\mathcal{E})$,

$$\text{Lan}_H(F)(e) = \text{Hom}_{\mathcal{E}}(H(-), e) \boxtimes_{\mathcal{C}} F.$$

Example 6.7 (Internal Geometric Realization). For a simplicial category \mathcal{M} tensored and cotensored over \mathbf{sSet} , given any simplicial object $X : \Delta^{\text{op}} \rightarrow \mathcal{M}$, the internal geometric realization is defined as

$$|X_*| := \Delta[\bullet]_* \boxtimes_{\Delta^{\text{op}}} X.$$

When $\mathcal{M} = \mathbf{Top}$, $\mathbf{sM} = \mathbf{sTop}$, if $X : \Delta^{\text{op}} \rightarrow \mathbf{Set}$ is viewed as a discrete simplicial space,

$$|X| \simeq \mathbf{Hom}_{\mathbf{sSet}}(\Delta[\bullet]_*, X) \boxtimes_{\Delta} \Delta^* \cong X \boxtimes_{\Delta} \Delta^*.$$

Exercise 6.8. For $\mathcal{M} = \mathbf{sSet}$, $X : \Delta^{\text{op}} \rightarrow \mathbf{sSet}$ a bisimplicial set,

$$|X| \cong \text{diag}(X) = \{X_{nn}\}_{n \geq 0}.$$

6.3 Two-sided Simplicial Bar Construction

Definition 6.9. Given $G : \mathcal{C}^{\text{op}} \rightarrow \mathbf{sSet}$ and $F : \mathcal{C} \rightarrow \mathcal{M}$, define

$$\begin{aligned} B_*(G, \mathcal{C}, F) : \Delta^{\text{op}} &\longrightarrow \mathcal{M} \\ [n] &\longmapsto B_n(G, \mathcal{C}, F) \end{aligned}$$

where

$$B_n(G, \mathcal{C}, F) = \coprod_{\left(c_n \xleftarrow{f_{n-1}} c_{n-1} \xleftarrow{\cdots} \xleftarrow{f_0} c_0\right) \in \mathcal{N}_n \mathcal{C}} G(c_n) \boxtimes F(c_0)$$

with face maps

$$\begin{aligned} d_i : B_n(G, \mathcal{C}, F) &\longrightarrow B_{n-1}(G, \mathcal{C}, F) \\ (y|f_{n-1}, \dots, f_0|x) &\mapsto d_i(y|f_{n-1}, \dots, f_0|x) \end{aligned}$$

defined by

$$d_i(y|f_{n-1}, \dots, f_0|x) = \begin{cases} (y|f_{n-1}, \dots, f_1|f_0 \cdot x), & i = 0, \\ (y|f_{n-1}, \dots, f_i \circ f_{i-1}, \dots, f_0|x), & 1 \leq i \leq n-1, \\ (y \cdot f_{n-1}|f_{n-2}, \dots, f_0|x), & i = n. \end{cases}$$

and degeneracy maps

$$\begin{aligned} s_j : B_n(G, \mathcal{C}, F) &\longrightarrow B_{n+1}(G, \mathcal{C}, F) \\ (y|f_{n-1}, \dots, f_0|x) &\mapsto (y|f_{n-1}, \dots, \text{Id}_{c_j}, \dots, f_0|x) \end{aligned}$$

Note

$$\operatorname{colim}_{\Delta^{\text{op}}} (B_*(G, \mathcal{C}, F)) \cong G \boxtimes_{\mathcal{C}} F.$$

Therefore $B_*(G, \mathcal{C}, F)$ is a fattened (thickened) tensor product. There is a canonical map

$$B_*(G, \mathcal{C}, F) \rightarrow G \boxtimes_{\mathcal{C}} F.$$

6.3.1 Examples

1. $\mathcal{M} = \mathbf{Set}$. $G : \mathcal{C}^{\text{op}} \rightarrow \mathbf{Set} \hookrightarrow \mathbf{sSet}$, $c \mapsto *$ and $F : \mathcal{C} \rightarrow \mathbf{Set}$, $c \mapsto *$ then

$$B_*(*, \mathcal{C}, *) = \mathcal{N}_* \mathcal{C}.$$

2. $\mathcal{M} = \mathbf{Set}$ or \mathbf{Top} with $G = *$ and $F : \mathcal{C} \rightarrow \mathbf{Set}$, then

$$B_*(*, \mathcal{C}, F) = \mathcal{N}_*(\mathcal{C}_F).$$

3. $\mathcal{M} = \mathbf{Set}$ and for $c \in \operatorname{Ob}(\mathcal{C})$, we have

$$\begin{aligned} h^c : \mathcal{C} &\longrightarrow \mathbf{Set} & h_c : \mathcal{C}^{\text{op}} &\longrightarrow \mathbf{Set} \\ d &\longmapsto \operatorname{Hom}_{\mathcal{C}}(c, d) & b &\longmapsto \operatorname{Hom}_{\mathcal{C}}(b, c) \end{aligned}$$

then

$$B_*(*, \mathcal{C}, h^c) \cong \mathcal{N}(c \downarrow \mathcal{C}), \quad B(h_c, \mathcal{C}, *) \cong \mathcal{N}_*(\mathcal{C} \downarrow c).$$

Varying $c \in \operatorname{Ob}(\mathcal{C})$ in the last two formulas gives us two functors

$$\begin{aligned} B_*(*, \mathcal{C}, \mathcal{C}) : \mathcal{C}^{\text{op}} &\longrightarrow \mathbf{sSet} & B_*(\mathcal{C}, \mathcal{C}, *) : \mathcal{C} &\longrightarrow \mathbf{sSet} \\ c &\longmapsto B_*(*, \mathcal{C}, h^c) & c &\longmapsto B(h_c, \mathcal{C}, *) \end{aligned}$$

and we have

$$B_*(*, \mathcal{C}, \mathcal{C}) \cong \mathcal{N}((-) \downarrow \mathcal{C}), \quad B(\mathcal{C}, \mathcal{C}, *) \cong \mathcal{N}_*(\mathcal{C} \downarrow (-)).$$

4. (Exercise) For $F : \mathcal{C} \rightarrow \mathcal{D}$,

$$B_*(*, \mathcal{C}, \mathcal{D}) \cong \mathcal{N}((-) \downarrow F), \quad B(\mathcal{D}, \mathcal{C}, *) \cong \mathcal{N}_*(F \downarrow (-)).$$

which is a generalization of the previous one.

5. Let G be a discrete group, $\mathcal{C} = \underline{G}$ where $\operatorname{Ob}(\underline{G}) = \{*_G\}$ then $\operatorname{Hom}_{\underline{G}}(*, *) = G$.

$$B_*(*, \underline{G}, \underline{G}) \cong E_*(G), \quad B(\underline{G}, \underline{G}, *) \cong E_* G.$$

where the first is as right G -action and the second is as left G -action.

Definition 6.10. The **bar construction** is

$$B(G, \mathcal{C}, F) := |B_*(G, \mathcal{C}, F)|_{\mathcal{M}} = \Delta[\bullet]_* \boxtimes_{\Delta^{\text{op}}} B_*(G, \mathcal{C}, F).$$

The canonical functor

$$\Delta[\bullet]_* \longrightarrow \Delta[0]_*$$

induces a functor

$$\varepsilon : B(G, \mathcal{C}, F) \longrightarrow G \boxtimes_{\mathcal{C}} F$$

in \mathcal{M} .

Lemma 6.11. For any $F : \mathcal{C} \rightarrow \mathcal{M}$,

$$B_*(*, \mathcal{C}, F) \cong B_*(*, \mathcal{C}, \mathcal{C}) \boxtimes_{\mathcal{C}} F.$$

Thus one-sided bar construction is a weight colimit.

Proof. Tensor products are all coends, and the following commutes.

$$\begin{aligned} B(*, \mathcal{C}, F) &:= \Delta[\bullet]_* \boxtimes_{\Delta^{\text{op}}} B_*(*, \mathcal{C}, F) \\ &\cong \Delta[\bullet]_* \boxtimes_{\Delta^{\text{op}}} B_*(*, \mathcal{C}, \mathcal{C} \boxtimes_{\mathcal{C}} F) \\ &\cong (\Delta[\bullet]_* \boxtimes_{\Delta^{\text{op}}} B_*(*, \mathcal{C}, \mathcal{C})) \boxtimes_{\mathcal{C}} F \\ &= B_*(*, \mathcal{C}, \mathcal{C}) \boxtimes_{\mathcal{C}} F. \end{aligned}$$

□

Definition 6.12 (BK formula for homotopy colimit). $\text{hocolim}_{\mathcal{C}}(F) := B(*, \mathcal{C}, F) \cong \mathcal{N}_*(-) \downarrow \mathcal{C} \boxtimes_{\mathcal{C}} F$.

Remark. The concrete formula is

$$\mathbb{L} \text{colim}_{\mathcal{C}}(F) \cong B(*, \mathcal{C}, QF)$$

where $Q : \mathcal{M} \rightarrow \mathcal{M}$ is a cofibrant replacement functor on \mathcal{M} .

6.4 Cyclic Bar Construction of a Discrete Group

Notation. We will use $\Gamma_* G = B_*^{\text{cyc}}(G)$ to denote the cyclic bar construction.

Definition 6.13. The cyclic bar construction of G is given by

$$\Gamma_*(G) : \Delta C^{\text{op}} \rightarrow \mathbf{Top}$$

$$[n] \longmapsto G^{n+1}$$

with

$$\begin{aligned} d_i : \Gamma_n G &\longrightarrow \Gamma_{n-1} G \\ (g_0, \dots, g_n) &\mapsto d_i(g_0, \dots, g_n) \end{aligned}$$

where

$$d_i(g_0, \dots, g_n) = \begin{cases} (g_0, \dots, g_i g_{i+1}, \dots, g_n), & 0 \leq i < n, \\ (g_n g_0, g_1, \dots, g_{n-1}), & i = n. \end{cases}$$

and

$$\begin{aligned} s_j : \Gamma_n G &\longrightarrow \Gamma_{n+1} G \\ (g_0, \dots, g_n) &\mapsto (g_0, \dots, g_{j-1}, 1, g_j, \dots, g_n) \end{aligned}$$

and

$$\begin{aligned} t : \Gamma_n G &\longrightarrow \Gamma_n G \\ (g_0, \dots, g_n) &\mapsto (g_n, g_0, g_1, \dots, g_{n-1}) \end{aligned}$$

Remark. If G is a discrete group, and let $A = k[G]$ be the group algebra, then

$$C_*^{\text{HH}}(A) \cong k[\Gamma_* G].$$

Remark. Fix $z \in \mathcal{Z}(G)$ a central element, can define a twisted cyclic map

$$\begin{aligned} t_n(z) : B_n G &\longrightarrow B_n G \\ (g_1, \dots, g_n) &\mapsto (z(g_1 \cdots g_n)^{-1}, g_1, \dots, g_{n-1}) \end{aligned}$$

then $(t_n(z))^{n+1} = \text{Id}_{B_n G}$.

Recall $BG = |B_* G|$.

Theorem 6.14. For any discrete simplicial/topological group G , there is a canonical S^1 equivariant homotopy equivalence

$$\gamma : |\Gamma_* G| \rightarrow \mathcal{L}(BG) = \text{Map}(S^1, BG)$$

where the free loop space $\mathcal{L}(BG)$ carries a natural S^1 -action.

Lemma 6.15. For any discrete group G ,

$$F : \text{hocolim}_{\underline{G}}(\text{Ad}) \cong \gamma_* G,$$

where

$$\begin{aligned} \text{Ad} : \underline{G} &\longrightarrow \mathbf{Set} \\ * &\longmapsto G \\ g &\longmapsto (\text{Ad}_g : G \rightarrow G, \gamma \mapsto g\gamma g^{-1}). \end{aligned}$$

Proof. By definition,

$$\begin{aligned} \operatorname{hocolim}_{\underline{G}}(\operatorname{Ad}) &= |B_*(\ast, \underline{G}, \underline{G}) \boxtimes_{\underline{G}} \operatorname{Ad}_G| \\ &\cong |E_* \underline{G} \boxtimes_{\underline{G}} \operatorname{Ad}_G| \\ &\cong \mathcal{N}_*(\underline{EG}) \boxtimes_{\underline{G}} \operatorname{Ad}_G \end{aligned}$$

where $E_* \underline{G} = \mathcal{N}_*(\ast \downarrow \underline{G}) = \mathcal{N}_*(\underline{EG})$.¹

Explicitly, $\underline{EG} = \ast \downarrow \underline{G}$ is the canonical groupoid attached to G , with

- objects are $g \in G$, and
- $\operatorname{Hom}_{\underline{EG}}(g_1, g_2) = \{h \in G \mid g_2 = hg_1\} = \{g_2 g_1^{-1}\}$.

The nerve of \underline{EG} is a simplicial set

$$E_* \underline{G} : \Delta^{\operatorname{op}} \longrightarrow \mathbf{Set}$$

$$[n] \longmapsto G^{n+1}$$

with face and degeneracy maps

$$d_i : G^{n+1} \longrightarrow G^n$$

$$s_j : G^{n+1} \longrightarrow G^{n+2}$$

$$(g_0, \dots, g_n) \mapsto (g_0, \dots, g_{i-1}, g_{i+1}, \dots, g_n) \quad (g_0, \dots, g_n) \mapsto (g_0, \dots, g_{j-1}, g_j, g_j, g_{j+1}, \dots, g_n)$$

Note we have the identification

$$\mathcal{N}_n(\underline{EG}) \xrightarrow{\cong} E_n G$$

$$(g_0 \xrightarrow{g_1 g_0^{-1}} g_1 \rightarrow \dots \rightarrow g_n) \mapsto (g_0, \cdot, g_n)$$

and the face map $d_i^{\operatorname{Nerve}}$ which skips g_i in an n -composable morphisms in $\mathcal{N}_n(\underline{EG})$ by composing morphisms can be identified with $d_i^{E_* \underline{G}}$ which skips g_i in the $n+1$ -tuple in $E_n G$.

Note $E_* \underline{G}$ is equipped with a right G -action given by

$$E_n G \times G \longrightarrow E_n G$$

$$((g_0, \dots, g_n), z) \mapsto (g_0 z, \dots, g_n z)$$

hence we can take the G -orbit of $E_* \underline{G} \times \operatorname{Ad}_G$ levelwise by

$$E_n G \times_G G \cong (G^{n+1} \times G) / G$$

¹In general, we have $B(\ast, \mathcal{C}, \mathcal{C}) \cong \mathcal{N}_*((-) \downarrow \mathcal{C}) : \mathcal{C}^{\operatorname{op}} \rightarrow \mathbf{Set}, c \mapsto \mathcal{N}_*(c \downarrow \mathcal{C})$.

where

$$z \cdot (g_0, \dots, g_n; g) = (g_0 z^{-1}, \dots, g_n z^{-1}, z g z^{-1}).$$

So we have

$$\begin{aligned} F_* : E_* G \times_G \text{Ad}_G &\longrightarrow \Gamma_* G \\ [g_0, \dots, g_n; g] &\longmapsto (g_n g g_0^{-1}, g_0 g_1^{-1}, \dots, g_{n-1} g_0^{-1}) \end{aligned}$$

which is compatible with face and degeneracy maps, and it has an inverse

$$\begin{aligned} F_*^{-1} : \Gamma_* G &\longrightarrow E_* G \times_G \text{Ad}_G \\ (\gamma_0, \dots, \gamma_n) &\mapsto [\gamma_1 \cdots \gamma_n, \gamma_2 \cdots \gamma_n, \dots, \gamma_n, 1, \gamma_0 \gamma_1 \cdots \gamma_n] \end{aligned}$$

□

Corollary 6.16. *There is a canonical decomposition of cyclic sets $\Gamma_* G = \coprod_{\langle z \rangle \in \langle G \rangle} \Gamma_*(G, z)$ where $\langle G \rangle$ is the set of conjugacy classes of elements of G and*

$$\Gamma_*(G, z) = \{ \Gamma_n(G, z) = \{ (\gamma_0, \dots, \gamma_n) \in \Gamma_n G : \gamma_0 \gamma_1 \cdots \gamma_n \in \langle z \rangle \} \}$$

Proof. Note

$$\begin{array}{ccc} \Gamma_* G \cong \text{hocolim}_{\underline{G}} \text{Ad} & \xrightarrow{\cong} & E_* G \times_G \text{Ad} \\ \downarrow & & \downarrow \\ \text{colim}_{\underline{G}} \text{Ad} & \xlongequal{\quad} & \{*\} \times_G \text{Ad} = \langle G \rangle \end{array}$$

Simplicially, in degree n , we have

$$\begin{aligned} \pi_0 : E_n G \times_G \text{Ad} &\rightarrow \langle G \rangle \\ [g_0, \dots, g_n; z] &\mapsto \langle z \rangle \end{aligned}$$

□

6.4.1 Simplicial Models of Circles

There are two simplicial models of S^1 .

1. $B_* C = \{\mathbb{Z}/(n+1)\}_{n \geq 0}$, and
2. $B_* \mathbb{Z} = \{\mathbb{Z}^n\}_{n \geq 0}$.

There is a natural homotopy equivalence $|C_*| \simeq |B_*\mathbb{Z}|$ induced by the following map of simplicial sets (**CHECK**)

$$\begin{aligned} f_n : C_n &\longrightarrow \mathbb{Z}^n \\ t_n^0 &\longmapsto (0, \dots, 0) \\ t_n^i &\longmapsto (0, \dots, 1, \dots, 0) \end{aligned}$$

Furthermore, it is a map of cyclic sets with

$$\begin{aligned} t_n : \mathbb{Z}^n &\longrightarrow \mathbb{Z}^n \\ (m_1, \dots, m_n) &\mapsto (1 - (m_1 + \dots + m_n), m_1, \dots, m_{n-1}). \end{aligned}$$

6.5 Nerves

6.5.1 Twisted Nerve of a Group

Let G be a discrete group G , fix $z \in \mathcal{Z}(G)$ an element in the center of G . We can define $B_*(G, z)$ by add a cyclic map to the nerve of G as follows

$$\begin{aligned} t_n : B_n G = G^n &\longrightarrow B_n G = G^n \\ (g_1, \dots, g_n) &\mapsto (z \cdot (g_1 \cdots g_n)^{-1}, g_1, \dots, g_{n-1}). \end{aligned}$$

then one can check

$$t_n^{n+1}(g_1, \dots, g_n) = (zg_1z^{-1}, \dots, zg_nz^{-1}) = (g_1, \dots, g_n).$$

Notation. When $z = 1$, we write $B_*G = B_*(G, 1)$.

Proposition 6.17. *Given G and $z \in \mathcal{Z}(G)$ as above. We can define a group homomorphism*

$$\begin{aligned} \gamma_z : \mathbb{Z} \times G &\longrightarrow G \\ (n, g) &\longmapsto z^n g = g z^n. \end{aligned}$$

Up to homotopy, the S^1 -action on $BG = |B_*(G, z)|$ is induced by the above cyclic structure, which is given by following map

$$S^1 \times BG \cong |B_*\mathbb{Z}| \times |B_*G| \cong |B_*(\mathbb{Z} \times G)| \xrightarrow{B\gamma_z} BG.$$

Recall the following adjunction

$$F : \mathbf{Set}^{\Delta^{\text{op}}} \rightleftarrows \mathbf{Set}^{\Delta^{\text{C}^{\text{op}}}} : U$$

Proof. It suffices to show on π_1 :

$$\begin{array}{ccc} |F(B_*G)| & \xrightarrow{|\text{ev}|} & |B_*G| \\ (p_1, p_2) \downarrow & & \parallel \\ |C_*| \times |B_*G| & \longrightarrow & |B_*G| \end{array}$$

Check $t_1^* = \tau_1$. □

Definition 6.18. $X(G, z) := ES^1 \times_{S^1} |B_*(G, z)| =: |B_*(G, z)|_{hS^1}$ is the S^1 -equivariant Borel homotopy quotient of $|BG|$ by S^1 .

For any S^1 -space X , we have the following fibration

$$\begin{array}{ccccc} S^1 & \longrightarrow & ES^1 \times X & \longrightarrow & ES^1 \times_{S^1} X \\ \parallel & & \downarrow \simeq & & \parallel \\ |B_*\mathbb{Z}| & \xrightarrow{\mu_*} & BG & \longrightarrow & X(G, z) \end{array}$$

where

$$\mu : \mathbb{Z} \rightarrow G$$

$$1 \mapsto z.$$

Claim 6.19. $X(G, z)$ is equivalent to the classifying space (nerve) of the crossed module defined by the map $\mu : \mathbb{Z} \rightarrow G$ together with the trivial action of G on \mathbb{Z} .

6.5.2 Crossed Modules

Digression (on crossed modules) Let G be a discrete group and A an abelian group.

Fact 6.20. $H^2(G, A) \cong \{\text{group extensions } 1 \rightarrow A \rightarrow E \rightarrow G \rightarrow 1\} / \sim$, which has 1-1 correspondence with maps

$$G \times G \longrightarrow A$$

$$(g_1, g_2) \mapsto s(g_1)s(g_2)s(g_1g_2)^{-1}.$$

Question 6.21. Is there a similar interpretation for $H^3(G, A)$?

Definition 6.22. A crossed module is a group homomorphism $\mu : M \rightarrow N$ together with

$$\rho : N \longrightarrow \text{Aut}(M)$$

$$n \mapsto (\rho(n) : m \mapsto {}^n m)$$

satisfying

1. $\mu(nm) = n\mu(m)n^{-1}$ for any $m \in M, n \in N$.
2. $\mu(m)(m') = m \cdot m' \cdot m^{-1}$ for any $m \in M, n \in N$.

Equivalently, given any $\mu : M \rightarrow N$, there is a commutative diagram

$$\begin{array}{ccc} M & \xrightarrow{\text{Ad}} & \text{Aut}(M) \\ \mu \downarrow & \nearrow \exists \rho & \downarrow \text{act} \\ N & \xrightarrow{\text{Ad}_\mu} & \text{Hom}(M, N) \end{array}$$

where

$$\text{Ad}(m)(m') = mm'm^{-1}$$

and

$$\text{Ad}_\mu(n) : M \longrightarrow N$$

$$m' \longmapsto n\mu(m')n^{-1}.$$

and

$$\text{act} : \text{Aut}(M) \longrightarrow \text{Hom}(M, N)$$

$$\sigma \longmapsto (\mu \circ \sigma : M \xrightarrow{\sigma} M \xrightarrow{\mu} N).$$

Then a crossed module structure on μ is equivalent to a lifting $\rho : N \rightarrow \text{Aut}(M)$.

Example 6.23. 1. Given $\mu : M \hookrightarrow N$, the normal morphisms

$$\rho : N \longrightarrow \text{Aut}(M)$$

$$n \mapsto M \rightarrow M, m \mapsto nmn^{-1}.$$

2. (Whitehead, 1949) Given a Serre fibration $F \rightarrow E \rightarrow B$ of pointed spaces, $\mu : \pi_1(F) \rightarrow \pi_1(E)$ is a crossed module.
3. (Loday) What are group objects in **Cat**? Suppose \underline{G} is a group object in **Cat**, then $\mathcal{N}_*\underline{G}$ is a group object in **sSet**, i.e. a simplicial group. Recall for any simplicial group G_* , its Moore complex is defined by $K_n = \bigcap_{i=1}^n \ker(d_i^{(n)}) \subseteq G_n$ with $\partial_n = d_0^{(n)}$. If we take $G_* = \mathcal{N}_*(\underline{G})$, then its Moore complex is a crossed module.

Lemma 6.24. Given a crossed module $(\mu : M \rightarrow N, \rho)$, consider $A = \ker(\mu)$ and $G = \text{coker}(\mu)$, then

$$1 \longrightarrow A \longrightarrow M \xrightarrow{\mu} N \longrightarrow G \longrightarrow 1$$

1. G is a group and A is a central subgroup of M .
2. the action map $N \times M \rightarrow M, (n, m) \mapsto {}^n m$ induces a well-defined G -structure on A .

Proof. Exercise. □

Fix G and A , define an equivalence relation on the set of all crossed modules $(\mu : M \rightarrow N, \rho)$ such that $A \cong \ker(\mu)$ and $G \cong \text{coker}(\mu)$. The equivalence relation is generated by a map of crossed modules (α, β) as follows.

$$\begin{array}{ccccccccc}
 1 & \longrightarrow & A & \longrightarrow & M & \xrightarrow{\mu} & N & \longrightarrow & G & \longrightarrow & 1 \\
 & & \parallel & & \downarrow \alpha & & \downarrow \beta & & \parallel & & \\
 1 & \longrightarrow & A & \longrightarrow & M' & \xrightarrow{\mu'} & N' & \longrightarrow & G & \longrightarrow & 1
 \end{array}$$

Proposition 6.25. *There is a canonical isomorphism*

$$\mathbf{CrMod}(G, A) / \sim \rightarrow H^3(G, A)$$

$$\left[(M \xrightarrow{\mu} N, \rho) \right]_{\sim} \mapsto k(M, N)$$

where $k(M, N)$ is called the **MacLane invariant of crossed modules**.

{thm:moore}

Theorem 6.26 (Loday). *The following are equivalent:*

1. a crossed module of groups $(\mu : M \rightarrow N, \rho : N \rightarrow \text{Aut}(M))$.
2. a simplicial group $G_* \in \mathbf{sGr}$ whose Moore (nonabelian chain) complex $K_* G$ has length 1.
3. a group object in **Cat**.
4. a “categorical group”, which is a pair (N, G) where $N \subseteq G$ together with two group homomorphisms $s, t : G \rightarrow N$ such that

$$(a) \quad s|_N = t|_N = \text{Id}_N, \text{ and}$$

$$(b) \quad [\ker(s), \ker(t)] = 1.$$

Moore Complex For a simplicial abelian group A_* , there are two chain complexes associated to it.

1. $C_*(A, \mathbb{Z})$ where $C_n(A, \mathbb{Z}) = A$ with differential $\partial_n = \sum_{i=0}^n (-1)^i d_i$.

2. $N_*(A)$ where $N_n(A) = \bigcap_{i=1}^n d_i^{(n)} \subseteq A_n$ with differential $\partial_n = d_0|_{N_n(A)}$.

For a simplicial group, its Moore complex is the nonabelian version of Dold-Kan normalization.

Definition 6.27. The homotopy groups of a simplicial group are defined by $\pi_n(G_*) := H_n(K_*(G), \partial) \cong \pi_n(|G_*|)$ for all $n \geq 0$.

Lemma 6.28. $\pi_n(B|G_*|) = \pi_{n-1}(|G_*|) = H_{n-1}(K_*(G), \partial)$.

Classifying Space of Crossed Modules Given a crossed module $(M \xrightarrow{\mu} N, \rho)$, define \mathcal{C} by

- $\text{Ob}(\mathcal{C}) = N$, and
- $\text{Mor}(\mathcal{C}) = M \rtimes N$ with

$$s : \text{Mor}(\mathcal{C}) \rightarrow \text{Ob}(\mathcal{C})$$

$$(m, n) \longmapsto n$$

and

$$s : \text{Mor}(\mathcal{C}) \rightarrow \text{Ob}(\mathcal{C})$$

$$(m, n) \longmapsto \mu(n) \cdot n$$

Therefore if we take the nerve of \mathcal{C} we will get $B_0\mathcal{C} = N$, $B_1\mathcal{C} = M \rtimes N$ and $B_2\mathcal{C} = M \rtimes (M \rtimes N)$, and $B_*\mathcal{C}$ is a simplicial group.

Recall there is a Kan loop group adjunction

$$G : \mathbf{sSet}_0 \rightleftarrows \mathbf{sGr} : \overline{W} :$$

which is a Quillen adjunction. Here \mathbf{sSet}_0 is the category of reduced simplicial sets X_* , i.e. $X_0 = \{\mathbf{pt}\}$. In particular $|\overline{W}G| \simeq \Omega X$.

Now we take $BB\mathcal{C} = |\overline{W}B_*\mathcal{C}|$ and

$$\pi_i BB\mathcal{C} = \begin{cases} \text{coker } \mu, & i = 1, \\ \ker \mu, & i = 2, \\ 0, & i \geq 3. \end{cases}$$

and $k \in H^3(B\pi_1, \pi_2) \cong H^3(BG, A)$.

Proof of Theorem 6.26. 1. $1 \implies 2$: take $B_*C(M, N)$ where $s_0 : N \hookrightarrow M \rtimes N$ is the natural inclusion.

2. $3 \implies 2$: take $K_*(B_*C) \cong \{N \xleftarrow{\partial_0=d_0} \ker d_1 \leftarrow 1\}$, where $d_0 s_0 = d_1 s_0 = \text{Id} \iff s|_N = t|_N = \text{Id}_N$.
3. $4 \implies 1$ comes from the observation that $[\ker(d_0), \ker(d_1)] = 1 \iff$ axiom (b) in the definition of crossed modules.

□

Question 6.29. What is a correct definition of normal maps of groups? What is the homotopy invariant version of normality?

Question 6.30. For a noncommutative ring spectrum, what does it mean to be simple?

Homotopy Normal Maps Let $\mu : M \rightarrow N$ be a map between (simplicial) groups in **sGr**.

In **Gr**, μ is normal if and only if

- μ is injective,
- $\text{Im}(\mu) \trianglelefteq N$.

This gives an exact sequence

$$M \longrightarrow N \longrightarrow N/M$$

which gives a homotopy fibration sequence

$$BM \longrightarrow BN \longrightarrow B(N/M)$$

Definition 6.31. A map of groups $\mu : M \rightarrow N$ is **homotopy normal** if there exists a pointed connected space with $\alpha : BN \rightarrow W$ such that

$$BM \longrightarrow BN \longrightarrow W$$

is a homotopy fibration sequence. Or equivalently, $B\mu \simeq \text{hofib}_*(\alpha)$.

Remark. For any pointed map of connected spaces, $f : X \rightarrow Y$, there is a Puppe sequence

$$\cdots \longrightarrow \Omega X \longrightarrow \Omega Y \longrightarrow \Omega X // \Omega Y \longrightarrow X \longrightarrow Y \xrightarrow{\alpha} W$$

Theorem 6.32 ([FH], [P]). (Higher normality) A map $\mu : M \rightarrow N$ is homotopy normal (in **sGr**) if there exists a simplicial group Γ_* with an isomorphism $\Gamma_0 \cong N$ which extends to a N -equivariant isomorphism of simplicial sets

$$\text{hocolim}_M(N) = N \times_M EM \xrightarrow{\cong} \Gamma_*$$

Theorem 6.33. *To give such an extension is equivalent to extend μ to a crossed module $(\mu : M \rightarrow N, \rho)$.*

Recall for a crossed module $(M \xrightarrow{\mu} N, \rho)$ we associate it with a simplicial group $B_*C(M, N)$, which gives us $BBC = B|B_*C|$ the classifying space of crossed modules.

$$\begin{aligned}\pi_1(BBC) &\cong \text{coker}(\mu), \\ \pi_2(BBC) &\cong \ker(\mu), \\ \pi_i(BBC) &= 0, \quad i \geq 3.\end{aligned}$$

Example 6.34. For a discrete group G together with $z \in \mathbb{Z}(G)$, we have $B_*(G, z)$ the twisted nerve of G and $X(G, z) := ES^1 \times_{S^1} |B_*(G, z)|$. To describe the homotopy type of this space depends on (G, z) .

Observe there is a sphere fibration

$$\begin{array}{ccccc} S^1 & \longrightarrow & ES^1 \times B(G, z) & \twoheadrightarrow & ES^1 \times_{S^1} B(G, z) \\ \parallel & & \downarrow \simeq & & \parallel \\ S^1 & \longrightarrow & \{*\} \times BG & \longrightarrow & X(G, z) \end{array}$$

Consider $\mu : \mathbb{Z} \rightarrow G, n \mapsto z^n$ together with the trivial G -action $\rho : G \rightarrow \text{Aut}(\mathbb{Z})$, this crossed module (μ, ρ) gives us a group homomorphism

$$\begin{aligned}\gamma_z : \mathbb{Z} \times G &\longrightarrow G \\ (n, g) &\longmapsto z^n g = g z^n\end{aligned}$$

which induces

$$S^1 \times BG \cong B(\mathbb{Z} \times G) \xrightarrow{B\gamma_z} BG$$

or equivalently, a fibration

$$B\mathbb{Z} \xrightarrow{\mu} BG \xrightarrow{\alpha} X(G, z).$$

By Proposition 1, $X(G, z)$ is the classifying space of $BBC(\mathbb{Z}, G, \rho)$, and

$$\begin{aligned}\pi_1(X(G, z)) &\cong \text{coker}(\mu) = G/\text{Im}(\mu), \\ \pi_2(X(G, z)) &\cong \ker(\mu), \\ \pi_i(X(G, z)) &= 0, \quad i \geq 3.\end{aligned}$$

Proposition 6.35. 1. *If z has infinite order, then $X(G, z) \cong B(G/\langle z \rangle)$.*

2. *If z has finite order, then*

$$\begin{aligned}\pi_1(X(G, z)) &\cong G/\langle z \rangle, \\ \pi_2(X(G, z)) &\cong \mathbb{Z}, \\ \pi_i(X(G, z)) &= 0, \quad i \geq 3.\end{aligned}$$

In the second case, π_1 acts trivially in π_2 ,

$$k(G, z) \in H^3(B\pi_1, \pi_2) = H^3(G/\langle z \rangle, \mathbb{Z}).$$

When $k(G, z) = 0$, by Postnikov decomposition,

$$X(G, z) = K(G/\langle z \rangle, 1) \times K(\mathbb{Z}, 1) = B(G/\langle z \rangle) \times BS^1.$$

This happens in two examples.

1. When G is any discrete group and $z = 1$, $B(G, 1) = BG$ and $X(G, 1) = BG \times BS^1$.
2. When G is abelian and $z \in G$ has finite order, $X(G, z) = B(G/\langle z \rangle) \times BS^1$.
Note in this case, G and \mathbb{Z} are abelian groups and G acts on \mathbb{Z} trivially, so $B_n \mathcal{C} = \mathbb{Z} \times G^n$ and $B_* \mathcal{C}$ is an abelian simplicial group. By a theorem in [M], any abelian simplicial group is the product of Eilenberg-MacLane spaces

$$|\Gamma_*| \simeq \prod_{i \geq 1} K(\pi_i(\Gamma_*), i)$$

or equivalently, Postnikov's invariants of $|\Gamma_*|$ are trivial.

Theorem 6.36. *For any group and any finite order $z \in \mathcal{Z}(G)$, there is a rational equivalence*

$$X(G, z) \simeq_{\mathbb{Q}} B(G/\langle z \rangle)_{\mathbb{Q}} \times BS^1_{\mathbb{Q}}.$$

6.6 Cyclic Nerve

Kapranov-Dyckenhoff 2015 [DK]

Definition 6.37. Let

$$\Psi : \Delta C \rightarrow \mathbf{Cat}$$

$$[n] \mapsto \Psi([n])$$

where $\Psi([n])$ is the path (free) category of the cyclic quiver

$$\begin{array}{ccccccc} 0 & \xleftarrow{x_1} & 1 & \xleftarrow{\quad} & \dots & \xleftarrow{x_n} & n \\ & & & & & & \nearrow x_0 \\ & & & & & & \end{array}$$

Then the cyclic nerve is

$$\mathcal{N}_*^{\text{cyc}} : \mathbf{Cat} \rightarrow \mathbf{Set}^{\Delta C^{\text{op}}}$$

$$C \longmapsto \mathcal{N}_*^{\text{cyc}} C$$

where

$$\mathcal{N}_n^{\text{cyc}} \mathcal{C} = \{c_0 \xleftarrow{x_1} c_1 \xleftarrow{x_2} c_2 \leftarrow \cdots \xleftarrow{x_n} c_n \xleftarrow{x_0} c_0\}$$

The structure maps are given as follows. For any $f : [m] \rightarrow [n]$, there is an unique factorization $f = \tau \circ \alpha$,

- For $\alpha \in \text{Hom}_{\Delta}([m], [n])$,

$$\mathcal{N}_{\alpha}^{\text{cyc}}(\mathcal{C}) : (c_0 \leftarrow c_1 \leftarrow \cdots \leftarrow c_n \leftarrow c_0) = (c_{\alpha(0)} \leftarrow c_{\alpha(1)} \leftarrow \cdots \leftarrow c_{\alpha(n)} \leftarrow c_{\alpha(0)}).$$

- For $\tau \in \text{Hom}_{\Delta C}([n], [n])$,

$$\mathcal{N}_{\tau}^{\text{cyc}}(\mathcal{C}) : (c_0 \leftarrow c_1 \leftarrow \cdots \leftarrow c_n \leftarrow c_0) = (c_n \leftarrow c_0 \leftarrow \cdots \leftarrow c_{n-1} \leftarrow c_n).$$

Lemma 6.38. 1. If G is a group, $\Gamma_*(G) = \mathcal{N}_*^{\text{cyc}}(\underline{G})$.

2. $\mathcal{N}_*^{\text{cyc}}$ is the restriction functor for certain map of small categories $b^* : \Delta C \rightarrow \mathcal{G} \subset \mathbf{Gr}$, i.e. $\mathcal{N}_*^{\text{cyc}} = (b^{\text{cyc}})^*$.

Proof. Consider the following adjunctions

$$\mathcal{U} : \mathbf{Cat} \rightleftarrows \mathbf{Grpd} \rightleftarrows \mathbf{Gr} : (-)$$

Define

$$b^{\text{cyc}} = \Psi : \Delta C \longrightarrow \mathbf{Cat}$$

$$[n] \longmapsto \mathbb{F}^{n+1} = \mathbb{F}\langle x_0, \dots, x_n \rangle$$

Check

$$\Gamma_n(G) = G^{n+1} \cong \text{Hom}_{\mathbf{Gr}}(b^{\text{cyc}}([n]), G) \cong \text{Hom}_{\mathbf{Gr}}(\mathcal{U}\Psi([n]), G) \cong \text{Hom}_{\mathbf{Cat}}(\Psi([n]), \underline{G}) = \mathcal{N}_n(\underline{G})$$

on morphisms. \square

Theorem 6.39. For any (simplicial topological) group G , there is a canonical homotopy equivalence

$$\gamma : |\Gamma_* G| \xrightarrow{\sim} \mathcal{L}(BG) = \text{Map}(S^1, BG)$$

which is S^1 -equivariant.

The left S^1 -action comes from cyclic structure and the right S^1 -action comes from the action of S^1 on itself.

Corollary 6.40. There are canonical equivalences

1. $\text{HH}_*(k[G]) \cong H_*(\mathcal{L}(BG); k)$.
2. $\text{HC}_*(k[G]) \cong H_*^{S^1}(\mathcal{L}(BG); k)$.

6.6.1 Burghelea Decomposition

Free Loop Space Let X be a pointed connected space. $\mathcal{L}X = \text{Map}(S^1, X)$.

The S^1 -action on $\mathcal{L}X$ is given by

$$\begin{aligned} S^1 \times \mathcal{L}X &\longrightarrow \mathcal{L}X \\ (z, \theta) &\longmapsto (z, \theta) : z' \mapsto \theta(z \cdot z') \end{aligned}$$

Theorem 6.41 (Goodwillie, Loday 2016). *For any topological group G , there is a canonical S^1 -equivariant homotopy equivalence*

$$|\Gamma_* G| \xrightarrow{\sim} \mathcal{L}BG$$

Proof. (Detailed sketch) Let G be a discrete group, there is a simplicial map

$$\begin{aligned} \text{pr}_* : \Gamma_* G &\longrightarrow B_* G = B_*(G, 1) \\ (g_0, \dots, g_n) &\longmapsto (g_1, \dots, g_n) \end{aligned}$$

which can be extended to a morphism of cyclic sets.

CHECK

Define

$$\gamma^\# : S^1 \times |\Gamma_* G| \xrightarrow{\text{act}} |\Gamma_* G| \xrightarrow{|\text{pr}_*|} |B_* G|$$

Note $\gamma^\#$ is S^1 -equivariant (because both are). γ is defined to be the adjoint map

$$\gamma : |\Gamma_* G| \longrightarrow \text{Map}(S^1, BG)$$

and γ is S^1 -equivariant because $\gamma^\#$ is. Recall $|B_* G| = |B_*(G, 1)|$ comes with trivial S^1 -action, so S^1 -action on $\mathcal{L}BG$ is the standard one.

Need to check γ is a homotopy equivalence.

If X is pointed, there is a natural fibration sequence

$$\Omega X \longrightarrow \mathcal{L}X \xrightarrow{\text{ev}} X$$

For $X = |B_* G|$, we have

$$\begin{array}{ccccc} G & \xrightarrow{\text{in}} & |\Gamma_* G| & \xrightarrow{|\text{pr}|} & |B_* G| \\ \sim \downarrow \overline{\gamma} & & \downarrow \gamma & & \\ \Omega BG & \longrightarrow & \mathcal{L}BG & \xrightarrow{\text{ev}} & BG \end{array}$$

where

$$\begin{aligned} G &\longrightarrow \Gamma_* G \\ g &\mapsto (g, 1, \dots, 1) \end{aligned}$$

Lemma 6.42. $\bar{\gamma} : G \rightarrow \Omega BG$ is the classical homotopy equivalence.

Remark. $\bar{\gamma}$ is called a **topological group completion**. Quillen shows that $\bar{\gamma}$ induces

$$\begin{array}{ccc} H_*(M, k) & \xrightarrow{\bar{\gamma}} & H_*(\Omega BM, k) \\ & \searrow & \nearrow \cong \\ & H_*(M, k)[\pi_0(M)^{-1}] & \end{array}$$

where $\pi_0(M)^{-1}$ is a 2-sided Ore set.

For M a topological monoid, $\bar{\gamma} : M \rightarrow \Omega BM$ is homotopy equivalence if and only if M is group like, $\pi_0(M)$ is a monoid group.

Proof. We will describe the classical map $\bar{\gamma}'$ simplicially and show that $\bar{\gamma}' = \bar{\gamma}$.

Recall that $\bar{\gamma}'$ is defined by looking at its adjoint

$$\begin{aligned} (\bar{\gamma}')^\# : S^1 \times G &\longrightarrow BG \\ C_1 \times G &\longrightarrow B_1 G = G \\ (t, g) &\longmapsto g \end{aligned}$$

Check there is a unique simplicial map that extends to a map $C_* \times G \rightarrow B_* G$.

On the other hand, by definition of S^1 -action on $|\Gamma_* G|$

$$\begin{array}{ccc} S^1 \times |\Gamma_* G| & \xrightarrow{(p_1, p_2)^{-1}} & |F(\Gamma_* G)| \xrightarrow{|ev|} |\Gamma G| \\ \uparrow |in| & & \\ S^1 \times G = |C_* \times G| & & \end{array}$$

and $\bar{\gamma}'$ is the realization of

$$C_* \times G \xrightarrow{in} C_* \times \Gamma_* G \xrightarrow{ev} \Gamma_* G \xrightarrow{pr} B_* G$$

$$(t, g) \longmapsto (t, (g, 1)) \longmapsto t_*(g, 1) = (1, g) \longmapsto g$$

which can be identified with $\bar{\gamma}$. □

□

Corollary 6.43. For any discrete group G and any commutative ring k ,

$$\begin{aligned} HH_*(k[G]) &\cong H_*^{\text{simp}}(\mathcal{L}(BG), k) \cong H_*^{\text{sing}}(\mathcal{L}(BG), k) \\ HC_*(k[G]) &\cong H_*^{S^1}(\mathcal{L}BG, k). \end{aligned}$$

Proof. Recall $k[\Gamma_* G] \cong C_*^{\text{Hoch}}(k[G])$ (by inspection), so

$$HH_*(k[G]) := H_*(C_*^{\text{Hoch}}(k[G]) \cong H_*(k[\Gamma_* G]) \cong H_*^{\text{simp}}(\mathcal{L}BG).$$

□

MacLane Isomorphism Let $A = k[G]$. Let M be a bimodule over A

$$HH_*(k[G], M) \xrightleftharpoons[\Phi^{-1}]{\Phi} H_*^{\text{EM}}(G, \tilde{M})$$

where \tilde{M} is the (right) G -module defined by $\tilde{M} = M$ as k -module.

$$\tilde{M} \times G \longrightarrow \tilde{M}$$

$$(m, g)n \mapsto m^g = g^{-1}mg$$

Proof. Let $C_n^{\text{Hoch}}(k[G], M) := (M \otimes_k k[G]^{\otimes n}, d^{\text{Hoch}})$ and $C_n^{\text{EM}}(G, \tilde{M}) := (\tilde{M} \times k[G^n], d^{\text{EM}})$ where

$$d_n^{\text{EM}} : C_n^{\text{EM}} \longrightarrow C_{n-1}^{\text{EM}}$$

$$(m, g_1, \dots, g_n) \mapsto (m^{g_1}, g_2, \dots, g_n) + \sum_{i=1}^{n-1} (-1)^i (m, g_1, \dots, g_i g_{i+1}, \dots, g_n) + (-1)^n (m, g_1, \dots, g_{n-1}).$$

For

$$\Phi_n : C_n^{\text{Hoch}}(k[G], M) \xrightleftharpoons{\quad} C_n^{\text{EM}}(k[G], \tilde{M}) : \Phi_n^{-1}$$

$$(m, g_1, \dots, g_n) \longmapsto (g_1 \cdots g_n \cdot m, g_1, \dots, g_n)$$

$$(g_n^{-1} \cdots g_1^{-1} \tilde{m}, g_1, \dots, g_n) \longleftarrow (\tilde{m}, g_1, \dots, g_n)$$

check $\phi_{n+1} \circ d^{\text{Hoch}} \cong d^{\text{EM}} \circ \Phi_n$, which induces isomorphism on cohomology.

□

Burghlelea decomposition Ref: Quillen diaries 1992 April-October, Clay Institute, Quillen notebooks, “cyclic torsors” over Segal groupoids.

Burghlelea, Goodwillie-Fiedorowicz

For any $z \in G$, $G_z = C_G(z) = \{g \in G : gz = zg\}$. Note $z \in G_z$, so we can define $B_*(G_z, z)$ and $X(G_z, z) = ES^1 \times_{S^1} |B(G_z, z)|$. And $\mathbb{Z} \rightarrow G_z, n \mapsto z^n$ defines a crossed module.

Note G_z and $X(G_z, z)$ depends only on the conjugacy class of z .

The set of conjugacy classes of all elements in G

$$\langle G \rangle = \langle G \rangle^{\text{Fin}} \coprod \langle G \rangle^{\infty}$$

consisting of finite order ones and infinite order ones.

Recall that $\Gamma_*(G, z)$ contains cyclic subsets $\Gamma_*(G, z)$ and we have a cyclic isomorphism $\Gamma_* G \cong \coprod_{\langle z \rangle} \Gamma(G, z)$, which comes from a homotopy decomposition, where

$$\Gamma_* G \cong \operatorname{hocolim}_G (\operatorname{Ad}) \xrightarrow{\pi_0} \operatorname{colim}_G (\operatorname{Ad}) = \langle G \rangle$$

and $\Gamma_*(G, z) = \operatorname{fiber}_{\langle z \rangle}(\pi_0)$, which implies

Lemma 6.44. $k[\Gamma_* G] \cong \bigoplus_{\langle z \rangle} k[\Gamma_*(G, z)]$.

Proof. Apply $k[-]$. □

Lemma 6.45 (Loday, Fiedorowicz). *For every $z \in G$, there is a natural inclusion*

{lem:classifying-cyclic}

$$i : B_*(G_z, z) \longrightarrow \Gamma_*(G, z)$$

$$(g_1, \dots, g_n) \mapsto ((g_1 \cdots g_n)^{-1} z, g_1, \dots, g_n)$$

with properties

1. i is a map of cyclic sets.
2. $i_* : k[B_*(G_z, z)] \rightarrow k[\Gamma_*(G, z)]$ induces an isomorphism on cyclic homology.

Proof. 1. Check

$$\begin{array}{ccc} B_n(G_z, z) & \xrightarrow{i_n} & \Gamma_n(G, z) \\ \downarrow t_n^B & & \downarrow t_n^\Gamma \\ B_n(G_z, z) & \xrightarrow{i_n} & \Gamma_n(G, z) \end{array}$$

where

$$\begin{aligned} t_n^B(g_1, \dots, g_n) &= ((g_1 \cdots g_n)^{-1} z, g_2, \dots, g_n) \\ t_n^\Gamma(g_0, \dots, g_n) &= (g_n, \dots, g_0, \dots, g_{n-1}) \end{aligned}$$

2. By the next lemma, it suffices to show that

$$i : k[B_*(G_z, z)] \hookrightarrow k[\Gamma_*(G, z)]$$

is Hochschild equivalence. Notice i factors as

$$\begin{aligned} k[B(G_z, z)] &= k[B_*(G_z)] \cong C_*^{\operatorname{Hoch}}(k[G_z] \xrightarrow{\alpha} C_*^{\operatorname{EM}}(G, k[\langle z \rangle])) \\ &\cong \downarrow \Phi^{-1} \\ &k[\Gamma_*(G, z)] \\ &\downarrow \\ &k[\Gamma_* G] \end{aligned}$$

where α is an isomorphism by Shapiro lemma: for $G_z \subseteq G$,

$$H_*(G_z, k) \cong H_*(G, \text{Ind}_{G_z}^G(k)),$$

$$\text{and } \text{Ind}_{G_z}^G(k) = k[\langle \rangle].$$

□

Lemma 6.46. *A map of cyclic k -modules $f : E \rightarrow E'$ is a cyclic equivalence*

$$f_* : \text{Tor}_*^{\Delta \text{C}^{\text{op}}}(k, E) \cong \text{Tor}_*^{\Delta \text{C}^{\text{op}}}(k, E')$$

if and only if

$$f_* : \text{Tor}_*^{\Delta \text{op}}(k, E) \cong \text{Tor}_*^{\Delta \text{op}}(k, E')$$

Proof. It follows from Connes' Periodicity sequence

$$\cdots \longrightarrow \text{HH}_n(E) \longrightarrow \text{HC}_n(E) \longrightarrow \text{HC}_{n-2}(E) \longrightarrow \text{HH}_{n-1}(E) \longrightarrow \cdots$$

which is the Gysin sequence of the following Borel fibration

$$\cdots \longrightarrow S^1 \longrightarrow X \longrightarrow \text{ES}^1 \times_{S^1} X \longrightarrow \text{BS}^1$$

□

6.6.2 Cyclic Homology of Group Algebras

Theorem 6.47. *Let G be a discrete group and k be a commutative ring.*

$$\text{HH}_*(k[G]) \cong \bigoplus_{\langle z \rangle \in \langle G \rangle} H_*(k[B_*(G_z, z)])$$

$$\text{HC}_*(k[G]) \cong \bigoplus_{\langle z \rangle \in \langle G \rangle} \text{HC}_*(k[B_*(G_z, z)])$$

$$\text{HC}_*(k[G]) \cong \bigoplus_{\langle z \rangle \in \langle G \rangle} H_*^{\text{top}}(X_*(G_z, z), k)$$

Notation For any $z \in G$, $G_z = C_G(z)$ and $z \in \mathcal{Z}(G_z)$. The twisted nerve of (G_z, z) is $B(G_z, z) \in \mathbf{Set}^{\Delta \text{C}^{\text{op}}}$.

Corollary 6.48. *If G is torsion-free,*

$$\begin{aligned} \text{HC}_*(k[G]) &\cong \bigoplus_{\langle z \rangle \in \langle G \rangle, z \neq 1} H_*(G_z / \langle z \rangle, k) \bigoplus H_*(\text{BS}^1 \times BG, k) \\ &\cong \bigoplus_{\langle z \rangle \in \langle G \rangle, z \neq 1} H_*(G_z / \langle z \rangle, k) \bigoplus \bigoplus_{i \in \mathbb{N}} H_{*-2i}(G, k) \end{aligned}$$

Corollary 6.49. *For abelian group G ,*

6.7 Homotopy Simplicial Groups

Example of homotopy simplicial \mathcal{T} -algebras over a (simplicial) algebraic theory \mathcal{T} of groups.

Definition 6.50. Let \mathcal{G} be the (skeleton) of the full subcategory of \mathbf{Gr} consisting of finitely generated free groups with

- objects $\{\langle n \rangle = \mathbb{F}\langle x_1, \dots, x_n \rangle\}_{n \geq 0}$, and
- morphisms $\mathrm{Hom}_{\mathcal{G}}(\langle n \rangle, \langle m \rangle) = \mathrm{Hom}_{\mathbf{Gr}}(\langle n \rangle, \langle m \rangle)$.

There is a strict monoidal structure on \mathcal{G} with $\langle n \rangle \coprod \langle m \rangle = \langle n \rangle * \langle m \rangle = \langle n + m \rangle$. $\mathcal{G}^{\mathrm{op}}$ has an opposite monoidal structure \coprod .

Starting Point. Consider the following composite

$$\mathbf{Gr} \hookrightarrow \mathrm{Fun}(\mathbf{Gr}^{\mathrm{op}}, \mathbf{Set}) \xrightarrow{\mathrm{Res}} \mathrm{Fun}(\mathcal{G}^{\mathrm{op}}, \mathbf{Set})$$

which gives

$$\mathbf{Gr} \xrightarrow{\cong} \mathrm{Fun}^{\otimes}(\mathcal{G}^{\mathrm{op}}, \mathbf{Set})$$

$$\Gamma \longmapsto \underline{\Gamma} : \langle n \rangle \mapsto \mathrm{Hom}_{\mathbf{Gr}}(\langle n \rangle, \Gamma) = \Gamma^n$$

Note for each $n \geq 0$ and $1 \leq k \leq n$, there are maps

$$i_{n,k} : \langle 1 \rangle \longrightarrow \langle n \rangle = \coprod_k \langle 1 \rangle$$

$$x \longmapsto x_k$$

which corresponds to opposite maps $p_{n,k}$ in $\mathcal{G}^{\mathrm{op}}$.

If we replace \mathbf{Gr} by \mathbf{sGr} and \mathbf{Set} by \mathbf{sSet} , we have similar argument for simplicial groups and given any functor $A : \mathcal{G}^{\mathrm{op}} \rightarrow \mathbf{sSet}$, we can patch maps $p_{n,k}$ to get

$$A(p_n) := \prod_{k=1}^n A(p_{n,k}) : A(\langle n \rangle) \longrightarrow \prod_{i=1}^k A(\langle 1 \rangle) = A(\langle n \rangle)^n$$

Definition 6.51. $A : \mathcal{G}^{\mathrm{op}} \rightarrow \mathbf{sSet}$ is **(strictly) product-preserving** if $A(p_n)$ are isomorphisms in \mathbf{sSet} for any $n \geq 1$. Then

$$\mathbf{sGr} \cong \mathrm{Fun}^{\otimes}(\mathcal{G}^{\mathrm{op}}, \mathbf{sSet})$$

and $A(p_0) = \mathbf{pt}$.

Definition 6.52 (B, Badzioch [B]). A **homotopy simplicial group** is a functor $A : \mathcal{G}^{\mathrm{op}} \rightarrow \mathbf{sSet}$ such that $A(p_n)$ are weak homotopy equivalences in \mathbf{sSet} .

Remark. The full subcategory of all weakly product-preserving functors cannot carry a model structure because it is not closed under colimits. (Examples?)

Badzioch Model Category is a model structure on $\mathbf{sSet}^{\mathcal{G}^{\text{op}}}$ in which homotopy simplicial groups are fibrant objects.

Definition 6.53. Let $\mathbf{sGr}^h := L_{\mathcal{S}}(\mathbf{sSet}^{\mathcal{G}^{\text{op}}})$ be the left Bousfield localization of $\mathbf{sSet}^{\mathcal{G}^{\text{op}}}$ with BK projective model structure with respect to

$$\mathcal{S} = \{j_n : \coprod_{k=1}^n \text{Hom}_{\mathcal{G}}(-, \langle 1 \rangle) \xrightarrow{\coprod_{k=1}^n (i_{n,k})^*} \text{Hom}_{\mathcal{G}}(-, \langle n \rangle)\}_{n \geq 1}$$

The localization induces a Quillen pair

$$j = \text{Id} : \mathbf{sSet}^{\mathcal{G}^{\text{op}}} \rightleftarrows \mathbf{sGr}^h : \text{Id} = q$$

which induces an adjunction on homotopy category

$$\overline{\text{Id}} : \text{Ho}(\mathbf{sSet}^{\mathcal{G}^{\text{op}}}) \rightleftarrows \text{Ho}(\mathbf{sGr}^h) : \mathbb{R}\text{Id}$$

Definition 6.54. The localization with respect to \mathcal{S} is the adjunction unit $\{L_{\mathcal{S}}(X) : X \rightarrow \mathbb{R}q \circ \bar{j}(X)\}_{X \in \text{Ob}(\text{Ho}(\mathbf{sSet}^{\mathcal{G}^{\text{op}}}))}$ of this pair

$$\bar{j} : \text{Ho}(\mathbf{sSet}^{\mathcal{G}^{\text{op}}}) \rightleftarrows \text{Ho}(\mathbf{sGr}^h) : \mathbb{R}q$$

Proposition 6.55 ([B]). *Homotopy simplicial groups are precisely the fibrant objects in \mathbf{sGr}^h (which are objectwise fibrant as simplicial set).*

The natural inclusion $J : \mathbf{sGr} \hookrightarrow \mathbf{sSet}^{\mathcal{G}^{\text{op}}}$ has a left adjoint

$$K : \mathbf{sSet}^{\mathcal{G}^{\text{op}}} \longrightarrow \mathbf{sGr}$$

where $J(\Gamma) = \underline{\Gamma}$ and

$$K(\Gamma) := \int^{n \in \text{Ob}(\mathcal{G})} \underline{\Gamma}(\langle n \rangle) \boxtimes \mathbb{F}\langle n \rangle.$$

Theorem 6.56 (Rigidification Theorem). 1. $\mathbb{L}K$ maps $\bar{S} \in \text{Ho}(\mathbf{sSet}^{\mathcal{G}^{\text{op}}})$ to isomorphisms in $\text{Ho}(\mathbf{sGr})$. Hence $K^h : \mathbf{sGr}^h \rightleftarrows \mathbf{sGr} : J^h$ is a Quillen pair.

2. (K^h, J^h) is a Quillen equivalence, $\text{Ho}(\mathbf{sGr}^h) \cong \text{Ho}(\mathbf{sGr})$.

Remark. 1. The above theorem generalizes to (multi-sorted) algebraic theories.

Remark. If we consider

$$\mathcal{G} \hookrightarrow \mathbf{Gr} \hookrightarrow \text{Ho}(\mathbf{Top}_{0,*})$$

$$\langle n \rangle \longmapsto B\langle n \rangle$$

in $\text{Ho}(\mathbf{Top}_{0,*})$, notice $\mathcal{G} \cong \mathcal{B}$ where $\text{Ob}(\mathcal{B}) = \{\bigvee_{i=1}^n S^1\}_{n \geq 0}$.

6.7.1 Cyclic Homology

Let Γ be a discrete group, we can define

$$\begin{aligned} B^{cyc}\Gamma : \Delta C^{op} &\rightarrow \mathbf{Set} \\ [n] &\longmapsto \Gamma^{n+1} \end{aligned}$$

and extend to a functor

$$\begin{aligned} k[-] : \mathbf{Set}^{\Delta C^{op}} &\longrightarrow \mathbf{Mod}_k^{\Delta C^{op}} \\ (\Delta C^{op} \rightarrow \mathbf{Set}) &\mapsto (\Delta C^{op} \rightarrow \mathbf{Set} \xrightarrow{k[-] \mathbf{Mod}_k}) \end{aligned}$$

and we have $k[B^{cyc}\Gamma] \cong C_*^{Hoch}(k[\Gamma])$.

Key Observation There is a functor

$$\begin{aligned} \Psi_{cyc} : \Delta C &\longrightarrow \mathcal{G} \\ [n] &\longmapsto \langle n+1 \rangle \end{aligned}$$

which induces

$$\Psi_{cyc}^*(\Gamma) : \Delta C^{op} \longrightarrow \mathcal{G}^{op} \xrightarrow{\Gamma} \mathbf{Set}$$

such that $B^{cyc}\Gamma \cong \Psi_{cyc}^*(\Gamma)$.

Define for any $\Gamma : \mathcal{G}^{op} \rightarrow \mathbf{sSet}$, we can define

$$B^{cyc}(\Gamma) := \Psi_{cyc}^*(\Gamma) : \Delta C^{op} \longrightarrow \mathbf{sSet}$$

and

$$k[B^{cyc}(\Gamma)] = k[\Psi_{cyc}^*(\Gamma)] : \Delta C^{op} \longrightarrow \mathbf{sMod}_k \xrightarrow{N} \mathbf{Ch}_k$$

Lemma 6.57. *If Γ is a homotopy simplicial group, then $HC_*(k[\Gamma])$ depends only on S -local weak equivalence class of Γ , i.e.*

$$HC_*(k[-]) : (\mathbf{sGr}^h)^{fib} \longrightarrow \mathbf{Mod}_k$$

Corollary 6.58. $HC_*(k[\Gamma])$ depends only on the homotopy type of the classifying space of Γ .

Question 6.59. *What is the center/centralizer of homotopy simplicial group? What is the twisted nerve?*

7 Realizable (Voedvodsky) Homotopy Colimits

Voedvodsky (2010): from BK construction to simplicial descent category, proposed axiomatics for simplicial descent categories.

Barwick-Kan (2012): the fifth model for $(\infty, 1)$ -categories.

B. Rodziguez-Gonzalez [G] (2016) used Barwick-Kan model **RelCat** (the fifth model for ∞ -category) to make it concrete.

Motivation For $f : \mathcal{J} \rightarrow \mathcal{J}$ in **Cat**. Let \mathcal{M} be a (nice) model category, there is a Quillen pair

$$f_! : \mathcal{M}^{\mathcal{J}} \longrightarrow \mathcal{M}^{\mathcal{J}} : f^*$$

where the left adjoint is defined by left Kan extension. Thus we have a (left) derived functor

$$\mathbb{L}f_! : \mathrm{Ho}(\mathcal{M}^{\mathcal{J}}) \longrightarrow \mathcal{M}^{\mathcal{J}}.$$

When $\mathcal{J} = \{*\}$, we have $p : \mathcal{J} \rightarrow \{*\}$ and $\mathbb{L}p_! : (X) = \mathrm{hocolim}_{\mathcal{J}} X$.

Note $\mathbb{L}f_!$ is defined on $\mathrm{Ho}(\mathcal{M}^{\mathcal{J}})$ but not $\mathrm{Ho}(\mathcal{M})^{\mathcal{J}}$, so we want to lift diagrams from $\mathrm{Ho}(\mathcal{M}^{\mathcal{J}})$ to $\mathcal{M}^{\mathcal{I}}$.

Want notion of “hocolim” at the level of the diagram categories.

Example 7.1 (Bousfield-Kan). $\mathrm{hocolim}_{\mathcal{J}}^{\mathrm{BK}} X := |B_*(\mathcal{J}, X)|_{\mathcal{M}}$.

Question 7.2. Replace B_* by “smaller resolutions”.

Theorem 7.3. Homotopy colimits are realizable if and only if they satisfy Voedvodsky’s axiomatics.

7.1 Relative Categories

Definition 7.4. A **relative category** is a pair $(\mathcal{C}, \mathcal{W})$ where

- \mathcal{C} is the underlying category, and
- $\mathcal{W} \subset \mathcal{C}$ is a (wide) subcategory (of weak equivalences in \mathcal{C}).

A **relative functor** $F : (\mathcal{C}, \mathcal{W}_{\mathcal{C}}) \rightarrow (\mathcal{D}, \mathcal{W}_{\mathcal{D}})$ is a functor $F : \mathcal{C} \rightarrow \mathcal{D}$ such that $F(\mathcal{W}_{\mathcal{C}}) \subseteq \mathcal{W}_{\mathcal{D}}$.

Denote by **RelCat** the category of all (small) relative categories and relative functors.

Theorem 7.5 ([BK]). There is a cofibrantly generated left proper model structure on **RelCat** that is Quillen equivalent to **CSS** and **QC**at.

Sketch of Construction

Definition 7.6. A relative category $(\mathcal{C}, \mathcal{W})$ is called

- **maximal** if $\mathcal{W} = \mathcal{C}$.
- **minimal** if $\mathcal{W} = \text{Ob}(\mathcal{C})$ has no identity morphisms.
- a **relative poset** if $\mathcal{C} = \mathcal{P}$ is a poset category.

RelPos is the full subcategory of **RelCat** with objects being relative posets.

Notation For a category \mathcal{C} , we denote by \mathcal{C}_{\max} and \mathcal{C}_{\min} the maximal (resp. minimal) relative model structure on \mathcal{C} .

Example 7.7. Let $\mathcal{C} = [n] = \{0 \rightarrow 1 \rightarrow \dots \rightarrow n\}$ be a poset category, $[n]_{\min}$ and $[n]_{\max}$ are distinguished relative posets.

Definition 7.8. Let $\mathcal{P} \in \mathbf{RelPos}$

- the **terminal subdivision** of \mathcal{P} is the relative poset $\xi_t \mathcal{P}$ with
 - objects are monomorphisms in **RelPos** $\chi : [n]_{\min} \rightarrow \mathcal{P}, \forall n \geq 0$.
 - morphisms are commutative diagrams of the form

$$\begin{array}{ccc} [n]_{\min} & \xrightarrow{\quad} & [m]_{\min} \\ & \searrow \chi_n & \swarrow \chi_m \\ & \mathcal{P} & \end{array}$$

- weak equivalences are commutative diagrams as above such that $\chi_n([n]) \xrightarrow{\sim} \chi_m([m]) \in \mathcal{W}(\mathcal{P})$ are weak equivalences in \mathcal{P} .
- the **initial subdivision** in **RelPos** of \mathcal{P} is given by the relative poset $\xi_i(\mathcal{P})$ such that $\xi_i \mathcal{P} := \xi_t(\mathcal{P}^{\text{op}})^{\text{op}}$. More explicitly, $\xi_i(\mathcal{P})$ is described by
 - objects are monomorphisms $\chi : [n]_{\min} \rightarrow \mathcal{P}^{\text{op}}, \forall n \geq 0$.
 - morphisms are commutative diagrams of the form

$$\begin{array}{ccc} [m]_{\min} & \xrightarrow{\quad} & [n]_{\min} \\ & \searrow \chi_m & \swarrow \chi_n \\ & \mathcal{P}^{\text{op}} & \end{array}$$

- weak equivalences are commutative diagrams as above such that $\chi_n([n]) \xrightarrow{\sim} \chi_m([m]) \in \mathcal{W}(\mathcal{P})$ are weak equivalences in \mathcal{P} .

Note $\xi_t(\mathcal{P})$ and $\xi_i(\mathcal{P})$ come with two projection functors

$$\pi_t : \xi_t(\mathcal{P}) \longrightarrow \mathcal{P}$$

$$(x_n \rightarrow x_m) \longmapsto x_n([n]) \rightarrow x_m([m])$$

$$\pi_i : \xi_t(\mathcal{P}) \longrightarrow \mathcal{P}$$

$$(x_n \rightarrow x_m) \longmapsto x_n([n]) \rightarrow x_m([m])$$

There are canonical extensions of ξ_t and ξ_i to endofunctors on **RelPos**.

$$\xi_t : \mathbf{RelPos} \longrightarrow \mathbf{RelPos}$$

- on objects $\xi_t(\mathcal{P})$
- on morphisms, for each $[n]_{\min} \rightarrow \mathcal{P}$ and $[m]_{\min} \rightarrow \mathcal{Q}$ there is a unique epi $[n]_{\min} \twoheadrightarrow [m]_{\min}$ such that

$$\begin{array}{ccc} [n]_{\min} & \twoheadrightarrow & [m]_{\min} \\ \downarrow & & \downarrow \\ \mathcal{P} & \xrightarrow{f} & \mathcal{Q} \end{array}$$

commutes.

The extension $\xi_i : \mathbf{RelPos} \rightarrow \mathbf{RelPos}$ is defined similarly.

Definition 7.9. The two-fold subdivision of \mathcal{P} is defined by $\xi\mathcal{P} := \xi_i \xi_t \mathcal{P}$.

Remark. $\xi_{\mathcal{P}^{\text{conj}}} = \xi_t \xi_i \mathcal{P} \neq \xi\mathcal{P}$, what would happen to the main theorem in this case?

Example 7.10. Let $\mathcal{P} = [2]_{\min} = \{0 \rightarrow 1 \rightarrow 2\}$. Then

$$\xi_t([2]_{\min}) = \left\{ \begin{array}{c} \begin{array}{ccccc} & & 1 & & \\ & \swarrow & \downarrow & \searrow & \\ 01 & \twoheadrightarrow & 012 & \leftarrow & 12 \\ \swarrow & \nearrow & \uparrow & \nwarrow & \swarrow \\ 0 & \twoheadrightarrow & 02 & \leftarrow & 2 \end{array} \end{array} \right\}$$

and $\xi_i[2]_{\min}$ is given by the opposite diagram.

Definition 7.11. The **classification diagram** (Rezk) of a relative category \mathcal{C}, \mathcal{W} is the simplicial space $\mathcal{N}\mathcal{C}_* \in \mathbf{sSet}^{\Delta^{\text{op}}}$ with

$$\begin{aligned} \mathcal{N}\mathcal{C}_* : \Delta^{\text{op}} &\longrightarrow \mathbf{sSet} \\ [n] &\longmapsto \mathcal{N}\mathcal{C}_n = \mathcal{N}_*(\mathcal{W}(\mathcal{C}^{[n]})) \end{aligned}$$

where \mathcal{N}_* is the usual simplicial nerve and $\mathcal{C}^{[n]} = \text{Fun}([n], \mathcal{C})$ and $\mathcal{W}(\mathcal{C}^{[n]}) := \text{We}(\mathcal{C}^{[n]})$ the natural transformations with objectwise weak equivalences.

Exercise 7.12. Check (by decoding notations) $\mathcal{N}\mathcal{C}_n([m]) = \{[n]_{\min} \times [m]_{\max} \longrightarrow \mathcal{C}\}$.

Lemma 7.13. The classification diagram functor \mathcal{N} has a left adjoint

$$\mathcal{K} : \mathbf{sSet}^{\Delta^{\text{op}}} \rightleftarrows \mathbf{RelCat} : \mathcal{N}$$

such that $\mathcal{K}(\Delta[n]_*^t \times \Delta[m]_*) = [n]_{\min} \times [m]_{\max}$, where $(-)^t$ is the transpose embedding.

In particular \mathcal{K} takes values in $\mathbf{RelPos} \subset \mathbf{RelCat}$. Hence we can define the two-subdivided functor

$$\mathcal{K}_\xi : \mathbf{sSet}^{\Delta^{\text{op}}} \xrightarrow{\mathcal{K}} \mathbf{RelPos} \xrightarrow{\xi = \xi_i \xi_t} \mathbf{RelPos} \hookrightarrow \mathbf{RelCat}$$

Proposition 7.14 ([BK], Section 3). \mathcal{K}_ξ has a right adjoint denoted by

$$\mathcal{N}_\xi : \mathbf{RelCat} \rightleftarrows \mathbf{sSet}^{\Delta^{\text{op}}} : \mathcal{K}_\xi \quad (7.1) \quad \{\text{eq:rel-adj}\}$$

Note there is a natural transformation $\pi : \mathcal{N} \Rightarrow \mathcal{N}_\xi$ induced by $\pi = \pi_t \pi_i : \xi \Rightarrow \text{Id}$.

Recall $\mathbf{sSet}^{\Delta^{\text{op}}}$ carry the Reedy model structure with weak equivalences being levelwise weak equivalences of simplicial sets.

Theorem 7.15 (Main Theorem, [BK], Theorem 6.1). The adjunction 7.1 induces the BK model structure on \mathbf{RelCat} with

- weak equivalences are maps $f : \mathcal{C} \rightarrow \mathcal{D}$ such that $\mathcal{N}_\xi(f)$ are Reedy equivalences in $\mathbf{sSet}^{\Delta^{\text{op}}}$.
- Fibrations are maps $f : \mathcal{C} \rightarrow \mathcal{D}$ such that $\mathcal{N}_\xi(f)$ are Reedy fibrations in $\mathbf{sSet}^{\Delta^{\text{op}}}$.
- cofibrant objects are relative posets.

7.2 2-Category

Assumptions (on relative categories): small

- $(\mathcal{C}, \mathcal{W})$ is saturated: in the formal localization $\gamma : \mathcal{C} \rightarrow \mathcal{C}[\mathcal{W}^{-1}]$, $\gamma^{-1}(\text{Iso}) = \mathcal{W}$. For instance, if $(\mathcal{C}, \mathcal{W})$ comes from a model category, then \mathcal{W} is saturated.
- $(\mathcal{C}, \mathcal{W})$ is closed under finite coproducts. That is, \mathcal{C} has an initial object 0 and both \mathcal{C} and \mathcal{W} are closed under finite coproducts.

Notation If \mathcal{C} is a category and $(\mathcal{D}, \mathcal{W})$ is a relative category, then $\mathcal{D}^{\mathcal{C}} = \text{Fun}(\mathcal{C}, \mathcal{D})$ has obvious relative structure with $\mathcal{W}^{\mathcal{C}} \subset \mathcal{D}^{\mathcal{C}}$ defined pointwise.

Definition 7.16. A **relative natural transformation** $\tau : F \Rightarrow G$ between two functors $F, G : (\mathcal{C}, \mathcal{W}_{\mathcal{C}}) \rightarrow (\mathcal{D}, \mathcal{W}_{\mathcal{D}})$ is a morphism of in $\mathcal{D}^{\mathcal{C}}[(\mathcal{W}^{\mathcal{D}})^{-1}] := \text{Fun}(\mathcal{C}, \mathcal{D})[(\mathcal{W}^{\mathcal{D}})^{-1}]$. More explicitly, τ is represented by zigzags of natural transformations with reversed arrows being weak equivalences.

Proposition 7.17. **RelCat** with relative functors and relative natural transformations is a 2-category.

Definition 7.18. A bicategory is a tuple $(\mathcal{B}, 2, c, a, l, r)$ consisting of the following data:

- Objects (0-cells) are $B_0 = \text{Ob}(\mathcal{B})$
- Hom categories: for any $X, Y \in \text{Ob}(\mathcal{B})$, $B(X, Y)$ is a category with
 - objects (1-cells) $B_1 = \text{Ob}(B(X, Y))$, and
 - morphisms (2-cells) $B_2 = \text{Ob}(B(X, Y))$
 - \circ and Id in $B(X, Y)$ are **vertical compositions** and identity 2-cells.
 - isomorphisms are invertible 2-cells.
- Identity 1-cells: for any $X \in B_0$, $1_X : * \rightarrow B(X, X)$ is a functor called the identity cell of X .
- Horizontal compositions: for any $X, Y, Z \in B_0$, $c_{X,Y,Z} : B(Y, Z) \times B(X, Y) \rightarrow B(X, Z)$.
- Associations: for any $W, X, Y, Z \in B_0$,

$$\alpha_{W,X,Y,Z} : c_{W,Y,Z}(c_{X,Y,Z} \times \text{Id}_{B(W,X)}) \xrightarrow{\sim} c_{W,Y,Z}(\text{Id}_{B(Y,Z)} \times c_{W,X,Y}).$$

a natural isomorphism of functors $B(Y, Z) \times B(X, Y) \times B(W, X) \rightarrow B(W, Z)$.

- Example 7.19.**
1. usual categories are discrete bicategories.
 2. monoidal categories are one object bicategories.
 3. bimodules: B_0 is the set of rings and B_1 are $R - S$ -bimodules and B_2 are bimodule morphisms.

Definition 7.20. A 2-category is a bicategory $(B, 1, c, a, l, r)$ where a, l, r are identity natural transformations.

$$\begin{array}{ccc}
 \mathbf{BiCat} & \xrightleftharpoons{\text{strict}} & 2\text{-Cat} \\
 \uparrow \text{one object} & & \uparrow \\
 \mathbf{MonCat} & \xrightleftharpoons{\text{strict}} & \mathbf{StrictMonCat}
 \end{array}$$

In \mathbf{RelCat} horizontal compositions are compositions of relative functors.

- Definition 7.21.**
- A **relative isomorphism** in \mathbf{RelCat} is an invertible 2-cell, i.e. a relative natural transformation $\tau : F \dashrightarrow G$ which is an isomorphism in $\mathbf{Fun}(F, G)[W^{-1}]$.
 - An equivalence of relative categories is a relative functor $F : (\mathcal{C}, \mathcal{W}_{\mathcal{C}}) \Rightarrow (\mathcal{D}, \mathcal{W}_{\mathcal{D}})$ such that there is a relative functor $G : (\mathcal{D}, \mathcal{W}) \rightarrow (\mathcal{C}, \mathcal{W})$ with relative isomorphisms

$$\tau : F \circ G \xrightarrow{\sim} \text{Id}_{\mathcal{D}}$$

$$\rho : G \circ F \xrightarrow{\sim} \text{Id}_{\mathcal{C}}.$$

Remark. Barwick-Kan (Chapter 33) define relative homotopy equivalence which is a relative equivalence, where ρ and τ are represented by zigzags with all arrows being natural isomorphisms in $\mathbf{Fun}(\mathcal{C}, \mathcal{D})[W^{-1}]$.

Lemma 7.22. *The localization is a 2-functor*

$$\begin{aligned}
 \text{loc} : \mathbf{RelCat} &\longrightarrow \mathbf{Cat} \\
 (\mathcal{C}, \mathcal{W}) &\longmapsto \mathcal{C}[W^{-1}]
 \end{aligned}$$

Lemma 7.23. *For any small $I \in \mathbf{Cat}$, the exponentiation functor is a 2-functor:*

$$\begin{aligned}
 (-)^I : \mathbf{RelCat} &\rightarrow \mathbf{RelCat} \\
 (\mathcal{C}, \mathcal{W}) &\longmapsto (\mathcal{C}^I, \mathcal{W}^I)
 \end{aligned}$$

Definition 7.24. A **relative adjunction** consists of

- relative functors $F : (\mathcal{C}, \mathcal{W}) \rightarrow (\mathcal{D}, \mathcal{W})$. and $G : (\mathcal{D}, \mathcal{W}) \rightarrow (\mathcal{C}, \mathcal{W})$

- relative natural transformations

$$\alpha : F \circ G \rightrightarrows \text{Id}_{\mathcal{D}}$$

$$\beta : \text{Id}_{\mathcal{C}} \rightrightarrows G \circ F$$

such that the triangle identities

$$F \xrightarrow{F\beta} FGF \xrightarrow{\alpha F} F$$

$$G \xrightarrow{\beta G} GFG \xrightarrow{G\alpha} G$$

are identity 1-cells.

Example 7.25. 1. Usual adjoint functors are relative adjunctions

$$F : (\mathcal{C}, \text{Iso}) \rightleftarrows (\mathcal{D}, \text{Iso}) : G$$

2. More generally, any relative adjoint functors for a relative adjunction.
3. A Quillen adjunction between two cofibrantly generated model categories

$$F : \mathcal{M} \rightleftarrows \mathcal{N} : G$$

with functorial fibrant and cofibrant replacements $Q : \mathcal{M} \rightarrow \mathcal{M}$ and $R : \mathcal{N} \rightarrow \mathcal{N}$, then

$$F \circ Q : (\mathcal{M}, \mathcal{W}) \rightleftarrows (\mathcal{N}, \mathcal{W}) : G \circ R$$

is a genuine relative adjunction.

4. (Conjecturally) Model approximation of Quillen Adjunctions

Example 7.26 (Thomason subdivision). $\mathcal{N}^{\text{op}} : \mathbf{Cat}^{\text{op}} \rightleftarrows \mathbf{sSet}^{\text{op}} : \mathcal{C}^{\text{op}}$ is a relative adjunction with respect to Thomason subdivision.

Example 7.27. Let G be an affine algebraic group (scheme) over a field k , e.g. $G = \text{GL}_n$. There is an adjunction

$$(-)_G : \mathbf{Gr} \rightleftarrows \mathbf{Comm}_k : G$$

$$\Gamma \longmapsto \mathcal{O}[\text{Rep}_G(\Gamma)]$$

where $\mathcal{O}[\text{Rep}_G(\Gamma)]$ is the affine scheme of G -representations of Γ $\rho_A : \Gamma \rightarrow G(A), \forall A \in \mathbf{Comm}_k$. This adjunction extends to an adjunction in simplicial setting

$$(-)_G : \mathbf{sGr} \rightleftarrows \mathbf{sComm}_k : G$$

which however is not a Quillen pair. Nevertheless, it behaves like a Quillen pair.

1. $\mathbb{L}(-)_G$ and $\mathbb{R}G$ exists.
2. $\mathbb{L}(-)_G : \mathbf{sGr} \rightleftarrows \mathbf{sComm}_k : \mathbb{R}G$ is an adjunction between homotopy categories.
3. $\mathbb{L}(-)_G$ commutes with homotopy colimits.

Conjecture 7.28. $\mathbb{L}(-)_G$ is a relative Quillen adjunction.

It then follows $\mathbb{L}(-)_G$ commutes with realizable colimits.

Lemma 7.29. • If

$$F : (\mathcal{C}, \mathcal{W}) \rightleftarrows (\mathcal{D}, \mathcal{W}) : G$$

is a relative adjunction, then

$$F : \mathcal{C}[\mathcal{W}^{-1}] \rightleftarrows \mathcal{D}[\mathcal{W}^{-1}] : G$$

is a classical adjunction.

- For any small category I , if

$$F : (\mathcal{C}, \mathcal{W}) \rightleftarrows (\mathcal{D}, \mathcal{W}) : G$$

is a relative adjunction, then

$$F^I : (\mathcal{C}^I, \mathcal{W}^I) \rightleftarrows (\mathcal{D}^I, \mathcal{W}^I) : G^I$$

is a relative adjunction.

Definition 7.30 (Rodügues-Gonzalez). **Realizable homotopy colimit** is a relative colimit.

Given $I \in \mathbf{Cat}$, and a relative category $(\mathcal{C}, \mathcal{W})$, consider the constant diagram functor

$$c_I : \mathcal{C} \longrightarrow \mathcal{C}^I$$

a realizable hocolimit is a relative adjunction

Model approximation Chacholski-Schrer (Dwyer) Homotopy theory of diagrams:

If $(\mathcal{C}, \mathcal{W})$ is a relative category, we define (left) model approximation of $(\mathcal{C}, \mathcal{W})$ to be an adjunction

$$l : \mathcal{M} \rightleftarrows \mathcal{C} : r$$

where \mathcal{M} is a model category.

1. r is a relative functor $r(\mathcal{W}_{\mathcal{C}}) \subseteq \mathcal{W}_{\mathcal{M}}$.
2. l is a relative functor on $\mathcal{W}_{\mathcal{M}}|_{\mathcal{M}^{\text{cof}}}$.
3. for any cofibrant $X \in \text{Ob}(\mathcal{M}^{\text{cof}})$ and any $A \in \text{Ob}(\mathcal{C})$, $f : X \rightarrow G(A)$ is a weak equivalence if and only if $f^{\#} : F(X) \rightarrow A$ is a weak equivalence.

Lemma 7.31. *There is a derived adjunction*

$$\mathbb{L}l : \text{Ho}(\mathcal{M}) \rightleftarrows \text{Ho}(\mathcal{C}) : \bar{r}$$

and $\bar{r} : \text{Ho}(\mathcal{C}) \hookrightarrow \text{Ho}(\mathcal{M})$ is fully faithful.

Example 7.32 (CS). Let \mathcal{M} be a(ny) model category and I a small category. \mathcal{M}^I is not a model category but a relative category with pointwise weak equivalences.

Theorem 7.33. *There is a natural left model approximation*

$$l : \mathcal{M}_b^{\Delta I} \rightleftarrows \mathcal{M}^I : r$$

where $\Delta I \downarrow \mathcal{N}_* I$ and $\mathcal{M}_b^{\Delta I} \subset \mathcal{M}^{\Delta I}$ subcategory of bounded diagrams (i.e. $F : \Delta I \rightarrow \mathcal{M}$ maps degeneracy maps s^j to weak equivalences in \mathcal{M}).

Question 7.34. *What left model approximation of an adjoint pair of functors*

$$F : \mathcal{C} \rightleftarrows \mathcal{D} : G$$

where \mathcal{C}, \mathcal{D} are relative functor but functors are not.

Definition 7.35 (B-Ramadoss). A **left model approximation** of (F, G) is a Quillen pair $\hat{F} : \mathcal{M} \rightleftarrows \mathcal{N} : \hat{G}$ such that

$$\begin{array}{ccc} \mathcal{M} & \xrightleftharpoons[r]{l} & \mathcal{C} \\ \hat{G} \uparrow \downarrow \hat{F} & & G \uparrow \downarrow F \\ \mathcal{N} & \xrightleftharpoons[q]{p} & \mathcal{D} \end{array}$$

1. $l : \mathcal{M} \rightleftarrows \mathcal{C} : r$ is a left model approximation.
2. $p : \mathcal{N} \rightleftarrows \mathcal{D} : q$ is a left model approximation.
3. $p \circ \hat{F} \dashrightarrow F \circ l$ is a relative isomorphism of functors such that “left-looking” arrows are natural weak equivalences on \mathcal{M}^{cof} , i.e. $\text{Im}(RG \circ \bar{q}) \subseteq \text{Im}(\bar{r})$.

Theorem 7.36. If $F : \mathcal{C} \rightleftarrows \mathcal{D} : G$ is an adjoint pair between two relative categories that admits a left model approximation $\widehat{F} : \mathcal{M} \rightleftarrows \mathcal{N} : \widehat{G}$ in the above sense. Then both F and G have total left (right) derived functors and they are adjoint

$$\mathbb{L}F : \mathrm{Ho}(\mathcal{C}) \rightleftarrows \mathrm{Ho}(\mathcal{D}) : \mathbb{R}G$$

given by

$$\begin{aligned} \mathbb{L}F &\cong \mathbb{L}p \circ \mathbb{L}\widehat{F} \circ \bar{r} \\ \mathbb{R}G &\cong \mathbb{L}l \circ \mathbb{R}\widehat{G} \circ \bar{q} \end{aligned}$$

Corollary 7.37. $\mathbb{L}F$ commutes with (small) colimits.

Question 7.38. In this setting, (F, G) defines a relative adjunction.

Example 7.39. Special case: $(-)_G : \mathbf{sGr} \rightleftarrows \mathbf{sComm} : G$.

8 Stable Homotopy Theory

8.1 Motivation Results

S. Schwede “Spectra in model categories”

Theorem 8.1. “Stabilization = Abelianization” Theorem (over \mathbb{Q})

Theorem 8.2. $\Sigma_+^\infty \cong \mathbb{L}(-)_{\mathrm{ab}}$ derived cotangent complex.

Let \mathcal{C} be a proper (simplicial) model category, for instance $\mathcal{C} = \mathbf{sComm}_k$. Define \mathcal{C}/B to be the augmented category over B with

- objects are $B \xleftarrow{s} X \xrightarrow{r} B$.
- morphisms are $f : X \rightarrow Y$ such that the following diagram commutes.

$$\begin{array}{ccccc} B & \xleftarrow{s} & X & \xrightarrow{r} & B \\ \parallel & & \downarrow f & & \parallel \\ B & \xleftarrow{s'} & Y & \xrightarrow{r'} & B \end{array}$$

Lemma 8.3. \mathcal{C}/B inherits naturally a model structure from \mathcal{C} .

Note \mathcal{C}/B is pointed with $\emptyset = * = \{B \xrightarrow{\mathrm{Id}} B \xrightarrow{\mathrm{Id}} B\}$. So $(\mathcal{C}/B)^\infty$ is the category of spectra over $\mathcal{C}/B (\simeq \mathbf{Top}_*)$.

There is a Quillen pair

$$\Sigma^\infty : \mathcal{C}/B \rightarrow (\mathcal{C}/b)^\infty : \Omega^\infty$$

$$X_0 \longleftarrow \{X_n, \sigma_n\} \tag{8.1} \quad \{\mathbf{eq:stab}\}$$

$$X \longmapsto \Sigma^\infty X = \{X, \mathrm{Id}\}$$

Factorization Category Fix $f : A \rightarrow B$ in \mathcal{C} , define the category $A \backslash \mathcal{C} / B$ with

- objects are $(A \xrightarrow{\alpha} P \xrightarrow{\beta} B)$ such that $f = \beta \circ \alpha$.

Note when $f = \text{Id}_B$, this is \mathcal{C} / B .

Lemma 8.4 (Schwede). $(A \backslash \mathcal{C} / B)_f$ is a simplicial category.

There is a natural pair of adjoint functors

$$\begin{aligned} f^* = (-)_+ : (A \backslash \mathcal{C} / B)_f &\longrightarrow \mathcal{C} / B : f_* \\ (A \xrightarrow{sf} X \xrightarrow{r} B) &\longleftarrow (B \xrightarrow{s} X \xrightarrow{r} B) \\ ((A \xrightarrow{\alpha} X \xrightarrow{\beta} B) &\longmapsto (B \hookrightarrow B \cup_A P \xrightarrow{(\text{Id}, \beta)} B) \end{aligned} \quad (8.2) \quad \{\text{eq:f}\}$$

Combining (8.1) and (8.2) define the Quillen pair

$$\Sigma_+^\infty : (A \backslash \mathcal{C} / B)_f \xrightleftharpoons[f_*]{(-)_+} (\mathcal{C} / B) \xrightleftharpoons[\Omega^\infty]{\Sigma^\infty} (\mathcal{C} / B)^\infty \quad (8.3) \quad \{\text{eq:stab-adj}\}$$

and we get an adjunction between homotopy categories

$$\mathbb{L}\Sigma_+^\infty : \text{Ho}((A \backslash \mathcal{C} / B)_f) \xrightleftharpoons{\quad} \text{Ho}((\mathcal{C} / B)^\infty) : \mathbb{R}f_{*,0}.$$

Definition 8.5. The suspension spectrum of unreduced suspension of A over B is given by

$$\Sigma_+^\infty(A \xrightarrow{f} B) := \mathbb{L}\Sigma_+^\infty(A \xrightarrow{f} B \xrightarrow{\text{Id}} B).$$

Suppose $\mathcal{C} = \mathbf{sComm}_k$ where $k \supseteq \mathbb{Q}$.

Theorem 8.6. For any $B \in \text{Ob}(\mathbf{sComm}_k)$ fixed, the composition

$$B - \mathbf{sMod}_k \xrightleftharpoons[\Omega^\infty]{\Sigma^\infty} (B - \mathbf{sMod}_k)^\infty \xrightleftharpoons[\mathbb{R}i^\infty]{\text{Sym}_B^\infty} \mathbf{sComm}_k / B)^\infty$$

is a Quillen equivalence.

Hence

$$\text{Ho}(B - \mathbf{sMod}_k) \xrightleftharpoons[\mathbb{R}\Omega^\infty]{\mathbb{L}\Sigma^\infty} \text{Ho}((B - \mathbf{sMod}_k)^\infty) \xrightleftharpoons[\mathbb{R}i^\infty]{\mathbb{L}\text{Sym}_B^\infty} \text{Ho}(\mathbf{sComm}_k / B)^\infty$$

are equivalences of homotopy categories.

Theorem 8.7. Given a map $f : A \rightarrow B$ in \mathbf{sComm}_k , the following diagram commutes.

$$\begin{array}{ccccc}
 (A \backslash \mathbf{sComm}_k / B)_f & \xrightarrow{(-)_+} & \mathbf{sComm}_k / B & \xrightarrow{\Sigma^\infty} & (\mathbf{sComm}_k / B)^\infty \\
 & \searrow \text{ab}_{A/B} & \downarrow \text{ab}_B = (B \backslash (-) / B)_{\text{ab}} & & \downarrow \text{ab}_B^\infty \\
 & & B - \mathbf{sMod}_k & \xrightarrow{\Sigma^\infty} & (B - \mathbf{sMod}_k)^\infty
 \end{array}$$

The Quillen derived cotangent functor

$$\mathbb{L}ab_{A/B} : \text{Ho}(A \backslash \mathbf{sComm}_k / B) \longrightarrow \text{Ho}(B - \mathbf{sMod}_k)$$

is isomorphic to $\mathbb{L}\Sigma_+^\infty(A \xrightarrow{f} B)$.

Recall Quillen abelianization (or Quillen homology, 1970)

$$\text{ab} : \mathcal{C} \rightleftarrows \mathcal{C}_{\text{ab}} : i$$

where \mathcal{C}_{ab} is the category of abelian group objects in a model category \mathcal{C} . Then $\mathbb{L}(-)_{\text{ab}} : \text{Ho}(\mathcal{C}) \rightarrow \text{Ho}(\mathcal{C}_{\text{ab}})$ gives Quillen homology.

Example 8.8. Consider

$$\Omega(-/B) : \mathbf{Alg}_k \downarrow B \rightleftarrows (\mathbf{Alg}_k \downarrow B)_{\text{ab}} \cong B - \mathbf{Bimod} : B \ltimes (-)$$

$$B \ltimes M \longleftarrow \longrightarrow M$$

$$(A \xrightarrow{f} B) \longmapsto \Omega_{\text{Kahl}}^1(A) \otimes_A B$$

then $\mathbb{L}\Omega(-/B)(A \xrightarrow{f} B) = \Omega_{\text{Kahl}}^1(P) \otimes_P B$ where P is a cofibrant replacement of A .

Question 8.9. What happens to $A \neq S$ Theorem if $k \not\supseteq \mathbb{Q}$.

S. Schwede Stable homotopy of Algebraic theories.
Feng, Galatius, Venkatesh

8.2 Γ -Spaces and Homotopy-theoretic generalization of Quillen homology

Γ -Spaces (G. Segal) are simplest models for connective spectra.
Survey [DGM].

Definition 8.10. Let Γ^{op} be the skeleton of the category of finite pointed sets.

- objects are $n_+ = \{0, 1, \dots, n\}_{n \geq 0}$ with basepoint 0.
- morphisms are based morphisms $f : n_+ \rightarrow m_+$ such that $f(0) = 0$.

A Γ -space is a pointed functor $X : \Gamma^{\text{op}} \rightarrow \mathbf{sSet}_*$ such that $X(0_+) = \Delta[0]_*$.

The idea is to replace simplicial abelian groups

Fix an algebraic theory \mathcal{T} and let $\mathcal{A}_{\mathcal{T}}$ be the category of \mathcal{T} -algebras. $\mathbf{s}\mathcal{A}_{\mathcal{T}}$ is the category of (strict) simplicial \mathcal{T} -algebras/

Quillen's abelianization (restricted to $\mathcal{A}_{\mathcal{T}}$) is

$$(\mathcal{A}_{\mathcal{T}})_{\text{ab}} \cong \text{Mod}(\mathcal{T}^{\text{ab}})$$

Theorem 8.11 (Schwede). *For any algebraic theory, there exists a Gamma-ring \mathcal{T}^s such that $\mathcal{A}_{\mathcal{T}}^{\infty} = \text{Sp}_{\mathcal{T}} \simeq \text{Mod}(\mathcal{T}^s)$.*

Γ -spaces as (simple) model for spectra

1. Quillen

2. THH

algebraic theories

(Schwede's) stabilization theorem v.s. (Quillen) abelianization (linearization) -theorem

The category $A \backslash \mathbf{sComm}_k / A$ of augmented algebras $\{A \rightarrow B \rightarrow A\}$ has a model structure with initial and terminal object $A \rightarrow A \rightarrow A$, thus a stable model category, and by Quillen's abelianization get a functor $A \backslash \mathbf{sComm}_k / A \rightarrow \mathbf{sMod}_k - A$.

$$\Sigma^{\infty} : A \backslash \mathbf{sComm}_k / A \hookrightarrow \mathbf{Sp}_0(\mathbf{sComm}_k).$$

References

- [B] Bernard Badzioch, *Algebraic theories in homotopy theory*, *Annals of Mathematics* **155** (2002), 895–913.
- [BK] C Barwick and DM Kan, *Relative categories: Another model for the homotopy theory of homotopy theories*, *arxiv.org math*, AT, November (2010).
- [DGM] Bjørn Ian Dundas, Thomas G Goodwillie, and Randy McCarthy, *The local structure of algebraic k -theory*, Vol. 18, Springer Science & Business Media, 2012.
- [DK] Tobias Dyckerhoff and Mikhail Kapranov, *Crossed simplicial groups and structured surfaces*, *Stacks and categories in geometry, topology, and algebra* **643** (2015), 37–110.
- [FH] Emmanuel D Farjoun and Kathryn Hess, *Normal and conormal maps in homotopy theory*, *Homology, Homotopy and Applications* **14** (2012), no. 1, 79–112.
- [G] Beatriz Rodriguez Gonzalez, *Realizable homotopy colimits*, arXiv preprint arXiv:1104.0646 (2011).
- [H] Philip S Hirschhorn, *Model categories and their localizations*, American Mathematical Soc., 2003.
- [J] John F Jardine, *Categorical homotopy theory*, *Homology, Homotopy and Applications* **8** (2006), no. 1, 71–144.
- [M] J Peter May, *Simplicial objects in algebraic topology*, Vol. 11, University of Chicago Press, 1992.
- [P] Matan Prezma, *Homotopy normal maps*, *Algebraic & Geometric Topology* **12** (2012), no. 2, 1211–1238.