


PHP 2550: Worksheet 4

Due: September 27th at 11:59pm

Reading Recap

1. Recap Chapters 3-6 of the “The 9 Pitfalls of Data Science” in three points and choose your favorite example. Each point could be summarizing a key takeaway, something that surprised you, or something that you want to remember.
 - In pitfall 4, worshipping computer, the example of “Seeing the World Through Pixels” surprises me a lot. This example talks about how differently computers and human “see” an object. When a picture of school bus was shown, the DNN (deep neural network) identified it as an airplane instead of a school bus. This occurs since, unlike humans, computers would not understand the picture but analyzes the pixel patterns. Human always identify the object by understanding the context and meaning, even the picture is blocked or unclear, we rely on our experiences to identify it. I choose this example since it helped me understand something I had been confused about. When trying to log in to some websites, I have encountered tasks asking me to recognize traffic lights, buses, or taxis and they said this task can prevent computer from logging in. I used to believe this should be a easy task for “smart computers”, but after reading this example, I understand the reason for that and I realize how important and sophisticated human perception really is. In addition, I would like to remember this tip and avoid overly trust computers.
 - In pitfall 5, Torturing Data, the example of “Aspartame doesn’t cause cancer” does catch my attention. I usually drink diet coke to avoid suger intake and I was concerned about this topic previously when there are people saying that Aspartame could cause cancer. A lot of studies have shown that aspartame does not cause cancer, and this result is confirmed by both the FDA and European Food Safety Authority. However, because of fears inflamed by hoax emails, people continued to test it. Because of over-testing, there a few occasional anomal cases popped up, suggesting possible links between brain tumors and aspartame. This p-hacking case shows that over-testing or over-analyzing is a way to torturing data and this would increase the probability of finding meaningless patterns which are not the result we would like to have. This is also a tip I would like to remind myself.

- In pitall 6, fooling yourself, the example of “Wishful Thinking” attracts my attention since I also have experiences of overestimating my math test scores during high school and this scenario is common in our class. This example talks about how people can overestimate their ability that the 100 high school students predicted a score of 75 for a math test on average while their mean score is actually 60. This discrepancy occurs due to bias of optimism or an underestimation of the math test’s difficulty, leading to wrong decision making or conclusion. Our human nature can lead us to be overly optimistic which is a way of fooling ourselves, letting our subjective biases and desires influence our judgement. This is also a tip I would remind myself in the future.
2. The questions below relate to the article [Personalized Research on Diet in Ulcerative Colitis and Crohn’s Disease: A Series of N-of-1 Diet Trials](#).
- Summarize the paper in 1-2 paragraphs using your own words (a good way to help this is to not look at the paper when writing your response). Use the six questions from last week to help guide your summary
 - What do the author(s) want to know (motivation)?
 - What did they do (approach/methods)?
 - Why was it done that way (context within the field)?
 - What do the results show (figures and data tables)?
 - How did the author(s) interpret the results (interpretation/discussion)?
 - What should be done next (discussion/own reflection)?
 - Reflecting on the replication crisis and documentation discussion from class, how easy would it be to try to replicate this study? Explain your response.
 - How was the data presented in the paper? What parts of an exploratory analysis were included?
 - Take a look at the visual below that did not make into the paper and was replaced with Figure 1 in the paper. How is the data presented in each visual? Why do you think the authors went with Figure 1?



prob, less CI, dominance color better diet, extra info: split people into disease groups vs. descriptive stat / uncertainty / CI include 0

3. The following questions relate to the reading “An Introduction to Modern Missing Data Analyses”.

- Explain the three missing data mechanisms introduced Missing Completely at Random (MCAR), Missing at Random (MAR), and Missing Not at Random (MNAR) and give an example of each. Describe a setting in which it would be difficult to tell which setting is appropriate for our data.
1. MCAR: MCAR, missing completely at random, is when the missing is completely random and it does not depend on any variables in the data. This means that the probability of missing data on a variable is unrelated to any observed or unobserved data. An example for MCAR is in a heart disease study, the technician accidentally loses some lab samples of some participants. Here, the missingness is not related to any participants' health related or other observed and unobserved variables, making it completely random.
 2. MAR: MAR, missing at random, is when the missingness is related to observed variables, but not on the missing variable itself. In this situation, the missingness

can be explained by other observed variables, but not the missing variable itself. An example of MAR is, in a study of depression, male participants are less likely to complete the survey questions asking about depression severity compared to the female participants, leading to missingness. This missingness is not related to the severity of participants' severity of depression, but is related to their gender which is observed.

3. MNAR: MNAR, missing not at random, is when the probability of missing data is related to the incompleted variables itself. An example for MNAR is that people with higher incomes may be more likely to leave the income question blank due to privacy concerns in a survey. Here, the missing of income response is related to participants' income level but not related to other observed variables.

In a longitudinal depression study, people complete surveys regularly through time and some participants might drop out from the study. This missingness might have various causes which might be difficult to distinguish among these three types. For example, if the participants move to other places where the reason for the moving is not related to any variables in our studies, this missingness is random and it should be MCAR. Or, if the participants are too old to make regular appointment and complete the survey, the missing data is related to their age, but not related to their severity of depression. This case is MAR. Or, if the participants' depression severity becomes much worse and they drop out since they are not willing to share their worse conditions, this missingness is MNAR where it is related to the missing variable itself. In this case, it would be difficult to tell which setting is appropriate for our data because in reality, researchers might not have clear information about the drop out reason and drop out might occur due to mix of reasons mentioned earlier.

- In this class, we will focus on exploring missing data, thinking through what could have contributed to missing data, and introduce using multiple imputation as a possible tool to use. Write a 3-5 sentence explanation of multiple imputation and then describe settings when multiple imputation might not be appropriate to use.

Multiple imputation is a modern statistical technique to handle missing data. The basic procedure for multiple imputation includes three steps: the imputation phase, the analysis phase, and the pooling phase. In the imputation phase, we generate a specified number of datasets with different estimates of missing values. Then, we perform analyses separately using the same methods and steps (with no change due to missingness) on each dataset generated during the imputation phase, yielding multiple sets of estimates of each parameter and standard error. Last, in the pooling phase, we combine these multiple sets of results from each dataset's analysis into a single set of results.

Multiple imputation might not be appropriate to use when MNAR occurs, that is, when the missingness is related to the missing variable itself. This is because mul-

multiple imputation assumes the missingness in the data can be explained by observed data. The implementation of multiple imputation on MNAR data would lead to biases. Or, if our data has small sample size, it would be challenged to do reliable imputation using limited observations. Moreover, if the proportion of missingness is extremely high, our imputation might also be unreliable due to lack of information which would introduce uncertainty.

Data Exploration and Visualization

The data (schmid_data.csv and schmid_codebook.xlsx) in this question comes from a previous PhD qualifying exam. The data is a national data set of demographic, diagnostic, and respiratory parameters of infants with severe bronchopulmonary dysplasia (sBPD) admitted to collaborative NICUs and with known respiratory support parameters at 36 weeks postmenstrual age (PMA). This data was used to develop a regression model to predict the composite outcome of tracheostomy/death to guide the indication criteria and timing of tracheostomy placement. For our purposes, we will focus on exploratory analysis of this data and you do not need to do any modeling.

Conduct an exploratory analysis of this data (3-5 pages). To guide yourself, think of at least three questions you want to answer using your EDA. You should also look at patterns of missing data. As part of your analysis, create a summary table and include at least two visualizations. When creating your visuals, you should think about what you want the reader to learn from your visual, making your visual clear and effective, and the data-to-ink ratio (i.e. how much information your visual contains and whether the visual is more effective than putting the same information in text). Your writing accompanying your analysis can be short and informal.

codebook - structure: longitudinal (36/44) correlation; missing at over time (a lot 44 wk are missing, healthier babies are more likely to lose data at 44 wk); explain missing: due to the center clinical setting: multiple centers: clustering, difference between centers; race: data quality issue (unable to answer); missing / center / relationship outcome


Baseline characteristics (birth, demographics, medications, and clinical measurements) are listed in Table 1. The outcomes we are interested in are binary variables indicating whether the infant received tracheotomy prior to discharge or whether the infant died. According to Table 1, we observe a large proportion of missing values for certain variables, such as complete prenatal steroids, maternal chorioamnionitis, hospital discharge. For example, 42% of the tracheotomy outcome 'Trach' and 39% of the death outcome 'Death' lacked complete prenatal steroid information. This pattern suggests that the exclusion of these variables needs to be considered when conducting multiple imputations.

To further explore the missing patterns in the dataset, we created Figure 1. Missing observations are shown in gray, with more gray shaded areas indicating more missing values. We can see from the figure that missing values increase over time, with more values missing at 44

weeks than at 36 weeks. This may be due to the fact that healthier infants are more likely to be discharged from the hospital and the researchers were unable to collect information on them after discharge.

To explore the association between outcomes and the medical centers where the data were collected, we created a new categorical variable that combined information on both outcomes into four categories: had tracheotomy and died (TD), did not have tracheotomy and died (NTD), had tracheotomy without death (TND), and had no tracheotomy without death (NTND). As we can see in Figure 2, more than half of the infants did not have a tracheotomy and did not die, except in centers 1, 12, and 21. Especially in center 21, all infants had tracheotomy without death. This implies that there may be potential differences in the characteristics and settings of infants admitted to the centers.


The longitudinal structure of the data also allows us to understand the distribution of infant weight over time, and how this may affect our new outcome of ‘tracheal intubation/death’. As shown in Figure 3, the distribution of infant weights is approximately normally distributed, with the distribution becoming more spread out over time. Looking at the stacked bars in the figure, we find that the light blue bars (NTND) consistently have the largest proportion, suggesting that the majority of infants did not require tracheotomy and survived during the study period. Notably, comparing the red bars (TD) between the three time points, we found that more newborns required tacheometry at baseline compared to 36 and 44 weeks but did not survive the study period compared.




Characteristic	Tracheostomy		Death	
	No, N = 853 ¹	Yes, N = 146 ¹	No, N = 943 ¹	Yes, N = 54 ¹
Gender				
Female	348 (41%)	60 (41%)	390 (41%)	17 (31%)
Male	501 (59%)	86 (59%)	549 (58%)	37 (69%)
Missing	4 (0.5%)	0 (0%)	4 (0.4%)	0 (0%)
Ethnicity				
Hispanic or Latino	66 (7.7%)	8 (5.5%)	71 (7.5%)	3 (5.6%)
Not Hispanic or Latino	740 (87%)	128 (88%)	818 (87%)	48 (89%)
Missing	47 (5.5%)	10 (6.8%)	54 (5.7%)	3 (5.6%)
Birth Weight	814 (295)	761 (303)	811 (288)	721 (406)
Gestational Age	26 (2)	26 (2)	26 (2)	26 (2)
Birth Length	33 (4)	32 (4)	33 (4)	30 (4)
Unknown	48	30	71	7
Birth Head Circumference	23.22 (2.71)	22.99 (3.07)	23.24 (2.70)	22.34 (3.54)
Unknown	46	31	72	5
Delivery Method				
Vaginal delivery	254 (30%)	31 (21%)	274 (29%)	10 (19%)
Cesarean section	597 (70%)	114 (78%)	666 (71%)	44 (81%)
Missing	2 (0.2%)	1 (0.7%)	3 (0.3%)	0 (0%)
Prenatal Corticosteroids				
Yes	715 (84%)	123 (84%)	792 (84%)	44 (81%)
No	118 (14%)	8 (5.5%)	121 (13%)	5 (9.3%)
Missing	20 (2.3%)	15 (10%)	30 (3.2%)	5 (9.3%)
Small for gestational age				
SGA	161 (19%)	42 (29%)	177 (19%)	26 (48%)
Not SGA	681 (80%)	100 (68%)	751 (80%)	28 (52%)
Missing	11 (1.3%)	4 (2.7%)	15 (1.6%)	0 (0%)
Complete Prenatal Steroids				
Yes	527 (62%)	86 (59%)	577 (61%)	34 (63%)
No	166 (19%)	27 (18%)	184 (20%)	9 (17%)
Missing	160 (19%)	33 (23%)	182 (19%)	11 (20%)
Maternal Chorioamnionitis				
Yes	138 (16%)	22 (15%)	153 (16%)	7 (13%)
No	665 (78%)	112 (77%)	732 (78%)	43 (80%)
Missing	50 (5.9%)	12 (8.2%)	58 (6.2%)	4 (7.4%)
Hospital Discharge Gestational Age	49 (24)	80 (30)	52 (27)	59 (24)
Unknown	86	38	116	7

¹Mean (SD) for continuous; n (%) for categorical

Outcome Proportion by Center



Birth Weight Distribution by Tracheostomy/Death Outcome




Missing Data and Imputation with MICE



Load in the data called `pain` from the `HDSinRdata` package and read the documentation. The data contains information from patient-reported pain assessments at baseline and at a 3-month follow-up.

1. First, describe the patterns of missing data observed in the data set overall. How would you present this information in an exploratory analysis?

I would present the missing data pattern using heat map. Base on the data structure, I plot three separate heat maps focusing on the body region variables, baseline characteristics, and the follow-up variable, respectively.

In the heat map of the body region variables, we see that most variables have nearly complete data since there is no visible grey area on the plot, indicating minimal or no missing values. Our baseline characteristics, including both continuous and categorical, exhibits some missingness among certain columns. Categorical variables, such as `PAT_RACE` and `MEDICAID_BIN`, having small portion of missing data, while continuous variables like `BMI` and `GH_PHYSICAL_SCORE` have larger proportion of missingness. To be noticed, the follow-up variable in the third heat map exhibits a significantly large portion of missingness, with over 50% of value missing, which might be a crucial issue.





2. If we were interested in analyzing the change in pain over time, it would be important to think about the missing data due to loss to follow-up. Compare the baseline characteristics between those with and without follow-up information. Comment on your results and discuss whether you think the data is MCAR, MAR, or MNAR.

To compare the baseline characteristics between those with and without follow-up information, we create a table summary of all baseline characteristics by follow-up presence. Here, participants without follow-up had significantly higher pain intensity score (7.00 vs. 6.00 with $p < 0.01$) compared to participants with follow-up. Moreover, in several PROMIS measures, patients with and without follow-up exhibits small but statistically significant differences with smaller p-value. For those demographic variables, age, BMI, gender, and race, significant difference between groups present as well. For example, younger people or patients with lower BMI value are more likely not to have the follow-up information. Moreover, observing the MEDICAID_BIN summary, medicaid recipients are more likely to skip the follow-up.

We think this data is more likely to be MAR where the follow-up variable missingness is relates to observed baseline characteristics variables. Since many baseline characteristics variables exhibits significant difference between people with and without follow-up information as we mentioned earlier, these characteristics seem associated with the probability of missingness which fits the assumption of MAR.

Characteristic	Follow-up, N = 7,138	No Follow-up, N = 14,521	p-value
PAIN_INTENSITY_AVERAGE	6.00 (5.00, 8.00)	7.00 (5.00, 8.00)	<0.001
PROMIS_PHYSICAL_FUNCTION	36 (31, 39)	35 (30, 39)	0.045
PROMIS_PAIN_BEHAVIOR	61.6 (59.0, 63.4)	61.7 (59.5, 63.4)	<0.001
PROMIS_DEPRESSION	55 (48, 62)	55 (48, 62)	0.009
PROMIS_ANXIETY	56 (49, 63)	58 (50, 63)	<0.001
PROMIS_SLEEP_DISTURBANCE	59 (54, 65)	60 (54, 65)	<0.001
PROMIS_PAIN_INTERFERENCE	62 (52, 70)	67 (63, 72)	<0.001
GH_MENTAL_SCORE	44 (36, 51)	44 (36, 51)	0.5
GH_PHYSICAL_SCORE	35 (30, 40)	35 (30, 40)	0.028
AGE_AT_CONTACT	58 (48, 67)	57 (45, 68)	0.001
BMI	30 (25, 35)	29 (25, 34)	<0.001
CCI_TOTAL_SCORE			0.7
0	6,405 (90%)	13,119 (90%)	
1	606 (8.5%)	1,149 (7.9%)	
2	104 (1.5%)	196 (1.3%)	
3	19 (0.3%)	45 (0.3%)	
4	4 (<0.1%)	10 (<0.1%)	
5	0 (0%)	1 (<0.1%)	
PAT_SEX			<0.001

Characteristic	Follow-up, N = 7,138	No Follow-up, N = 14,521	p-value
female	4,431 (62%)	8,671 (60%)	<0.001
male	2,707 (38%)	5,849 (40%)	
PAT_RACE			
ALASKA NATIVE	1 (<0.1%)	1 (<0.1%)	
AMERICAN INDIAN	29 (0.4%)	29 (0.2%)	
BLACK	934 (13%)	2,295 (16%)	
CHINESE	4 (<0.1%)	17 (0.1%)	
DECLINED	28 (0.4%)	93 (0.6%)	
FILIPINO	3 (<0.1%)	3 (<0.1%)	
GUAM/CHAMORRO	0 (0%)	1 (<0.1%)	
HAWAIIAN	1 (<0.1%)	0 (0%)	
INDIAN (ASIAN)	13 (0.2%)	36 (0.2%)	
JAPANESE	1 (<0.1%)	8 (<0.1%)	
KOREAN	3 (<0.1%)	7 (<0.1%)	
NOT SPECIFIED	2 (<0.1%)	2 (<0.1%)	
OTHER	0 (0%)	1 (<0.1%)	
OTHER ASIAN	8 (0.1%)	39 (0.3%)	
OTHER PACIFIC ISLANDER	1 (<0.1%)	11 (<0.1%)	
VIETNAMESE	3 (<0.1%)	3 (<0.1%)	
WHITE	6,080 (86%)	11,860 (82%)	
CCI_BIN			0.2
Any comorbidity	733 (10%)	1,401 (9.6%)	<0.001
No comorbidity	6,405 (90%)	13,119 (90%)	
MEDICAID_BIN			
Missing	92 (1.3%)	209 (1.4%)	
no	5,708 (80%)	11,049 (76%)	<0.001
yes	1,338 (19%)	3,263 (22%)	

3. Now suppose we are interested in mental and physical function at baseline and how that is associated with pain intensity. We will use the `mice` package to perform multiple imputation on this data. First, drop the variable `PAIN_INTENSITY_AVERAGE.FOLLOW_UP` and the body map variables `X101:X238`.

```
pain_mod <- pain[, -c(1:75)] %>%
  select(-PAIN_INTENSITY_AVERAGE.FOLLOW_UP)

pain_mod$PAT_SEX <- as.factor(pain_mod$PAT_SEX)
pain_mod$PAT_RACE <- as.factor(pain_mod$PAT_RACE)
```

```
pain_mod$CCI_BIN <- as.factor(pain_mod$CCI_BIN)
pain_mod$MEDICAID_BIN <- as.factor(pain_mod$MEDICAID_BIN)
```

4. Use the `mice()` function to perform multiple imputation to create five imputed data sets and save the result as `pain_mice`. You should read the documentation on this function to understand how it is implementing this and what arguments you might want to set. What is the structure of the returned object?

The returned object contains the following parts:

- **data:** The original dataset with missing data.
- **imp:** A list of imputed values for variables with missingness.
- **m:** The number of imputed dataset which is 5 here.
- **where:** A matrix presents the location of missing values in our original dataset where missing values are coded as TRUE.
- **blocks:** A list of variable names. Variables in one block would be imputed together.
- **call:** The command used to run the `mice()` function, including information of data, number of iterations, random seed, and any set parameters.
- **nmis:** A numeric vector presents number of missing values for each variable.
- **method:** The imputation method used.
- **predictorMatrix:** A numeric matrix specifying which variables are used as predictors in the imputation models.
- **visitSequence:** The order variables are imputed in mice function.
- **formulas:** A list of formulas used for imputation of each variable.
- **post:** Post-processing rules applied to the imputed data.
- **blots:** Contains information of additional processing for certain variables during imputation.
- **ignore:** A vector specifying whether specific rows were ignored during imputation. False represents the observation was included in the process while TRUE represents it is ignored.
- **seed:** The random seed set in parameter.
- **iteration:** Number of iteration for the imputation.
- **lastSeedValue:** This component stores the final state of the random number generator after all imputations.

- **chainMean:** The mean of the imputed values for each variable at each iteration and for each imputation chain.
- **chainVar:** The variance of the imputed values for each variable at each iteration and for each imputation chain.
- **loggedEvents:** A data frame that records unusual events occurred during imputation, like warnings and errors.
- **version:** The version of mice package used for the imputation.
- **date:** The date when the imputation performed.

```
# pain_mice <- mice(pain_mod, maxit = 5, seed = (2550))
# saveRDS(pain_mice, file = "mice_result.rds")
pain_mice <- readRDS("mice_result.rds")

str(pain_mice)
```

List of 22

```
$ data          : 'data.frame':   21659 obs. of  16 variables:
..$ PAIN_INTENSITY_AVERAGE : num [1:21659] 7 5 4 7 8 8 7 7 5 6 ...
..$ PROMIS_PHYSICAL_FUNCTION : num [1:21659] 39.4 36.3 41.5 28.9 36.5 ...
..$ PROMIS_PAIN_BEHAVIOR : num [1:21659] 60 60.9 60.4 62.5 61.7 ...
..$ PROMIS_DEPRESSION : num [1:21659] 64.2 51.3 55.2 57.6 58.8 ...
..$ PROMIS_ANXIETY : num [1:21659] 61.5 51.2 56.5 57.7 59.5 ...
..$ PROMIS_SLEEP_DISTURB_V1_0: num [1:21659] 59.5 59.4 55.8 62 65.2 ...
..$ PROMIS_PAIN_INTERFERENCE : num [1:21659] 61.5 66.9 60.1 68.8 64.2 ...
..$ GH_MENTAL_SCORE : num [1:21659] 43.5 53.3 48.3 NA 45.8 31.3 43.5 41.1 56 41.1
..$ GH_PHYSICAL_SCORE : num [1:21659] 29.6 37.4 37.4 NA 34.9 26.7 34.9 37.4 50.8 32
..$ AGE_AT_CONTACT : num [1:21659] 78 70 75 61 63 54 32 93 51 69 ...
..$ BMI : num [1:21659] 33.7 36.3 27.2 NA 28.9 ...
..$ CCI_TOTAL_SCORE : num [1:21659] 0 0 0 0 0 0 0 0 0 0 ...
..$ PAT_SEX : Factor w/ 2 levels "female","male": 2 1 2 2 2 2 1 1 2 2 ...
..$ PAT_RACE : Factor w/ 17 levels "ALASKA NATIVE",...: 17 17 17 17 17 17 17 17 17 17
..$ CCI_BIN : Factor w/ 2 levels "Any comorbidity",...: 2 2 2 2 2 2 2 2 2 2
..$ MEDICAID_BIN : Factor w/ 2 levels "no","yes": 1 1 1 1 1 NA 2 1 1 1 ...

$ imp          :List of 16
..$ PAIN_INTENSITY_AVERAGE : 'data.frame':   1 obs. of  5 variables:
.. ..$ 1: num 4
.. ..$ 2: num 4
.. ..$ 3: num 10
.. ..$ 4: num 10
.. ..$ 5: num 6
```

```

..$ PROMIS_PHYSICAL_FUNCTION : 'data.frame':  1 obs. of  5 variables:
.. ..$ 1: num 24.7
.. ..$ 2: num 38.6
.. ..$ 3: num 39.5
.. ..$ 4: num 36.5
.. ..$ 5: num 35.6
..$ PROMIS_PAIN_BEHAVIOR      : 'data.frame': 6371 obs. of  5 variables:
.. ..$ 1: num [1:6371] 62.5 60.8 55.2 60 61 ...
.. ..$ 2: num [1:6371] 56.3 60.7 58.5 57.3 60 ...
.. ..$ 3: num [1:6371] 63.4 58.5 56.3 60.2 62.5 ...
.. ..$ 4: num [1:6371] 58.5 60.8 51.2 63.4 59 ...
.. ..$ 5: num [1:6371] 60.7 59 51.2 66 62.5 ...
..$ PROMIS_DEPRESSION        : 'data.frame':  88 obs. of  5 variables:
.. ..$ 1: num [1:88] 79.3 36.7 62.2 48.6 53.3 ...
.. ..$ 2: num [1:88] 60.5 41.2 42.9 57.1 62.2 ...
.. ..$ 3: num [1:88] 61.8 51.2 49.9 65.7 55.4 ...
.. ..$ 4: num [1:88] 49.9 50.8 59.6 57.6 57.3 ...
.. ..$ 5: num [1:88] 56.3 67.7 45.8 57.5 71.4 ...
..$ PROMIS_ANXIETY           : 'data.frame':  88 obs. of  5 variables:
.. ..$ 1: num [1:88] 70.5 36.1 65.2 56.5 53.5 ...
.. ..$ 2: num [1:88] 64.8 43 40 61.6 65.1 ...
.. ..$ 3: num [1:88] 65.6 56.5 45.7 51.4 46 ...
.. ..$ 4: num [1:88] 55.7 44.3 52.9 63.4 52.1 ...
.. ..$ 5: num [1:88] 58.5 66.9 44.3 59.5 63.4 ...
..$ PROMIS_SLEEP_DISTURB_V1_0: 'data.frame':  88 obs. of  5 variables:
.. ..$ 1: num [1:88] 57.9 54.3 62 79 55.8 ...
.. ..$ 2: num [1:88] 65.5 63 48.9 63.9 63 ...
.. ..$ 3: num [1:88] 52.7 43.2 54.3 54.3 39.5 ...
.. ..$ 4: num [1:88] 64.1 63 72.7 59.4 61.2 ...
.. ..$ 5: num [1:88] 52.7 63 59.4 54.3 63 ...
..$ PROMIS_PAIN_INTERFERENCE : 'data.frame': 152 obs. of  5 variables:
.. ..$ 1: num [1:152] 59.1 66.9 65.7 66.9 73.7 ...
.. ..$ 2: num [1:152] 61.5 74.1 66.9 66.9 73 ...
.. ..$ 3: num [1:152] 61.5 70.3 65.5 73.7 73.7 ...
.. ..$ 4: num [1:152] 61.5 68.2 66.9 64.2 69.9 ...
.. ..$ 5: num [1:152] 62.7 68.2 66.9 66.9 74.1 ...
..$ GH_MENTAL_SCORE          : 'data.frame': 2947 obs. of  5 variables:
.. ..$ 1: num [1:2947] 41.1 56 56 48.3 31.3 28.4 31.3 33.8 41.1 48.3 ...
.. ..$ 2: num [1:2947] 43.5 59 59 45.8 41.1 38.8 33.8 33.8 41.1 43.5 ...
.. ..$ 3: num [1:2947] 48.3 56 43.5 45.8 28.4 33.8 41.1 36.3 48.3 43.5 ...
.. ..$ 4: num [1:2947] 41.1 59 48.3 50.8 21.2 31.3 48.3 33.8 50.8 50.8 ...
.. ..$ 5: num [1:2947] 48.3 45.8 56 36.3 36.3 33.8 45.8 31.3 43.5 43.5 ...
..$ GH_PHYSICAL_SCORE        : 'data.frame': 2947 obs. of  5 variables:

```

```

.. ..$ 1: num [1:2947] 34.9 34.9 44.9 34.9 26.7 39.8 39.8 29.6 47.7 32.4 ...
.. ..$ 2: num [1:2947] 34.9 50.8 50.8 32.4 34.9 34.9 39.8 29.6 37.4 39.8 ...
.. ..$ 3: num [1:2947] 34.9 47.7 54.1 42.3 34.9 34.9 29.6 26.7 47.7 29.6 ...
.. ..$ 4: num [1:2947] 26.7 47.7 44.9 44.9 29.6 29.6 44.9 26.7 34.9 34.9 ...
.. ..$ 5: num [1:2947] 32.4 42.3 50.8 32.4 32.4 29.6 39.8 29.6 37.4 32.4 ...
..$ AGE_AT_CONTACT          : 'data.frame':  1 obs. of  5 variables:
.. ..$ 1: num 50
.. ..$ 2: num 44
.. ..$ 3: num 63
.. ..$ 4: num 41
.. ..$ 5: num 86
..$ BMI                    : 'data.frame':  5633 obs. of  5 variables:
.. ..$ 1: num [1:5633] 25.1 21.8 26.6 29.4 24 ...
.. ..$ 2: num [1:5633] 36.9 21.9 33.8 27.3 25.8 ...
.. ..$ 3: num [1:5633] 37.3 38.6 26.7 21.9 40.7 ...
.. ..$ 4: num [1:5633] 25.8 25.3 26.2 22.5 28.4 ...
.. ..$ 5: num [1:5633] 23.2 26.3 22.6 54.7 29.1 ...
..$ CCI_TOTAL_SCORE        : 'data.frame':  1 obs. of  5 variables:
.. ..$ 1: num 1
.. ..$ 2: num 0
.. ..$ 3: num 0
.. ..$ 4: num 0
.. ..$ 5: num 1
..$ PAT_SEX                : 'data.frame':  1 obs. of  5 variables:
.. ..$ 1: Factor w/ 2 levels "female","male": 1
.. ..$ 2: Factor w/ 2 levels "female","male": 2
.. ..$ 3: Factor w/ 2 levels "female","male": 1
.. ..$ 4: Factor w/ 2 levels "female","male": 1
.. ..$ 5: Factor w/ 2 levels "female","male": 2
..$ PAT_RACE               : 'data.frame':  142 obs. of  5 variables:
.. ..$ 1: Factor w/ 17 levels "ALASKA NATIVE",...: 17 17 3 17 17 17 3 3 17 17 ...
.. ..$ 2: Factor w/ 17 levels "ALASKA NATIVE",...: 17 17 17 17 3 17 17 17 3 17 ...
.. ..$ 3: Factor w/ 17 levels "ALASKA NATIVE",...: 17 17 17 17 17 13 16 17 17 17 ...
.. ..$ 4: Factor w/ 17 levels "ALASKA NATIVE",...: 3 17 17 17 12 17 17 17 17 17 ...
.. ..$ 5: Factor w/ 17 levels "ALASKA NATIVE",...: 17 17 17 3 1 17 8 17 17 17 ...
..$ CCI_BIN               : 'data.frame':  1 obs. of  5 variables:
.. ..$ 1: Factor w/ 2 levels "Any comorbidity",...: 2
.. ..$ 2: Factor w/ 2 levels "Any comorbidity",...: 2
.. ..$ 3: Factor w/ 2 levels "Any comorbidity",...: 2
.. ..$ 4: Factor w/ 2 levels "Any comorbidity",...: 1
.. ..$ 5: Factor w/ 2 levels "Any comorbidity",...: 1
..$ MEDICAID_BIN          : 'data.frame':  301 obs. of  5 variables:
.. ..$ 1: Factor w/ 2 levels "no","yes": 1 1 1 1 1 1 1 1 2 1 ...

```



```

.. ..$ 2: Factor w/ 2 levels "no","yes": 2 1 1 2 2 1 1 1 1 2 ...
.. ..$ 3: Factor w/ 2 levels "no","yes": 2 1 1 2 1 1 1 1 1 1 ...
.. ..$ 4: Factor w/ 2 levels "no","yes": 1 1 1 2 1 1 1 1 1 2 ...
.. ..$ 5: Factor w/ 2 levels "no","yes": 1 1 1 2 1 2 1 1 1 1 ...
$ m                : num 5
$ where            : logi [1:21659, 1:16] FALSE FALSE FALSE FALSE FALSE FALSE ...
..- attr(*, "dimnames")=List of 2
.. ..$ : chr [1:21659] "1" "2" "3" "4" ...
.. ..$ : chr [1:16] "PAIN_INTENSITY_AVERAGE" "PROMIS_PHYSICAL_FUNCTION" "PROMIS_PAIN_BEHAVI
$ blocks          :List of 16
..$ PAIN_INTENSITY_AVERAGE : chr "PAIN_INTENSITY_AVERAGE"
..$ PROMIS_PHYSICAL_FUNCTION : chr "PROMIS_PHYSICAL_FUNCTION"
..$ PROMIS_PAIN_BEHAVIOR    : chr "PROMIS_PAIN_BEHAVIOR"
..$ PROMIS_DEPRESSION      : chr "PROMIS_DEPRESSION"
..$ PROMIS_ANXIETY         : chr "PROMIS_ANXIETY"
..$ PROMIS_SLEEP_DISTURB_V1_0: chr "PROMIS_SLEEP_DISTURB_V1_0"
..$ PROMIS_PAIN_INTERFERENCE : chr "PROMIS_PAIN_INTERFERENCE"
..$ GH_MENTAL_SCORE        : chr "GH_MENTAL_SCORE"
..$ GH_PHYSICAL_SCORE      : chr "GH_PHYSICAL_SCORE"
..$ AGE_AT_CONTACT        : chr "AGE_AT_CONTACT"
..$ BMI                    : chr "BMI"
..$ CCI_TOTAL_SCORE       : chr "CCI_TOTAL_SCORE"
..$ PAT_SEX               : chr "PAT_SEX"
..$ PAT_RACE              : chr "PAT_RACE"
..$ CCI_BIN               : chr "CCI_BIN"
..$ MEDICAID_BIN          : chr "MEDICAID_BIN"
..- attr(*, "calltype")= Named chr [1:16] "type" "type" "type" "type" ...
.. ..- attr(*, "names")= chr [1:16] "PAIN_INTENSITY_AVERAGE" "PROMIS_PHYSICAL_FUNCTION" "PROMIS_PAIN_BEHAVI
$ call            : language mice(data = pain_mod, maxit = 5, seed = (2550))
$ nmis            : Named int [1:16] 1 1 6371 88 88 88 152 2947 2947 1 ...
..- attr(*, "names")= chr [1:16] "PAIN_INTENSITY_AVERAGE" "PROMIS_PHYSICAL_FUNCTION" "PROMIS_PAIN_BEHAVI
$ method          : Named chr [1:16] "pmm" "pmm" "pmm" "pmm" ...
..- attr(*, "names")= chr [1:16] "PAIN_INTENSITY_AVERAGE" "PROMIS_PHYSICAL_FUNCTION" "PROMIS_PAIN_BEHAVI
$ predictorMatrix: num [1:16, 1:16] 0 1 1 1 1 1 1 1 1 1 ...
..- attr(*, "dimnames")=List of 2
.. ..$ : chr [1:16] "PAIN_INTENSITY_AVERAGE" "PROMIS_PHYSICAL_FUNCTION" "PROMIS_PAIN_BEHAVI
.. ..$ : chr [1:16] "PAIN_INTENSITY_AVERAGE" "PROMIS_PHYSICAL_FUNCTION" "PROMIS_PAIN_BEHAVI
$ visitSequence  : chr [1:16] "PAIN_INTENSITY_AVERAGE" "PROMIS_PHYSICAL_FUNCTION" "PROMIS_PAIN_BEHAVI
$ formulas       :List of 16
..$ PAIN_INTENSITY_AVERAGE :Class 'formula' language PAIN_INTENSITY_AVERAGE ~ PROMIS_PHYSICAL_FUNCTION
.. ..- attr(*, ".Environment")=<environment: 0x000001a06ddd9998>
..$ PROMIS_PHYSICAL_FUNCTION :Class 'formula' language PROMIS_PHYSICAL_FUNCTION ~ PAIN_INTENSITY_AVERAGE
.. ..- attr(*, ".Environment")=<environment: 0x000001a06ddd9998>

```

```

..$ PROMIS_PAIN_BEHAVIOR      :Class 'formula' language PROMIS_PAIN_BEHAVIOR ~ PAIN_INTENS
.. .. ..- attr(*, ".Environment")=<environment: 0x000001a06ddd9998>
..$ PROMIS_DEPRESSION        :Class 'formula' language PROMIS_DEPRESSION ~ PAIN_INTENSITY
.. .. ..- attr(*, ".Environment")=<environment: 0x000001a06ddd9998>
..$ PROMIS_ANXIETY           :Class 'formula' language PROMIS_ANXIETY ~ PAIN_INTENSITY_AVI
.. .. ..- attr(*, ".Environment")=<environment: 0x000001a06ddd9998>
..$ PROMIS_SLEEP_DISTURB_V1_0:Class 'formula' language PROMIS_SLEEP_DISTURB_V1_0 ~ PAIN_I
.. .. ..- attr(*, ".Environment")=<environment: 0x000001a06ddd9998>
..$ PROMIS_PAIN_INTERFERENCE :Class 'formula' language PROMIS_PAIN_INTERFERENCE ~ PAIN_IN
.. .. ..- attr(*, ".Environment")=<environment: 0x000001a06ddd9998>
..$ GH_MENTAL_SCORE          :Class 'formula' language GH_MENTAL_SCORE ~ PAIN_INTENSITY_A
.. .. ..- attr(*, ".Environment")=<environment: 0x000001a06ddd9998>
..$ GH_PHYSICAL_SCORE        :Class 'formula' language GH_PHYSICAL_SCORE ~ PAIN_INTENSITY
.. .. ..- attr(*, ".Environment")=<environment: 0x000001a06ddd9998>
..$ AGE_AT_CONTACT           :Class 'formula' language AGE_AT_CONTACT ~ PAIN_INTENSITY_AVI
.. .. ..- attr(*, ".Environment")=<environment: 0x000001a06ddd9998>
..$ BMI                      :Class 'formula' language BMI ~ PAIN_INTENSITY_AVERAGE + PRO
.. .. ..- attr(*, ".Environment")=<environment: 0x000001a06ddd9998>
..$ CCI_TOTAL_SCORE          :Class 'formula' language CCI_TOTAL_SCORE ~ PAIN_INTENSITY_A
.. .. ..- attr(*, ".Environment")=<environment: 0x000001a06ddd9998>
..$ PAT_SEX                  :Class 'formula' language PAT_SEX ~ PAIN_INTENSITY_AVERAGE +
.. .. ..- attr(*, ".Environment")=<environment: 0x000001a06ddd9998>
..$ PAT_RACE                  :Class 'formula' language PAT_RACE ~ PAIN_INTENSITY_AVERAGE
.. .. ..- attr(*, ".Environment")=<environment: 0x000001a06ddd9998>
..$ CCI_BIN                   :Class 'formula' language CCI_BIN ~ PAIN_INTENSITY_AVERAGE +
.. .. ..- attr(*, ".Environment")=<environment: 0x000001a06ddd9998>
..$ MEDICAID_BIN             :Class 'formula' language MEDICAID_BIN ~ PAIN_INTENSITY_AVER
.. .. ..- attr(*, ".Environment")=<environment: 0x000001a06ddd9998>
$ post                        : Named chr [1:16] "" "" "" "" "" ...
..- attr(*, "names")= chr [1:16] "PAIN_INTENSITY_AVERAGE" "PROMIS_PHYSICAL_FUNCTION" "PROM
$ blots                       :List of 16
..$ PAIN_INTENSITY_AVERAGE   : list()
..$ PROMIS_PHYSICAL_FUNCTION : list()
..$ PROMIS_PAIN_BEHAVIOR      : list()
..$ PROMIS_DEPRESSION         : list()
..$ PROMIS_ANXIETY           : list()
..$ PROMIS_SLEEP_DISTURB_V1_0: list()
..$ PROMIS_PAIN_INTERFERENCE : list()
..$ GH_MENTAL_SCORE          : list()
..$ GH_PHYSICAL_SCORE        : list()
..$ AGE_AT_CONTACT           : list()
..$ BMI                      : list()
..$ CCI_TOTAL_SCORE          : list()

```

```

..$ PAT_SEX                : list()
..$ PAT_RACE                : list()
..$ CCI_BIN                : list()
..$ MEDICAID_BIN           : list()
$ ignore                   : logi [1:21659] FALSE FALSE FALSE FALSE FALSE FALSE ...
$ seed                     : num 2550
$ iteration                : num 5
$ lastSeedValue            : int [1:626] 10403 485 -442884590 937493873 1097172893 669123325 11446045
$ chainMean                : num [1:16, 1:5, 1:5] 8 40.7 60.6 56 56.4 ...
..- attr(*, "dimnames")=List of 3
.. ..$ : chr [1:16] "PAIN_INTENSITY_AVERAGE" "PROMIS_PHYSICAL_FUNCTION" "PROMIS_PAIN_BEHAV
.. ..$ : chr [1:5] "1" "2" "3" "4" ...
.. ..$ : chr [1:5] "Chain 1" "Chain 2" "Chain 3" "Chain 4" ...
$ chainVar                 : num [1:16, 1:5, 1:5] NA NA 11.8 74.4 88.7 ...
..- attr(*, "dimnames")=List of 3
.. ..$ : chr [1:16] "PAIN_INTENSITY_AVERAGE" "PROMIS_PHYSICAL_FUNCTION" "PROMIS_PAIN_BEHAV
.. ..$ : chr [1:5] "1" "2" "3" "4" ...
.. ..$ : chr [1:5] "Chain 1" "Chain 2" "Chain 3" "Chain 4" ...
$ loggedEvents             : 'data.frame': 375 obs. of 5 variables:
..$ it : int [1:375] 1 1 1 1 1 1 1 1 1 1 ...
..$ im : int [1:375] 1 1 1 1 1 1 1 1 1 1 ...
..$ dep : chr [1:375] "PAIN_INTENSITY_AVERAGE" "PROMIS_PHYSICAL_FUNCTION" "PROMIS_PAIN_BEH
..$ meth: chr [1:375] "pmm" "pmm" "pmm" "pmm" ...
..$ out : chr [1:375] "PAT_RACEGUAM/CHAMORRO, PAT_RACEHAWAIIAN, PAT_RACEOTHER" "PAT_RACEGU
$ version                  :Classes 'package_version', 'numeric_version' hidden list of 1
..$ : int [1:3] 3 16 0
$ date                    : Date[1:1], format: "2024-09-26"
- attr(*, "class")= chr "mids"

```

5. For one imputed data set, find the average mental and physical health score for each possible pain intensity level (0-10). To access the first imputed data set you can use the code below.


```
mice::complete(pain_mice,1)
```

Table 3: Average Summary of Pain Intensity Score

PAIN_INTENSITY_AVERAGE	avg_mental_health	avg_physical_health
0	49.12553	49.43191
1	47.97703	45.07432
2	48.68802	44.81221
3	46.97733	43.19800
4	45.47257	39.30918

5	44.58407	38.27742
6	43.67522	37.04332
7	42.71223	33.86519
8	41.94154	32.67854
9	42.12265	32.38978
10	41.34510	28.63168

6. Repeat this for the other four data sets and then use Rubin's rules to plot the results.
<https://bookdown.org/mwheymans/bookmi/rubins-rules.html>



Appendix

```
library(tidyverse)
library(ggplot2)
library(visdat)
library(gt)
schmid_data <- read.csv("schmid_data.csv")
schmid_data_m <- schmid_data
schmid_data_m$center <- as.factor(schmid_data_m$center)
schmid_data_m$mat_race <- as.factor(schmid_data_m$mat_race)
schmid_data_m$mat_ethn <- as.factor(schmid_data_m$mat_ethn)
```

```

schmid_data_m$del_method <- as.factor(schmid_data_m$del_method)
schmid_data_m$prenat_ster <- as.factor(schmid_data_m$prenat_ster)
schmid_data_m$com_prenat_ster <- as.factor(schmid_data_m$com_prenat_ster)
schmid_data_m$mat_chorio <- as.factor(schmid_data_m$mat_chorio)
schmid_data_m$gender <- as.factor(schmid_data_m$gender)
schmid_data_m$any_surf <- as.factor(schmid_data_m$any_surf)
schmid_data_m$ventilation_support_level.36 <- as.factor(schmid_data_m$ventilation_support_level.36)
schmid_data_m$ventilation_support_level_modified.44 <- as.factor(schmid_data_m$ventilation_support_level_modified.44)
schmid_data_m$Trach <- as.factor(schmid_data_m$Trach)
schmid_data_m$Death <- as.factor(schmid_data_m$Death)
schmid_data_m$sga <- as.factor(schmid_data_m$sga)

vis_dat(schmid_data_m[, -1]) +
  theme(axis.text.x = element_text(angle = 90, size = 5))
library(tidyr)
library(gtsummary)
library(ggpubr)

schmid_trach <- schmid_data %>%
  select(gender, mat_ethn, bw, ga, blength, birth_hc, del_method,
         prenat_ster, sga, com_prenat_ster, mat_chorio, hosp_dc_ga, Trach) %>%
  mutate(Trach = recode(Trach, `0` = "No", `1` = "Yes")) %>%
  mutate(
    # mat_race = case_when(mat_race == 1 ~ "American Indian or Alaskan Native",
    #                       mat_race == 2 ~ "Asian",
    #                       mat_race == 3 ~ "Black or African American",
    #                       mat_race == 4 ~ "Native Hawaiian or Other Pacific Islander",
    #                       mat_race == 5 ~ "White",
    #                       mat_race == 6 ~ "Other",
    #                       TRUE ~ "Missing")
    gender = case_when(gender == "Female" ~ "Female",
                      gender == "Male" ~ "Male",
                      TRUE ~ "Missing"),
    mat_ethn = case_when(mat_ethn == 1 ~ "Hispanic or Latino",
                        mat_ethn == 2 ~ "Not Hispanic or Latino",
                        TRUE ~ "Missing"),
    del_method = case_when(del_method == 1 ~ "Vaginal delivery",
                          del_method == 2 ~ "Cesarean section",
                          TRUE ~ "Missing"),
    sga = case_when(sga == "SGA" ~ "SGA",
                   sga == "Not SGA" ~ "Not SGA",

```

```

      TRUE ~ "Missing"),
prenat_ster = case_when(prenat_ster == "Yes" ~ "Yes",
                        prenat_ster == "No" ~ "No",
                        TRUE ~ "Missing"),
com_prenat_ster = case_when(com_prenat_ster == "Yes" ~ "Yes",
                            com_prenat_ster == "No" ~ "No",
                            TRUE ~ "Missing"),
mat_chorio = case_when(mat_chorio == "Yes" ~ "Yes",
                       mat_chorio == "No" ~ "No",
                       TRUE ~ "Missing")) %>%
mutate(mat_ethn = factor(mat_ethn, levels = c('Hispanic or Latino',
                                              'Not Hispanic or Latino', 'Missing')),

       del_method = factor(del_method,
                           levels = c('Vaginal delivery',
                                       'Cesarean section', 'Missing')),
       sga = factor(sga, levels = c('SGA', 'Not SGA', 'Missing')),
       prenat_ster = factor(prenat_ster, levels = c('Yes', 'No', 'Missing')),
       com_prenat_ster = factor(com_prenat_ster, levels = c('Yes', 'No', 'Missing')),
       mat_chorio = factor(mat_chorio, levels = c('Yes', 'No', 'Missing'))) %>%
tbl_summary(by=Trach,
            label = list(gender ~ "Gender",
                         #mat_race ~ "Race",
                         mat_ethn ~ "Ethnicity",
                         bw ~ "Birth Weight",
                         ga ~ " Gestational Age",
                         blength ~ "Birth Length",
                         birth_hc ~ "Birth Head Circumference",
                         del_method ~ "Delivery Method",
                         prenat_ster ~ "Prenatal Corticosteroids",
                         sga ~ "Small for gestational age",
                         com_prenat_ster ~ "Complete Prenatal Steroids",
                         mat_chorio ~ "Maternal Chorioamnionitis",
                         hosp_dc_ga ~ "Hospital Discharge Gestational Age"
                        ),
            statistic = all_continuous() ~ "{mean} ({sd})" %>%
modify_spanning_header(update = all_stat_cols() ~ "**Trachostomy**") %>%
modify_footnote(update = all_stat_cols() ~ "Mean (SD) for continuous; n (%) for categori
bold_labels()

schmid_death <- schmid_data %>%

```

```

select(gender, mat_ethn, bw, ga, blength, birth_hc, del_method,
       prenat_ster, sga, com_prenat_ster, mat_chorio, hosp_dc_ga, Death) %>%
mutate(Death = recode(Death, `0` = "No", `1` = "Yes")) %>%
mutate(
  # mat_race = case_when(mat_race == 1 ~ "American Indian or Alaskan Native",
  #                       mat_race == 2 ~ "Asian",
  #                       mat_race == 3 ~ "Black or African American",
  #                       mat_race == 4 ~ "Native Hawaiian or Other Pacific Islande",
  #                       mat_race == 5 ~ "White",
  #                       mat_race == 6 ~ "Other",
  #                       TRUE ~ "Missing")
  gender = case_when(gender == "Female" ~ "Female",
                     gender == "Male" ~ "Male",
                     TRUE ~ "Missing"),
  mat_ethn = case_when(mat_ethn == 1 ~ "Hispanic or Latino",
                       mat_ethn == 2 ~ "Not Hispanic or Latino",
                       TRUE ~ "Missing"),
  del_method = case_when(del_method==1 ~ 'Vaginal delivery',
                         del_method==2 ~ 'Cesarean section',
                         TRUE ~ "Missing"),
  sga = case_when(sga=="SGA" ~ "SGA",
                  sga=="Not SGA" ~ "Not SGA",
                  TRUE ~ "Missing"),
  prenat_ster = case_when(prenat_ster == "Yes" ~ "Yes",
                          prenat_ster == "No" ~ "No",
                          TRUE ~ "Missing"),
  com_prenat_ster = case_when(com_prenat_ster == "Yes" ~ "Yes",
                              com_prenat_ster == "No" ~ "No",
                              TRUE ~ "Missing"),
  mat_chorio = case_when(mat_chorio == "Yes" ~ "Yes",
                         mat_chorio == "No" ~ "No",
                         TRUE ~ "Missing")) %>%
mutate(mat_ethn = factor(mat_ethn, levels = c('Hispanic or Latino',
                                              'Not Hispanic or Latino', 'Missing')),
       del_method = factor(del_method, levels = c('Vaginal delivery', 'Cesarean section',
                                                  'Missing')),
       sga = factor(sga, levels = c('SGA', 'Not SGA', 'Missing')),
       prenat_ster = factor(prenat_ster, levels = c('Yes', 'No', 'Missing')),
       com_prenat_ster = factor(com_prenat_ster, levels = c('Yes', 'No', 'Missing')),
       mat_chorio = factor(mat_chorio, levels = c('Yes', 'No', 'Missing'))) %>%
tbl_summary(by=Death,

```

```

    label = list(gender ~ "Gender",
                  #mat_race ~ "Race",
                  mat_ethn ~ "Ethnicity",
                  bw ~ "Birth Weight",
                  ga ~ " Gestational Age",
                  blength ~ "Birth Length",
                  birth_hc ~ "Birth Head Circumference",
                  del_method ~ "Delivery Method",
                  prenat_ster ~ "Prenatal Corticosteroids",
                  sga ~ "Small for gestational age",
                  com_prenat_ster ~ "Complete Prenatal Steroids",
                  mat_chorio ~ "Maternal Chorioamnionitis",
                  hosp_dc_ga ~ "Hospital Discharge Gestational Age"
                  ),
    statistic = all_continuous() ~ "{mean} ({sd})" %>%
  modify_spanning_header(update = all_stat_cols() ~ "**Death**") %>%
  modify_footnote(update = all_stat_cols() ~ "Mean (SD) for continuous; n (%) for categori

schimd_tbl <- tbl_merge(list(schmid_trach, schmid_death),
                           tab_spanner = c("**Trachostomy**", "**Death**"))

schimd_tbl %>%
  as_gt() %>%
  cols_align(aligned = "center", columns = everything()) %>%
  tab_options(
    table.width = pct(120),
    table.align = "center"
  )

schmid_data <- schmid_data %>%
  mutate(Outcome = case_when(Trach == 1 & Death == "Yes" ~ "Trach/Death",
                             Trach == 1 & Death == "No" ~ "Trach/No Death",
                             Trach == 0 & Death == "Yes" ~ "No Trach/Death",
                             Trach == 0 & Death == "No" ~ "No Trach/No Death"))

schmid_count <- schmid_data %>%
  filter(!is.na(center)) %>%
  group_by(center) %>%
  count(Outcome) %>%
  spread(Outcome, n, fill = 0)
schmid_count <- schmid_count %>%
  gather(`Trach/Death`, `No Trach/Death`, `No Trach/No Death`, `Trach/No Death`, `<NA>`,
        key = "Outcomes", value = "Count") %>%

```



```

mutate(Outcomes = factor(Outcomes,
                          levels = c("Trach/Death", "No Trach/Death", "Trach/No Death",
                                       "No Trach/No Death", "<NA>"))) %>%

group_by(center) %>%
mutate(Proportion = Count/sum(Count))

ggplot(schmid_count, aes(x = Proportion, y = factor(center), fill = Outcomes)) +
  geom_bar(stat = "identity", position = "stack", color = "black") +
  labs(title = "Outcome Proportion by Center", x = "Proportion", y = "Center") +
  theme_minimal() +
  theme(legend.position = "bottom") +
  scale_fill_manual(values = c("Trach/Death" = "#EF8100",
                               "No Trach/Death" = "#FFA842",
                               "Trach/No Death" = "#FFD925",
                               "No Trach/No Death" = "#FFEB8C",
                               "<NA>" = "gray")) +
  theme(legend.text = element_text(size = 6),
        plot.title = element_text(size = 10),
        axis.title = element_text(size = 8)
        )

schmid_data$ventilation_support_level.36 <-
  as.factor(schmid_data$ventilation_support_level.36)
schmid_data$ventilation_support_level_modified.44 <-
  as.factor(schmid_data$ventilation_support_level_modified.44)

schmid_data$Trach <- as.factor(schmid_data$Trach)
schmid_data$Death <- as.factor(schmid_data$Death)

schmid_long <- schmid_data %>%
  rename(ventilation_support_level.44 = 'ventilation_support_level_modified.44') %>%
  pivot_longer(cols = c('ventilation_support_level.36',
                        'ventilation_support_level.44',
                        'p_delta.36',
                        'p_delta.44'),
              names_to = c(".value", "time"),
              names_pattern = "(ventilation_support_level|p_delta).(\\d+)")
schmid_long$time <- as.factor(schmid_long$time)
schmid_long$Trach <- as.factor(schmid_long$Trach)
schmid_long$ventilation_support_level <- as.factor(schmid_long$ventilation_support_level)
weight_df <- schmid_data %>%
  pivot_longer(cols = c("bw", "weight_today.36", "weight_today.44"),

```

```

      names_to = "Time",
      values_to = "Weight")

weight_df <- weight_df %>%
  mutate(Time = case_when(Time == "bw" ~ "Baseline",
                           Time == "weight_today.36" ~ "Week 36",
                           Time == "weight_today.44" ~ "Week 44",
                           TRUE ~ NA))
options(repr.plot.width = 10, repr.plot.height = 10)

weight_df %>%
  group_by(Time) %>%
  ggplot(aes(x = Weight, fill = Outcome)) +
  geom_histogram(bins = 20, position = "stack", color = "black", alpha = 0.7, size = 0.1)
  labs(title = "Birth Weight Distribution by Tracheostomy/Death Outcome",
        x = "Birth Weight (g)", y = "Count") +
  facet_wrap(~Time, ncol = 3) +
  theme_minimal() +
  theme(legend.position = "bottom") +
  scale_fill_manual(values = c("#F48126", "skyblue", "red", "greenyellow")) +
  theme(legend.text = element_text(size = 6),
        legend.title = element_text(size = 8),
        axis.text = element_text(size = 6),
        plot.title = element_text(size = 10),
        axis.title = element_text(size = 8)
  )

library(HDSinRdata)
library(mice)
library(dplyr)
library(ggplot2)
library(kableExtra)
library(visdat)
library(gtsummary)
data(pain)

vis_dat(pain[, 2:75], warn_large_data = FALSE) +
  theme(axis.text.x = element_text(angle = 90, size = 5)) +
  theme(legend.position = "bottom")

vis_dat(pain[, c(76:87, 89:92)]) +

```

```

theme(axis.text.x = element_text(angle = 90, size = 5)) +
theme(legend.position = "bottom")

vis_dat(pain[, 88]) +
  theme(axis.text.x = element_text(angle = 90, size = 5)) +
  theme(legend.position = "bottom")
pain_followup <- pain %>%
  mutate(follow_up_status = ifelse(is.na(PAIN_INTENSITY_AVERAGE.FOLLOW_UP),
                                   "No Follow-up", "Follow-up")) %>%
  mutate(MEDICAID_BIN = ifelse(is.na(MEDICAID_BIN), "Missing", MEDICAID_BIN))

tbl <- pain_followup %>%
  select(follow_up_status, colnames(pain)[c(76:87, 89:92)]) %>%
  tbl_summary(
    by = follow_up_status,
    missing = "no"
  ) %>%
  add_p(
    test = list(
      all_categorical() ~ "fisher.test",
      all_continuous() ~ "t.test"
    ),
    test.args = list(
      all_categorical() ~ list(simulate.p.value = TRUE)
    )
  ) %>%
  modify_table_body(~ .x %>% dplyr::mutate_at(vars(label), as.character))

tbl
pain_mod <- pain[, -c(1:75)] %>%
  select(-PAIN_INTENSITY_AVERAGE.FOLLOW_UP)

pain_mod$PAT_SEX <- as.factor(pain_mod$PAT_SEX)
pain_mod$PAT_RACE <- as.factor(pain_mod$PAT_RACE)
pain_mod$CCI_BIN <- as.factor(pain_mod$CCI_BIN)
pain_mod$MEDICAID_BIN <- as.factor(pain_mod$MEDICAID_BIN)
# pain_mice <- mice(pain_mod, maxit = 5, seed = (2550))
# saveRDS(pain_mice, file = "mice_result.rds")
pain_mice <- readRDS("mice_result.rds")

str(pain_mice)

```

```

score_list <- list()

for (i in 1:5) {
  mice_imp <- mice::complete(pain_mice, i)

  score <- mice_imp %>%
    group_by(PAIN_INTENSITY_AVERAGE) %>%
    summarize(
      avg_mental_health = mean(GH_MENTAL_SCORE),
      se_mental_health = sd(GH_MENTAL_SCORE)/sqrt(n()),
      avg_physical_health = mean(GH_PHYSICAL_SCORE),
      se_physical_health = sd(GH_PHYSICAL_SCORE)/sqrt(n())
    )

  score_list[[i]] <- score
}

score1_avg <- score_list[[1]][, c("PAIN_INTENSITY_AVERAGE", "avg_mental_health",
                                "avg_physical_health")]

knitr::kable(score1_avg,
              caption = "Average Summary of Pain Intensity Score") %>%
  kable_styling(latex_options = "HOLD_position")
mice_pool <- do.call(rbind, score_list)

score_pool <- mice_pool %>%
  group_by(PAIN_INTENSITY_AVERAGE) %>%
  summarize(
    avg_mental_health_pool = mean(avg_mental_health),

    se_mental_health_pool = sqrt(mean(se_mental_health^2)
  + (sum((avg_mental_health - mean(avg_mental_health))^2))/4
  + (sum((avg_mental_health - mean(avg_mental_health))^2))/20),

    avg_physical_health_pool = mean(avg_physical_health),

    se_physical_health_pool = sqrt(mean(se_mental_health^2)
  + (sum((avg_physical_health - mean(avg_physical_health))^2))/4
  + (sum((avg_physical_health - mean(avg_physical_health))^2))/20)
  )

ggplot(score_pool, aes(x = factor(PAIN_INTENSITY_AVERAGE))) +

```

```

geom_point(aes(y = avg_mental_health_pool, color = "Mental Health")) +
geom_line(aes(y = avg_mental_health_pool, color = "Mental Health"), group = 1) +
geom_errorbar(aes(ymin = avg_mental_health_pool - se_mental_health_pool,
                  ymax = avg_mental_health_pool + se_mental_health_pool,
                  color = "Mental Health",), alpha = 0.6) +
geom_point(aes(y = avg_physical_health_pool, color = "Physical Health")) +
geom_line(aes(y = avg_physical_health_pool, color = "Physical Health"), group = 1) +
geom_errorbar(aes(ymin = avg_physical_health_pool - se_physical_health_pool,
                  ymax = avg_physical_health_pool + se_physical_health_pool,
                  color = "Physical Health"), alpha = 0.6) +
labs(x = "Pain Intensity", y = "Average Score",
      title = "Average Mental and Physical Health by Pain Intensity with SE",
      color = "Score Type") +
theme_minimal()

```