

Pointwise Complexity for Gaussian Fields: Upper Envelopes, Algorithmic Lower Bounds, and Finite-Cutoff Renormalization

Yunbei Xu

National University of Singapore

yunbei@nus.edu.sg

Abstract

We prove a variance-aware pointwise majorizing-measure theorem for centered Gaussian processes. Classical generic chaining characterizes the scalar quantity $\mathbb{E} \sup_{x \in T} X_x$; the theorem here gives a simultaneous high-probability envelope for the entire field. For an ambient prior μ , the envelope at x is governed by a localized Fernique–Talagrand functional $\Phi_\mu(x)$, truncated at the variance scale $\sigma(x) = (\mathbb{E} X_x^2)^{1/2}$, together with the corresponding Gaussian tail term. Optimizing the resulting in-expectation bound recovers the anchored Fernique–Talagrand scale up to universal constants. The theorem is intended as a standard reference whenever one needs a field-level refinement of classical generic chaining, and as the Gaussian-process counterpart of pointwise empirical-process bounds for deep neural networks (Li and Xu, 2026).

We also derive a single-radius Bayesian algorithmic lower envelope from the interactive Fano/data-processing principle (Chen et al., 2024). For a known prior, an observation channel, and a concrete estimator $\hat{t}(Y)$, the lower bound is expressed through the exact ghost small-ball mass $\mathbb{E}_{Y \sim Q} \pi(B_d(\hat{t}(Y), \Delta))$, rather than a worst-case covering number. In Gaussian location experiments, comparison decoders convert Bayes location error into lower bounds on decision-aligned Gaussian ranges. This includes ordinary nearest-neighbor decoding and, importantly in overparameterized ambient classes, unrestricted norm-regularized nearest-neighbor rules. The resulting lower bound is valid for arbitrary priors, with magnitude controlled by the ghost-mass-induced pointwise dimension. For hard priors localized at the tested radius, a class-supremum relaxation recovers a Sudakov-type single-radius obstruction. We then construct an elementary heterogeneous-orthogonal-basis example separating the usual Fano relaxation for a fixed prior, the Bayesian algorithmic lower envelope, the pointwise Gaussian envelope on the estimator-selected subatlas, and the full-class minimax risk/global Gaussian scale. Together, these results show that Bayesian algorithmic lower bounds can serve as local-geometric certificates of pointwise complexity for fixed estimators in overparameterized ambient classes.

We further isolate a pointwise-complexity principle for statistical field theory and finite-cutoff renormalization, guided by probabilistic and physical motivation. For renormalization maps and interacting fields, we prove finite-cutoff theorems and construct a nonconvex phase-atlas example, illustrating a route to controlling coarse marginals of globally non-log-concave and anisotropic measures beyond the scope of traditional convex geometry. An exact finite-step telescoping identity replaces one renormalization level at a time; the resulting secant response Gram defines a pointwise ellipsoidal metric; and a hierarchical prior over phase labels and active subspaces splits the complexity into a local effective dimension and a global atlas cost. This makes precise the structural analogy between exponential-family RG and nonlinear feature learning in fully connected DNNs (Li and Xu, 2026): both settings exploit a composite linear structure. For free fields and local time, we specialize the pointwise Gaussian theorem to finite pinned Gaussian free fields. The resulting target-set local-time envelopes explicitly separate direct field-level control from reductions through full cover time. On phase-star graphs, a selected subatlas pays its own atlas code length and chart cost, while any full-cover route must pay for all phases.

Contents

1	Introduction and Main Results	2
1.1	Main pointwise Gaussian theorem	4
1.2	Main Bayesian algorithmic lower bounds and consequences	6
1.3	Separating example construction and pointwise certification	9
1.4	Main finite-cutoff renormalization theorem	12
1.5	Main graph local-time application	13
1.6	Organization	13
2	Proof of the Pointwise Gaussian Theorem	13
3	Proofs and Consequences of the Bayesian algorithmic lower bounds	17
3.1	Proof of the Bayesian algorithmic lower and upper bounds	17
3.2	An algorithmic relaxation route to Sudakov-type comparison	18
4	A separating example: overparameterized class and pointwise certification	20
5	Finite-Cutoff Renormalization and Pointwise Dimension	25
5.1	Pointwise complexity for nonlinear RG maps	26
5.2	A finite nonconvex phase-atlas example	30
6	Graph Local-Time Application versus Cover Time	33
6.1	Target local times versus cover-time reduction	34
7	Conclusion	39

1 Introduction and Main Results

Classical generic chaining controls the scalar quantity $\mathbb{E} \sup_{x \in T} X_x$ for a Gaussian process $\{X_x\}_{x \in T}$. For many field-level questions, however, this scalar is not the first object one wants to estimate. In the graph setting, the second Ray–Knight theorem identifies a local-time field at an inverse-local-time stopping rule with a shifted square of a Gaussian free field. In constructive and lattice field theory, one similarly studies local observables before passing to global maxima or continuum limits. This suggests that the Gaussian input should itself be pointwise: one should first prove a simultaneous field-level envelope and only afterwards take suprema or threshold events.

A further motivation comes from recent work on pointwise generalization for deep neural networks (Li and Xu, 2026). There the central object is not a class-wide supremum but a hypothesis-dependent finite-scale complexity. The proof is structural: an exact non-perturbative telescoping expansion exposes learned feature Gram matrices, and a hierarchical local-chart/global-atlas prior converts the resulting data-dependent active subspaces into a data-independent bound. The present paper develops the Gaussian-process counterpart of that viewpoint. Theorem 1.1 can serve as a standard reference whenever one needs a field-level refinement of classical generic chaining.

We also record a single-radius Bayesian lower-envelope counterpart. It is not a pathwise lower bound on each Gaussian coordinate. Rather, it is a data-processing Bayes-risk lower bound for a specified prior, observation channel, and estimator. In Gaussian location experiments, comparison decoders turn Bayes location error into an algorithmic lower bound on a decision-aligned Gaussian range. Conversely, a simultaneous pointwise Gaussian envelope yields an algorithmic upper bound

on the same decoding risk. Ordinary nearest-neighbor decoding is the simplest instance; the separating example in Theorem 1.7 uses an unrestricted norm-regularized nearest-neighbor rule over the full ambient class, reflecting the inductive-bias/interpolation principle that good empirical fit must be paired with a locality or complexity bias (Cover and Hart, 1967; Belkin et al., 2018). This conversion is variance-normalized and valid for arbitrary priors, with magnitude controlled by the ghost-mass-induced pointwise dimension. When applied to hard priors localized at the tested radius, a class-supremum relaxation yields a Sudakov-type single-radius obstruction. We then construct an elementary heterogeneous-orthogonal-basis “hub–leaves–cloud” example in which the fixed-prior Fano relaxation is vacuous, the algorithmic ghost-mass lower envelope is sharp for the chosen Bayes problem and for the pointwise upper envelope on the estimator-selected subatlas, while the full-class minimax risk and the global Gaussian supremum scale are much larger.

These results clarify why a Bayesian algorithmic lower bound can provide the right certificate for a pointwise upper envelope under a specified prior, even when the ambient class is overparameterized. In this sense, the pointwise and algorithmic viewpoint developed here is analogous to probabilistic refinements of worst-case analysis in theoretical computer science (Yao, 1977; Levin, 1986; Spielman and Teng, 2004). Just as distributional or smoothed input models can make the effective computational complexity of an algorithm much smaller than its worst-case complexity, a concrete estimator and reference law can make the effective probabilistic or information-theoretic complexity of a rich field much smaller than what is suggested by a global supremum or class-wide relaxation.¹

The paper then records a finite-cutoff renormalization analogue of this pointwise viewpoint. The analogy with deep feature learning is not simply that layers resemble levels. Rather, the essential common structure is the internal linear *feature/operator inner product* that appears within each composite layer or level. The shared proof architecture is exact finite-step telescoping, a pointwise ellipsoidal metric induced by a learned or renormalized response Gram, and a hierarchical atlas separating local effective dimension from global prior-independence cost. The practical significance in statistical field theory is that RG is fundamentally about controlling coarse marginals after integrating out fluctuations. For globally non-log-concave and anisotropic interacting fields, a direct scalar Lipschitz or global convex-geometric bound is usually too crude. The finite-cutoff viewpoint instead allows one to work on stable local charts where the effective action has a curvature lower bound; directions below the finite resolution are absorbed as approximation error, and the global nonconvexity is paid through an atlas cost. We state this part as an abstract deterministic complexity theorem, together with a chartwise nonconvex example. The graph application later in the paper uses the same local-to-global logic to compare two probabilistic questions: controlling a local-time field on a specified target set versus reducing the problem to global cover time. The phase-star example shows that retaining the target field can give a strictly smaller scale than any argument that first proves full cover.

Terminology for radii, resolutions, and RG levels. The word “scale” appears in several neighboring literatures, so we keep the terminology separate. The Bayesian lower envelope is a *single-radius* statement at a testing radius Δ , whereas generic chaining is multiresolution and integrates over radii. The Gaussian upper theorem also uses neighborhood radii, such as ε , r_0 , and s_0 , only as metric resolutions. In the renormalization section we use *RG level* for the index of a block-spin or fluctuation integration, and *finite-cutoff* or *finite-step* for the non-infinitesimal nature

¹Here “algorithmic” is used in the decision-theoretic sense of bounds depending on a concrete estimator. This is distinct from classical algorithmic information theory, which is traditionally formulated for finite strings or countable description spaces. For related notions of pointwise dimension and their connections to algorithmic and effective statistical dimension, see Li and Xu (2026); Lutz (2016).

of the telescoping argument. Finally, in the separating example the numbers R and S are norms of two different parts of the ambient class. These are related by analogy but are not the same mathematical object.

1.1 Main pointwise Gaussian theorem

Let $\{X_x\}_{x \in T}$ be a centered Gaussian process on a finite index set T . Fix an anchor $x_0 \in T$ such that $X_{x_0} = 0$ almost surely, and define

$$d(x, y) := (\mathbb{E}|X_x - X_y|^2)^{1/2}, \quad \sigma(x) := (\mathbb{E}X_x^2)^{1/2} = d(x, x_0), \quad R_* := \sup_{x \in T} \sigma(x).$$

For an ambient prior $\mu \in \Delta(T)$, where $\Delta(T)$ denotes the set of probability measures on T , define the localized pointwise Fernique–Talagrand functional

$$\Phi_\mu(x) := \int_0^{4\sigma(x)} \sqrt{\log \frac{1}{\mu(B_d(x, \varepsilon))}} d\varepsilon, \quad B_d(x, \varepsilon) := \{y \in T : d(x, y) \leq \varepsilon\}. \quad (1.1)$$

Here the upper limit is local: it is proportional to the standard deviation $\sigma(x)$, rather than the global radius. The factor 4 is inessential; it absorbs the two constant-factor losses coming from anchoring and from the nearest-point pushforward used in the ambient-equivalence step. If one instead works with the non-local full-radius integral, then the upper limits R_* and $2R_*$ are equivalent up to universal constants, since the entropy integrand is nonincreasing in ε .

Write

$$\Phi_{\mu,*} := \sup_{x \in T} \Phi_\mu(x), \quad \log_2^+(u) := \max\{\log_2 u, 0\}.$$

The degenerate case $R_* = 0$ is trivial, and the statements below are understood to be vacuous if $\Phi_{\mu,*} = +\infty$.

For $r_0 > 0$, $s_0 \in (0, R_*]$, and $\delta \in (0, 1)$, set

$$M_{r_0, s_0} := (1 + \lceil \log_2^+(\Phi_{\mu,*}/r_0) \rceil)(1 + \lceil \log_2(R_*/s_0) \rceil), \quad (1.2)$$

and

$$\mathcal{E}_{\mu, r_0, s_0}(x; \delta) := \max\{\Phi_\mu(x), r_0\} + \max\{\sigma(x), s_0\} \sqrt{\log \left(\frac{eM_{r_0, s_0}}{\delta} \right)}. \quad (1.3)$$

Thus all logarithmic losses from the two successive pointwise-peeling steps are isolated in M_{r_0, s_0} .

We state the main result as a variance-aware pointwise Gaussian majorizing-measure theorem in high probability.

Theorem 1.1 (Pointwise Gaussian majorizing-measure bound). *Assume $0 < R_* < \infty$ and $\Phi_{\mu,*} < \infty$. There exists an absolute constant $C > 0$ such that, for every ambient prior $\mu \in \Delta(T)$, every $r_0 > 0$, every $s_0 \in (0, R_*]$, and every $\delta \in (0, 1)$, with probability at least $1 - \delta$,*

$$\forall x \in T : \quad X_x \leq C \mathcal{E}_{\mu, r_0, s_0}(x; \delta). \quad (1.4)$$

Consequently, applying the same bound to X and $-X$ and taking a union bound, with probability at least $1 - \delta$,

$$\forall x \in T : \quad |X_x| \leq A_{\mu, r_0, s_0}(x; \delta), \quad A_{\mu, r_0, s_0}(x; \delta) := C \mathcal{E}_{\mu, r_0, s_0}(x; \delta/2). \quad (1.5)$$

When r_0, s_0 are fixed, we abbreviate A_{μ, r_0, s_0} to A_μ .

Besides providing the Gaussian-process counterpart of the pointwise empirical-process bound in Li and Xu (2026, Theorem1), the present result makes a key technical refinement: the global boundedness parameter for the metric is replaced by the local variance scale $4\sigma(x)$. Moreover, in the Gaussian-process setting we do not need the advanced symmetrization machinery or the mixed empirical-ghost metric used in Li and Xu (2026, Theorem1).

Before presenting the in-expectation consequence, we record the corresponding anchored form of the classical generic-chaining scale. This is a localized reformulation of the usual FerniqueTalagrand theorem: since the cutoff $4\sigma(x)$ is not the standard full-radius presentation, the global radius R_* must remain part of the anchored scale.

Proposition 1.2 (Classical scale in anchored local form). *There exist absolute constants $0 < c < C < \infty$ such that*

$$c \left(R_* + \inf_{\mu \in \Delta(T)} \Phi_{\mu,*} \right) \leq \mathbb{E} \sup_{x \in T} X_x \leq C \left(R_* + \inf_{\mu \in \Delta(T)} \Phi_{\mu,*} \right). \quad (1.6)$$

Moreover, since the process is centered and anchored,

$$\mathbb{E} \sup_{x \in T} |X_x| \asymp \mathbb{E} \sup_{x \in T} X_x.$$

Therefore

$$\mathbb{E} \sup_{x \in T} |X_x| \asymp R_* + \inf_{\mu \in \Delta(T)} \Phi_{\mu,*}. \quad (1.7)$$

We now record the in-expectation consequence of Theorem 1.1. In particular, after optimizing over the ambient prior, the resulting global in-expectation bound is sharp up to universal constants. The variance-radius term is not an additional loss: it is part of the anchored generic-chaining scale in (1.7), while the peeling logarithms disappear by taking $r_0 \geq \Phi_{\mu,*}$ and $s_0 = R_*$.

Corollary 1.3 (In-expectation consequence and sharp global scale). *Under the assumptions of Theorem 1.1, there exists an absolute constant $C > 0$ such that*

$$\mathbb{E} \left[\sup_{x \in T} \frac{(|X_x| - C \max\{\Phi_{\mu}(x), r_0\})_+}{\max\{\sigma(x), s_0\}} \right] \leq C \sqrt{\log(eM_{r_0, s_0})}. \quad (1.8)$$

Consequently, for every $x \in T$,

$$\mathbb{E} |X_x| \leq C \max\{\Phi_{\mu}(x), r_0\} + C \max\{\sigma(x), s_0\} \sqrt{\log(eM_{r_0, s_0})}. \quad (1.9)$$

At the global level, taking $r_0 \geq \Phi_{\mu,*}$ and $s_0 = R_*$ gives $M_{r_0, s_0} = 1$, hence

$$\mathbb{E} \sup_{x \in T} |X_x| \leq C(r_0 + R_*). \quad (1.10)$$

Letting $r_0 \downarrow \Phi_{\mu,*}$, then optimizing over μ , and using (1.7), we obtain the sharp equivalence

$$\mathbb{E} \sup_{x \in T} |X_x| \asymp R_* + \inf_{\mu \in \Delta(T)} \Phi_{\mu,*}. \quad (1.11)$$

Thus the optimized expectation version is optimal up to absolute constants.

Remark (variance term versus chaining complexity). The term in Theorem 1.1 involving $\sigma(x)$ is a variance-sensitive Gaussian tail term. It is necessary for high-probability control, already for a single non-anchor point $X_x \sim N(0, \sigma(x)^2)$, but it is not a replacement for the chaining complexity $\Phi_\mu(x)$, which carries the local geometry of the index set. For a fixed marginal, $\mathbb{E}|X_x| = \sqrt{\frac{2}{\pi}} \sigma(x)$, so the extra factor $\sqrt{\log(eM_{r_0, s_0})}$ in (1.9) is not intrinsic to $\mathbb{E}|X_x|$. It is the price of extracting a simultaneous pointwise envelope by peeling first over the local Fernique–Talagrand complexity and then over the local variance. At the global in-expectation level, this peeling factor disappears, yielding the sharp scale in (1.11).

Remark (pointwise high probability). For Gaussian processes, subset-level in-expectation bounds transfer to high-probability bounds through Borell–Tsirelson concentration, with variance scale $\sigma_H = \sup_{x \in H} (\mathbb{E}X_x^2)^{1/2}$ for a fixed subset H . The genuinely pointwise statement is not obtained from this transfer alone: after the subset-homogeneous estimate is established, one first applies uniform pointwise peeling to localize the Fernique–Talagrand term $\Phi_\mu(x)$, and then applies the same peeling mechanism a second time to improve the tail scale from the global radius to the local variance $\sigma(x)$. Conversely, the resulting high-probability envelope implies the stated in-expectation consequence by integrating the tail.

1.2 Main Bayesian algorithmic lower bounds and consequences

The pointwise upper envelope is non-decision-theoretic, simultaneous, multiscale, and pathwise in the Gaussian field. A lower-bound analogue should therefore be stated with care. We first record a one-scale Bayesian and estimator-dependent primitive. Let (T, d) be a finite metric space, let $\pi \in \Delta(T)$ be a known prior, and let $\{P_t : t \in T\}$ be an experiment on an observation space $\mathcal{Y}$. For an estimator $\hat{t} : \mathcal{Y} \rightarrow T$, a reference law $Q \in \Delta(\mathcal{Y})$, and a scale $\Delta > 0$, define

$$\rho_{\Delta, Q}(\hat{t}) := \mathbb{P}_{t \sim \pi, Y \sim Q} \{d(t, \hat{t}(Y)) < \Delta\} = \mathbb{E}_{Y \sim Q} \pi(B_d(\hat{t}(Y), \Delta)),$$

and

$$\mathcal{I}_\pi(Q) := \mathbb{E}_{t \sim \pi} D_{\text{KL}}(P_t \| Q).$$

The following result is the Gaussian-process and KL specialization of the interactive Fano method of Chen et al. (2024, Theorem 2). The original theorem provides a unifying information-theoretic principle for proving lower bounds in learning and decision-making problems.

Proposition 1.4 (Bayesian algorithmic lower envelope). *For every estimator $\hat{t}$, every $\Delta > 0$, and every reference law Q with $0 < \rho_{\Delta, Q}(\hat{t}) < 1$,*

$$\mathbb{P}_{t \sim \pi, Y \sim P_t} \{d(t, \hat{t}(Y)) \geq \Delta\} \geq \left(1 - \frac{\mathcal{I}_\pi(Q) + \log 2}{\log(1/\rho_{\Delta, Q}(\hat{t}))}\right)_+. \quad (1.12)$$

Consequently,

$$\mathbb{E}_{t \sim \pi, Y \sim P_t} d(t, \hat{t}(Y)) \geq \sup_{\Delta > 0, Q} \Delta \left(1 - \frac{\mathcal{I}_\pi(Q) + \log 2}{\log(1/\rho_{\Delta, Q}(\hat{t}))}\right)_+. \quad (1.13)$$

The key point is that $\rho_{\Delta, Q}(\hat{t})$ is not immediately replaced by the class-wide quantity

$$q_\pi(\Delta) := \sup_{a \in T} \pi(B_d(a, \Delta)).$$

Keeping the exact ghost mass $\mathbb{E}_{Y \sim Q} \pi(B_d(\hat{t}(Y), \Delta))$ makes the lower bound algorithmic: it is attached to the image of the actual estimator under ghost data. Replacing it by $q_\pi(\Delta)$ gives the class-wide small-ball term used in classical Fano-type lower bounds (Zhang, 2006) and in the fractional-covering specialization of Chen et al. (2024). In the Gaussian nearest-neighbor comparison below, this small-ball term must still be balanced against the information radius of the Gaussian channel.

We next present a lower-bound comparison for the Gaussian process itself, rather than only for its induced distance. The result is stated for a comparison decoder. This includes ordinary nearest neighbor and the norm-regularized rule used in the separating example in Theorem 1.7.

Theorem 1.5 (Algorithmic lower bound for comparison decoders). *Let $v : T \rightarrow \mathcal{H}$ be an embedding into a finite-dimensional Hilbert space and set $d(s, t) = \|v_s - v_t\|_{\mathcal{H}}$. Let*

$$Y = v_\Theta + \tau Z, \quad \Theta \sim \pi, \quad Z \sim N(0, I_{\mathcal{H}}),$$

and let $m \in \mathcal{H}$ be any reference center. Set

$$Q_{\tau, m} := N(m, \tau^2 I_{\mathcal{H}}), \quad \mathcal{I}_{\pi, m} := \frac{1}{2\tau^2} \int_T \|v_t - m\|^2 \pi(dt).$$

With $\bar{v}_\pi := \int_T v_t \pi(dt)$, the centered choice $m = \bar{v}_\pi$ gives

$$\mathcal{I}_{\pi, \bar{v}_\pi} = \frac{V_\pi}{4\tau^2}, \quad V_\pi := \iint d(s, t)^2 \pi(ds) \pi(dt).$$

Let $\hat{\Theta}(Y)$ be an estimator with values in T . Assume that there is a penalty $\Omega : T \rightarrow \mathbb{R}$ such that, under the true joint law,

$$\|Y - v_{\hat{\Theta}}\|^2 + \Omega(\hat{\Theta}) \leq \|Y - v_\Theta\|^2 + \Omega(\Theta), \quad \Omega(\hat{\Theta}) \geq \Omega(\Theta) \quad a.s. \quad (1.14)$$

If, for some $c \in (0, 1)$ and $\Delta > 0$,

$$\mathcal{I}_{\pi, m} + \log 2 \leq (1 - c) \log \frac{1}{\rho_{\Delta, Q_{\tau, m}}(\hat{\Theta})}, \quad (1.15)$$

then

$$\mathbb{E}_{\Theta, Y} d(\Theta, \hat{\Theta}(Y)) \geq c\Delta, \quad (1.16)$$

and, for the Gaussian process $X_t := \langle Z, v_t \rangle$ built from the same Gaussian vector Z ,

$$\mathbb{E}_{\Theta, Z} [X_{\hat{\Theta}} - X_\Theta] \geq \frac{c^2 \Delta^2}{2\tau}. \quad (1.17)$$

Thus the exact ghost mass gives an algorithmic lower bound on the Gaussian range sampled by the comparison decoder.

Ordinary nearest neighbor corresponds to $\Omega \equiv 0$. The unrestricted norm-regularized nearest-neighbor rule

$$\hat{\Theta}_\lambda(Y) \in \arg \min_{a \in T} \{\|Y - v_a\|^2 + \lambda \|v_a\|^2\}$$

is also covered whenever the realized outputs have norm at least the norm of the true parameter, as in the separating example in Theorem 1.7. This is a canonical minimum-complexity inductive bias consist with modern perspectives, rather than an explicit restriction of the hypothesis class.

The same pathwise comparison underlying Theorem 1.5, together with the pointwise Gaussian upper envelope, gives an algorithmic upper bound for Gaussian location decoding. Thus, when the two estimates match, the Bayesian algorithmic lower bound and the pointwise upper envelope certify the sharpness of one another.

Corollary 1.6 (Algorithmic pointwise upper bound for estimator decoding). *Work in the Gaussian location setup of Theorem 1.5. Thus T is embedded as $\{v_t : t \in T\}$ in a Hilbert space,*

$$d(s, t) = \|v_s - v_t\|, \quad Y = v_\Theta + \tau Z, \quad X_t := \langle Z, v_t \rangle,$$

where Z is standard Gaussian. Let $\hat{\Theta} = \hat{\Theta}(Y)$ be a decoder satisfying the comparison condition (1.14); in particular, the condition implies

$$\|Y - v_{\hat{\Theta}(Y)}\|^2 \leq \|Y - v_\Theta\|^2 \quad (1.18)$$

under the joint law of (Θ, Y) .

Suppose that a deterministic envelope $A : T \rightarrow [0, \infty)$ satisfies

$$\mathbb{P}_Z \{ \forall t \in T : |X_t| \leq A(t) \} \geq 1 - \delta. \quad (1.19)$$

Then, with probability at least $1 - \delta$ under the joint law of (Θ, Y) ,

$$d(\Theta, \hat{\Theta}(Y))^2 \leq 2\tau(A(\Theta) + A(\hat{\Theta}(Y))). \quad (1.20)$$

Consequently, if

$$D_T := \sup_{s, t \in T} d(s, t) < \infty,$$

then

$$\mathbb{E} d(\Theta, \hat{\Theta}(Y)) \leq \mathbb{E} \sqrt{2\tau(A(\Theta) + A(\hat{\Theta}(Y)))} + \delta D_T. \quad (1.21)$$

In particular, after adjoining an anchor point t_0 with $v_{t_0} = 0$ and $X_{t_0} = 0$ if necessary, one may take

$$A(t) = A_{\mu, r_0, s_0}(t; \delta)$$

from (1.5) in Theorem 1.1, with the prior extended to $T \cup \{t_0\}$. This yields a pointwise-complexity upper bound for the estimation risk.

We defer to Section 3.2 a route from the algorithmic bound to a Sudakov-type comparison via supremum relaxation, which consists of several technical results.

Why keep the Bayesian and algorithmic form. The Bayesian algorithmic form is crucial because a full-class minimax benchmark can be the wrong certificate for an overparameterized decision problem, while a restricted minimax benchmark can be *oracle-dependent*. Full minimax over T can be too pessimistic, because it is governed by the worst point of the ambient class rather than by the geometry selected by the prior and the algorithm; maximizing over priors can similarly erase the structure that makes a particular estimator appropriate. On the other hand, a restricted minimax benchmark is useful only when the correct restricted class is known in advance. In the separating example in Theorem 1.7, the relevant class is the hub-leaf subatlas H_M , but identifying H_M as the right estimator-selected subatlas is precisely the decision-dependent information revealed by the ghost image $Y \mapsto \hat{t}(Y)$. Along the sequence $N = M^K$ with $K \rightarrow \infty$, the norm-regularized decoder uses a fixed numerical penalty λ , independent of the prior π and the problem

parameters M, N, R, S, τ , and has Bayes risk $\asymp R$, whereas the full-class minimax risk is $\asymp S$, with $\frac{S}{R} \asymp \sqrt{\frac{\log N}{\log M}} = \sqrt{K} \rightarrow \infty$. Thus the full-class benchmark misses the decision-aligned scale, while the restricted benchmark recovers it only after an oracle has selected H_M .

The same issue is compounded in deep networks: the effective subatlas is rarely a fixed combinatorial class and may depend on the architecture, data distribution, learned representation, and compressed feature spectrum (Li and Xu, 2026). Moreover, computationally efficient procedures may also operate through relaxations or implicit regularization rather than by solving an exact restricted-class problem, as in sparse regression. The algorithmic ghost mass $\rho_{\Delta, Q}(\hat{t})$ is designed for this regime: it evaluates the estimator actually used, under the intended prior geometry, and therefore can certify a pointwise upper envelope without requiring a separate restricted class to be specified in advance.

The Bayesian *algorithmic* lower bounds in Proposition 1.4 and Theorem 1.5 should also not be read as ordinary Bayesian lower bounds whose purpose is to optimize over all estimators given a specific prior. The prior is an averaging and comparison device, while the certified object is the performance of a concrete estimator that is of practical interests regardless of knowledge of the prior. This distinction is important because the usual Fano relaxation may be vacuous for a fixed nonuniform prior even though the estimator of interest remains hard on the subgeometry selected by the prior and the ghost law. Such vacuity does not mean that the Bayesian decision problem is easy (in the separating example in Theorem 1.7, the Bayes-optimal risk for this prior is also $\asymp R$, despite the oracle estimator being highly non-analytical). It means only that the particular worst-case small-ball relaxation $\sup_a \pi(B_d(a, \Delta))$ has discarded the algorithmic geometry of the estimator.

Relationship to the pointwise upper theorem and existing lower bounds. The Bayesian algorithmic lower bound also clarifies the relation to the pointwise upper theorem. The upper envelope is simultaneous, multiscale, and high-probability: it is pathwise in the Gaussian field and integrates local prior masses across scales. By contrast, the lower bound is Bayesian, one-scale, and estimator-dependent: it averages over $\Theta \sim \pi$ and $Y \sim P_\Theta$ for a fixed estimator, while the additional ghost/reference law $Y \sim Q$ enters through the quantile term $\rho_{\Delta, Q}(\hat{t})$. This parallels the information-theoretic upper–lower duality emphasized by Zhang (2006), but with the additional point that the lower bound is a data-processing statement valid for every estimator.

DEC-type lower bounds (Foster et al., 2021, 2023) provide an independent motivation for incorporating algorithmic dependence into the fundamental limits of online regret, exploration, and interactive decision making. This perspective further motivates the Bayesian algorithmic lower bound of Chen et al. (2024) and related variants (Xu and Zeevi, 2025b). The present algorithmic lower-bound perspective isolates a distinct but complementary principle: a prior-mass lower-bound analogue of pointwise complexity, in the spirit of the pointwise generalization framework of Li and Xu (2026) for deep representation learning.

1.3 Separating example construction and pointwise certification

The next theorem is the formal version of the separating example proved in Section 4. It shows that, in an overparameterized ambient class, a fixed-prior worst-case Fano relaxation, a Bayesian algorithmic lower envelope, a pointwise Gaussian upper envelope, and the full minimax/global Sudakov scale can all have different meanings and different orders. The estimator is unrestricted over the full ambient class; the relevant subatlas is selected by a norm-regularized inductive bias and by the ghost law, rather than imposed as a constraint.

Theorem 1.7 (Decision-aligned separation and pointwise certification). *There exist universal constants $0 < c < C < \infty$ with the following property. Let $M, N \geq 2$ satisfy $\log N \geq 1024 \log M$. Let*

$$g_0, e_1, \dots, e_M, f_1, \dots, f_N$$

be orthonormal vectors in $\mathbb{R}^{M+N+1}$. Fix $R > 0$, define

$$\tau := \frac{R}{\sqrt{\log M}}, \quad S := \frac{1}{8} \tau \sqrt{\log N} = \frac{R}{8} \sqrt{\frac{\log N}{\log M}},$$

and set

$$H_M := \{Rg_0, Re_1, \dots, Re_M\}, \quad C_N := \{Sf_1, \dots, Sf_N\}, \quad T_{M,N} := H_M \cup C_N.$$

Let $d(s, t) = \|s - t\|_2$, let $\Delta = R/3$, and let the prior be

$$\pi(Rg_0) = \frac{1}{2}, \quad \pi(Re_i) = \frac{1}{2M}, \quad i = 1, \dots, M, \quad \pi(Sf_j) = 0, \quad j = 1, \dots, N.$$

Consider the Gaussian location experiment

$$Y = \Theta + \tau Z, \quad \Theta \sim \pi, \quad Z \sim N(0, I_{M+N+1}),$$

and the unrestricted norm-regularized decoder

$$\hat{\Theta}_\lambda(Y) \in \arg \min_{a \in T_{M,N}} \{\|Y - a\|^2 + \lambda \|a\|^2\}, \quad \lambda = 64. \quad (1.22)$$

Let $Q_\tau = N(0, \tau^2 I_{M+N+1})$. Then, for all sufficiently large M, N , the following hold.

- (i) Fixed-prior Fano relaxation is vacuous. *Since every Δ -ball in $T_{M,N}$ is a singleton,*

$$q_\pi(\Delta) := \sup_{a \in T_{M,N}} \pi(B_d(a, \Delta)) = \frac{1}{2}.$$

Moreover the usual data-processing lower envelope obtained by replacing the exact ghost mass by $q_\pi(\Delta)$ gives zero at the reference law Q_τ .

- (ii) Exact algorithmic ghost mass is small. *The estimator-dependent ghost mass satisfies*

$$\rho_{\Delta, Q_\tau}(\hat{\Theta}_\lambda) = \mathbb{E}_{Y \sim Q_\tau} \pi(B_d(\hat{\Theta}_\lambda(Y), \Delta)) \leq \frac{C}{M}.$$

Consequently the exact ghost entropy is of order $\log M$, whereas the worst-case Fano small-ball entropy for the same prior is only $\log 2$.

- (iii) Sharp Bayes risk for the actual unrestricted estimator.

$$cR \leq \mathbb{E}_{\Theta, Y} d(\Theta, \hat{\Theta}_\lambda(Y)) \leq CR.$$

The lower bound follows from Proposition 1.4 using the exact ghost mass; the upper bound is the actual Bayes risk of the same unrestricted estimator.

Moreover, the the Bayes-optimal risk is also $\asymp R$, despite the oracle estimator being highly non-analytical. Thus the class-wide Fano relaxation below is vacuous only as a certificate; it does not reflect an easy Bayes problem.

(iv) Full-class minimax risk is much larger. *If*

$$\mathfrak{R}_{\text{mm}}(T_{M,N}, \tau) := \inf_{\hat{\theta}} \sup_{t \in T_{M,N}} \mathbb{E}_t d(t, \hat{\theta}(Y)),$$

where the infimum is over all measurable estimators with values in the ambient Euclidean space, then

$$cS \leq \mathfrak{R}_{\text{mm}}(T_{M,N}, \tau) \leq CS.$$

Thus, whenever $\log N / \log M \rightarrow \infty$, the full-class minimax risk is asymptotically larger than the Bayes risk under π .

(v) Algorithmically induced Gaussian certificate matches the selected subatlas. *For the Gaussian process*

$$X_t := \langle Z, t \rangle, \quad t \in T_{M,N},$$

where Z is the same standard Gaussian vector as in the location experiment $Y = \Theta + \tau Z$, define

$$\mathcal{G}_{\text{alg}} := \mathbb{E}_{\Theta, Z} [X_{\hat{\Theta}_\lambda(\Theta + \tau Z)} - X_\Theta].$$

Then

$$\mathcal{G}_{\text{alg}} \asymp \mathbb{E} \sup_{s, u \in H_M} (X_s - X_u) \asymp R \sqrt{\log M}.$$

Moreover, under the ghost/reference law $Y \sim Q_\tau = N(0, \tau^2 I)$,

$$Q_\tau \{\hat{\Theta}_\lambda(Y) \in C_N\} \leq N^{-2}$$

for all sufficiently large N , and hence the ghost-selected subatlas is H_M up to negligible probability. Conditional on $\hat{\Theta}_\lambda(Y) \in H_M$, the ghost selection law is uniform on H_M . Therefore, after adjoining the zero anchor, the pointwise Gaussian upper theorem applied to the restricted process on H_M gives the matching high-probability envelope

$$\mathbb{P} \left(\forall t \in H_M : |X_t| \leq CR \sqrt{\log \frac{M}{\delta}} \right) \geq 1 - \delta.$$

(vi) Global Gaussian scale can be arbitrarily larger.

$$\mathbb{E} \sup_{s, u \in T_{M,N}} (X_s - X_u) \geq cS \sqrt{\log N} = \frac{c}{8} R \frac{\log N}{\sqrt{\log M}}.$$

Thus the global Gaussian scale exceeds the decision-aligned scale $R \sqrt{\log M}$ by a factor of order $\log N / \log M$, which can be made arbitrarily large by taking $N = M^K$ with $K \rightarrow \infty$.

Consequently the same explicit problem sequence separates

$$\begin{aligned} & \text{fixed-prior Fano relaxation} \\ & \ll \text{algorithmic ghost-mass lower envelope} \\ & \asymp \text{pointwise upper envelope on the selected subatlas} \\ & \ll \text{full-class minimax/global Gaussian scale.} \end{aligned}$$

1.4 Main finite-cutoff renormalization theorem

We next state the finite-dimensional renormalization theorem in the form used later. Its main role is to provide an analogue of the pointwise complexity theory for deep neural networks developed in Li and Xu (2026). This analogy relies not only on the correspondence between composite layers and RG levels, but more essentially on the common *feature/operator inner-product* structure that appears within each composite layer or level. Let a depth- J RG trajectory be parametrized by level variables

$$g = (g_0, \dots, g_{J-1}), \quad g_j \in B_2(R_j) \subset \mathbb{R}^{p_j},$$

and let $S_J(g)$ denote the final effective action, or any finite-dimensional observable extracted from the final effective action, equipped with a semimetric ρ . The theorem is conditional on a finite-scale domination statement of the form

$$\rho(S_J(g'), S_J(g))^2 \leq \sum_{j=0}^{J-1} (g'_j - g_j)^\top G_j(g, \varepsilon) (g'_j - g_j) \quad (1.23)$$

for all g' in a finite ρ -neighborhood of g , where the secant Grams G_j are defined via the renormalized operators for exponential-family RG steps; see Section 5 for the details. This defines the pointwise ellipsoidal RG metric

$$\rho_{G, g, \varepsilon}(g, g')^2 := \sum_{j=0}^{J-1} (g'_j - g_j)^\top G_j(g, \varepsilon) (g'_j - g_j),$$

which is the RG analogue of the pointwise ellipsoidal metric induced by learned feature Gram matrices in the DNN theory. The matrices $G_j(g, \varepsilon)$ are not infinitesimal Hessians; they are finite-scale secant response Grams produced by replacing one RG level at a time.

For a positive semidefinite matrix G , radius R , and resolution $u > 0$, define

$$r_{\text{eff}}(G, R, u) := \max\{k : \lambda_k(G)R^2 \geq u^2\}, \quad d_{\text{ell}}(G, R, u) := \frac{1}{2} \sum_{k=1}^{r_{\text{eff}}} \log \left(\frac{CR^2 \lambda_k(G)}{u^2} \right), \quad (1.24)$$

with eigenvalues in nonincreasing order. Let $V_j(g, \varepsilon)$ be the active eigenspace of $G_j(g, \varepsilon)$, and let $\mathfrak{A}_j(g, \varepsilon, u_j)$ be the atlas cost of choosing the phase label, effective rank, and a reference subspace close to $V_j(g, \varepsilon)$. A concrete Grassmannian expression is given in Section 5.

Theorem 1.8 (Finite-scale RG pointwise dimension, main form). *Assume the finite-scale domination (1.23). Choose numbers $u_j > 0$ such that $\sum_j u_j^2 \leq c\varepsilon^2$. Then there exists a trajectory-independent hierarchical prior Π over phase labels, active ranks, reference subspaces, and local scale couplings such that*

$$\log \frac{1}{\Pi(B_\rho(g, \varepsilon))} \leq C \sum_{j=0}^{J-1} [d_{\text{ell}}(G_j(g, \varepsilon), R_j, u_j) + \mathfrak{A}_j(g, \varepsilon, u_j)]. \quad (1.25)$$

Thus finite-cutoff renormalization has the same complexity structure as nonlinear feature-learning DNN: a local ellipsoidal effective dimension plus a global atlas cost at each scale.

We then apply the theorem to a nonconvex phase-atlas construction, showcasing how finite-cutoff geometry can control coarse marginals of globally non-log-concave and anisotropic measures. The theorem and the example should be read as abstract finite-cutoff results, not as a shortcut to continuum constructive QFT. To prove a scaling-limit theorem such as four-dimensional marginal triviality, one also needs a model-specific contraction or Gaussianization estimate for non-Gaussian connected correlations. The pointwise-dimension theorem supplies the local finite-scale complexity layer once such RG or correlation estimates are available.

1.5 Main graph local-time application

We give a specialized application of the pointwise-complexity principle to the control of local times in Gaussian free fields. The results separate two distinct ways of using Gaussian information, highlighting the advantage of a field-level route. In the field-level route, one fixes a target set, controls the pinned GFF on that set by a pointwise envelope, and then transfers the resulting estimate through the RayKnight theorem to a local-time statement. In the global route, one first establishes that the entire graph has been covered and only afterwards deduces that the target set has been visited. These two stopping-time arguments may occur at different scales.

Section 6.1 proves this separation on phase-star hierarchies. A target subatlas I is encoded by an atlas prior q , and the same ambient pointwise envelope can be evaluated on the selected target. The exact law

$$\mathbb{P}\{\tau_I \leq \tau(t)\} = (1 - e^{-t})^{|I|}, \quad \mathbb{P}\{\tau_{\text{cov}} \leq \tau(t)\} = (1 - e^{-t})^N$$

shows that target cover has intrinsic inverse-root-local-time scale $\log |I|$, whereas any route through full cover pays $\log N$. The pointwise envelope pays the target code length $\log(|I|/q(I))$, and the chart-decorated phase-star adds a compressed within-chart term. This gives a concrete finite-graph analogue of the phase/subspace atlas cost in Section 5.

1.6 Organization

Section 2 proves the pointwise Gaussian theorem. Section 3 proves the Bayesian algorithmic lower bounds and their consequences. Section 4 constructs an important example separating the fixed-prior Fano relaxation, the algorithmic ghost-mass lower bound, the pointwise Gaussian envelope, and the full-class minimax/global Gaussian scale. Section 5 develops the finite-cutoff renormalization pointwise-complexity theory. Section 6 gives the graph and local-time applications. Section 7 summarizes the pointwise/algorithmic scope and the main lessons.

2 Proof of the Pointwise Gaussian Theorem

The proof follows a subset-homogeneity and peeling strategy. First, for each fixed subset $H \subseteq T$, we combine an anchored form of Fernique’s majorizing-measure upper bound (Fernique, 1975; Talagrand, 1987), the ambient equivalence principle of pointwise dimension, and Borell–Tsirelson concentration. Second, we apply the one-parameter uniform pointwise peeling lemma twice: first to localize the Fernique–Talagrand integral $\Phi_\mu(x)$, and then to localize the variance scale from R_* to $\sigma(x)$. This avoids introducing a separate two-dimensional peeling corollary.

The following uniform pointwise peeling lemma was originally developed for empirical processes in Xu and Zeevi (2025a). It identifies minimal conditions for obtaining pointwise envelopes, up to negligible double-logarithmic factors, without imposing unrealistic Bernstein-type or rigid sub-root assumptions. Necessary and sufficient justifications and discussions of data dependence are provided in Li and Xu (2026). Here we extend the lemma to general indexed processes, including Gaussian processes.

Lemma 2.1 (Uniform pointwise peeling for indexed processes). *Let $\{Z_x\}_{x \in T}$ be random variables, let $d_0 : T \rightarrow [0, R]$, and let $\psi(r; \delta)$ be nondecreasing in r and nonincreasing in δ . Assume that for every fixed subset $H \subseteq T$ and every $\delta \in (0, 1)$,*

$$\mathbb{P}\left(\sup_{x \in H} Z_x \leq \sup_{x \in H} \psi(d_0(x); \delta)\right) \geq 1 - \delta. \quad (2.1)$$

Then for every $r_0 > 0$, with probability at least $1 - \delta$, uniformly over all $x \in T$,

$$Z_x \leq \psi\left(\max\{2d_0(x), r_0\}; \frac{\delta}{1 + \lceil \log_2^+(R/r_0) \rceil}\right), \quad (2.2)$$

where $\log_2^+(u) = \max\{\log_2 u, 0\}$.

Proof. Let $m := \lceil \log_2^+(R/r_0) \rceil$. The sets

$$H_0 := \{x : d_0(x) \leq r_0\}, \quad H_j := \{x : 2^{j-1}r_0 < d_0(x) \leq 2^j r_0\} \quad (1 \leq j \leq m)$$

cover T . Apply (2.1) to each H_j with confidence level $\delta/(m+1)$, and take a union bound. On the resulting event, if $x \in H_0$, then

$$Z_x \leq \psi\left(r_0; \frac{\delta}{m+1}\right) \leq \psi\left(\max\{2d_0(x), r_0\}; \frac{\delta}{m+1}\right),$$

and if $x \in H_j$ for $j \geq 1$, then $2^j r_0 \leq 2d_0(x)$, so

$$Z_x \leq \psi\left(2^j r_0; \frac{\delta}{m+1}\right) \leq \psi\left(\max\{2d_0(x), r_0\}; \frac{\delta}{m+1}\right).$$

This proves the claim. $\square$

We also use the following localized version of the ambient equivalence principle of pointwise dimension from Lemma 7 of [Li and Xu \(2026\)](#).

Lemma 2.2 (Ambient equivalence for the localized integral). *Let (T, d) be a finite metric space and let $H \subseteq T$. Let $p : T \rightarrow H$ be a nearest-point selector, so that $d(y, p(y)) = \min_{h \in H} d(y, h)$, and let $\mu_H = p_{\#}\mu$ be the pushforward of an ambient prior $\mu \in \Delta(T)$. Then, for every $x \in H$ and every $\varepsilon > 0$,*

$$\mu_H(B_d(x, 2\varepsilon)) \geq \mu(B_d(x, \varepsilon)).$$

Consequently, for the localized functional Φ in (1.1),

$$\Phi_{\mu_H}(x) \leq 2\Phi_{\mu}(x), \quad x \in H. \quad (2.3)$$

Proof. If $y \in B_d(x, \varepsilon)$, then

$$d(p(y), x) \leq d(p(y), y) + d(y, x) \leq 2d(y, x) \leq 2\varepsilon,$$

because $x \in H$ and $p(y)$ is a nearest point in H . Hence $p(B_d(x, \varepsilon)) \subseteq B_d(x, 2\varepsilon)$, which gives the ball-mass inequality. Therefore

$$\Phi_{\mu_H}(x) = \int_0^{4\sigma(x)} \sqrt{\log \frac{1}{\mu_H(B_d(x, u))}} du \leq 2 \int_0^{2\sigma(x)} \sqrt{\log \frac{1}{\mu(B_d(x, \varepsilon))}} d\varepsilon \leq 2\Phi_{\mu}(x),$$

where we used the change of variables $u = 2\varepsilon$. $\square$

Lemma 2.3 (Anchored subset majorizing-measure bound). *For every fixed subset $H \subseteq T$, every ambient prior $\mu \in \Delta(T)$, and*

$$\sigma_H := \sup_{x \in H} \sigma(x),$$

there is an absolute constant $C > 0$ such that

$$\mathbb{E} \sup_{x \in H} X_x \leq C \sup_{x \in H} \Phi_{\mu}(x) + C\sigma_H. \quad (2.4)$$

Proof. Let $H_0 := H \cup \{x_0\}$, and let $p : T \rightarrow H_0$ be a nearest-point selector. Define the probability measure

$$\bar{\mu}_H := \frac{1}{2}\delta_{x_0} + \frac{1}{2}p_{\#}\mu$$

on H_0 . Fernique's majorizing-measure upper bound (Fernique, 1975; Talagrand, 1987), applied to the restricted process on H_0 , gives

$$\mathbb{E} \sup_{x \in H} X_x \leq C \sup_{x \in H_0} \int_0^{2\sigma_H} \sqrt{\log \frac{1}{\bar{\mu}_H(B_d(x, u))}} du,$$

since $\text{diam}(H_0, d) \leq 2\sigma_H$. For $x = x_0$, the integral is at most $2\sigma_H \sqrt{\log 2}$, because $\bar{\mu}_H\{x_0\} \geq 1/2$. For $x \in H$, split the integral at $4\sigma(x)$. On $[4\sigma(x), 2\sigma_H]$, the ball $B_d(x, u)$ contains x_0 , hence has $\bar{\mu}_H$ -mass at least $1/2$, so this part is bounded by $C\sigma_H$. On $[0, 4\sigma(x)]$, Lemma 2.2 and the extra factor $1/2$ in $\bar{\mu}_H$ give a bound of order $\Phi_\mu(x) + \sigma(x)$. Taking the supremum over $x \in H$ proves (2.4). $\square$

Lemma 2.4 (Subset-homogeneous Gaussian estimate). *For every fixed subset $H \subseteq T$, every ambient prior $\mu \in \Delta(T)$, and every $\delta \in (0, 1)$,*

$$\mathbb{P} \left(\sup_{x \in H} X_x \leq C \sup_{x \in H} \Phi_\mu(x) + C\sigma_H \sqrt{\log(e/\delta)} \right) \geq 1 - \delta, \quad (2.5)$$

where $\sigma_H = \sup_{x \in H} \sigma(x)$.

Proof. By Lemma 2.3,

$$\mathbb{E} \sup_{x \in H} X_x \leq C \sup_{x \in H} \Phi_\mu(x) + C\sigma_H.$$

Realize $X_x = \langle g, v_x \rangle$ as an isonormal process on a Hilbert space. The map $g \mapsto \sup_{x \in H} \langle g, v_x \rangle$ is σ_H -Lipschitz. Borell–Tsirelson concentration therefore gives the desired inequality, after absorbing the expectation term $C\sigma_H$ into $C\sigma_H \sqrt{\log(e/\delta)}$. $\square$

Proof of Theorem 1.1. Fix an arbitrary subset $H \subseteq T$. Applying Lemma 2.1 on the index set H , with $Z_x = X_x$, $d_0(x) = \Phi_\mu(x)$, $R = \Phi_{\mu,*}$, and using the subset estimate (2.5), yields the following: for every $r_0 > 0$ and every $\delta \in (0, 1)$, with probability at least $1 - \delta$,

$$\forall x \in H : \quad X_x \leq C \max\{\Phi_\mu(x), r_0\} + C\sigma_H \sqrt{\log\left(\frac{eM_\Phi}{\delta}\right)}, \quad (2.6)$$

where

$$M_\Phi := 1 + \lceil \log_2^+(\Phi_{\mu,*}/r_0) \rceil.$$

Since H was arbitrary, (2.6) is now a subset-homogeneous estimate for the centered residual

$$Y_x := X_x - C \max\{\Phi_\mu(x), r_0\}$$

with complexity $\sigma(x)$. Applying Lemma 2.1 a second time, now with $Z_x = Y_x$, $d_0(x) = \sigma(x)$, $R = R_*$, and

$$\psi(s; \delta) := Cs \sqrt{\log\left(\frac{eM_\Phi}{\delta}\right)},$$

gives, with probability at least $1 - \delta$,

$$\forall x \in T : \quad Y_x \leq C \max\{\sigma(x), s_0\} \sqrt{\log\left(\frac{eM_\Phi M_\sigma}{\delta}\right)},$$

where

$$M_\sigma := 1 + \lceil \log_2(R_*/s_0) \rceil.$$

Combining the last display with the definition of Y_x proves (1.4), since $M_{r_0, s_0} = M_\Phi M_\sigma$. The absolute-value estimate (1.5) follows by applying the same argument to $-X$ and taking a union bound. $\square$

Proof of Proposition 1.2. The upper bound follows from Lemma 2.3 with $H = T$, followed by optimization over μ . For the lower bound, choose x_* with $\sigma(x_*) = R_*$. Since $X_{x_0} = 0$,

$$\sup_{x \in T} X_x \geq (X_{x_*})_+, \quad \mathbb{E}(X_{x_*})_+ = \frac{R_*}{\sqrt{2\pi}}.$$

It remains to lower-bound the optimized local functional. Let

$$\Phi_\mu^{\text{full}}(x) := \int_0^{2R_*} \sqrt{\log \frac{1}{\mu(B_d(x, \varepsilon))}} d\varepsilon.$$

The classical majorizing-measure theorem gives

$$\inf_{\mu \in \Delta(T)} \sup_x \Phi_\mu^{\text{full}}(x) \lesssim \mathbb{E} \sup_{x \in T} X_x.$$

Since $\Phi_\mu(x) \leq \Phi_\mu^{\text{full}}(x)$ for every x , we have

$$\inf_\mu \Phi_{\mu,*} \leq \inf_\mu \sup_x \Phi_\mu^{\text{full}}(x) \lesssim \mathbb{E} \sup_{x \in T} X_x.$$

Together with the radius lower bound this proves the left side of (1.6). The absolute-value statement follows from

$$\sup_x X_x \leq \sup_x |X_x| \leq \sup_x X_x + \sup_x (-X_x)$$

and symmetry. $\square$

Proof of Corollary 1.3. Let

$$Y := \sup_{x \in T} \frac{(|X_x| - C \max\{\Phi_\mu(x), r_0\})_+}{\max\{\sigma(x), s_0\}}.$$

By the two-sided form of Theorem 1.1, after increasing C if necessary,

$$\mathbb{P}\left(Y > C \sqrt{\log\left(\frac{eM_{r_0, s_0}}{\delta}\right)}\right) \leq \delta, \quad \delta \in (0, 1).$$

Integrating this tail gives

$$\mathbb{E}Y \leq C \sqrt{\log(eM_{r_0, s_0})},$$

which proves (1.8). The pointwise bound (1.9) follows by dropping the supremum, and the global bound (1.10) follows by taking $r_0 \geq \Phi_{\mu,*}$, $s_0 = R_*$, and then letting $r_0 \downarrow \Phi_{\mu,*}$. Optimizing over μ and applying Proposition 1.2 proves (1.11). $\square$

3 Proofs and Consequences of the Bayesian algorithmic lower bounds

3.1 Proof of the Bayesian algorithmic lower and upper bounds

Proof of Proposition 1.4. The result is the Gaussian-process and KL specialization of the interactive Fano method of [Chen et al. \(2024, Theorem 2\)](#); we give a self-contained proof. Fix $\hat{t}$, Q , and Δ , and let

$$E := \{(t, Y) : d(t, \hat{t}(Y)) < \Delta\}.$$

Under the true joint law $P := \int \pi(dt) P_t(dY)$, write $p = P(E)$. Under the ghost law $P_0 := \pi \otimes Q$, write

$$q = P_0(E) = \rho_{\Delta, Q}(\hat{t}).$$

By data processing,

$$\mathcal{I}_\pi(Q) = D_{\text{KL}}(P \| P_0) \geq D_{\text{KL}}(\text{Bern}(p) \| \text{Bern}(q)).$$

The elementary inequality

$$D_{\text{KL}}(\text{Bern}(p) \| \text{Bern}(q)) \geq p \log \frac{1}{q} - \log 2$$

gives

$$p \leq \frac{\mathcal{I}_\pi(Q) + \log 2}{\log(1/q)}.$$

Therefore

$$P\{d(t, \hat{t}(Y)) \geq \Delta\} = 1 - p \geq 1 - \frac{\mathcal{I}_\pi(Q) + \log 2}{\log(1/\rho_{\Delta, Q}(\hat{t}))},$$

with the positive part inserted to make the bound nonnegative. Multiplying by Δ and optimizing over Δ, Q proves (1.13). $\square$

Proof of Theorem 1.5. For the Gaussian location experiment and the reference law $Q_{\tau, m}$,

$$\mathcal{I}_\pi(Q_{\tau, m}) = \int_T D_{\text{KL}}(N(v_t, \tau^2 I) \| N(m, \tau^2 I)) \pi(dt) = \frac{1}{2\tau^2} \int_T \|v_t - m\|^2 \pi(dt) = \mathcal{I}_{\pi, m}.$$

Applying Proposition 1.4 to $\hat{\Theta}$ and $Q_{\tau, m}$ gives (1.16) under condition (1.15).

It remains to relate this Bayes location error to a Gaussian range. By the comparison-decoder condition (1.14),

$$\|Y - v_{\hat{\Theta}}\|^2 - \|Y - v_{\Theta}\|^2 \leq \Omega(\Theta) - \Omega(\hat{\Theta}) \leq 0.$$

Since $Y = v_{\Theta} + \tau Z$, expanding and cancelling $\tau^2 \|Z\|^2$ gives

$$\|v_{\Theta} - v_{\hat{\Theta}}\|^2 + 2\tau \langle Z, v_{\Theta} - v_{\hat{\Theta}} \rangle \leq 0,$$

and hence

$$\|v_{\Theta} - v_{\hat{\Theta}}\|^2 \leq 2\tau \langle Z, v_{\hat{\Theta}} - v_{\Theta} \rangle = 2\tau (X_{\hat{\Theta}} - X_{\Theta}).$$

Taking expectations and using Jensen's inequality,

$$c^2 \Delta^2 \leq \left(\mathbb{E} d(\Theta, \hat{\Theta}) \right)^2 \leq \mathbb{E} d(\Theta, \hat{\Theta})^2 \leq 2\tau \mathbb{E} [X_{\hat{\Theta}} - X_{\Theta}].$$

This proves (1.17). $\square$

Proof of Corollary 1.6. By the comparison condition (1.18) and the identity $Y = v_\Theta + \tau Z$,

$$\|v_\Theta + \tau Z - v_{\hat{\Theta}}\|^2 \leq \|\tau Z\|^2.$$

Expanding the left-hand side and cancelling the common term $\tau^2\|Z\|^2$, we obtain the pathwise inequality

$$d(\Theta, \hat{\Theta})^2 = \|v_\Theta - v_{\hat{\Theta}}\|^2 \leq 2\tau\langle Z, v_{\hat{\Theta}} - v_\Theta \rangle = 2\tau(X_{\hat{\Theta}} - X_\Theta).$$

On the event in (1.19), which holds with probability at least $1 - \delta$ and is uniform over all $t \in T$,

$$X_{\hat{\Theta}} - X_\Theta \leq |X_{\hat{\Theta}}| + |X_\Theta| \leq A(\hat{\Theta}) + A(\Theta).$$

Hence, on the same event,

$$d(\Theta, \hat{\Theta})^2 \leq 2\tau(A(\Theta) + A(\hat{\Theta})),$$

which proves (1.20).

For the expectation bound, let E denote the event $\{\forall t \in T : |X_t| \leq A(t)\}$. On E , the preceding display gives

$$d(\Theta, \hat{\Theta}) \leq \sqrt{2\tau(A(\Theta) + A(\hat{\Theta}))}.$$

On E^c , we use the trivial bound

$$d(\Theta, \hat{\Theta}) \leq D_T.$$

Therefore

$$\begin{aligned} \mathbb{E}d(\Theta, \hat{\Theta}) &\leq \mathbb{E}\left[\sqrt{2\tau(A(\Theta) + A(\hat{\Theta}))} \mathbf{1}_E\right] + D_T \mathbb{P}(E^c) \\ &\leq \mathbb{E}\sqrt{2\tau(A(\Theta) + A(\hat{\Theta}))} + \delta D_T, \end{aligned}$$

which proves (1.21). The final claim follows by applying the absolute pointwise Gaussian envelope (1.5) to the anchored process $(X_t)_{t \in T \cup \{t_0\}}$ and then restricting back to T . $\square$

3.2 An algorithmic relaxation route to Sudakov-type comparison

By replacing the exact ghost mass with a class-wide small-ball mass and optimizing this small-ball term over hard priors in Theorem 1.5, we obtain a nearest-neighbor route to a Sudakov-type comparison for priors localized at the tested radius.

Corollary 3.1 (Sudakov-type comparison through supremum relaxation). *Under the setup of Theorem 1.5 with the ordinary nearest-neighbor decoder $\Omega \equiv 0$ and the centered reference choice $m = \bar{v}_\pi$, set*

$$q_\pi(\Delta) := \sup_{a \in T} \pi(B_d(a, \Delta)), \quad L_\pi(\Delta) := \log \frac{1}{q_\pi(\Delta)}.$$

Since $\rho_{\Delta, Q_\tau, \bar{v}_\pi}(\hat{\Theta}_{\text{nn}}) \leq q_\pi(\Delta)$, if $L_\pi(\Delta)$ is larger than a universal constant, one may choose $\tau^2 \asymp V_\pi/L_\pi(\Delta)$ so that (1.15) holds with an absolute $c \in (0, 1)$. Relaxing the nearest-neighbor lower bound (1.17) to supremum over the Gaussian process yields the variance-normalized single-radius obstruction

$$\mathbb{E} \sup_{s, u \in T} (X_s - X_u) \gtrsim \Delta^2 \sqrt{\frac{L_\pi(\Delta)}{V_\pi}}. \quad (3.1)$$

In particular, if the hard prior is localized at scale Δ , meaning $V_\pi \leq C_0 \Delta^2$, then

$$\mathbb{E} \sup_{s, u \in T} (X_s - X_u) \gtrsim_{C_0} \Delta \sqrt{L_\pi(\Delta)}. \quad (3.2)$$

Equivalently, with

$$N_{\text{spread}}^{\text{loc}}(T, d, \Delta; C_0) := \sup_{\pi: V_\pi \leq C_0 \Delta^2} \frac{1}{\sup_{a \in T} \pi(B_d(a, \Delta))}, \quad (3.3)$$

one obtains, whenever the logarithm is above the universal threshold required by the information condition,

$$\mathbb{E} \sup_{s, u \in T} (X_s - X_u) \gtrsim_{C_0} \Delta \sqrt{\log N_{\text{spread}}^{\text{loc}}(T, d, \Delta; C_0)}. \quad (3.4)$$

Proof of Corollary 3.1. For the variance-normalized small-ball statement, use the centered reference choice $m = \bar{v}_\pi$, for which $\mathcal{I}_{\pi, m} = V_\pi / (4\tau^2)$, and the bound $\rho_{\Delta, Q_\tau, \bar{v}_\pi}(\hat{\Theta}_{\text{nn}}) \leq q_\pi(\Delta)$. If $L_\pi(\Delta)$ is above a universal constant, choosing $\tau^2 \asymp V_\pi / L_\pi(\Delta)$ with a sufficiently large absolute constant makes (1.15) hold for some absolute $c \in (0, 1)$. Since $X_{\hat{\Theta}_{\text{nn}}} - X_\Theta \leq \sup_{s, u \in T} (X_s - X_u)$ pointwise in Z , (1.17) gives

$$\mathbb{E} \sup_{s, u \in T} (X_s - X_u) \gtrsim \Delta^2 \sqrt{\frac{L_\pi(\Delta)}{V_\pi}}.$$

If $V_\pi \leq C_0 \Delta^2$, this becomes $\Delta \sqrt{L_\pi(\Delta)}$, with a constant depending only on C_0 . Optimizing over such localized priors gives (3.4). $\square$

The preceding corollary directly recovers a Sudakov-scale lower bound under a natural localized hard-prior condition. This is the precise mild condition needed by the nearest-neighbor route: the prior used to certify small balls must also have second moment comparable to the tested radius.

Corollary 3.2 (Localized packing form). *Let $S \subset T$ be 2Δ -separated and assume that, for some $t_0 \in T$ and $C_0 < \infty$,*

$$S \subset B_d(t_0, C_0 \Delta).$$

If $|S| = M \geq M_0$, where M_0 is a universal constant, then

$$\mathbb{E} \sup_{s, u \in T} (X_s - X_u) \gtrsim_{C_0} \Delta \sqrt{\log M}.$$

Proof of Corollary 3.2. Apply Theorem 1.5 with π equal to the uniform distribution on S . Since S is 2Δ -separated, every Δ -ball contains at most one point of S , so

$$q_\pi(\Delta) = \sup_{a \in T} \pi(B_d(a, \Delta)) \leq M^{-1}, \quad L_\pi(\Delta) \geq \log M.$$

The localization assumption gives

$$V_\pi = \iint d(s, t)^2 \pi(ds) \pi(dt) \leq 4C_0^2 \Delta^2.$$

Equation (3.2) yields the claim. $\square$

For comparison with classical covering notation, we record the finite-set fractional-covering duality used in Chen et al. (2024). Recall that the fractional covering number is equivalent to the canonical covering number, up to absolute-constant changes in the covering radius.

Lemma 3.3 (Fractional-covering duality). *For a finite metric space (T, d) and any $\Delta > 0$,*

$$N_{\text{frac}}(T, d, \Delta) := \inf_{\mu \in \Delta(T)} \sup_{x \in T} \frac{1}{\mu(B_d(x, \Delta))} = \sup_{\pi \in \Delta(T)} \frac{1}{\sup_{a \in T} \pi(B_d(a, \Delta))}. \quad (3.5)$$

Proof of Lemma 3.3. Let

$$r_* := \sup_{\mu \in \Delta(T)} \inf_{x \in T} \mu(B_d(x, \Delta)).$$

Then

$$N_{\text{frac}}(T, d, \Delta) = \frac{1}{r_*}.$$

Since T is finite, von Neumann's minimax theorem gives

$$\begin{aligned} r_* &= \sup_{\mu \in \Delta(T)} \inf_{x \in T} \sum_{a \in T} \mu(a) \mathbf{1}\{a \in B_d(x, \Delta)\} \\ &= \inf_{\pi \in \Delta(T)} \sup_{\mu \in \Delta(T)} \sum_{x, a \in T} \pi(x) \mu(a) \mathbf{1}\{a \in B_d(x, \Delta)\} \\ &= \inf_{\pi \in \Delta(T)} \sup_{a \in T} \pi(B_d(a, \Delta)). \end{aligned}$$

Taking reciprocals yields (3.5). $\square$

The equality (3.5) is an entropy duality. It optimizes the small-ball term, but it does not remove the factor V_π in (3.1), which comes from the information radius of the Gaussian location channel. There is therefore no contradiction between the fractional-covering lower bounds of [Chen et al. \(2024\)](#) and the localization requirement above: the fractional-covering duality identifies the optimal hard prior for the small-ball obstruction, while the nearest-neighbor Gaussian comparison also requires that this hard prior have controlled second moment at scale Δ . Thus the comparison directly gives the localized obstruction (3.4) and the localized packing form in Corollary 3.2.

The unrestricted classical Sudakov minoration is a separate Gaussian comparison theorem, not a consequence of linear-programming duality alone. Standard proofs use nontrivial Gaussian comparison and convex-geometric duality arguments; see, for example, the expositions of [Pollard \(2001\)](#) and [Ledoux and Talagrand \(1991, Chapter 3\)](#). The full lower half of the Fernique–Talagrand majorizing-measure theorem is stronger and multiscale ([Talagrand, 2005](#)). In the present paper we therefore keep the roles separate: the Bayesian algorithmic envelope supplies a one-scale, decision-theoretic, and prior-agnostic primitive, while the unrestricted Sudakov and generic-chaining lower bounds require the classical Gaussian comparison or multiscale localization machinery.

4 A separating example: overparameterized class and pointwise certification

This section proves Theorem 1.7. The example is a finite-dimensional heterogeneous-orthogonal-basis construction. It is intentionally elementary, but it captures the benchmark separation needed for the paper: the full ambient class is overparameterized, the correct subatlas is decision-dependent, and the actual estimator is unrestricted over the full class but uses a minimum-norm inductive bias. The proof keeps the discussion in separate paragraphs because each paragraph isolates one benchmark in the separation.

Proof of Theorem 1.7. Let $M, N \geq 2$ satisfy $\log N \geq 1024 \log M$. Let

$$g_0, e_1, \dots, e_M, f_1, \dots, f_N$$

be orthonormal vectors in $\mathbb{R}^{M+N+1}$, and define

$$\tau := \frac{R}{\sqrt{\log M}}, \quad S := \frac{1}{8}\tau\sqrt{\log N} = \frac{R}{8}\sqrt{\frac{\log N}{\log M}}.$$

The assumption on M, N implies $S \geq 4R$. Set

$$H_M := \{Rg_0, Re_1, \dots, Re_M\}, \quad C_N := \{Sf_1, \dots, Sf_N\}, \quad T_{M,N} := H_M \cup C_N,$$

with Euclidean metric $d(s, t) = \|s - t\|_2$. The prior is

$$\pi(Rg_0) = \frac{1}{2}, \quad \pi(Re_i) = \frac{1}{2M}, \quad i = 1, \dots, M, \quad \pi(Sf_j) = 0, \quad j = 1, \dots, N,$$

and we take $\Delta = R/3$. Since $S \geq 4R$, every Δ -ball in $T_{M,N}$ is a singleton. Hence

$$q_\pi(\Delta) = \sup_{a \in T_{M,N}} \pi(B_d(a, \Delta)) = \frac{1}{2}.$$

This proves the fixed-prior small-ball claim in part (i).

Algorithmic ghost mass versus worst-case Fano mass. Let

$$Y = \Theta + \tau Z, \quad \Theta \sim \pi, \quad Z \sim N(0, I_{M+N+1}),$$

and use the ghost/reference law $Q_\tau = N(0, \tau^2 I_{M+N+1})$. Under Q_τ , the scores of the $M+1$ lower-norm points in H_M are exchangeable. We first show that the cloud is selected with negligible ghost probability. If a cloud point Sf_j beats the hub point Rg_0 , then

$$2\langle Y, Sf_j \rangle - (1 + \lambda)S^2 \geq 2\langle Y, Rg_0 \rangle - (1 + \lambda)R^2.$$

Under Q_τ , the random part of the difference is centered Gaussian with variance at most $4\tau^2(S^2 + R^2)$. Since $S \geq 4R$ and $\lambda = 64$, a Gaussian union bound gives

$$Q_\tau(\hat{\Theta}_\lambda \in C_N) \leq N^{-2} \tag{4.1}$$

for all sufficiently large N . Conditional on $\hat{\Theta}_\lambda \in H_M$, exchangeability gives

$$Q_\tau(\hat{\Theta}_\lambda = Rg_0 \mid \hat{\Theta}_\lambda \in H_M) = \frac{1}{M+1}.$$

Consequently,

$$Q_\tau(\hat{\Theta}_\lambda = Rg_0) \leq \frac{1}{M+1} + N^{-2}.$$

Since every Δ -ball is a singleton and π vanishes on the cloud,

$$\begin{aligned} \rho_{\Delta, Q_\tau}(\hat{\Theta}_\lambda) &= \mathbb{E}_{Y \sim Q_\tau} \pi(B_d(\hat{\Theta}_\lambda(Y), \Delta)) \\ &\leq \frac{1}{2}Q_\tau(\hat{\Theta}_\lambda = Rg_0) + \frac{1}{2M}Q_\tau(\hat{\Theta}_\lambda \in \{Re_1, \dots, Re_M\}) \leq \frac{2}{M} \end{aligned}$$

for all large M, N . Hence

$$\log \frac{1}{\rho_{\Delta, Q_\tau}(\hat{\Theta}_\lambda)} \geq \log M - O(1),$$

whereas the worst-case relaxation for the same prior gives only $\log 2$. This proves part (ii) and the entropy separation in part (i).

Sharp algorithmic lower bound for the actual unrestricted estimator. The information radius under Q_τ is

$$I_{\pi, Q_\tau} = \mathbb{E}_{\Theta \sim \pi} D_{\text{KL}}(N(\Theta, \tau^2 I) \| N(0, \tau^2 I)) = \frac{\mathbb{E}_\pi \|\Theta\|^2}{2\tau^2} = \frac{R^2}{2\tau^2} = \frac{1}{2} \log M.$$

Combining this with the ghost-mass bound, for all large M ,

$$I_{\pi, Q_\tau} + \log 2 \leq \frac{3}{4} \log \frac{1}{\rho_{\Delta, Q_\tau}(\hat{\Theta}_\lambda)}.$$

Proposition 1.4 therefore yields

$$\mathbb{E}_{\Theta, Y} d(\Theta, \hat{\Theta}_\lambda(Y)) \geq c\Delta \asymp R. \quad (4.2)$$

This lower bound is sharp for the actual Bayesian decision problem. When $\Theta \in H_M$, all true parameters have norm R . The same score comparison as above, now under $Y = \Theta + \tau Z$, gives

$$\mathbb{P}_{\Theta, Y}(\hat{\Theta}_\lambda \in C_N) \leq N^{-2}.$$

On the complementary event, both Θ and $\hat{\Theta}_\lambda$ lie in H_M , whose diameter is at most $\sqrt{2}R$. Hence

$$\mathbb{E}_{\Theta, Y} d(\Theta, \hat{\Theta}_\lambda(Y)) \leq \sqrt{2}R + N^{-2}S \lesssim R. \quad (4.3)$$

Combining the algorithmic lower bound (4.2) and the algorithmic upper bound (4.3) proves the Bayes risk statement for the estimator $\hat{\Theta}_\lambda$:

$$E_{\Theta, Y} d(\Theta, \hat{\Theta}_\lambda(Y)) \asymp R.$$

This proves the statement about the Bayes risk of the actual estimator in part (iii).

Bayes-optimal risk is of the same order. The Bayes-optimal risk for this prior is also $\asymp R$, despite the oracle estimator being highly non-analytical. The upper bound follows from (4.3). For the lower bound, condition on the event $\mathcal{L}$ that Θ is one of the leaves. Under this conditional experiment, write

$$J \sim \text{Unif}\{1, \dots, M\}, \quad \Theta = Re_J, \quad Y = Re_J + \tau Z, \quad Z \sim N(0, I).$$

Given any estimator $\hat{\theta}(Y)$, define the induced leaf classifier

$$\hat{J}(Y) \in \arg \min_{1 \leq j \leq M} \|\hat{\theta}(Y) - Re_j\|,$$

with ties broken arbitrarily. Since the leaves are separated by $\sqrt{2}R$, the event $\|\hat{\theta}(Y) - Re_J\| < R/2$ implies $\hat{J} = J$. Hence

$$\mathbb{E} \left[\|\hat{\theta}(Y) - \Theta\| \mid \mathcal{L} \right] \geq \frac{R}{2} \mathbb{P}(\hat{J} \neq J \mid \mathcal{L}).$$

Let $P_j = N(Re_j, \tau^2 I)$, let $\bar{P} = M^{-1} \sum_{j=1}^M P_j$, and take the reference law $Q = N(0, \tau^2 I)$. Then

$$I(J; Y) = \frac{1}{M} \sum_{j=1}^M D_{\text{KL}}(P_j \| \bar{P}) \leq \frac{1}{M} \sum_{j=1}^M D_{\text{KL}}(P_j \| Q) = \frac{R^2}{2\tau^2} = \frac{1}{2} \log M,$$

where the last equality uses $\tau = R/\sqrt{\log M}$. Fano's inequality therefore gives

$$\mathbb{P}(\hat{J} \neq J \mid \mathcal{L}) \geq 1 - \frac{I(J; Y) + \log 2}{\log M} \geq c_0$$

for a universal constant $c_0 > 0$, for all sufficiently large M . Since $\mathbb{P}(\mathcal{L}) = 1/2$ under the original prior,

$$\mathbb{E}\|\hat{\theta}(Y) - \Theta\| \geq \frac{1}{2} \mathbb{E}\left[\|\hat{\theta}(Y) - \Theta\| \mid \mathcal{L}\right] \geq cR.$$

Taking the infimum over $\hat{\theta}$ gives the Bayes-risk lower bound cR . This proves the statement about the Bayes-optimal risk in part (iii). Thus the class-wide Fano relaxation below is vacuous only as a certificate; it does not reflect an easy Bayes problem.

Usual Fano relaxation is vacuous for the fixed prior. Replacing the exact ghost mass by the worst-case relaxation $q_\pi(\Delta) = 1/2$ gives

$$\Delta \left[1 - \frac{\mathfrak{l}_{\pi, Q_\tau} + \log 2}{\log(1/q_\pi(\Delta))} \right]_+ = \Delta \left[1 - \frac{\mathfrak{l}_{\pi, Q_\tau} + \log 2}{\log 2} \right]_+ = 0.$$

Thus, for the same prior, channel, and estimator, the usual fixed-prior Fano relaxation is vacuous, while the exact algorithmic ghost mass gives a sharp Bayes estimation lower bound.

Full-class minimax risk has arbitrarily large gap. Let

$$\mathfrak{R}_{\text{mm}}(T_{M,N}, \tau) := \inf_{\hat{\theta}} \sup_{\theta \in T_{M,N}} \mathbb{E}_\theta d(\theta, \hat{\theta}(Y)),$$

where the infimum is over all measurable estimators with values in the ambient Euclidean space. The trivial estimator $\hat{\theta} \equiv 0$ gives $\mathfrak{R}_{\text{mm}}(T_{M,N}, \tau) \leq S$. Conversely, restrict the parameter to the cloud C_N and put the uniform prior on C_N . For any estimator $\hat{\theta}$, define

$$\hat{J}(\hat{\theta}) \in \arg \min_{1 \leq j \leq N} \|\hat{\theta} - Sf_j\|_2.$$

If $d(Sf_j, \hat{\theta}) < S/2$, then $\hat{J}(\hat{\theta}) = j$, since the balls $B(Sf_j, S/2)$ are disjoint. Therefore

$$\mathbb{E}_J d(Sf_J, \hat{\theta}(Y)) \geq \frac{S}{2} \mathbb{P}(\hat{J} \neq J).$$

By Fano's inequality with reference law $N(0, \tau^2 I_{M+N+1})$,

$$I(J; Y) \leq \frac{S^2}{2\tau^2} = \frac{1}{128} \log N.$$

Thus, for all sufficiently large N ,

$$\mathbb{P}(\hat{J} \neq J) \geq 1 - \frac{I(J; Y) + \log 2}{\log N} \geq \frac{1}{2}.$$

It follows that

$$\mathfrak{R}_{\text{mm}}(T_{M,N}, \tau) \geq cS.$$

Together with the trivial upper bound, this gives $\mathfrak{R}_{\text{mm}}(T_{M,N}, \tau) \asymp S$, proving part (iv). Since $S/R = (1/8)\sqrt{\log N/\log M}$, the full-class minimax risk is much larger than the Bayes risk whenever $\log N/\log M \rightarrow \infty$. The minimax scale is governed by an ambient cloud that is irrelevant to the specified Bayesian decision problem.

Algorithmically induced Gaussian certificate matches the selected subatlas. Now define

$$X_t := \langle Z, t \rangle, \quad t \in T_{M,N},$$

using the same standard Gaussian vector Z as in the location experiment. We verify the comparison condition in Theorem 1.5. The decoder minimizes $\|Y - a\|^2 + \lambda\|a\|^2$, so (1.14) holds with $\Omega(t) = \lambda\|t\|^2$ once $\Omega(\hat{\Theta}_\lambda) \geq \Omega(\Theta)$ is checked. But $\Theta \in H_M$ almost surely, so $\|\Theta\| = R$, whereas every output of $\hat{\Theta}_\lambda$ has norm either R or $S \geq R$. Hence $\Omega(\hat{\Theta}_\lambda) \geq \Omega(\Theta)$ almost surely. Applying Theorem 1.5 gives

$$\mathcal{G}_{\text{alg}} := \mathbb{E}_{\Theta, Y} [X_{\hat{\Theta}_\lambda(Y)} - X_\Theta] \gtrsim R\sqrt{\log M}.$$

Conversely, on the event $\hat{\Theta}_\lambda \in H_M$, both Θ and $\hat{\Theta}_\lambda$ lie in H_M , and

$$\mathbb{E} \sup_{s, u \in H_M} (X_s - X_u) \asymp R\sqrt{\log M}.$$

The event $\hat{\Theta}_\lambda \in C_N$ has probability at most N^{-2} under the true prior. By Cauchy–Schwarz and the standard Gaussian maximal bound on $T_{M,N}$, its contribution is negligible. Therefore

$$\mathcal{G}_{\text{alg}} \asymp \mathbb{E} \sup_{s, u \in H_M} (X_s - X_u) \asymp R\sqrt{\log M},$$

which proves the first assertion in part (v).

This also matches the pointwise Gaussian upper envelope on the selected subatlas. By (4.1), we already have

$$Q_\tau\{\hat{\Theta}_\lambda(Y) \in C_N\} \leq N^{-2}.$$

for all sufficiently large N . Let

$$\nu_Q^H(a) := Q_\tau\{\hat{\Theta}_\lambda(Y) = a \mid \hat{\Theta}_\lambda(Y) \in H_M\}, \quad a \in H_M,$$

be the ghost selection law conditioned on the lower-norm subatlas. By exchangeability, ν_Q^H is uniform on H_M . Since the restricted process on H_M consists of $M + 1$ Gaussian variables of variance at most R^2 , Theorem 1.1 gives, up to universal constants and confidence logarithms,

$$\mathbb{P}\left(\forall t \in H_M : |X_t| \leq CR\sqrt{\log \frac{M}{\delta}}\right) \geq 1 - \delta.$$

This proves the pointwise-envelope assertion in part (v).

Global Gaussian supremum has arbitrarily large gap. The full ambient Gaussian field has a larger global scale because of the cloud. Restricting only to C_N ,

$$\mathbb{E} \sup_{s, u \in T_{M,N}} (X_s - X_u) \geq \mathbb{E} \max_{1 \leq j \leq N} X_{Sf_j} = S \mathbb{E} \max_{1 \leq j \leq N} Z_j \gtrsim S\sqrt{\log N}.$$

Using the definition of S ,

$$S\sqrt{\log N} = \frac{1}{8}R \frac{\log N}{\sqrt{\log M}}.$$

This proves part (vi). Comparing with the decision-aligned scale $R\sqrt{\log M}$, the gap is of order $\log N / \log M$, and taking $N = M^K$ makes the ratio of order K . □

Summary and interpretation. The construction gives a fully explicit sequence of finite-dimensional Gaussian location problems for which

- fixed-prior Fano relaxation
- $\ll$ algorithmic ghost-mass lower envelope
- $\asymp$ pointwise upper envelope on the selected subatlas
- $\ll$ full-class minimax/global Gaussian scale.

The example does not claim that minimax theory is wrong or fundamentally insufficient. Rather, it shows that a full-class minimax benchmark can be the wrong certificate for a pointwise decision problem, while a restricted minimax benchmark can be oracle-dependent. The norm-regularized decoder uses a fixed numerical penalty λ , independent of the prior π and the problem parameters M, N, R, S, τ , while identifying the hub-leaf subatlas H_M as the relevant subatlas is precisely the decision-dependent information revealed by the ghost image $Y \mapsto \hat{\Theta}_\lambda(Y)$. The algorithmic ghost-mass lower bound supplies the missing certificate: it is tied to the fixed prior, the actual estimator, and the subgeometry selected by that estimator. This is the finite-dimensional analogue of the pointwise-complexity viewpoint in deep networks, where the effective subatlas is induced by the learned representation and compressed spectrum rather than specified as a fixed restricted class in advance.

Here we keep the cleanest finite formulation. Finite sets are already the standard setting for Sudakov, Fano, and packing lower bounds, and they make the quantifiers transparent: the full ambient class, the oracle restricted subatlas, the fixed prior, and the actual estimator are all explicit. Extensions to continuous settings, as well as computational-eciency questions concerning the full class versus selected subatlases, are left to future work.

5 Finite-Cutoff Renormalization and Pointwise Dimension

This section makes precise the analogy between finite-cutoff renormalization and nonlinear feature learning in [Li and Xu \(2026\)](#). The analogy does not rest merely on the correspondence between composite layers and RG levels. More essentially, both settings exhibit a common *feature/operator inner-product* structure within each composite layer of a fully connected DNN, or within each exponential-family RG level. This correspondence leads to the same proof mechanism: exact finite-step telescoping, a learned or renormalized secant Gram matrix, and a hierarchical prior over local charts and a global atlas.

This is the mechanism that makes the DNN pointwise theory computable, and it is the same mechanism that can be used in a finite-cutoff nonlinear RG analysis. The language is compatible with rigorous Wilsonian RG frameworks such as [Brydges and Slade \(2015\)](#); [Bauerschmidt et al. \(2019\)](#), but the theorem below is an abstract finite-dimensional statement.

Its practical significance in statistical field theory is that RG is fundamentally about controlling coarse marginals after integrating out fluctuations. For globally non-log-concave and anisotropic interacting fields, direct scalar Lipschitz bounds or global convex-geometric arguments are often too crude. The finite-cutoff viewpoint instead allows one to work on stable local charts where the effective action has a curvature lower bound: directions below the finite resolution are absorbed as approximation error, while global nonconvexity is paid for through an atlas cost. We formalize this mechanism as an abstract deterministic complexity theorem and illustrate it through a chartwise nonconvex phase-atlas example, as discussed at the end of [Section 5.2](#).

Deep-network pointwise theory	Finite-cutoff RG analogue
Layer index ℓ	RG level index j
Weights W_ℓ	Level coupling vector g_j
Feature matrix $F_{\ell-1}(W, X)$	Renormalized operator family $O_j(S_j)$
Feature Gram $F_{\ell-1}^\top F_{\ell-1}$	Secant response Gram Γ_j^{sec}
Exact layer telescoping	Exact replacement over RG levels
Outer Lipschitz $M_{\ell \rightarrow L}$	Stability of later RG maps $M_{j \rightarrow J}$
Active feature eigenspace	Relevant/marginal active operator subspace
Grassmannian prior	Phase/subspace atlas prior
Feature-rank compression	Contraction or truncation of irrelevant directions
Kernel/NTK baseline	Free or massive GFF baseline

Limiting scope relative to constructive QFT, RG approximation, and gauge fields.

All statements below are finite-dimensional and finite-cutoff: they provide a structural complexity layer, not a replacement for model-specific constructive-QFT estimates. In the four-dimensional Ising and lattice Φ_4^4 setting, [Aizenman and Duminil-Copin \(2021\)](#) prove marginal triviality by controlling deviations from Wick’s rule through random currents and multiscale estimates; the issue is not amplitude control, since Ising spins are already uniformly bounded, but Gaussianization of macroscopic observables through decay of connected non-Gaussian remainders. Regarding the technical difficulty of Gaussianization, while universal approximation theorems ([Cybenko, 1989](#); [Hornik et al., 1989](#)) can approximate a finite-dimensional RG map on a compact set at fixed cutoff, they do not provide scale-uniform error bounds, preservation of locality or reflection positivity, or control of Wick-factorization errors. Such conclusions require model-specific RG or correlation estimates. For Yang–Mills, an analogous finite-cutoff program would have to be formulated for lattice gauge fields and gauge-invariant observables such as plaquettes, Wilson loops, and character observables; a mass-gap theorem would still require continuum construction, Osterwalder–Schrader positivity and reconstruction, and uniform exponential clustering of gauge-invariant correlations ([Jaffe and Witten, 2000](#); [Chatterjee, 2018](#)). Thus scaling-limit Gaussianity, nontrivial continuum interaction, and gauge-theoretic mass gaps remain separate model-specific problems.

5.1 Pointwise complexity for nonlinear RG maps

Exact finite-step telescoping for multi-level RG maps. Let $\mathcal{S}_0, \dots, \mathcal{S}_J$ be finite-dimensional normed spaces. For each RG level j , let

$$\mathcal{R}_j^{g_j} : \mathcal{S}_j \rightarrow \mathcal{S}_{j+1}, \quad g_j \in B_2(R_j) \subset \mathbb{R}^{p_j},$$

be a nonlinear coarse-graining map. Starting from a fixed $S_0 \in \mathcal{S}_0$, define the RG trajectory

$$S_{j+1}(g) := \mathcal{R}_j^{g_j}(S_j(g)), \quad S_J(g) = \mathcal{R}_{J-1}^{g_{J-1}} \circ \dots \circ \mathcal{R}_0^{g_0}(S_0).$$

This notation covers, for example, the exact Wilsonian identity

$$e^{-S_{j+1}(\Phi)} = \int e^{-S_j(\Phi + \psi_j)} d\gamma_j(\psi_j).$$

This is the finite-dimensional analogue of integrating out one fluctuation field at one RG level. Perturbative RG expands this map; here we keep the finite map itself.

For two coupling sequences g, g' , define the hybrid trajectory

$$S_j^{(g', g; m)} := \mathcal{R}_{j-1}^{g'_{j-1}} \circ \dots \circ \mathcal{R}_m^{g'_m} \circ \mathcal{R}_{m-1}^{g_{m-1}} \circ \dots \circ \mathcal{R}_0^{g_0}(S_0),$$

with the evident convention at the endpoints. Then the following identity is exact.

Lemma 5.1 (Exact RG replacement telescoping). *For every g, g' ,*

$$S_J(g') - S_J(g) = \sum_{j=0}^{J-1} \left[\mathcal{R}_{J-1:j+1}^{g'}(\mathcal{R}_j^{g'_j}(S_j^{(j)})) - \mathcal{R}_{J-1:j+1}^{g'}(\mathcal{R}_j^{g_j}(S_j^{(j)})) \right], \quad (5.1)$$

where $S_j^{(j)} := \mathcal{R}_{j-1}^{g_{j-1}} \circ \dots \circ \mathcal{R}_0^{g_0}(S_0)$ and $\mathcal{R}_{J-1:j+1}^{g'} := \mathcal{R}_{J-1}^{g'_{J-1}} \circ \dots \circ \mathcal{R}_{j+1}^{g'_{j+1}}$.

Proof. Insert the hybrid trajectories that use g up to level $j - 1$ and g' from level j onward. The difference between two consecutive hybrids changes only the scale- j map. Summing these consecutive differences gives (5.1). $\square$

Let ρ be a semimetric on $\mathcal{S}_J$. The next result is the RG analogue of the non-perturbative DNN feature expansion in Li and Xu (2026): the finite replacement identity induces an ellipsoidal metric whose matrices are finite-scale secant response Grams.

Theorem 5.2 (Secant-Gram metric domination for nonlinear RG). *Fix g and a finite neighborhood $\mathcal{N}(g, \varepsilon)$. Assume that for each j :*

(i) *the later RG flow is locally stable: for all states U, V arising in $\mathcal{N}(g, \varepsilon)$,*

$$\rho(\mathcal{R}_{J-1:j+1}^{g'}(U), \mathcal{R}_{J-1:j+1}^{g'}(V)) \leq M_{j \rightarrow J}(g, \varepsilon) \|U - V\|_{j+1};$$

(ii) *the scale- j finite replacement is dominated by a positive semidefinite secant Gram $\Gamma_j(g, \varepsilon)$:*

$$\|\mathcal{R}_j^{g'_j}(S_j^{(j)}) - \mathcal{R}_j^{g_j}(S_j^{(j)})\|_{j+1}^2 \leq (g'_j - g_j)^\top \Gamma_j(g, \varepsilon) (g'_j - g_j).$$

Then all $g' \in \mathcal{N}(g, \varepsilon)$ satisfy

$$\rho(S_J(g'), S_J(g))^2 \leq \sum_{j=0}^{J-1} (g'_j - g_j)^\top G_j(g, \varepsilon) (g'_j - g_j), \quad G_j := J M_{j \rightarrow J}(g, \varepsilon)^2 \Gamma_j(g, \varepsilon). \quad (5.2)$$

Proof. Apply Lemma 5.1, the triangle inequality for ρ , the local stability assumption, and the one-step secant-Gram bound. Cauchy-Schwarz gives a factor J when the J increments are summed; this factor is included in the definition of G_j . $\square$

How the secant Gram is verified in exponential-family RG steps. Condition (ii) in Theorem 5.2 is a finite-dimensional secant calculation. Fix the state $S_j^{(j)}$ and set

$$F_j(g_j) := \mathcal{R}_j^{g_j}(S_j^{(j)}).$$

If F_j is continuously Frechet differentiable on the finite neighborhood under consideration and the next-scale norm is Hilbertian, then for $\Delta_j = g'_j - g_j$ and $g_{j,t} = g_j + t\Delta_j$,

$$F_j(g'_j) - F_j(g_j) = \int_0^1 DF_j(g_{j,t})[\Delta_j] dt,$$

and Jensen's inequality gives

$$\|F_j(g'_j) - F_j(g_j)\|_{j+1}^2 \leq \Delta_j^\top \left(\int_0^1 DF_j(g_{j,t})^* DF_j(g_{j,t}) dt \right) \Delta_j. \quad (5.3)$$

Thus condition (ii) holds with the pairwise secant Gram in parentheses, or with any positive-semidefinite Loewner envelope that dominates it uniformly over the finite neighborhood. For an exponential-family RG step

$$\mathcal{R}_j^g(S)(\Phi) = -\log \int \exp \left\{ -S(\Phi + \psi) - \sum_{a=1}^{p_j} g_a O_{j,a}(\Phi + \psi) \right\} d\gamma_j(\psi),$$

one has

$$\partial_{g_a} \mathcal{R}_j^g(S)(\Phi) = \mathbb{E}_{j,\Phi,g} [O_{j,a}(\Phi + \psi)].$$

Consequently the secant Gram is the Gram matrix of the finite-scale response functions

$$\Gamma_{j,ab}^{\text{sec}}(g, g') = \int_0^1 \langle m_{j,a}^t, m_{j,b}^t \rangle_{j+1} dt, \quad m_{j,a}^t(\Phi) := \mathbb{E}_{j,\Phi,g,t} [O_{j,a}(\Phi + \psi)].$$

This is the precise RG analogue of the learned feature Gram matrix in the DNN pointwise metric.

Compressed response Grams. The scalar Lipschitz choice $\Gamma_j = L_j^2 I$ verifies condition (ii), but it gives no pointwise gain. A useful pointwise-complexity theorem requires an anisotropic response estimate. Let $K_{j+1 \rightarrow J}$ be a positive-semidefinite quadratic form measuring the sensitivity of the later RG flow from level $j+1$ to the terminal observable. The propagated response Gram is then

$$G_j^{\text{RG}}(g, g') := \int_0^1 DF_j(g_{j,t})^* K_{j+1 \rightarrow J} DF_j(g_{j,t}) dt. \quad (5.4)$$

Suppose that the coupling space decomposes as $\mathbb{R}^{p_j} = U_j \oplus U_j^\perp$. Let P_j be an orthonormal basis matrix for U_j . Assume that, uniformly over the finite neighborhood,

$$\|DF_j(\theta)P_j a\|_{K_{j+1 \rightarrow J}}^2 \leq a^\top A_j a \quad (a \in \mathbb{R}^{\dim U_j}), \quad \|DF_j(\theta)v\|_{K_{j+1 \rightarrow J}}^2 \leq \tau_j \|v\|^2 \quad (v \in U_j^\perp).$$

Then, for every g' in the neighborhood,

$$G_j^{\text{RG}}(g, g') \preceq 2P_j A_j P_j^\top + 2\tau_j I. \quad (5.5)$$

Indeed, write any direction $h \in \mathbb{R}^{p_j}$ as $h = P_j a + v$, with $v \in U_j^\perp$, and use $\|\xi + \zeta\|^2 \leq 2\|\xi\|^2 + 2\|\zeta\|^2$. Integrating along the secant path gives (5.5) in the Loewner order. After diagonalizing A_j , the active eigenvalues give the local ellipsoidal effective dimension, while directions with $\tau_j R_j^2 \lesssim u_j^2$ are below finite resolution and do not contribute. In perturbative RG language, U_j is the finite-resolution span of relevant and marginal operator directions, while irrelevant operators contribute only the residual term. In a nonconvex model this compression is imposed chartwise, and the phase/subspace prior below pays the cost of selecting the chart and active subspace.

Hierarchical phase/subspace prior. For a positive semidefinite matrix $G \in \mathbb{R}^{p \times p}$, a Euclidean radius R , and a resolution $u > 0$, set

$$r_{\text{eff}}(G, R, u) := \max\{k : \lambda_k(G)R^2 \geq u^2\}, \quad V_{\text{eff}}(G, R, u) := \text{span}\{\text{top } r_{\text{eff}} \text{ eigenvectors}\},$$

and

$$d_{\text{ell}}(G, R, u) := \frac{1}{2} \sum_{k=1}^{r_{\text{eff}}(G, R, u)} \log \left(\frac{CR^2 \lambda_k(G)}{u^2} \right). \quad (5.6)$$

The following elementary estimate is the local-chart calculation used throughout. For subspaces $V, \bar{V} \subset \mathbb{R}^p$, write

$$\rho_{\text{proj}, G}(V, \bar{V}) := \|G^{1/2}(P_V - P_{\bar{V}})\|_{\text{op}}.$$

Lemma 5.3 (Ellipsoidal local chart). *Let $r = r_{\text{eff}}(G, R, u)$, let $V = V_{\text{eff}}(G, R, u)$, and let $\bar{V} \in \text{Gr}(p, r)$ satisfy*

$$\rho_{\text{proj}, G}(V, \bar{V}) \leq \frac{u}{4R}.$$

Let $\pi_{\bar{V}}$ be the uniform law on $B_2(2R) \cap \bar{V}$. Then, after changing the universal numerical constant in the radius, uniformly over $g \in B_2(R)$,

$$\log \frac{1}{\pi_{\bar{V}}\{g' : (g' - g)^\top G(g' - g) \leq Cu^2\}} \leq C + d_{\text{ell}}(G, R, u).$$

Equivalently, replacing u by a constant multiple gives the displayed local-chart bound used in Theorem 5.4.

Proof. The point is that the prior may be supported on the reference subspace $\bar{V}$ even though the center g need not belong to $\bar{V}$. Since $r = r_{\text{eff}}(G, R, u)$, the component of any vector in the inactive subspace $V^\perp$ has G -length at most u on the Euclidean ball $B_2(R)$. The projection-distance assumption transfers the active subspace V to $\bar{V}$ with additional G -error at most $u/4$. Hence the G -ball centered at g , with radius enlarged by a universal constant, contains a translate inside $\bar{V}$ of an r -dimensional ellipsoid whose principal semi-axes are comparable to

$$\min \left\{ R, \frac{u}{\sqrt{\lambda_k(G)}} \right\}, \quad k = 1, \dots, r.$$

Because the support of $\pi_{\bar{V}}$ is $B_2(2R) \cap \bar{V}$, this translated ellipsoid lies in the support up to another universal loss. The volume ratio between $B_2(2R) \cap \bar{V}$ and this ellipsoid is therefore bounded by

$$C \prod_{k=1}^r \left(\frac{CR\sqrt{\lambda_k(G)}}{u} \right),$$

with factors below one omitted precisely by the definition of r_{eff} . Taking logarithms gives $C + d_{\text{ell}}(G, R, u)$. This is the same finite-dimensional volume-ratio argument used in the DNN local-chart lemma. $\square$

Let $\mathcal{A}_j$ be a finite set of phase labels at level j , with prior weights $q_j(a) > 0$. Conditional on a phase label a and rank r , let $\nu_{j,a,r}$ be a prior on the Grassmannian $\text{Gr}(p_j, r)$. For $G_j = G_j(g, \varepsilon)$, define the atlas cost

$$\mathfrak{A}_j(g, \varepsilon, u_j) := \log \frac{1}{q_j(a_j(g))} + \log(1 + p_j) + \log \frac{1}{\nu_{j,a_j,r_j}(B_{\rho_{\text{proj}, G_j}}(V_j, u_j/(4R_j)))}, \quad (5.7)$$

where $r_j = r_{\text{eff}}(G_j, R_j, u_j)$, $V_j = V_{\text{eff}}(G_j, R_j, u_j)$, and $a_j(g)$ is any chart containing the trajectory at level j . If $\nu_{j,a,r}$ is Haar-uniform on $\text{Gr}(p_j, r)$, the last term is bounded by the usual Grassmannian entropy; in an ellipsoidal projection metric one may use the anisotropic Grassmannian bound from the DNN pointwise theory (Li and Xu, 2026).

Theorem 5.4 (Finite-scale RG pointwise dimension). *Assume the metric domination (5.2). Choose resolutions $u_j > 0$ such that $\sum_{j=0}^{J-1} u_j^2 \leq c\varepsilon^2$. Construct a trajectory-independent hierarchical prior Π by independently, at each level j , sampling a phase label a , an effective rank r , a reference subspace $\bar{V} \in \text{Gr}(p_j, r)$, and then sampling g_j uniformly in $B_2(2R_j) \cap \bar{V}$. Then, uniformly for every trajectory g ,*

$$\log \frac{1}{\Pi(B_\rho(g, \varepsilon))} \leq C \sum_{j=0}^{J-1} [d_{\text{ell}}(G_j(g, \varepsilon), R_j, u_j) + \mathfrak{A}_j(g, \varepsilon, u_j)]. \quad (5.8)$$

Proof. For each scale choose the phase label $a_j(g)$, the active rank r_j , and a reference subspace $\bar{V}_j$ within projection distance $u_j/(4R_j)$ of V_j . Lemma 5.3 gives the local prior mass of the ellipsoidal u_j -ball inside this chart. Multiplying the scale-wise prior masses and including the phase, rank, and subspace probabilities gives the right-hand side of (5.8). Finally, if each scale perturbation lies in its ellipsoidal u_j -ball, then (5.2) and $\sum_j u_j^2 \leq c\varepsilon^2$ imply $g' \in B_\rho(g, \varepsilon)$. Hence the product event is contained in the ρ -ball, and its prior mass lower-bounds $\Pi(B_\rho(g, \varepsilon))$. $\square$

Interpretation. The theorem is the RG counterpart of the DNN Riemannian-dimension calculation. The local term d_{ell} is the effective dimension of the finite-scale response Gram. The atlas term is the price of making the prior independent of the realized RG trajectory. Irrelevant directions do not contribute once their eigenvalues fall below the finite resolution u_j^2/R_j^2 ; relevant and marginal directions are precisely the active directions that remain in the sum.

Free Gaussian RG as the fixed-kernel baseline Let $E = E_C \oplus E_F$ be finite-dimensional and let a centered Gaussian field have precision matrix

$$Q = \begin{pmatrix} Q_{CC} & Q_{CF} \\ Q_{FC} & Q_{FF} \end{pmatrix}, \quad Q_{FF} \succ 0.$$

Define the Schur-complement precision

$$Q_C^{\text{RG}} := Q_{CC} - Q_{CF} Q_{FF}^{-1} Q_{FC}.$$

Then marginalizing the fluctuation variables gives $\phi_C \sim N(0, (Q_C^{\text{RG}})^{-1})$, and conditionally

$$\phi_F = -Q_{FF}^{-1} Q_{FC} \phi_C + \zeta, \quad \zeta \sim N(0, Q_{FF}^{-1})$$

independently of ϕ_C . This is a fixed-kernel RG step: all pointwise complexity is determined by the Schur-complement covariance and the fluctuation covariance. It is therefore analogous to a kernel, GP, or NTK model rather than to nonlinear feature learning.

5.2 A finite nonconvex phase-atlas example

The previous theorem is abstract. We now give a concrete finite-volume chartwise example. The point is not to construct a full continuum double-well field, but to show how a globally non-log-concave finite measure can be controlled after it is decomposed into stable phase charts. Let $E = E_C \oplus E_F$, let $\mathcal{A}$ be a finite set of phases, and let

$$\nu(d\phi_C, d\phi_F) = \sum_{a \in \mathcal{A}} w_a \nu_a(d\phi_C, d\phi_F), \quad \nu_a \propto e^{-S_a(\phi_C, \phi_F)} \mathbf{1}_{\Omega_{C,a} \times \Omega_{F,a}} d\phi_C d\phi_F,$$

where the sets $\Omega_{C,a}, \Omega_{F,a}$ are closed convex sets with nonempty interior. The mixture may be globally non-log-concave and multimodal. Assume that, on each chart,

$$\nabla^2 S_a(\phi_C, \phi_F) \succeq H_a = \begin{pmatrix} A_a & B_a \\ B_a^\top & D_a \end{pmatrix}, \quad D_a \succ 0, \quad H_a^{\text{RG}} := A_a - B_a D_a^{-1} B_a^\top \succ 0.$$

Theorem 5.5 (One-step nonconvex phase-atlas RG stability). *Under the preceding assumptions, the coarse marginal is again a finite phase-atlas mixture*

$$\nu_C(d\phi_C) = \sum_{a \in \mathcal{A}} w_a^+ \nu_{C,a}(d\phi_C), \quad \nu_{C,a} \propto e^{-S_a^{\text{RG}}(u)} \mathbf{1}_{\Omega_{C,a}} du,$$

where the weights w_a^+ are the corresponding normalized marginal weights and

$$S_a^{\text{RG}}(u) := -\log \int_{\Omega_{F,a}} e^{-S_a(u,v)} dv.$$

Moreover, S_a^{RG} is H_a^{RG} -strongly convex. Hence, for any finite family of coarse linear observables $Y_x(u) = \langle v_x, u \rangle$, the centered chart process $Y_x - \mathbb{E}_{\nu_{C,a}} Y_x$ is sub-Gaussian with canonical metric

$$d_a(x, y)^2 = \langle v_x - v_y, (H_a^{\text{RG}})^{-1} (v_x - v_y) \rangle.$$

Consequently, the standard sub-Gaussian chaining and variance peeling arguments applied chart by chart, with the comparison metric induced by $(H_a^{\text{RG}})^{-1}$, yields a simultaneous envelope for the mixture, with an additional phase-atlas cost $\log(1/\min_a w_a^+)$, or $\log |\mathcal{A}|$ for equal weights.

Proof. The marginal formula is the definition of integration over E_F . The Schur-complement lower bound follows by completing the square:

$$\begin{pmatrix} u \\ v \end{pmatrix}^\top H_a \begin{pmatrix} u \\ v \end{pmatrix} = u^\top H_a^{\text{RG}} u + (v + D_a^{-1} B_a^\top u)^\top D_a (v + D_a^{-1} B_a^\top u).$$

Moreover,

$$H_a - \begin{pmatrix} H_a^{\text{RG}} & 0 \\ 0 & 0 \end{pmatrix} = \begin{pmatrix} B_a D_a^{-1} B_a^\top & B_a \\ B_a^\top & D_a \end{pmatrix} \succeq 0.$$

Hence $S_a(u, v) - \frac{1}{2} u^\top H_a^{\text{RG}} u$ is jointly convex on the product chart. Including the indicators of the convex sets $\Omega_{C,a}$ and $\Omega_{F,a}$, the Prékopa–Leindler theorem implies that $S_a^{\text{RG}}(u) - \frac{1}{2} u^\top H_a^{\text{RG}} u$ is convex on $\Omega_{C,a}$. Thus the coarse chart is H_a^{RG} -strongly log-concave. Brascamp–Lieb and Herbst then give the stated sub-Gaussian bound for linear observables. The envelope follows from the usual generic-chaining upper bound for sub-Gaussian increments, followed by the same pointwise peeling used in the Gaussian proof and a union bound over $a \in \mathcal{A}$. $\square$

Corollary 5.6 (Pure-phase charts for a finite-volume double-well Φ^4 model). *Let $G = (V, E)$ be finite and*

$$S(\phi) = \frac{1}{2} \sum_{\{u,v\} \in E} c_{uv} (\phi_u - \phi_v)^2 + \lambda \sum_{u \in V} (\phi_u^2 - a^2)^2.$$

Fix $m > a/\sqrt{3}$ and define phase charts

$$\Omega_+ = \{\phi : \phi_u \geq m \ \forall u\}, \quad \Omega_- = -\Omega_+.$$

The restrictions of $e^{-S(\phi)}d\phi$ to Ω_+ and Ω_- define two stable phase charts. A finite mixture of these chart measures is generally non-log-concave, but every chart has Hessian bounded below by

$$L_G + 4\lambda(3m^2 - a^2)I.$$

Therefore Theorem 5.5 applies to every finite decomposition of variables $V = C \sqcup F$, and one nonlinear coarse-graining step preserves a two-chart pointwise-complexity envelope for these pure-phase restrictions. To cover the full double-well Gibbs measure one would need additional charts, for instance droplet or interface charts, and corresponding chartwise Hessian bounds.

Proof. On either chart, $|\phi_u| \geq m$, and

$$\frac{d^2}{dt^2}\lambda(t^2 - a^2)^2 = 4\lambda(3t^2 - a^2) \geq 4\lambda(3m^2 - a^2) > 0.$$

Adding the graph Dirichlet Hessian gives the displayed lower bound. The result follows from Theorem 5.5. $\square$

Interpretation of the non-convex phase-atlas example. The finite nonconvex phase-atlas example should be read as a finite-cutoff bridge between convex concentration theory, high-dimensional geometry, and interacting-field intuition. Within a single stable phase chart, the measure is strongly log-concave: a uniform Hessian lower bound gives Brascamp–Lieb covariance domination and, through logarithmic-Sobolev/Herbst arguments, Gaussian-type concentration for linear and Lipschitz observables (Brascamp and Lieb, 1976; Bakry and Emery, 1985). A mixture of several such charts, however, is generally not log-concave and may be genuinely multimodal, so there need not be a single global convex potential or a single global Gaussian comparison. The phase-atlas decomposition restores tractability by separating these effects. Chartwise, the Schur-complement Hessian gives the local Gaussian comparison metric and hence the pointwise ellipsoidal complexity; globally, one pays an explicit atlas cost for selecting the relevant phase and active subspace.

This distinction is also natural from the viewpoint of convex geometry. Ordinary log-concavity gives powerful isoperimetric and tail estimates, and one-dimensional marginals of isotropic log-concave measures have universal sub-exponential tails (Borell, 1975; Lovasz and Vempala, 2007). It does not, however, by itself provide dimension-free Gaussian-type control in all directions. The Kannan–Lovasz–Simonovits conjecture is a central benchmark for dimension-free isoperimetric and Poincare-type control of isotropic log-concave measures (Kannan et al., 1995). The finite-cutoff viewpoint used here is different. Rather than trying to prove such global dimension-free control for an arbitrary log-concave measure, we work on local charts where the cutoff and the quadratic part of the effective action give a uniform curvature lower bound. On each chart this strong log-concavity yields Gaussian-type comparison in the anisotropic Hessian metric. Global non-log-concavity is then handled by the phase atlas, and directions below the finite resolution are paid for as approximation error rather than controlled through a global Cheeger constant.

In this sense, the pointwise-complexity theorem gives an explicit spectral version of a local convex-geometric principle with global, finite-cutoff extension: the effective spectrum of the chartwise Hessian determines the local ellipsoidal dimension, while the phase atlas records the cost of moving between globally non-log-concave charts. This is parallel to the local-chart/global-atlas construction in the DNN pointwise theory (Li and Xu, 2026), but expressed here in the language of finite-cutoff interacting fields.

6 Graph Local-Time Application versus Cover Time

This section specializes the pointwise Gaussian theorem to pinned Gaussian free fields and then, through Ray–Knight, to random-walk local times. The distinction from cover-time theory is clearest for target questions. A scalar cover-time theorem concerns the event that every vertex has been visited. The field-level Ray–Knight estimate can instead be evaluated on a target field before one passes to any full-cover event. Local time is a standard tool towards cutoff-uniform local integrability control in constructive QFT (Dynkin, 1984).

The phase-star examples below make this distinction quantitative. A target subatlas I is assigned a code length through an atlas prior q . The pointwise local-time route pays this target code length, plus any within-chart fluctuation cost; a full-cover route pays the cost of all N phases. Thus the graph examples make explicit the same local-chart/global-atlas principle that underlies the finite-scale ellipsoidal theory in Section 5.

Graph setup, pinned GFF, and Ray–Knight. Let $G = (V, E, c)$ be a finite connected weighted graph with symmetric conductances $c_{xy} = c_{yx} \geq 0$, where $c_{xy} > 0$ exactly when $\{x, y\} \in E$. Write

$$c_x := \sum_{y \in V} c_{xy}.$$

The weighted graph Laplacian acts on functions $f : V \rightarrow \mathbb{R}$ by

$$(Lf)(x) = \sum_y c_{xy}(f(x) - f(y)).$$

We consider the continuous-time random walk, started at a distinguished root v_0 , that jumps from x to y at rate c_{xy}/c_x . Its normalized local time is

$$L_t(x) = \frac{1}{c_x} \int_0^t \mathbf{1}\{X_s = x\} ds, \quad \tau(t) := \inf\{s \geq 0 : L_s(v_0) > t\}.$$

The pinned Gaussian free field with root v_0 is the centered Gaussian vector $\eta = \{\eta_x\}_{x \in V}$ such that $\eta_{v_0} = 0$ almost surely and whose covariance is the killed Green function

$$\mathbb{E}[\eta_x \eta_y] = \Gamma_{v_0}(x, y) := \mathbb{E}_x[L_{T_{v_0}}(y)],$$

where T_{v_0} is the hitting time of v_0 . Equivalently, this covariance is the inverse of the pinned Laplacian on $V \setminus \{v_0\}$. Its canonical metric is

$$d(x, y) = (\mathbb{E}|\eta_x - \eta_y|^2)^{1/2} = \sqrt{R_{\text{eff}}(x, y)},$$

where R_{eff} is the effective-resistance metric of the conductance network.

We use the following finite-graph form of the second Ray–Knight theorem (Dynkin, 1984). Let η' be an independent copy of the pinned GFF.

Proposition 6.1 (Second Ray–Knight isomorphism on a finite graph). *With the above normalization, under the product law of the walk and the independent fields (η, η') ,*

$$\left(L_{\tau(t)}(x) + \frac{1}{2}\eta_x^2\right)_{x \in V} \stackrel{d}{=} \left(\frac{1}{2}(\eta'_x + \sqrt{2t})^2\right)_{x \in V}. \quad (6.1)$$

In particular, $L_{\tau(t)}(v_0) = t$ almost surely.

6.1 Target local times versus cover-time reduction

Target local-time envelopes. Let $H \subseteq V$ be a target set. The proof below only requires a deterministic envelope for the pinned GFF on H . Such an envelope may be obtained either by applying Theorem 1.1 to the restricted field on $H \cup \{v_0\}$, or by applying it once to a larger ambient field and then evaluating the resulting bound on H . This distinction is useful in the phase-star examples, where one ambient atlas prior is fixed before the target subatlas is selected.

Corollary 6.2 (Target-set pointwise local-time envelope). *Let $A_H(\cdot; \delta)$ be any deterministic function satisfying*

$$\mathbb{P}\{|\eta_x| \leq A_H(x; \delta) \text{ for all } x \in H\} \geq 1 - \delta$$

for the pinned GFF. Then, under the law of the walk, with probability at least $1 - \delta$,

$$\forall x \in H : \quad [t - \sqrt{2t} A_H(x; \delta/2)]_+ \leq L_{\tau(t)}(x) \leq t + \sqrt{2t} A_H(x; \delta/2) + \frac{1}{2} A_H(x; \delta/2)^2. \quad (6.2)$$

Let

$$\tau_H := \inf\{s \geq 0 : L_s(x) > 0 \text{ for every } x \in H\}$$

be the target-cover time of H . The preceding envelope gives the following target-cover and target-blanket criterion.

Corollary 6.3 (Target cover and target blanket criterion). *Assume that for some $t > 0$, $H \subseteq V$, and $\theta \in (0, 1/\sqrt{2})$,*

$$\sup_{x \in H} A_H(x; \delta/2) \leq \theta \sqrt{t}.$$

Then, under the law of the walk, with probability at least $1 - \delta$,

$$(1 - \sqrt{2}\theta)t \leq L_{\tau(t)}(x) \leq \left(1 + \sqrt{2}\theta + \frac{1}{2}\theta^2\right)t \quad \forall x \in H.$$

In particular, $\tau_H \leq \tau(t)$ on this event, and

$$\max_{u, v \in H} \frac{L_{\tau(t)}(u)}{L_{\tau(t)}(v)} \leq \frac{1 + \sqrt{2}\theta + \theta^2/2}{1 - \sqrt{2}\theta}.$$

Proof of Corollary 6.2. Set

$$A_x := A_H(x; \delta/2), \quad U_x := L_{\tau(t)}(x) + \frac{1}{2}\eta_x^2, \quad V_x := \frac{1}{2}(\eta'_x + \sqrt{2t})^2, \quad x \in H.$$

By Theorem 6.1, $(U_x)_{x \in H} \stackrel{d}{=} (V_x)_{x \in H}$. Define

$$D_x := \frac{1}{2}(\sqrt{2t} - A_x)_+^2, \quad B_x := t + \sqrt{2t} A_x + \frac{1}{2} A_x^2.$$

By the envelope assumption applied to the independent copy η' , the event $\{|\eta'_x| \leq A_x \text{ for all } x \in H\}$ has probability at least $1 - \delta/2$. On this event, $D_x \leq V_x \leq B_x$ for all $x \in H$. Equality in distribution therefore gives

$$\mathbb{P}\{D_x \leq U_x \leq B_x \text{ for all } x \in H\} \geq 1 - \delta/2.$$

Applying the same envelope to η itself and taking a union bound, with probability at least $1 - \delta$, we also have $|\eta_x| \leq A_x$ for all $x \in H$. On the intersection of these events,

$$L_{\tau(t)}(x) = U_x - \frac{1}{2}\eta_x^2 \leq B_x,$$

and

$$L_{\tau(t)}(x) \geq D_x - \frac{1}{2}A_x^2.$$

Using nonnegativity of local time and the identity

$$\max \left\{ 0, \frac{1}{2}(\sqrt{2t} - A_x)_+^2 - \frac{1}{2}A_x^2 \right\} = [t - \sqrt{2t} A_x]_+$$

gives (6.2). The event in the conclusion depends only on the walk, so the same probability bound holds under the walk law. $\square$

Proof of Corollary 6.3. Under the displayed small-envelope assumption, Corollary 6.2 gives, simultaneously for all $x \in H$,

$$L_{\tau(t)}(x) \geq (1 - \sqrt{2}\theta)t > 0$$

and

$$L_{\tau(t)}(x) \leq \left(1 + \sqrt{2}\theta + \frac{1}{2}\theta^2\right)t.$$

Thus every vertex in H has been visited by time $\tau(t)$, and the displayed ratio bound follows by dividing the upper estimate by the lower estimate. $\square$

Phase-star hierarchies. We now give a finite graph model that separates direct target-local-time control from a reduction through full cover time. If a deterministic target set is fixed in advance and one is allowed to redo the entire Gaussian analysis on that target alone, then classical generic chaining on the restricted field has the correct target scale. The point here is different. We fix one ambient atlas prior before the target is selected, and then evaluate the same pointwise envelope on any target subatlas. The target scale includes the code length of the selected subatlas under this prior. A proof that first establishes full cover loses this target dependence, because it must cover every phase.

Theorem 6.4 (Phase-star target subatlas versus full cover). *Let S_N be the star graph with root v_0 , leaves $\{1, \dots, N\}$, and unit conductances. Let $\mathcal{I}$ be a collection of nonempty subsets of $\{1, \dots, N\}$, and let $q \in \Delta(\mathcal{I})$. For $I \in \mathcal{I}$, set*

$$H_I := I, \quad \tau_I := \inf\{s \geq 0 : L_s(i) > 0 \text{ for every } i \in I\}.$$

Let τ_{cov} be the cover time of the whole star. Define the ambient prior

$$\mu_q := \frac{1}{2}\delta_{v_0} + \frac{1}{4}\text{Unif}\{1, \dots, N\} + \frac{1}{4} \sum_{I \in \mathcal{I}} q(I) \text{Unif}(I). \quad (6.3)$$

Let $A_q(i; \delta)$ be the two-sided pointwise Gaussian envelope from (1.5) applied to the full pinned GFF on S_N , with prior μ_q , with $r_0 = s_0 = 1$, and with constants enlarged if necessary. Then the following hold.

- (i) *The pinned GFF satisfies $\eta_1, \dots, \eta_N$ independent $N(0, 1)$. Moreover, with probability at least $1 - \delta$, simultaneously for all $I \in \mathcal{I}$,*

$$\sup_{i \in I} |\eta_i| \leq C \left[\sqrt{\log \frac{4|I|}{q(I)}} + \sqrt{\log \left(\frac{2 + \log \log(N + 3)}{\delta} \right)} \right]. \quad (6.4)$$

Consequently, the same deterministic right-hand side may be used as $\sup_{i \in I} A_q(i; \delta)$ in the Ray–Knight local-time envelope.

(ii) For every $\theta \in (0, 1/\sqrt{2})$, there is $C_\theta < \infty$ such that, for each $I \in \mathcal{I}$,

$$t \geq C_\theta \left[\log \frac{4|I|}{q(I)} + \log \left(\frac{2 + \log \log(N+3)}{\delta} \right) \right] \implies \mathbb{P}\{\tau_I \leq \tau(t)\} \geq 1 - \delta. \quad (6.5)$$

The probability statement is for the selected target I . The prior μ_q and the Gaussian envelope are fixed independently of that selection.

(iii) The exact inverse-root-local-time laws are

$$\mathbb{P}\{\tau_I \leq \tau(t)\} = (1 - e^{-t})^{|I|}, \quad \mathbb{P}\{\tau_{\text{cov}} \leq \tau(t)\} = (1 - e^{-t})^N. \quad (6.6)$$

Thus the $(1 - \delta)$ -quantile for covering k prescribed leaves is

$$q_k(\delta) := -\log \left(1 - (1 - \delta)^{1/k} \right) = \log k + O_\delta(1). \quad (6.7)$$

In particular, $q_N(\delta) - q_{|I|}(\delta) = \log(N/|I|) + O_\delta(1)$. Hence, whenever

$$\log \frac{|I|}{q(I)} = o(\log N),$$

the direct target-local-time route has an inverse-root-local-time scale asymptotically smaller than the scale required by any argument that certifies the same target event only through full cover.

Proof. For the star graph, the pinned Laplacian on the leaves is the identity matrix, so the pinned GFF has covariance I_N . For a leaf i , $\sigma(i) = d(i, v_0) = 1$, and the star metric satisfies $d(i, j) = \sqrt{2}$ for distinct leaves. If $i \in I$, then (6.3) gives

$$\mu_q(\{i\}) \geq \frac{q(I)}{4|I|}.$$

Since the metric diameter of the star leaves is bounded by an absolute constant, the localized Fernique–Talagrand functional satisfies

$$\Phi_{\mu_q}(i) \leq C \sqrt{\log \frac{4|I|}{q(I)}} \quad (i \in I).$$

The uniform component gives $\mu_q(\{i\}) \geq 1/(4N)$ for every leaf. Hence $\Phi_{\mu_q,*} \leq C\sqrt{\log(N+1)}$. With $r_0 = s_0 = 1$, the peeling multiplicity in Theorem 1.1 is bounded by a constant times $2 + \log \log(N+3)$. The two-sided pointwise Gaussian theorem gives (6.4). Applying Corollary 6.3, with the usual confidence split, gives (6.5) after increasing C_θ .

It remains to prove (6.6). During visits to the root, the walk waits an exponential time of rate one before departing and then chooses one of the N leaves uniformly. Since $c_{v_0} = N$, accumulated root occupation time at $\tau(t)$ is Nt . Therefore, per unit normalized root local time, departures from the root form a Poisson process of rate N with independent uniform leaf marks. By Poisson thinning, for each leaf i , the number of excursions from the root to i before $\tau(t)$ is Poisson with mean t , and these counts are independent over leaves. A leaf has positive local time by $\tau(t)$ if and only if it has been chosen at least once. This gives the two identities in (6.6). Solving $(1 - e^{-t})^k \geq 1 - \delta$ gives (6.7), and the asymptotic formula follows from $(1 - \delta)^{1/k} = 1 + k^{-1} \log(1 - \delta) + o(k^{-1})$. $\square$

The collapsed star has no within-phase geometry. The next theorem adds a local chart to each phase. It is a graph analogue of a phase/subspace atlas: the prior first pays for selecting a target subatlas, and the effective resistance inside each dense chart suppresses the within-chart fluctuation.

Theorem 6.5 (Decorated phase-star subatlas theorem). *Fix $N, K \geq 1$ and $\kappa > 0$ with $\kappa(K+1) \geq 1$. Let $G_{N,K,\kappa}$ be the conductance graph with root v_0 , phase centers $c_1, \dots, c_N$, and local chart vertices $z_{a,1}, \dots, z_{a,K}$ for each phase a . The root is connected to each c_a by an edge of conductance one. For each a ,*

$$C_a := \{c_a, z_{a,1}, \dots, z_{a,K}\}$$

is a complete graph with edge conductance κ , and there are no other edges. Let $\mathcal{I}$ be a collection of nonempty subsets of $\{1, \dots, N\}$, and let $q \in \Delta(\mathcal{I})$. For $I \in \mathcal{I}$, define the decorated target

$$H_{I,K} := \bigcup_{a \in I} C_a.$$

Define the ambient prior

$$\mu_{q,K} := \frac{1}{2} \delta_{v_0} + \frac{1}{4} \text{Unif} \left(\bigcup_{a=1}^N C_a \right) + \frac{1}{4} \sum_{I \in \mathcal{I}} q(I) \text{Unif} \left(\bigcup_{a \in I} C_a \right). \quad (6.8)$$

Then the following hold.

(i) *The effective resistance within a phase is compressed:*

$$R_{\text{eff}}(x, y) \leq \frac{2}{\kappa(K+1)}, \quad x, y \in C_a, \quad (6.9)$$

while

$$R_{\text{eff}}(v_0, c_a) = 1, \quad R_{\text{eff}}(v_0, x) \leq 1 + \frac{2}{\kappa(K+1)}, \quad x \in C_a. \quad (6.10)$$

(ii) *Let $A_{q,K}(x; \delta)$ be the two-sided pointwise Gaussian envelope from (1.5) applied to the full pinned GFF on $G_{N,K,\kappa}$, with prior $\mu_{q,K}$, with $r_0 = s_0 = 1$, and with constants enlarged if necessary. Then, with probability at least $1 - \delta$, simultaneously for all $I \in \mathcal{I}$,*

$$\begin{aligned} \sup_{x \in H_{I,K}} |\eta_x| \leq C & \left[\sqrt{\log \frac{4|I|}{q(I)}} + \sqrt{\frac{1}{\kappa(K+1)} \log \frac{4|I|(K+1)}{q(I)}} \right. \\ & \left. + \sqrt{\log \left(\frac{2 + \log \log((N+3)(K+3))}{\delta} \right)} \right]. \end{aligned} \quad (6.11)$$

Consequently, for every $\theta \in (0, 1/\sqrt{2})$, there is $C_\theta < \infty$ such that, for the selected I ,

$$\begin{aligned} t \geq C_\theta & \left[\log \frac{4|I|}{q(I)} + \frac{1}{\kappa(K+1)} \log \frac{4|I|(K+1)}{q(I)} \right. \\ & \left. + \log \left(\frac{2 + \log \log((N+3)(K+3))}{\delta} \right) \right]. \end{aligned} \quad (6.12)$$

implies

$$\mathbb{P}\{\tau_{H_{I,K}} \leq \tau(t)\} \geq 1 - \delta.$$

(iii) Let τ_{cov} be the cover time of the full decorated graph, and set $C_N^{\text{cen}} := \{c_1, \dots, c_N\}$.

$$\mathbb{P}\{C_N^{\text{cen}} \subseteq X[0, \tau(t)]\} = (1 - e^{-t})^N. \quad (6.13)$$

Since full cover implies that all phase centers have been visited,

$$\mathbb{P}\{\tau_{\text{cov}} \leq \tau(t)\} \leq (1 - e^{-t})^N. \quad (6.14)$$

Thus any argument that proves target cover only by first proving full cover at confidence $1 - \delta$ must use inverse-root-local-time at least $q_N(\delta)$ from (6.7). If, for fixed $\delta \in (0, 1)$,

$$\log \frac{|I|}{q(I)} + \frac{1}{\kappa(K+1)} \log \frac{|I|(K+1)}{q(I)} + \log \log((N+3)(K+3)) = o(\log N),$$

then the sufficient scale in (6.12) is $o(q_N(\delta))$, while $\mathbb{P}\{\tau_{\text{cov}} \leq \tau(t)\} \rightarrow 0$ along that scale.

Proof. The resistance estimate inside C_a is the standard effective resistance of a complete graph with $K+1$ vertices and edge conductance κ , namely $2/[\kappa(K+1)]$. Since c_a is the only vertex of C_a connected to the root, attaching the remaining clique to c_a does not change the effective resistance between v_0 and c_a ; hence $R_{\text{eff}}(v_0, c_a) = 1$. The triangle inequality for effective resistance gives (6.10).

Put

$$\alpha_{K,\kappa} := \sqrt{\frac{2}{\kappa(K+1)}}.$$

For $x \in C_a$ and $a \in I$, the ball $B_d(x, \alpha_{K,\kappa})$ contains C_a by (6.9). Under $\mu_{q,K}$, this ball has mass at least $q(I)/(4|I|)$. For smaller radii, the singleton mass is at least $q(I)/[4|I|(K+1)]$. Also $\sigma(x) \leq 2$ by (6.10). Therefore, for $x \in C_a$ with $a \in I$,

$$\Phi_{\mu_{q,K}}(x) \leq C \sqrt{\log \frac{4|I|}{q(I)}} + C \alpha_{K,\kappa} \sqrt{\log \frac{4|I|(K+1)}{q(I)}}.$$

The uniform component in $\mu_{q,K}$ gives singleton mass at least $1/[4N(K+1)]$, and hence $\Phi_{\mu_{q,K},*} \leq C \sqrt{\log((N+1)(K+1))}$. With $r_0 = s_0 = 1$, the peeling multiplicity contributes only the displayed $\log \log((N+3)(K+3))$ term. This proves (6.11). The target-cover implication follows from Corollary 6.3 as in the proof of Theorem 6.4.

Finally, each departure from the root chooses one of the phase centers uniformly, and the same Poisson-thinning argument used in Theorem 6.4 gives (6.13). Since full cover requires every phase center to have been visited, (6.14) follows. The final asymptotic statement follows from $q_N(\delta) = \log N + O_\delta(1)$. $\square$

Relation to classical cover-time theory. For a finite connected conductance graph $G = (V, E, c)$, let τ_{cov} be the first time every vertex has been visited, and write

$$t_{\text{cov}}(G) := \max_{v \in V} \mathbb{E}_v[\tau_{\text{cov}}]. \quad (6.15)$$

A blanket time strengthens cover time by requiring that the walk has accumulated roughly stationary proportions of visits at all vertices. One convenient strong version leads to

$$t_{\text{bl}}(G, \delta) := \max_{v \in V} \mathbb{E}_v[\tau_{\text{bl}}(\delta)]. \quad (6.16)$$

Let $H(u, v)$ be the expected hitting time from u to v , and define the commute-time metric

$$\kappa(u, v) := H(u, v) + H(v, u).$$

Let $\mathcal{C} := \sum_{x \in V} c_x$ be the total conductance. The commute-time identity gives

$$\kappa(u, v) = \mathcal{C} R_{\text{eff}}(u, v).$$

For an unweighted graph, $\mathcal{C} = 2|E|$. If $\eta = \{\eta_v\}_{v \in V}$ denotes the pinned GFF, then $\mathbb{E}(\eta_u - \eta_v)^2 = R_{\text{eff}}(u, v)$.

Proposition 6.6 (Ding–Lee–Peres (Ding et al., 2012)). *For any connected conductance graph $G = (V, E, c)$ and any fixed $0 < \delta < 1$,*

$$t_{\text{cov}}(G) \asymp \gamma_2(V, \sqrt{\kappa})^2 = \mathcal{C} \cdot \gamma_2(V, \sqrt{R_{\text{eff}}})^2 \asymp_{\delta} t_{\text{bl}}(G, \delta). \quad (6.17)$$

Equivalently,

$$t_{\text{cov}}(G) \asymp \mathcal{C} \cdot \left(\mathbb{E} \max_{v \in V} \eta_v \right)^2. \quad (6.18)$$

Proposition 6.6 is a global theorem: it identifies the scale of the time needed to cover every vertex. The target local-time estimates above answer a different question. They keep a single ambient field-level envelope and evaluate it on a selected subatlas before any full-cover event is invoked. Theorems 6.4 and 6.5 show that this distinction can be strict. On the star, a selected target I has inverse-root-local-time quantile $\log |I| + O_{\delta}(1)$, while full cover has quantile $\log N + O_{\delta}(1)$. The pointwise target envelope pays the atlas code length $\log(|I|/q(I))$; in the decorated star it also pays the compressed within-chart term $(\kappa(K+1))^{-1} \log(|I|(K+1)/q(I))$. Any argument that first passes through the full-cover event necessarily pays the coupon-collector cost associated with all N phase centers. If a target set is fixed in advance, and the Gaussian analysis is allowed to be restricted to that target, then classical generic chaining applied to the restricted field can recover the corresponding target scale. The purpose of Theorems 6.4 and 6.5, however, is different: they construct a simultaneous envelope valid uniformly over all target subsets I . The pointwise formulation is what makes this possible. It retains the dependence on the selected subatlas under a single ambient prior, rather than collapsing the analysis at the outset to the full-cover event.

7 Conclusion

The paper has three linked pointwise/algorithmic and finite-cutoff messages. First, for Gaussian processes, the natural refinement of generic chaining is a simultaneous pointwise envelope for the field, not only a scalar estimate on $\mathbb{E} \sup_x X_x$. This envelope is sharp in expectation after optimizing over the prior and recovers the anchored Fernique–Talagrand scale. It should be useful whenever one wants to retain local field information before taking a final supremum.

Second, the Bayesian algorithmic lower envelope gives a single-radius lower-bound counterpart to pointwise complexity. It is an information-theoretic Bayes-risk bound for every estimator $\hat{t}(Y)$, expressed through the exact ghost small-ball mass $\mathbb{E}_Q \pi(B_d(\hat{t}(Y), \Delta))$. The comparison-decoder theorem turns this Bayes distance obstruction into a lower bound on a decision-aligned Gaussian range; conversely, a simultaneous pointwise Gaussian envelope yields an algorithmic upper bound on the same decoding risk. Importantly, we construct a heterogeneous-orthogonal-basis “hub–leaves–cloud” example for norm-regularized nearest-neighbor decoder where the fixed-prior Fano relaxation is vacuous, the algorithmic ghost-mass lower envelope and pointwise upper envelope

match on the estimator-selected subatlas, while the full-class minimax risk and global Gaussian scale are much larger. Keeping the algorithmic mass is the key to certifying pointwise complexity locally.

Third, finite-cutoff renormalization illustrates the same structural mechanism in a nonlinear setting. Exact telescoping exposes a response Gram, and a hierarchical atlas converts trajectory-dependent active directions into a trajectory-independent prior. The practical reason to keep this theorem finite-cutoff is that it gives a way to control coarse marginals of globally non-log-concave, anisotropic interacting systems through local strong log-concavity, spectral ellipsoidal complexity, and an explicit phase/subspace atlas. More concretely, the local-time application shows how the pointwise upper envelope can be used before reducing the problem to cover times of finite graphs via the Gaussian free field. Ray–Knight transfers pointwise GFF control to target-set local-time control, and the phase-star examples demonstrate that a selected subatlas can have a smaller intrinsic scale than any route through full cover. Across these settings, the contribution is the same finite-scale complexity layer: expose the correct ellipsoidal metric, keep the relevant local prior mass, and pay the atlas cost only when global uniformity requires it.

Taken together, the results suggest a unified viewpoint. The pointwise Gaussian upper theorem shows that a rich field should be controlled through local envelopes before it is collapsed to a global supremum. The Bayesian algorithmic upper and lower bounds show that, for a fixed prior, reference law, and estimator, the relevant information-theoretic complexity may be the estimator-selected subgeometry rather than the whole ambient class. The finite-cutoff RG framework identifies the analogous issue for interacting fields: before a covering argument is meaningful, the nonlinear dynamics must first expose the appropriate response Gram and phase/subspace atlas. Thus the same principle appears in three forms: global worst-case objects capture genuine hardness of the full class, while pointwise and algorithmic objects can certify tractability on the local geometry selected by a field, an estimator, or a renormalization trajectory. This is the sense in which the paper connects field-level generic chaining, algorithmic complexity certificates, and finite-cutoff renormalization.

References

- Michael Aizenman and Hugo Duminil-Copin. Marginal triviality of the scaling limits of critical 4D Ising and ϕ_4^4 models. *Annals of Mathematics*, 194(1):163–235, 2021.
- Dominique Bakry and Michel Emery. Diffusions hypercontractives. In *Séminaire de probabilités XIX*, Lecture Notes in Mathematics, vol. 1123, pages 177–206. Springer, 1985.
- Roland Bauerschmidt, David C. Brydges, and Gordon Slade. Introduction to a Renormalisation Group Method. *Lecture Notes in Mathematics*, vol. 2242. Springer, 2019.
- Mikhail Belkin, Daniel Hsu, and Partha P. Mitra. Overfitting or perfect fitting? Risk bounds for classification and regression rules that interpolate. In *Advances in Neural Information Processing Systems 31*, 2018.
- Adam Block, Yuval Dagan, and Alexander Rakhlin. Majorizing measures, sequential complexities, and online learning. In *Proceedings of the 34th Annual Conference on Learning Theory*, vol. 134, pages 1–4, 2021.
- Christer Borell. Convex set functions in d -space. *Periodica Mathematica Hungarica*, 6:111–136, 1975.

- Herm Jan Brascamp and Elliott H. Lieb. On extensions of the Brunn–Minkowski and Prekopa–Leindler theorems, including inequalities for log-concave functions, and with an application to the diffusion equation. *Journal of Functional Analysis*, 22(4):366–389, 1976.
- David C. Brydges and Gordon Slade. A renormalisation group method. V. A single renormalisation group step. *Journal of Statistical Physics*, 159:589–667, 2015.
- Sourav Chatterjee. Yang–Mills for probabilists. arXiv:1803.01950, 2018.
- Fan Chen, Dylan J. Foster, Yanjun Han, Jian Qian, Alexander Rakhlin, and Yunbei Xu. Assouad, Fano, and Le Cam with interaction: A unifying lower bound framework and characterization for bandit learnability. arXiv:2410.05117, 2024.
- Thomas M. Cover and Peter E. Hart. Nearest neighbor pattern classification. *IEEE Transactions on Information Theory*, 13(1):21–27, 1967.
- George Cybenko. Approximation by superpositions of a sigmoidal function. *Mathematics of Control, Signals and Systems*, 2:303–314, 1989.
- Jian Ding, James R. Lee, and Yuval Peres. Cover times, blanket times, and majorizing measures. *Annals of Mathematics*, 175(3):1409–1471, 2012.
- E. B. Dynkin. Local times and quantum fields. In *Seminar on Stochastic Processes, 1983*. Birkhäuser, 1984.
- Xavier Fernique. Régularité des trajectoires des fonctions aléatoires gaussiennes. In *École d’Été de Probabilités de Saint-Flour IV–1974*, Lecture Notes in Mathematics, vol. 480. Springer, 1975.
- Dylan J. Foster, Sham M. Kakade, Jian Qian, and Alexander Rakhlin. The statistical complexity of interactive decision making. arXiv:2112.13487, 2021.
- Dylan J. Foster, Noah Golowich, and Yanjun Han. Tight guarantees for interactive decision making with the decision-estimation coefficient. In *Proceedings of the 36th Annual Conference on Learning Theory*, vol. 195, pages 1–75, 2023.
- Kurt Hornik, Maxwell Stinchcombe, and Halbert White. Multilayer feedforward networks are universal approximators. *Neural Networks*, 2(5):359–366, 1989.
- Arthur Jaffe and Edward Witten. Quantum Yang–Mills theory. Clay Mathematics Institute Millennium Prize Problem description, 2000.
- Ravi Kannan, Laszlo Lovasz, and Miklos Simonovits. Isoperimetric problems for convex bodies and a localization lemma. *Discrete & Computational Geometry*, 13:541–559, 1995.
- Shaojie Li and Yunbei Xu. Pointwise generalization in deep neural networks. arXiv:2605.18598, 2026.
- Michel Ledoux and Michel Talagrand. *Probability in Banach Spaces: Isoperimetry and Processes*. Springer, 1991.
- Leonid A. Levin. Average case complete problems. *SIAM Journal on Computing*, 15(1):285–286, 1986.

- Laszlo Lovasz and Santosh Vempala. The geometry of logconcave functions and sampling algorithms. *Random Structures & Algorithms*, 30(3):307–358, 2007.
- Neil Lutz. A note on pointwise dimensions. arXiv:1612.05849, 2016.
- David Pollard. The duality proof of the Sudakov minoration. Lecture notes, Yale University, available at <http://www.stat.yale.edu/~pollard/Notes/Sudakov.pdf>, 2001.
- Daniel A. Spielman and Shang-Hua Teng. Smoothed analysis of algorithms: Why the simplex algorithm usually takes polynomial time. *Journal of the ACM*, 51(3):385–463, 2004.
- Michel Talagrand. *The Generic Chaining: Upper and Lower Bounds of Stochastic Processes*. Springer, 2005.
- Michel Talagrand. Regularity of Gaussian processes. *Acta Mathematica*, 159:99–149, 1987.
- Yunbei Xu and Assaf Zeevi. Towards optimal problem dependent generalization error bounds in statistical learning theory. *Mathematics of Operations Research*, 2025.
- Yunbei Xu and Assaf Zeevi. Bayesian design principles for frequentist sequential learning. *Journal of the ACM*, 2025.
- Andrew C. Yao. Probabilistic computations: Toward a unified measure of complexity. In *Proceedings of the 18th Annual Symposium on Foundations of Computer Science*, pages 222–227, 1977.
- Tong Zhang. Information theoretical upper and lower bounds for statistical estimation. *IEEE Transactions on Information Theory*, 52(4):1307–1321, 2006.