

# Introduction to Rigid Analytic Geometry

Yunhai Xiang

June 7, 2024

Motivations

Affinoid Domains

Tate Uniformization

For an algebraic variety  $X/\mathbb{C}$ , one associates its analytification

$$X^{\text{an}} = X(\mathbb{C})$$

a complex analytic space (a complex manifold when  $X$  smooth).

## Example

Suppose  $X/\mathbb{C}$  a smooth algebraic curve of genus  $g$ , then  $X^{\text{an}}$  is a complex manifold of dimension 1, i.e. a Riemann surface, of genus  $g$ . In particular, when  $X$  is an elliptic curve,  $X^{\text{an}}$  is a torus.

## Theorem (Uniformization theorem)

*The only simply connected Riemann surfaces are the Riemann sphere  $\mathbb{CP}^1$ , the complex plane  $\mathbb{C}$ , and the upper-half plane  $\mathbb{H}$ .*

Thus, we have the correspondence

| genus    | $X$            | $X^{\text{an}}$      |
|----------|----------------|----------------------|
| 0        | conic section  | $\mathbb{CP}^1$      |
| 1        | elliptic curve | $\mathbb{C}/\Lambda$ |
| $\geq 2$ | modular curve  | $\mathbb{H}/\Gamma$  |

where  $\Lambda$  a lattice, and  $\Gamma$  a congruence subgroup (congruent to  $I$  mod some  $N$ ) of  $\text{SL}_2(\mathbb{Z})$  acting by the Mobius transformation.

Thus, we have the correspondence

| genus    | $X$            | $X^{\text{an}}$      |
|----------|----------------|----------------------|
| 0        | conic section  | $\mathbb{CP}^1$      |
| 1        | elliptic curve | $\mathbb{C}/\Lambda$ |
| $\geq 2$ | modular curve  | $\mathbb{H}/\Gamma$  |

where  $\Lambda$  a lattice, and  $\Gamma$  a congruence subgroup (congruent to  $I$  mod some  $N$ ) of  $\text{SL}_2(\mathbb{Z})$  acting by the Mobius transformation.

Moreover, Serre showed a collection of results (GAGA) relating coherent sheaves on a variety  $X/\mathbb{C}$  to that of  $X^{\text{an}}$ .

The complex numbers  $\mathbb{C} = \overline{\mathbb{R}}$  has a non-archimedean analogue  $\mathbb{C}_p$ .



The complex numbers  $\mathbb{C} = \overline{\mathbb{R}}$  has a non-archimedean analogue  $\mathbb{C}_p$ .

### Theorem (Ostrowski)

*The nontrivial norms on  $\mathbb{Q}$  are precisely the archimedean norm  $|\cdot|$  and the nonarchimedean  $p$ -adic norms  $|\cdot|_p$ . Thus the completions of  $\mathbb{Q}$  are precisely  $\mathbb{R}$  and  $\mathbb{Q}_p$  for each prime  $p$ .*

The complex numbers  $\mathbb{C} = \overline{\mathbb{R}}$  has a non-archimedean analogue  $\mathbb{C}_p$ .

### Theorem (Ostrowski)

*The nontrivial norms on  $\mathbb{Q}$  are precisely the archimedean norm  $|\cdot|$  and the nonarchimedean  $p$ -adic norms  $|\cdot|_p$ . Thus the completions of  $\mathbb{Q}$  are precisely  $\mathbb{R}$  and  $\mathbb{Q}_p$  for each prime  $p$ .*

However, the algebraic closure  $\overline{\mathbb{Q}_p}$  is not complete, so one has to complete it again  $\mathbb{C}_p = \widehat{\overline{\mathbb{Q}_p}}$  to get a complete and algebraically closed non-archimedean field (Krasner theorem).

Is there a non-archimedean analogue of complex analytic space?

Is there a non-archimedean analogue of complex analytic space?

Is there a uniformization theorem or GAGA for this analogue?

There are many such analogous theories!

There are many such analogous theories!

1. Tate's theory of rigid analytic spaces,

There are many such analogous theories!

1. Tate's theory of rigid analytic spaces,
2. Berkovich's theory of Berkovich spaces,

There are many such analogous theories!

1. Tate's theory of rigid analytic spaces,
2. Berkovich's theory of Berkovich spaces,
3. Huber's theory of Adic spaces,



There are many such analogous theories!

1. Tate's theory of rigid analytic spaces,
2. Berkovich's theory of Berkovich spaces,
3. Huber's theory of Adic spaces,
4. Raynaud's theory of formal schemes,

There are many such analogous theories!

1. Tate's theory of rigid analytic spaces,
2. Berkovich's theory of Berkovich spaces,
3. Huber's theory of Adic spaces,
4. Raynaud's theory of formal schemes,
5. Other approaches: Scholze, Fujiwara-Kato, etc.

There are many such analogous theories!

1. Tate's theory of rigid analytic spaces,
2. Berkovich's theory of Berkovich spaces,
3. Huber's theory of Adic spaces,
4. Raynaud's theory of formal schemes,
5. Other approaches: Scholze, Fujiwara-Kato, etc.

We will only talk about Tate's approach.

The function

$$f(z) = \begin{cases} 1 & \text{if } |z|_p \leq 1 \\ 0 & \text{if } |z|_p > 1 \end{cases}$$

is locally constant.

The function

$$f(z) = \begin{cases} 1 & \text{if } |z|_p \leq 1 \\ 0 & \text{if } |z|_p > 1 \end{cases}$$

is locally constant.

This is bad! Many complex analysis properties fail.

The function

$$f(z) = \begin{cases} 1 & \text{if } |z|_p \leq 1 \\ 0 & \text{if } |z|_p > 1 \end{cases}$$

is locally constant.

This is bad! Many complex analysis properties fail.

1. no analytic continuation,

The function

$$f(z) = \begin{cases} 1 & \text{if } |z|_p \leq 1 \\ 0 & \text{if } |z|_p > 1 \end{cases}$$

is locally constant.

This is bad! Many complex analysis properties fail.

1. no analytic continuation,
2. no maximum modulus principle,

The function

$$f(z) = \begin{cases} 1 & \text{if } |z|_p \leq 1 \\ 0 & \text{if } |z|_p > 1 \end{cases}$$

is locally constant.

This is bad! Many complex analysis properties fail.

1. no analytic continuation,
2. no maximum modulus principle,
3. no global power series expansion.



The function

$$f(z) = \begin{cases} 1 & \text{if } |z|_p \leq 1 \\ 0 & \text{if } |z|_p > 1 \end{cases}$$

is locally constant.

This is bad! Many complex analysis properties fail.

1. no analytic continuation,
2. no maximum modulus principle,
3. no global power series expansion.

We must eliminate these bad functions.

## Definition

Let  $K$  a (complete) non-archimedean field, define the Tate algebra

$$K\langle X_1, \dots, X_r \rangle = \left\{ \sum_{n \in \mathbb{N}^r} a_n \underline{X}^n : a_n \rightarrow 0 \text{ as } |n| \rightarrow \infty \right\}$$

An affinoid  $K$ -algebra is a quotient  $K\langle X_1, \dots, X_r \rangle / I$ .

Let  $A$  be an affinoid  $K$ -algebra.

Let  $A$  be an affinoid  $K$ -algebra.

An affinoid domain  $\mathrm{Sp}(A)$  is a  $G$ -ringed space which is,

Let  $A$  be an affinoid  $K$ -algebra.

An affinoid domain  $\mathrm{Sp}(A)$  is a  $G$ -ringed space which is,

1. built from maximal ideals of  $A$ ,

Let  $A$  be an affinoid  $K$ -algebra.

An affinoid domain  $\mathrm{Sp}(A)$  is a  $G$ -ringed space which is,

1. built from maximal ideals of  $A$ ,
2. equipped with a  $G$ -topology called the admissible topology,

Let  $A$  be an affinoid  $K$ -algebra.

An affinoid domain  $\mathrm{Sp}(A)$  is a  $G$ -ringed space which is,

1. built from maximal ideals of  $A$ ,
2. equipped with a  $G$ -topology called the admissible topology,
3. equipped with a sheaf of functions  $\mathcal{O}_{\mathrm{Sp}(A)}$ .

Let  $A$  be an affinoid  $K$ -algebra.

An affinoid domain  $\mathrm{Sp}(A)$  is a  $G$ -ringed space which is,

1. built from maximal ideals of  $A$ ,
2. equipped with a  $G$ -topology called the admissible topology,
3. equipped with a sheaf of functions  $\mathcal{O}_{\mathrm{Sp}(A)}$ .

A rigid analytic space defined by gluing affinoid domains, i.e. it is a  $G$ -ringed space locally isomorphic to affinoid domains.



Completely analogous to the archimedean case, every algebraic variety  $X/\mathbb{C}_p$  has an analytification  $X^{\text{an}}$  as a rigid analytic space.

Completely analogous to the archimedean case, every algebraic variety  $X/\mathbb{C}_p$  has an analytification  $X^{\text{an}}$  as a rigid analytic space.

However, there is generally no universal covers in rigid geometry, and almost nothing is simply connected: there is no uniformization!

For an elliptic curve  $E/\mathbb{C}$ ,

$$E \cong \mathbb{C}/\Lambda \cong \mathbb{C}/(\mathbb{Z} \oplus \tau\mathbb{Z})$$

For an elliptic curve  $E/\mathbb{C}$ ,

$$E \cong \mathbb{C}/\Lambda \cong \mathbb{C}/(\mathbb{Z} \oplus \tau\mathbb{Z})$$

The universal cover factors

$$\mathbb{C} \xrightarrow{e^{2\pi iz}} \mathbb{C}^\times \rightarrow \mathbb{C}^\times/q^\mathbb{Z} \cong E$$

where  $q = e^{2\pi i\tau}$  satisfies  $0 < |q| < 1$ .

Analogously, if  $q \in \mathbb{C}_p$  satisfies  $0 < |q|_p < 1$ , then the quotient

$$E = \mathbb{C}_p^\times / q^{\mathbb{Z}}$$

is an elliptic curve.

Analogously, if  $q \in \mathbb{C}_p$  satisfies  $0 < |q|_p < 1$ , then the quotient

$$E = \mathbb{C}_p^\times / q^{\mathbb{Z}}$$

is an elliptic curve. Moreover, by a theorem of Tate, an elliptic curve can be obtained this way iff it has bad reduction.

Rigid analytic geometry has been used in the following:

1. Moduli of curves,

Rigid analytic geometry has been used in the following:

1. Moduli of curves,
2. Mirror symmetry,



Rigid analytic geometry has been used in the following:

1. Moduli of curves,
2. Mirror symmetry,
3. Raynaud's proof of Abhyankar's conjecture (constructing finite étale covers of  $\mathbb{A}_{\mathbb{F}_p}^1$  with given Galois group),

Rigid analytic geometry has been used in the following:

1. Moduli of curves,
2. Mirror symmetry,
3. Raynaud's proof of Abhyankar's conjecture (constructing finite étale covers of  $\mathbb{A}_{\mathbb{F}_p}^1$  with given Galois group),

and many more.