Introduction

On March 28th, 2019, San Francisco based ride hailing company Lyft became a public company via IPO. It is expected that Lyft's major competitor Uber will follow the suit to be IPO soon. Peer-to-peer ridesharing become an important part of people's life everywhere. Ridesharing company implements a dynamic pricing model when calculating the trip cost. As an active ridesharing service customer, I have been interested in the pricing model those companies implemented for a while. The goal of this project is to develop a ridesharing service pricing model that allows customer like me to estimate the trip cost.

Dataset

During my research for related data, I was lucky enough to discover that City of Chicago is the first city to release the ride hailing related data to public. Therefore, I was able to obtain Trip information from Chicago Data Portal. The actual dataset contains 17.4 million rows as the portal started to collect data from November, 2018. For this project, a random sample of 1 million rows were selected and loaded into Jupyter notebook. Dataset contains the following 21 features:

Trip ID: A unique identifier for the trip.

Trip Start Timestamp: When the trip started, rounded to the nearest 15 minutes.

Trip End Timestamp: When the trip ended, rounded to the nearest 15 minutes.

Trip Seconds: Time of the trip in seconds. **Trip Miles:** Distance of the trip in miles.

Pickup Census Tract: The Census Tract where the trip began. This column often will be blank for locations outside Chicago.

Dropoff Census Tract: The Census Tract where the trip ended. This column often will be blank for locations outside Chicago.

Pickup Community Area: The Community Area where the trip began. This column will be blank for locations outside Chicago.

Dropoff Community Area: The Community Area where the trip ended. This column will be blank for locations outside Chicago.

Fare: The fare for the trip, rounded to the nearest \$2.50.

Tip: The tip for the trip, rounded to the nearest \$1.00. Cash tips will not be recorded.

Additional Charges: The taxes, fees, and any other charges for the trip.

Trip Total: Total cost of the trip. This is calculated as the total of the previous columns, including rounding.

Shared Trip Authorized: Whether the customer agreed to a shared trip with another customer, regardless of whether the customer was actually matched for a shared trip.

Trips Pooled: If customers were matched for a shared trip, how many trips, including this one, were pooled. All customer trips from the time the vehicle was empty until it was empty again contribute to this count, even if some customers were never present in the vehicle at the same time. Each trip making up the overall shared trip will have a separate record in this dataset, with the same value in this column.

Pickup Centroid Latitude: The latitude of the center of the pickup census tract or the community area if the census tract has been hidden for privacy. This column often will be blank for locations outside Chicago.

Pickup Centroid Longitude: The longitude of the center of the pickup census tract or the community area if the census tract has been hidden for privacy. This column often will be blank for locations outside Chicago.

Pickup Centroid Location: The location of the center of the pickup census tract or the community area if the census tract has been hidden for privacy. This column often will be blank for locations outside Chicago.

Dropoff Centroid Latitude: The latitude of the center of the dropoff census tract or the community area if the census tract has been hidden for privacy. This column often will be blank for locations outside Chicago.

Dropoff Centroid Longitude: The longitude of the center of the dropoff census tract or the community area if the census tract has been hidden for privacy. This column often will be blank for locations outside Chicago.

Dropoff Centroid Location: The location of the center of the dropoff census tract or the community area if the census tract has been hidden for privacy. This column often will be blank for locations outside Chicago.

Since position features are outside the project scope, they will not be included in the analysis. Trip Total is the sum of Fare, Tip, Additional Charge, however, the goal of this project is to analyze the pricing model, therefore, only Fare is the related feature that is needed in the analysis. A modified dataset is created by eliminating Trip ID, position features and money features except Fare. This narrows down the dataset to 9 features: Trip Start Timestamp, Trip End Timestamp, Trip Seconds, Trip Miles, Pickup Community Area, Dropoff Community Area, Fare, Shared Trip Authorized, Trips Pooled.

Data Cleaning

The initial data type of the modified dataset is shown as below:

Trip Start Timestamp obiect **Trip End Timestamp** object Trip Seconds float64 **Trip Miles** float64 Pickup Community Area float64 **Dropoff Community Area** float64 Fare float64 **Shared Trip Authorized** bool **Trips Pooled** int64

Since Pickup Community Area and Dropoff Community Area are actually categorical variables, those two features were converted to categorical with astype ('category'). Trip Start Timestamp and Trip End Timestamp were converted to datetime variables with to_datetime. The updated data type of the modified dataset is shown here:

Trip Start Timestampdatetime64[ns]Trip End Timestampdatetime64[ns]

Trip Seconds float64
Trip Miles float64
Pickup Community Area category
Dropoff Community Area float64
Fare float64
Shared Trip Authorized bool
Trips Pooled int64

After checking the missing data, Trip Seconds, Pickup Community Area, Dropoff Community Area has 155, 75005, 84427 missing data respectively. Based on the definition of Pickup and Dropoff Community Area, all the blanks indicated the area outside Chicago. Blanks will be replaced by 0 to represent area outside Chicago. Imputation was performed with missing values of Fare feature with average fare amount.

A new column which convert the dates to day of the week was created to find some interesting patterns throughout the week.

Exploratory Data Analysis

One of the important features of the dataset is whether the trip is carpooled or not. For this random selected dataset, there were more trips not carpooled as shown below graph.




Figure 1. Count of whether the trip was shared trip or not.

Fare, Trip Seconds, and Trip Miles has been drawn against the day of week to find some patterns in below three boxplots with adjusted y scale (Figure 2, Figure 3, Figure 4). No conclusion regarding the mean of the fare, trip seconds and trip miles against day of week can be drawn from below boxplot. More sophisticated statistical test will be needed to determine whether the difference among the mean is significant against day of week.

One-way ANOVA test has been applied to those pairs of features via below result tables (Table 1, Table 2, Table 3). For Fare against day of the week, overall the model is significant as F = 227.1 and p-value = $5.00e-291 \approx 0$. This indicated that there is significant difference among the average Fare for each day of the week. For Trip miles against day of the week, overall the model is also significant as F = 374.8 and p-value = 0. This indicated that there is significant difference among the average trip distance for each day of the week. Last, for Trip seconds against day of the week, overall the model is significant as F = 227.1 and p-value = $5.00e-291 \approx 0$. This indicated that there is significant difference among the average trip duration for each day of the week.




Figure 2. Boxplot of Fare in day of week

OLS Regression Results

Dep. Variable: Fare R-squared: 0.001 0.001 Model: OLS Adj. R-squared: Method: Least Squares F-statistic: 227.1 Date: Mon, 27 May 2019 Prob (F-statistic): 5.00e-291 Time: 18:27:07 Log-Likelihood: -3.7242e+06 1000000 AIC: No. Observations: 7.448e+06 Df Residuals: 999993 BIC: 7.449e+06

Df Model: 6
Covariance Type: nonrobust

	coef st	d err	t P> t	[0.02	5 0.975	5]	
Intercept	11.470	7 0.024	473.17	7 0.00	0 11.42	 23 11.5	18
C(weekday)[T.Mo	nday]	0.4088	0.039	10.605	0.000	0.333	0.484
C(weekday)[T.Sa	turday]	-0.6592	0.033	-19.684	0.000	-0.725	-0.594
C(weekday)[T.Su	nday]	-0.3159	0.035	-8.969	0.000	-0.385	-0.247
C(weekday)[T.Th	ursday]	0.3578	0.036	10.008	0.000	0.288	0.428
C(weekday)[T.Tue	esday]	0.1588	0.039	4.086	0.000	0.083	0.235
C(weekday)[T.We	ednesday]	0.1155	0.038	3.060	0.002	0.042	0.190
Omnibus: 896728.677 Durbin-Watson:						====== 26	======
Prob(Omnibus):		0.000 Jar	que-Bera	(JB):	1336419	85.008	
Skew:	3.78	0 Prob(JE	3):	(0.00		

Kurtosis: 59.127 Cond. No. 7.31

Table 1. ANOVA for Fare Against Weekday




Figure 3. Boxplot of Trip Miles in day of week

OLS Regression Results

Dep. Variable: Trip_miles R-squared: 0.002					
Model: OLS Adj. R-squared: 0.002					
Method: Least Squares F-statistic: 374.8					
Date: Mon, 27 May 2019 Prob (F-statistic): 0.00					
Time: 18:28:28 Log-Likelihood: -3.3731e+06					
No. Observations: 1000000 AIC: 6.746e+06					
Df Residuals: 999993 BIC: 6.746e+06					
Df Model: 6					
Covariance Type: nonrobust					
coef std err t P> t [0.025 0.975]					
Intercept 6.0293 0.017 353.325 0.000 5.996 6.063					
C(weekday)[T.Monday] 0.7158 0.027 26.379 0.000 0.663 0.769					
C(weekday)[T.Saturday] -0.3712 0.024 -15.747 0.000 -0.417 -0.325					

C(weekday)[T.Sunday] 0.1690 0.025 6.817 0.000 0.120 C(weekday)[T.Thursday] 0.3988 0.025 15.845 0.000 0.349 0.448 0.4363 0.490 C(weekday)[T.Tuesday] 0.027 15.949 0.000 0.383 C(weekday)[T.Wednesday] 0.3384 0.027 12.734 0.000 0.286 0.391

 Omnibus:
 840651.198
 Durbin-Watson:
 1.906

 Prob(Omnibus):
 0.000
 Jarque-Bera (JB):
 72648318.970

 Skew:
 3.567
 Prob(JB):
 0.00

 Kurtosis:
 44.142
 Cond. No.
 7.31

Table 2. ANOVA for Trip Distance against Weekday




Figure 4. Boxplot of Trip Seconds in day of week

OLS Regression Results

Dep. Variable: Trip seconds R-squared: 0.007 Model: OLS Adj. R-squared: 0.007 Method: Least Squares F-statistic: 1150. Date: Mon, 27 May 2019 Prob (F-statistic): 0.00 Time: 18:27:56 Log-Likelihood: -8.0925e+06 No. Observations: 1000000 AIC: 1.618e+07 Df Residuals: 999993 BIC: 1.619e+07

Df Model: 6
Covariance Type: nonrobust

coef std err t P>|t| [0.025 0.975]

Intercept 1127.6919 1.913 589.537 0.000 1123.943 1131.441 C(weekday)[T.Monday] -1.6480 3.042 -0.542 0.588 -7.610 4.314 C(weekday)[T.Saturday] -116.0739 2.642 -43.926 0.000 -121.253 -110.895

```
C(weekday)[T.Sunday] -136.0845
                                  2.779
                                        -48.973
                                                   0.000 -141.531 -130.638
C(weekday)[T.Thursday] 27.8426
                                                          22.313
                                                                    33.372
                                  2 821
                                          9 869
                                                  0.000
C(weekday)[T.Tuesday]
                                                          12.243
                                                                   24.263
                       18.2530
                                  3.066
                                          5.953
                                                  0.000
C(weekday)[T.Wednesday] 13.9122
                                   2.979
                                            4.670
                                                    0.000
                                                            8.073
                                                                    19.751
                                                      1.954
Omnibus:
                  612885 938 Durbin-Watson:
Prob(Omnibus):
                       0.000 Jarque-Bera (JB):
                                                  30620121.981
Skew:
                   2.287 Prob(JB):
                                                0.00
Kurtosis:
                   29.720 Cond. No.
                                                 7.31
```

Table 2 ANOVA for Trip Duration excinct Models


Table 3. ANOVA for Trip Duration against Weekday

Correlation Analysis

A scatter plot with regression line between Fare and Trip Miles was created to verify the linearity between those two variables. Day of the week was incorporated with different colors. From the graph shown below, there was no significant difference among day of the week. The linear relationship between those two variables could be verified. The Pearson Correlation Coefficient was calculated and shown here:

The Pearson Correlation Coefficient is: 0.8929456863732214
The p-value is: 0.0

As the p-value is very small < α = 0.05, we could reject the null hypothesis. The Pearson Correlation Coefficient showed a strong positive correlation between Fare and Trip Miles which was consistent with the regression plot. Generally as customer travel longer distance, the trip fare also goes up. Since there was strong evidence showing the correlation between Fare and Trip Miles, Trip Miles needed to be added to the model




Graph 5. Regression plot between Fare and Trip Miles. Day of Week in colors

Similar to Fare and Trip Miles, the linear relationship could also be observed between Fare and Trip Seconds from the regression plot below. Again, the plot incorporated day of week as different color to find any potential difference among the days. From the graph, there was no big difference among them. They all showed a positive correlation which could be verified with the Pearson Correlation Coefficient calculation results shown here.

The Pearson Correlation Coefficient is: 0.7850275437258065
The p-value is: 0.0

As the p-value is very small < α = 0.05, we could reject the null hypothesis. The Pearson Correlation Coefficient showed a strong positive correlation between Fare and Trip Seconds which was consistent with the regression plot. Generally as trip times goes up, the trip fare also goes up. Since there was strong evidence showing the correlation between Fare and Trip Seconds, Trip Seconds needed to be added to the model.




Graph 6. Regression plot between Fare and Trip Seconds. Day of Week in colors

To analyze the correlation between the categorical variables, chi-square test has been implemented among the three categorical variables (Shared Trip Authorized, Pickup Community Area, and Dropoff Community Area) verse weekday. The results has been summarized in below table. The p-values for all the pairs were 0, therefore, we can reject the null hypothesis. Those pairs of variables are indeed dependent on one another.


Weekday vs	Chi-Square Test value	p-value	Degree of Freedom
Shared Authorized Trip	3884.10	0.0	6
Pickup Community Area	16563.67	0.0	462
Dropoff Community Area	20776.08	0.0	462

Table 4. Chi-Square Test Results


To analyze the impact of time on Fare, Trip Seconds and Trip Miles, those variables are plotted against time that aggregated by 'Day' frequency using Python <u>resample</u> attribute. The result graphs are shown in Graph 7, Graph 8, Graph 9 respectively. For fare over time, no overall downward or upward trend was observed. The fluctuation of the fare over time seems random. However, there was a big spike right after Christmas Eve. For trip duration over time, a slightly overall downward trend was observed. Like fare over time, the trip duration also showed a random fluctuation throughout the period. For trip distance over time, no overall downward or upward trend can be found from the graph. Again, the fluctuation of the trip distance also seems random. However, the same big spike was also found right after Christmas Eve.



Graph 7. Fare over time



Graph 8. Trip Duration over time



Graph 9. Trip Distance over time