

Notes on ML Models

Part I

Linear Models and Neural Networks

1 Generalized Linear Models

A GLM is a conditional version of an exponential family distribution $p(y | \theta) = h(y) \exp(\theta^\top T(y) - A(\theta))$, in which the natural parameters are a linear function of the input. More precisely, the model has the following form:

$$p(y_n | \mathbf{x}_n, \mathbf{w}, \sigma^2) = \exp \left[\frac{y_n \eta_n - A(\eta_n)}{\sigma^2} + \log h(y_n, \sigma^2) \right]$$

where

- y_n is the output/response variable, $\mathbf{x}_n$ is the input/feature vector, $\mathbf{w}$ and σ is the learnable model parameter
- $\eta_n \triangleq \mathbf{w}^\top \mathbf{x}_n$ is the (input dependent) natural parameter or canonical parameter
- $A(\eta_n)$ is the log normalizer
- $\mathcal{T}(y_n) = y_n$ is the sufficient statistic, which is the smallest summary of the output data y that retains everything relevant for estimating the parameter θ – while it may still lose other information about y itself.
- σ^2 is the dispersion term
- $h(y_n, \sigma^2)$ is the scaling constant or base measure

We can show that the mean and variance of the response variable are as follows:

$$\int p(y_n | \eta_n, \sigma^2) dy_n = 1 \quad \text{for all } \eta_n$$

To derive the mean, differentiate the equation w.r.t to η_n

$$\begin{aligned} \int \frac{\partial p(y_n | \eta_n, \sigma^2)}{\partial \eta_n} dy_n &= 0 \\ \int p(y_n | \eta_n, \sigma^2) \cdot \frac{y_n - A'(\eta_n)}{\sigma^2} dy_n &= 0 \\ \frac{1}{\sigma^2} \left[\int y_n p(y_n | \eta_n, \sigma^2) dy_n - A'(\eta_n) \right] &= 0 \end{aligned}$$

thus, the **mean** is:

$$\mathbb{E}[y_n | \mathbf{x}_n, \mathbf{w}, \sigma^2] = A'(\eta_n) \triangleq \ell^{-1}(\eta_n)$$

where the mapping from the linear inputs to the mean of the output using $\mu_n = \ell^{-1}(\eta_n)$, and the function ℓ is known as the link function, and ℓ^{-1} is known as the mean function.

To derive the variance, differentiate the above equation w.r.t η_n again,

$$\begin{aligned} \int \left[-A''(\eta_n) \cdot p + (y_n - A'(\eta_n)) \cdot p \cdot \frac{y_n - A'(\eta_n)}{\sigma^2} \right] dy_n &= 0 \\ -A''(\eta_n) \int p dy_n + \frac{1}{\sigma^2} \int (y_n - A'(\eta_n))^2 p dy_n &= 0 \end{aligned}$$

thus, the **variance** is:

$$\mathbb{V}[y_n | \mathbf{x}_n, \mathbf{w}, \sigma^2] = A''(\eta_n) \sigma^2$$

1.1 Logistic regression

Binary logistic regression is used when $y_n \in \{0, 1\}$. It has the form

$$p(y_n | \mathbf{x}_n, \mathbf{w}) = \mu_n^{y_n} (1 - \mu_n)^{1-y_n}$$

where

$$\mu_n = \sigma(\eta_n) = \frac{1}{1 + \exp(-\eta_n)}, \quad \eta_n = \mathbf{w}^\top \mathbf{x}_n.$$

Hence

$$\log p(y_n | \mathbf{x}_n, \mathbf{w}) = y_n \log \mu_n + (1 - y_n) \log(1 - \mu_n)$$

Using $\mu_n = \exp(\eta_n)/(1 + \exp(\eta_n))$, in GLM form:

$$\log p(y_n | \mathbf{x}_n, \mathbf{w}) = y_n \eta_n - \log(1 + \exp(\eta_n))$$

We see that $A(\eta_n) = \log(1 + \exp(\eta_n))$ and hence

$$\mathbb{E}[y_n] = A'(\eta_n) = \frac{1}{1 + \exp(-\eta_n)}, \quad \mathbb{V}[y_n] = A''(\eta_n) = \mu_n(1 - \mu_n)$$

Thus the link function is the logit function, $\ell(\mu_n) = \log\left(\frac{\mu_n}{1-\mu_n}\right)$, and the mean function is the sigmoid function, $\ell^{-1}(\eta_n) = \sigma(\eta_n)$.

1.2 Linear regression

Linear regression has the form with $y_n \in \mathbb{R}$

$$p(y_n | \mathbf{x}_n, \mathbf{w}, \sigma^2) = \frac{1}{\sqrt{2\pi\sigma^2}} \exp\left(-\frac{1}{2\sigma^2}(y_n - \mathbf{w}^\top \mathbf{x}_n)^2\right)$$

Hence

$$\log p(y_n | \mathbf{x}_n, \mathbf{w}, \sigma^2) = -\frac{1}{2\sigma^2}(y_n - \eta_n)^2 - \frac{1}{2} \log(2\pi\sigma^2)$$

where $\eta_n = \mathbf{w}^\top \mathbf{x}_n$. In GLM form:

$$\log p(y_n | \mathbf{x}_n, \mathbf{w}, \sigma^2) = \frac{y_n \eta_n - \frac{\eta_n^2}{2}}{\sigma^2} - \frac{1}{2} \left(\frac{y_n^2}{\sigma^2} + \log(2\pi\sigma^2) \right)$$

We see that $A(\eta_n) = \eta_n^2/2$ and hence

$$\mathbb{E}[y_n] = \eta_n = \mathbf{w}^\top \mathbf{x}_n, \quad \mathbb{V}[y_n] = \sigma^2$$

1.3 Binomial regression

If the response variable is the number of successes in N_n trials, $y_n \in \{0, \dots, N_n\}$, we can use binomial regression, which is defined by

$$p(y_n | \mathbf{x}_n, N_n, \mathbf{w}) = \text{Bin}(y_n | \sigma(\mathbf{w}^\top \mathbf{x}_n), N_n)$$

note that binary logistic regression is the special case when $N_n = 1$.

The log pdf is given by

$$\begin{aligned}\log p(y_n \mid \mathbf{x}_n, N_n, \mathbf{w}) &= y_n \log \mu_n + (N_n - y_n) \log(1 - \mu_n) + \log \binom{N_n}{y_n} \\ &= y_n \log \left(\frac{\mu_n}{1 - \mu_n} \right) + N_n \log(1 - \mu_n) + \log \binom{N_n}{y_n}\end{aligned}$$

where $\mu_n = \sigma(\eta_n)$. To rewrite this in GLM form, let us define

$$\eta_n \triangleq \log \left(\frac{\mu_n}{1 - \mu_n} \right) = \log \left[\frac{1}{1 + e^{-\mathbf{w}^\top \mathbf{x}_n}} \frac{1 + e^{-\mathbf{w}^\top \mathbf{x}_n}}{e^{-\mathbf{w}^\top \mathbf{x}_n}} \right] = \log \left(\frac{1}{e^{-\mathbf{w}^\top \mathbf{x}_n}} \right) = \mathbf{w}^\top \mathbf{x}_n$$

Hence we can write binomial regression in GLM form as follows:

$$\log p(y_n \mid \mathbf{x}_n, N_n, \mathbf{w}) = y_n \eta_n - A(\eta_n) + h(y_n)$$

where

$$h(y_n) = \log \binom{N_n}{y_n}, \quad A(\eta_n) = -N_n \log(1 - \mu_n) = N_n \log(1 + e^{\eta_n})$$

Hence,

$$\mathbb{E}[y_n] = \frac{dA}{d\eta_n} = \frac{N_n e^{\eta_n}}{1 + e^{\eta_n}} = N_n \mu_n, \quad \mathbb{V}[y_n] = \frac{d^2 A}{d\eta_n^2} = N_n \mu_n (1 - \mu_n)$$

1.4 Poisson regression

If the response variable is an integer count, $y_n \in \{0, 1, \dots\}$, we can use Poisson regression, which is defined by

$$p(y_n \mid \mathbf{x}_n, \mathbf{w}) = \text{Poi}(y_n \mid \exp(\mathbf{w}^\top \mathbf{x}_n))$$

where

$$\text{Poi}(y \mid \mu) = e^{-\mu} \frac{\mu^y}{y!}$$

is the Poisson distribution.

The log pdf is given by

$$\log p(y_n \mid \mathbf{x}_n, \mathbf{w}) = y_n \log \mu_n - \mu_n - \log(y_n!)$$

where $\mu_n = \exp(\mathbf{w}^\top \mathbf{x}_n)$. Hence in GLM form we have

$$\log p(y_n \mid \mathbf{x}_n, \mathbf{w}) = y_n \eta_n - A(\eta_n) + h(y_n)$$

where $\eta_n = \log(\mu_n) = \mathbf{w}^\top \mathbf{x}_n$, $A(\eta_n) = \mu_n = e^{\eta_n}$, and $h(y_n) = -\log(y_n!)$. Hence

$$\mathbb{E}[y_n] = \frac{dA}{d\eta_n} = e^{\eta_n} = \mu_n, \quad \mathbb{V}[y_n] = \frac{d^2 A}{d\eta_n^2} = e^{\eta_n} = \mu_n$$

1.5 Canonical and non-canonical link functions

Link functions connect three quantities:

- θ_n — the **natural parameter** of the *exponential family distribution*.
- $\mu_n = \mathbb{E}[y_n]$ — the **mean** of the output distribution.
- $\eta_n = \mathbf{w}^\top \mathbf{x}_n$ — the **linear predictor**, built from the *inputs* and weights.

The two connecting functions:

$$\theta_n \xrightarrow{A'} \mu_n \xrightarrow{\ell} \eta_n$$

- A' (derivative of the log-partition function) always maps $\theta_n \rightarrow \mu_n$. This relationship is fixed by the choice of exponential-family distribution (Bernoulli, Gaussian, Poisson, ...) and never changes.
- ℓ , the **link function**, maps $\mu_n \rightarrow \eta_n$. This is the function *you choose* when designing the GLM. Its inverse, ℓ^{-1} , is the **mean function**: $\mu_n = \ell^{-1}(\eta_n)$ — the direction actually used to generate predictions from η_n .

Canonical link. Choosing $\ell = (A')^{-1}$ makes the link exactly undo A' :

$$\eta_n = g(\mu_n) = (A')^{-1}(A'(\theta_n)) = \theta_n$$

so the natural parameter and linear predictor coincide: $\theta_n = \eta_n$.

Non-canonical link. Choosing any $g \neq (A')^{-1}$ gives

$$\eta_n = g(A'(\theta_n)) \neq \theta_n \quad \text{in general.}$$

The mean is still recovered correctly via $\mu_n = g^{-1}(\eta_n)$, but the natural parameter θ_n is no longer equal to the linear predictor η_n —recovering it requires the separate map $(A')^{-1}(\mu_n)$.

Examples of links:

- Logit: canonical link $\theta_n = \ell(\mu_n) = \log \frac{\mu_n}{1-\mu_n}$.
- Complementary log-log: non-canonical link $\eta_n = \ell(\mu_n) = \log(-\log(1 - \mu_n))$

1.6 Maximum likelihood estimation

GLMs can be fit the negative log-likelihood as the objective function (ignoring constant term h):

$$\text{NLL}(\mathbf{w}) = -\log p(\mathcal{D} \mid \mathbf{w}) = -\frac{1}{\sigma^2} \sum_{n=1}^N \ell_n = -\frac{1}{\sigma^2} \sum_{n=1}^N [\eta_n y_n - A(\eta_n)]$$

where $\eta_n = \mathbf{w}^\top \mathbf{x}_n$.

To find the MLE, we must solve $\nabla \text{NLL}(\mathbf{w}) = \mathbf{g}(\mathbf{w}) = 0$. For notational simplicity, we will assume $\sigma^2 = 1$. The gradient for a single term is:

$$\begin{aligned} \mathbf{g}_n &= -\frac{1}{\sigma^2} \frac{\partial \ell_n}{\partial \mathbf{w}} = -\frac{1}{\sigma^2} \frac{\partial \ell_n}{\partial \eta_n} \frac{\partial \eta_n}{\partial \mathbf{w}} \\ &= -\frac{1}{\sigma^2} (y_n - A'(\eta_n)) \mathbf{x}_n = -\frac{1}{\sigma^2} (y_n - \mu_n) \mathbf{x}_n \end{aligned}$$

where $\mu_n = f(\mathbf{w}^\top \mathbf{x}_n)$, and f is the inverse link function that maps from canonical parameters to mean parameters. If we interpret $(y_n - \mu_n)$ as an error signal, then the gradient $-\frac{1}{\sigma^2} \sum_{n=1}^N (y_n - \mu_n) \mathbf{x}_n$ weights each input $\mathbf{x}_n$ by its error, and averages the result.

Gradient-based optimization will find a stationary points where $\mathbf{g}(\mathbf{w}) = 0$, we need to examine the Hessian matrix to determine whether it is the global optimum. The Hessian is given by

$$\begin{aligned} \mathbf{H} &= \frac{\partial^2}{\partial \mathbf{w} \partial \mathbf{w}^\top} \text{NLL}(\mathbf{w}) \\ &= - \sum_{n=1}^N \frac{\partial \mathbf{g}_n}{\partial \mathbf{w}^\top} = - \sum_{n=1}^N \frac{\partial \mathbf{g}_n}{\partial \mu_n} \frac{\partial \mu_n}{\partial \mathbf{w}^\top} = - \sum_{n=1}^N \mathbf{x}_n f'(\mathbf{w}^\top \mathbf{x}_n) \mathbf{x}_n^\top \end{aligned}$$

In general, the Hessian is positive definite, since $f'(\mathbf{w}^\top \mathbf{x}_n) > 0$, hence the negative log-likelihood is convex, so the MLE for a GLM is unique given $f(\mathbf{w}^\top \mathbf{x}_n) > 0$ for all n .

Based on these results, gradient-based optimizers can be used to estimate the model weights.

1.6.1 Stochastic gradient descent

The goal is to solve

$$\hat{\mathbf{w}} \triangleq \underset{\mathbf{w}}{\operatorname{argmin}} \mathcal{L}(\mathbf{w}) = \underset{\mathbf{w}}{\operatorname{argmin}} \text{NLL}(\mathbf{w})$$

One of the methods is stochastic gradient descent: we use a minibatch from the whole dataset of size N , then we get the following update equation:

$$\mathbf{w}_{t+1} = \mathbf{w}_t - \alpha_t \nabla_{\mathbf{w}} \text{NLL}(\mathbf{w}_t) = \mathbf{w}_t - \alpha_t (\mu_n - y_n) \mathbf{x}_n$$

where we replaced the average over all N samples with a stochastically chosen minibatch size at each step t .

Since we know the objective is convex, then it can be shown that this will converge to a global optimum given the learning rate α_t is decayed appropriately at each step t .

Part II

Dimentionality Reduction

2 Principal Component Analysis

PCA algorithm is to find a **linear and orthogonal** projection of the high dimensional data $x \in \mathbb{R}^D$, to a low dimensional subspace $z \in \mathbb{R}^L$, such that the low dimensional representation is a "good approximation" to the original data. If we project or encode $\mathbf{x}$ to get $\mathbf{z} = \mathbf{W}^T \mathbf{x}$, and then unproject or decode $\mathbf{z}$ to get $\hat{\mathbf{x}} = \mathbf{W} \mathbf{z}$, then we want $\hat{\mathbf{x}}$ to be close to $\mathbf{x}$ in ℓ_2 distance. In particular, we can define the following **reconstruction error**:

$$\mathcal{L}(\mathbf{W}) \triangleq \frac{1}{N} \sum_{n=1}^N \|\mathbf{x}_n - \text{decode}(\text{encode}(\mathbf{x}_n; \mathbf{W}); \mathbf{W})\|_2^2$$

PCA algorithm states that this objective can be minimized by setting $\mathbf{W} = \mathbf{U}_L$, where $\mathbf{U}_L$ contains the the L eigenvectors with lages eigenvalues of the empirical covariance matrix

$$\hat{\Sigma} = \frac{1}{N} \sum_{n=1}^N (\mathbf{x}_n - \bar{\mathbf{x}})(\mathbf{x}_n - \bar{\mathbf{x}})^T = \frac{1}{N} \mathbf{X}_c^T \mathbf{X}_c$$

where $\mathbf{X}_c$ is a centered version of the $N \times D$ design matrix.

2.1 PCA setup

Suppose we have an (unlabeled) dataset $\mathcal{D} = \{\mathbf{x}_n : n = 1 : N\}$, where $\mathbf{x}_n \in \mathbb{R}^D$. We can represent this as an $N \times D$ data matrix $\mathbf{X}$. We will assume $\bar{\mathbf{x}} = \frac{1}{N} \sum_{n=1}^N \mathbf{x}_n = \mathbf{0}$, which we can ensure by centering the data.

We would like to approximate each $\mathbf{x}_n$ by a low dimensional representation, $\mathbf{z}_n \in \mathbb{R}^L$. We assume that each $\mathbf{x}_n$ can be explained in terms of a weighted combination of basis functions $\mathbf{w}_1, \dots, \mathbf{w}_L$, where each $\mathbf{w}_k \in \mathbb{R}^D$, and where the weights are given by $\mathbf{z}_n \in \mathbb{R}^L$, i.e.

$$\mathbf{x}_n \approx \sum_{k=1}^L z_{nk} \mathbf{w}_k$$

The vector $\mathbf{z}_n$ is the low dimensional representation of $\mathbf{x}_n$, and is known as the **latent vector**, the collection of these latent variables are called the **latent factors**.

The error produced by this approximation:

$$\mathcal{L}(\mathbf{W}, \mathbf{Z}) = \frac{1}{N} \|\mathbf{X} - \mathbf{Z} \mathbf{W}^T\|_F^2 = \frac{1}{N} \|\mathbf{X}^T - \mathbf{W} \mathbf{Z}^T\|_F^2 = \frac{1}{N} \sum_{n=1}^N \|\mathbf{x}_n - \mathbf{W} \mathbf{z}_n\|^2$$

where

- $\mathbf{X} \in \mathbb{R}^{N \times D}$: data matrix, rows are centered data points $\mathbf{x}_n^T$
- $\mathbf{Z} \in \mathbb{R}^{N \times L}$: latent code matrix, rows are centered data points $\mathbf{z}_n^T$
- $\mathbf{W} \in \mathbb{R}^{D \times L}$: basis matrix, columns $\mathbf{w}_1, \dots, \mathbf{w}_L$ are orthonormal
- approximating each $\mathbf{x}_n$ by $\hat{\mathbf{x}}_n = \mathbf{W} \mathbf{z}_n$; or, $\hat{\mathbf{X}} = \mathbf{Z} \mathbf{W}^T$.
- the minimization of $\mathcal{L}$ is subject to constraints $\mathbf{w}_k^T \mathbf{w}_j = 0$ for $k \neq j$; 1 for $k = j$, i.e., $\mathbf{W}$ is orthogonal

2.2 Algorithm derivation

Base case

Starting by considering the best 1d solution, $\mathbf{w}_1 \in \mathbb{R}^D$, letting the coefficients for each data-points associated with the first basis vector be denoted by $\tilde{\mathbf{z}}_1 = [z_{11}, \dots, z_{N1}] \in \mathbb{R}^N$. The reconstruction error is

$$\begin{aligned}\mathcal{L}(\mathbf{w}_1, \tilde{\mathbf{z}}_1) &= \frac{1}{N} \sum_{n=1}^N \|\mathbf{x}_n - z_{n1} \mathbf{w}_1\|^2 = \frac{1}{N} \sum_{n=1}^N (\mathbf{x}_n - z_{n1} \mathbf{w}_1)^\top (\mathbf{x}_n - z_{n1} \mathbf{w}_1) \\ &= \frac{1}{N} \sum_{n=1}^N [\mathbf{x}_n^\top \mathbf{x}_n - 2z_{n1} \mathbf{w}_1^\top \mathbf{x}_n + z_{n1}^2 \mathbf{w}_1^\top \mathbf{w}_1] \\ &= \frac{1}{N} \sum_{n=1}^N [\mathbf{x}_n^\top \mathbf{x}_n - 2z_{n1} \mathbf{w}_1^\top \mathbf{x}_n + z_{n1}^2] \quad \text{since } \mathbf{w}_1^\top \mathbf{w}_1 = 1\end{aligned}$$

Taking derivatives wrt z_{n1} and equating to zero gives

$$\frac{\partial}{\partial z_{n1}} \mathcal{L}(\mathbf{w}_1, \tilde{\mathbf{z}}_1) = \frac{1}{N} [-2\mathbf{w}_1^\top \mathbf{x}_n + 2z_{n1}] = 0 \Rightarrow z_{n1} = \mathbf{w}_1^\top \mathbf{x}_n$$

So the optimal embedding is obtained by orthogonally projecting the data onto $\mathbf{w}_1$. Plugging this back in gives the loss for the weights:

$$\mathcal{L}(\mathbf{w}_1) = \mathcal{L}(\mathbf{w}_1, \tilde{\mathbf{z}}_1^*(\mathbf{w}_1)) = \frac{1}{N} \sum_{n=1}^N [\mathbf{x}_n^\top \mathbf{x}_n - z_{n1}^2] = \text{const} - \frac{1}{N} \sum_{n=1}^N z_{n1}^2$$

To solve for $\mathbf{w}_1$, note that

$$\mathcal{L}(\mathbf{w}_1) = -\frac{1}{N} \sum_{n=1}^N z_{n1}^2 = -\frac{1}{N} \sum_{n=1}^N \mathbf{w}_1^\top \mathbf{x}_n \mathbf{x}_n^\top \mathbf{w}_1 = -\mathbf{w}_1^\top \hat{\Sigma} \mathbf{w}_1$$

where Σ is the empirical covariance matrix (since we assumed the data is centered). We can trivially optimize this by letting $\|\mathbf{w}_1\| \rightarrow \infty$, so we impose the constraint $\|\mathbf{w}_1\| = 1$ and instead optimize

$$\tilde{\mathcal{L}}(\mathbf{w}_1) = \mathbf{w}_1^\top \hat{\Sigma} \mathbf{w}_1 - \lambda_1 (\mathbf{w}_1^\top \mathbf{w}_1 - 1)$$

where λ_1 is a Lagrange multiplier. Taking derivatives and equating to zero we have

$$\frac{\partial}{\partial \mathbf{w}_1} \tilde{\mathcal{L}}(\mathbf{w}_1) = 2\hat{\Sigma} \mathbf{w}_1 - 2\lambda_1 \mathbf{w}_1 = 0$$

$$\hat{\Sigma} \mathbf{w}_1 = \lambda_1 \mathbf{w}_1$$

Hence the optimal direction onto which we should project the data is an eigenvector of the covariance matrix. Left multiplying by $\mathbf{w}_1^\top$ (and using $\mathbf{w}_1^\top \mathbf{w}_1 = 1$) we find

$$\mathbf{w}_1^\top \hat{\Sigma} \mathbf{w}_1 = \lambda_1$$

Since we want to maximize this quantity (minimize the loss), we pick the eigenvector which corresponds to the largest eigenvalue.

Induction Step

For another direction $\mathbf{w}_2$, the reconstruction error is

$$\mathcal{L}(\mathbf{w}_1, \tilde{\mathbf{z}}_1, \mathbf{w}_2, \tilde{\mathbf{z}}_2) = \frac{1}{N} \sum_{n=1}^N \|\mathbf{x}_n - z_{n1}\mathbf{w}_1 - z_{n2}\mathbf{w}_2\|^2$$

Optimizing wrt $\mathbf{w}_1$ and $\mathbf{z}_1$ gives the same solution as before. $\frac{\partial \mathcal{L}}{\partial \mathbf{z}_2} = 0$ yields $z_{n2} = \mathbf{w}_2^\top \mathbf{x}_n$. Substituting in yields

$$\mathcal{L}(\mathbf{w}_2) = \frac{1}{N} \sum_{n=1}^N [\mathbf{x}_n^\top \mathbf{x}_n - \mathbf{w}_1^\top \mathbf{x}_n \mathbf{x}_n^\top \mathbf{w}_1 - \mathbf{w}_2^\top \mathbf{x}_n \mathbf{x}_n^\top \mathbf{w}_2] = \text{const} - \mathbf{w}_2^\top \hat{\Sigma} \mathbf{w}_2$$

Dropping the constant term, plugging in the optimal $\mathbf{w}_1$ and adding the constraints yields

$$\tilde{\mathcal{L}}(\mathbf{w}_2) = -\mathbf{w}_2^\top \hat{\Sigma} \mathbf{w}_2 + \lambda_2(\mathbf{w}_2^\top \mathbf{w}_2 - 1) + \lambda_{12}(\mathbf{w}_2^\top \mathbf{w}_1 - 0)$$

with the extra constraint term, it can be shown that the solution is given by the eigenvector with the second largest eigenvalue:

$$\hat{\Sigma} \mathbf{w}_2 = \lambda_2 \mathbf{w}_2$$

The proof continues in this way to show that $\hat{\mathbf{W}} = \mathbf{U}_L$.

2.3 Computational tricks

1. **High-dimensional data.** PCA finds the eigenvectors of the $D \times D$ covariance matrix $\mathbf{X}^\top \mathbf{X}$. If $D > N$, it is faster to work with the $N \times N$ matrix $\mathbf{X}\mathbf{X}^\top$. Let $\mathbf{U}$ be an orthonormal matrix containing the eigenvectors of $\mathbf{X}\mathbf{X}^\top$ with corresponding eigenvalues in Λ . By definition we have $(\mathbf{X}\mathbf{X}^\top)\mathbf{U} = \mathbf{U}\Lambda$. Pre-multiplying by $\mathbf{X}^\top$ gives

$$(\mathbf{X}^\top \mathbf{X})(\mathbf{X}^\top \mathbf{U}) = (\mathbf{X}^\top \mathbf{U})\Lambda$$

from which we see that the eigenvectors of $\mathbf{X}^\top \mathbf{X}$ are $\mathbf{V} = \mathbf{X}^\top \mathbf{U}$, with eigenvalues Λ as before. However, these eigenvectors are not normalized, since $\|\mathbf{v}_j\|^2 = \mathbf{u}_j^\top \mathbf{X}\mathbf{X}^\top \mathbf{u}_j = \lambda_j \mathbf{u}_j^\top \mathbf{u}_j = \lambda_j$. The normalized eigenvectors are given by

$$\mathbf{V} = \mathbf{X}^\top \mathbf{U} \Lambda^{-\frac{1}{2}}$$

2. **Computing PCA using SVD.** Let $\mathbf{U}_\Sigma \Lambda_\Sigma \mathbf{U}_\Sigma^\top$ be the top L eigendecomposition of the covariance matrix $\Sigma \propto \mathbf{X}^\top \mathbf{X}$ (assume $\mathbf{X}$ is centered). Recall the optimal estimate of the projection weights $\mathbf{W}$ is given by the top L eigenvalues, so $\mathbf{W} = \mathbf{U}_\Sigma$. Now let $\mathbf{U}_X \mathbf{S}_X \mathbf{V}_X^\top \approx \mathbf{X}$ be the L -truncated SVD approximation to the data matrix $\mathbf{X}$. By SVD definition, the columns of $\mathbf{V}_X$ are the eigenvectors of $\mathbf{X}^\top \mathbf{X}$, so

$$\mathbf{V}_X = \mathbf{U}_\Sigma = \mathbf{W}.$$

In addition, the eigenvalues of the covariance matrix are related to the singular values of the data matrix via $\lambda_k = s_k^2/N$. Now suppose we are interested in the principal components (or PC scores), rather than the projection matrix. We have

$$\mathbf{Z} = \mathbf{X}\mathbf{W} = \mathbf{U}_X \mathbf{S}_X \mathbf{V}_X^\top \mathbf{V}_X = \mathbf{U}_X \mathbf{S}_X$$

Finally, if we want to approximately reconstruct the data, we have

$$\hat{\mathbf{X}} = \mathbf{Z}\mathbf{W}^\top = \mathbf{U}_X \mathbf{S}_X \mathbf{V}_X^\top$$

This is precisely the same as a **truncated SVD approximation**.

3. **Latent dimentions selection.** Recall that the reconstruction error can be written in terms of eigenvalues

$$\mathcal{L}_L = \frac{1}{|\mathcal{D}|} \sum_{n \in \mathcal{D}} \|\mathbf{x}_n - \hat{\mathbf{x}}_n\|^2 = \sum_{j=L+1}^D \lambda_j$$

where $\hat{\mathbf{x}}_n = \mathbf{W}\mathbf{z}_n + \mu$ and μ is the empirical mean. Thus, as the number of dimentions incrcases, the eigenvalues get smaller, and so does the reconstruction error. Therefore, if we give it more latent dimenions, it will be able to approximate the test date more accurately.

Thus, PCA can be performed either using an eigendecomposition of $\mathbf{\Sigma}$ or an SVD decomposition of $\mathbf{X}$. The latter is often preferable, for computational reasons.

3 Factor Analysis

Factor analysis is a generalization of PCA corresponds to the following linear-Gaussian latent variable probabilistic **generative** model:

$$\mathbf{x} = \mathbf{W}\mathbf{z} + \boldsymbol{\mu} + \boldsymbol{\epsilon}, \quad \boldsymbol{\epsilon} \sim \mathcal{N}(\mathbf{0}, \boldsymbol{\Psi})$$

$$p(\mathbf{z}) = \mathcal{N}(\mathbf{z} \mid \boldsymbol{\mu}_0, \boldsymbol{\Sigma}_0)$$

$$p(\mathbf{x} \mid \mathbf{z}, \boldsymbol{\theta}) = \mathcal{N}(\mathbf{x} \mid \mathbf{W}\mathbf{z} + \boldsymbol{\mu}, \boldsymbol{\Psi})$$

where $\mathbf{W}$ is a $D \times L$ matrix, known as the **factor loading matrix**, $\boldsymbol{\Psi}$ is a diagonal $D \times D$ noise covariance matrix, and $\boldsymbol{\mu}$ is a $D \times 1$ mean of observed data

FA can be thought of as a low-rank version of a Gaussian distribution. To see this, note that the induced marginal distribution $p(\mathbf{x} \mid \boldsymbol{\theta})$ is a Gaussian:

$$\begin{aligned} p(\mathbf{x} \mid \boldsymbol{\theta}) &= \int \mathcal{N}(\mathbf{x} \mid \mathbf{W}\mathbf{z} + \boldsymbol{\mu}, \boldsymbol{\Psi}) \mathcal{N}(\mathbf{z} \mid \boldsymbol{\mu}_0, \boldsymbol{\Sigma}_0) d\mathbf{z} \\ &= \mathcal{N}(\mathbf{x} \mid \mathbf{W}\boldsymbol{\mu}_0 + \boldsymbol{\mu}, \boldsymbol{\Psi} + \mathbf{W}\boldsymbol{\Sigma}_0\mathbf{W}^\top) \end{aligned}$$

Hence $\mathbb{E}[\mathbf{x}] = \mathbf{W}\boldsymbol{\mu}_0 + \boldsymbol{\mu}$ and $\text{Cov}[\mathbf{x}] = \mathbf{W}\text{Cov}[\mathbf{z}]\mathbf{W}^\top + \boldsymbol{\Psi} = \mathbf{W}\boldsymbol{\Sigma}_0\mathbf{W}^\top + \boldsymbol{\Psi}$. Without loss of generality, we can set $\boldsymbol{\mu}_0 = \mathbf{0}$ and $\boldsymbol{\Sigma}_0 = \mathbf{I}$. Thus,

$$p(\mathbf{z}) = \mathcal{N}(\mathbf{z} \mid \mathbf{0}, \mathbf{I})$$

$$p(\mathbf{x} \mid \mathbf{z}) = \mathcal{N}(\mathbf{x} \mid \mathbf{W}\mathbf{z} + \boldsymbol{\mu}, \boldsymbol{\Psi})$$

$$p(\mathbf{x}) = \mathcal{N}(\mathbf{x} \mid \boldsymbol{\mu}, \mathbf{W}\mathbf{W}^\top + \boldsymbol{\Psi})$$

where FA approximates the covariance matrix of the visible vector using a low-rank decomposition $\mathbf{C} = \text{Cov}[\mathbf{x}] = \mathbf{W}\mathbf{W}^\top + \boldsymbol{\Psi}$. The marginal variance of each visible variable is given by $\mathbb{V}[x_d] = \sum_{k=1}^L w_{dk}^2 + \psi_d$ where the first term is the variance due to the common factors, and the second ψ_d term is dimension-specific uniqueness.

Using EM algorithm to estimate the parameters $\mathbf{W}, \boldsymbol{\mu}, \boldsymbol{\Psi}$, then we can compute probabilistic latent embeddings using $p(\mathbf{z} \mid \mathbf{x})$ using Bayes rule for Gaussians,

$$p(\mathbf{z} \mid \mathbf{x}) = \mathcal{N}(\mathbf{z} \mid \mathbf{W}^\top \mathbf{C}^{-1}(\mathbf{x} - \boldsymbol{\mu}), \mathbf{I} - \mathbf{W}^\top \mathbf{C}^{-1} \mathbf{W})$$

3.1 Prerequisite: evidence lower bound (ELBO)

Let data x been generated by some latent variable z through a process $p(z) \rightarrow p(x \mid z)$.

- To compute the marginal likelihood:

$$p(x) = \int p(x \mid z) p(z) dz$$

which tells how well the model explains the data, and should be maximized during training. However, the integral is usually intractable (eg. $p(x \mid z)$ is a neural network).

- To compute posterior $p(z \mid x)$ (i.e. given this data point, what latent value likely produced it). By Bayes' rule,

$$p(z \mid x) = \frac{p(x \mid z)p(z)}{p(x)}$$

and this also requires the intractable $p(x)$ in the denominator.

Since $p(z | x)$ cannot be computed exactly, an approximate posterior $q(z | x)$ is chosen and try to make it as close as possible to the true posterior, measured by KL divergence:

$$D_{KL}(q(z | x) \| p(z | x)) \geq 0$$

expand from the definition of KL divergence and re-arrange,

$$\begin{aligned} \log p(x) &= \underbrace{\mathbb{E}_{q(z|x)}[\log p(x | z)] - D_{KL}(q(z | x) \| p(z))}_{\text{ELBO}} + D_{KL}(q(z | x) \| p(z | x)) \\ &= \mathcal{L}(x) + D_{KL}(q(z|x) \| p(z|x)) \end{aligned}$$

for which,

$$\log p(x) \geq \mathcal{L}(x) \quad \text{since } D_{KL}(\cdot \| \cdot) \geq 0$$

ELBO is an useful variational trick,

- $\log p(x)$ itself is intractable, but $\mathcal{L}(x)$ only involves $q(z | x)$, $p(z)$, and $p(x | z)$ — all things you choose/model directly.
- Maximizing the ELBO does two things simultaneously: it pushes $\log p(x)$ up (better data likelihood), and it pushes $q(z | x)$ closer to the true posterior $p(z | x)$ (since the gap between ELBO and $\log p(x)$ is that KL term).

In the EM algorithm for FA below, expressing ELBO equivalently as $\mathcal{L} = \mathbb{E}_q[\log p(x, z)] - \mathbb{E}_q[\log q(z)]$, according to the lower bound theorem above, maximizing $\log p(x)$ can be done by maximizing $\mathbb{E}_q[\log p(x, z)]$.

3.2 EM algorithm for FA

The FA generative model:

$$p(\mathbf{z}_i) = \mathcal{N}(\mathbf{z}_i | \mathbf{0}, \mathbf{I}), \quad p(\mathbf{x}_i | \mathbf{z}_i) = \mathcal{N}(\mathbf{x}_i | \mathbf{W}\mathbf{z}_i + \boldsymbol{\mu}, \boldsymbol{\Psi})$$

with parameters $\boldsymbol{\theta} = (\mathbf{W}, \boldsymbol{\mu}, \boldsymbol{\Psi})$ shared across all datapoints $i = 1, \dots, N$.

E-step: compute the posterior $p(\mathbf{z}_i | \mathbf{x}_i, \boldsymbol{\theta})$. Because both distributions above are Gaussian and linearly related, this posterior is available in closed form by completing the square:

$$p(\mathbf{z}_i | \mathbf{x}_i, \boldsymbol{\theta}) = \mathcal{N}(\mathbf{z}_i | \mathbf{m}_i, \boldsymbol{\Sigma}_i)$$

$$\boldsymbol{\Sigma}_i = (\mathbf{I}_L + \mathbf{W}^T \boldsymbol{\Psi}^{-1} \mathbf{W})^{-1} \in \mathbb{R}^{L \times L}, \quad \mathbf{m}_i = \boldsymbol{\Sigma}_i \mathbf{W}^T \boldsymbol{\Psi}^{-1} (\mathbf{x}_i - \boldsymbol{\mu}) \in \mathbb{R}^{L \times 1}$$

Note $\boldsymbol{\Sigma}_i$ does not depend on i in practice (no $\mathbf{x}_i$ term appears), so it can be computed once per iteration.

M-step: maximize the expected complete-data log-likelihood $\mathbb{E}[\log p(\mathbf{x}_i, \mathbf{z}_i | \boldsymbol{\theta})]$. It is convenient to estimate $\mathbf{W}$ and $\boldsymbol{\mu}$ jointly by defining

$$\tilde{\mathbf{W}} = (\mathbf{W}, \boldsymbol{\mu}), \quad \tilde{\mathbf{z}} = (\mathbf{z}, 1), \quad \text{so that } \mathbf{x} = \mathbf{W}\mathbf{z} + \boldsymbol{\mu} = \tilde{\mathbf{W}}\tilde{\mathbf{z}}$$

Define:

$$\begin{aligned} \mathbf{b}_i &\triangleq \mathbb{E}[\tilde{\mathbf{z}}_i | \mathbf{x}_i] = [\mathbf{m}_i; 1] \\ \mathbf{C}_i &\triangleq \mathbb{E}[\tilde{\mathbf{z}}_i \tilde{\mathbf{z}}_i^T | \mathbf{x}_i] = \begin{pmatrix} \mathbb{E}[\mathbf{z}_i \mathbf{z}_i^T | \mathbf{x}_i] & \mathbb{E}[\mathbf{z}_i | \mathbf{x}_i] \\ \mathbb{E}[\mathbf{z}_i | \mathbf{x}_i]^T & 1 \end{pmatrix} = \begin{pmatrix} \boldsymbol{\Sigma}_i + \mathbf{m}_i \mathbf{m}_i^T & \mathbf{m}_i \\ \mathbf{m}_i^T & 1 \end{pmatrix} \end{aligned}$$

Maximizing the expected complete-data log-likelihood with respect to $\tilde{\mathbf{W}}$ and Ψ (a weighted-least-squares-type problem, since it is quadratic in $\tilde{\mathbf{W}}$ given $\mathbf{b}_i, \mathbf{C}_i$) yields:

$$\begin{aligned} \arg \max_{\boldsymbol{\theta}} \sum_i \mathbb{E}[\log p(\mathbf{x}_i, \mathbf{z}_i \mid \boldsymbol{\theta})] &= \arg \max_{\boldsymbol{\theta}} \sum_i \mathbb{E}[\log p(\mathbf{z}_i) + \log p(\mathbf{x}_i \mid \mathbf{z}_i, \boldsymbol{\theta})] \\ &= \arg \max_{\boldsymbol{\theta}} \sum_i \mathbb{E}[\log p(\mathbf{x}_i \mid \mathbf{z}_i, \boldsymbol{\theta})] \\ &= \arg \max_{\boldsymbol{\theta}} \left\{ -\frac{N}{2} \log |\Psi| - \frac{1}{2} \sum_{i=1}^N \left[\mathbf{x}_i^\top \Psi^{-1} \mathbf{x}_i - 2 \mathbf{x}_i^\top \Psi^{-1} \tilde{\mathbf{W}} \mathbf{b}_i + \text{tr}(\tilde{\mathbf{W}}^\top \Psi^{-1} \tilde{\mathbf{W}} \mathbf{C}_i) \right] \right\} \end{aligned}$$

Solve for $\boldsymbol{\theta} = (\tilde{\mathbf{W}}, \Psi)$ by setting $\partial Q / \partial \tilde{\mathbf{W}} = 0$ to get $\hat{\tilde{\mathbf{W}}}$, then substituting back and setting $\partial Q / \partial \Psi^{-1} = 0$ to get $\hat{\Psi}$,

$$\begin{aligned} \hat{\tilde{\mathbf{W}}} &= \left[\sum_i \mathbf{x}_i \mathbf{b}_i^\top \right] \left[\sum_i \mathbf{C}_i \right]^{-1} \\ \hat{\Psi} &= \frac{1}{N} \text{diag} \left\{ \sum_i \left(\mathbf{x}_i - \hat{\tilde{\mathbf{W}}} \mathbf{b}_i \right) \mathbf{x}_i^\top \right\} \end{aligned}$$

the $\text{diag}\{\cdot\}$ enforces the modeling assumption that Ψ remains diagonal.

The loop. Alternate E-step (recompute $\mathbf{m}_i, \Sigma_i, \mathbf{b}_i, \mathbf{C}_i$ using current $\boldsymbol{\theta}$) and M-step (recompute $\mathbf{W}, \boldsymbol{\mu}, \Psi$ using current $\mathbf{b}_i, \mathbf{C}_i$) until the marginal log-likelihood converges.

3.3 EM algorithm for PPCA

PPCA is the special case of FA with $\Psi = \sigma^2 \mathbf{I}$. Substituting into the general FA-EM equations:

E-step.

$$\Sigma = (\mathbf{I}_L + \sigma^{-2} \mathbf{W}^T \mathbf{W})^{-1} = \sigma^2 \mathbf{M}^{-1}, \quad \mathbf{M} := \mathbf{W}^T \mathbf{W} + \sigma^2 \mathbf{I}$$

$$\mathbf{m}_i = \mathbf{M}^{-1} \mathbf{W}^T (\mathbf{x}_i - \boldsymbol{\mu})$$

(Σ has no subscript i : it does not depend on $\mathbf{x}_i$, only on $\mathbf{W}, \sigma^2$.)

M-step. The $\tilde{\mathbf{W}}$ update is unchanged in form:

$$\hat{\tilde{\mathbf{W}}} = \left[\sum_i \mathbf{x}_i \mathbf{b}_i^\top \right] \left[\sum_i \mathbf{C}_i \right]^{-1}$$

Since noise is isotropic, σ^2 is a single scalar (average over all D dimensions, rather than a diagonal Ψ):

$$\hat{\sigma}^2 = \frac{1}{ND} \sum_{i=1}^N \text{tr} \left[(\mathbf{x}_i - \hat{\tilde{\mathbf{W}}} \mathbf{b}_i) \mathbf{x}_i^\top \right]$$

3.4 Limiting case: EM algorithm for PCA

Noise-free limit ($\sigma^2 \rightarrow 0$): *EM for PCA.* The posterior collapses to a point estimate ($\Sigma \rightarrow \mathbf{0}$), reducing to alternating least squares. With data centered and column-stacked as $\tilde{\mathbf{X}} \in \mathbb{R}^{D \times N}$:

1. E-step: given current $\mathbf{W}$, project data onto it via least squares $\rightarrow \tilde{\mathbf{Z}}$

$$\tilde{\mathbf{Z}} = (\mathbf{W}^T \mathbf{W})^{-1} \mathbf{W}^T \tilde{\mathbf{X}}$$

2. M-step: given current $\tilde{\mathbf{Z}}$, re-fit $\mathbf{W}$ via least-squares regression of $\mathbf{X}$ onto $\mathbf{Z}$

$$\mathbf{W} = \tilde{\mathbf{X}}\tilde{\mathbf{Z}}^T(\tilde{\mathbf{Z}}\tilde{\mathbf{Z}}^T)^{-1}$$

Why use EM instead of the closed-form eigendecomposition? Each EM iteration only requires $L \times L$ inversions ($L \ll D$ typically), costing $O(NDL + L^3)$ per iteration, versus $O(D^3)$ for eigendecomposing the full covariance matrix $\mathbf{S}$. This matters when D is large (e.g. many neurons) but L is small. EM also extends naturally to missing data, unlike the closed-form solution.

Sanity check. As $\sigma^2 \rightarrow 0$, both EM-derived quantities converge to the standard PCA solution: $\hat{\mathbf{W}} \rightarrow \mathbf{U}_L \mathbf{L}_L^{1/2}$ and $\mathbb{E}[\mathbf{z} \mid \mathbf{x}] \rightarrow (\mathbf{W}^T \mathbf{W})^{-1} \mathbf{W}^T (\mathbf{x} - \bar{\mathbf{x}})$, matching the closed-form Tipping & Bishop result.

3.5 Special case: Probabilistic PCA

Probabilistic PCA (PPCA) is a special case of the factor analysis model in which $\mathbf{W}$ has orthonormal columns and $\Psi = \sigma^2 \mathbf{I}$. Then, the full-form marginal distribution on the visible variables reduces to:

$$\begin{aligned} p(\mathbf{x} \mid \boldsymbol{\mu}, \mathbf{W}, \sigma^2) &= \int \mathcal{N}(\mathbf{x} \mid \mathbf{W}\mathbf{z} + \boldsymbol{\mu}, \Psi) \mathcal{N}(\mathbf{z} \mid \boldsymbol{\mu}_0, \Sigma_0) d\mathbf{z} \\ &= \int \mathcal{N}(\mathbf{x} \mid \mathbf{W}\mathbf{z}, \sigma^2 \mathbf{I}) \mathcal{N}(\mathbf{z} \mid \mathbf{0}, \mathbf{I}) d\mathbf{z} \\ &= \mathcal{N}(\mathbf{x} \mid \boldsymbol{\mu}, \mathbf{W}\mathbf{W}^T + \sigma^2 \mathbf{I}) = \mathcal{N}(\mathbf{x} \mid \boldsymbol{\mu}, \mathbf{C}) \end{aligned}$$

Then, the log-likelihood for PPCA is:

$$\begin{aligned} \log p(\mathbf{X} \mid \boldsymbol{\mu}, \mathbf{W}, \sigma^2) &= \sum_{n=1}^N \log p(\mathbf{x}_n \mid \boldsymbol{\mu}, \mathbf{W}, \sigma^2) \\ &= -\frac{ND}{2} \log(2\pi) - \frac{N}{2} \log |\mathbf{C}| - \frac{1}{2} \sum_{n=1}^N (\mathbf{x}_n - \boldsymbol{\mu})^T \mathbf{C}^{-1} (\mathbf{x}_n - \boldsymbol{\mu}) \end{aligned}$$

Solve for $\boldsymbol{\mu}$

Differentiate the log-likelihood w.r.t. $\boldsymbol{\mu}$ gives

$$\hat{\boldsymbol{\mu}} = \bar{\mathbf{x}} = \frac{1}{N} \sum_{n=1}^N \mathbf{x}_n$$

Substituting this back gives the simplified objective in terms of the sample covariance matrix $\mathbf{S} = \frac{1}{N} \sum_{n=1}^N (\mathbf{x}_n - \bar{\mathbf{x}})(\mathbf{x}_n - \bar{\mathbf{x}})^T$:

$$\log p(\mathbf{X} \mid \boldsymbol{\mu}, \mathbf{W}, \sigma^2) = -\frac{N}{2} [D \log(2\pi) + \log |\mathbf{C}| + \text{tr}(\mathbf{C}^{-1} \mathbf{S})]$$

Solve for $\mathbf{W}$ and σ^2 jointly

Rather than iterative fitting, the maximizer is derived directly (Tipping & Bishop 1999):

$$\mathbf{W} = \mathbf{U}_L (\mathbf{L}_L - \sigma^2 \mathbf{I})^{1/2} \mathbf{R}$$

where:

- $\mathbf{U}_L$: $D \times L$ matrix of the top L eigenvectors of $\mathbf{S}$
- $\mathbf{L}_L$: $L \times L$ diagonal matrix of the corresponding top L eigenvalues
- $\mathbf{R}$: arbitrary $L \times L$ rotation matrix (WLOG set $\mathbf{R} = \mathbf{I}$)

and the noise variance:

$$\hat{\sigma}^2 = \frac{1}{D-L} \sum_{i=L+1}^D \lambda_i$$

i.e., the *average of the discarded eigenvalues* (the variance in directions not kept as signal).

Both $\hat{\mathbf{W}}$ and $\hat{\sigma}^2$ come from *one* eigendecomposition of $\mathbf{S}$: compute all D eigenvalues/eigenvectors once, keep the top L for $\hat{\mathbf{W}}$, and average the rest for $\hat{\sigma}^2$.

Sanity check

As $\sigma^2 \rightarrow 0$:

$$\hat{\mathbf{W}} \rightarrow \mathbf{U}_L \mathbf{L}_L^{1/2}$$

which is proportional to the PCA solution.

Getting the latent representation

Unlike PCA's deterministic projection, PPCA computes the *posterior* $p(\mathbf{z} \mid \mathbf{x})$ via Bayes' rule:

$$p(\mathbf{z} \mid \mathbf{x}) = \frac{p(\mathbf{x} \mid \mathbf{z})p(\mathbf{z})}{\int p(\mathbf{x} \mid \mathbf{z}')p(\mathbf{z}') d\mathbf{z}'} = \mathcal{N}(\mathbf{z} \mid \mathbf{M}^{-1}\mathbf{W}^T(\mathbf{x} - \boldsymbol{\mu}), \sigma^2\mathbf{M}^{-1})$$

where $p(\mathbf{z}) = \mathcal{N}(\mathbf{z} \mid \mathbf{0}, \mathbf{I})$, $p(\mathbf{x} \mid \mathbf{z}) = \mathcal{N}(\mathbf{x} \mid \mathbf{W}\mathbf{z} + \boldsymbol{\mu}, \sigma^2\mathbf{I})$ and $\mathbf{M} := \mathbf{W}^T\mathbf{W} + \sigma^2\mathbf{I}$.

The posterior mean serves as the latent embedding (PPCA's analogue of PCA's $\mathbf{z}_n = \mathbf{W}^T\mathbf{x}_n$). In the $\sigma^2 \rightarrow 0$ limit:

$$\mathbb{E}[\mathbf{z} \mid \mathbf{x}] = (\mathbf{W}^T\mathbf{W})^{-1}\mathbf{W}^T(\mathbf{x} - \bar{\mathbf{x}})$$

which reduces exactly to the orthogonal projection used in standard PCA.

3.6 Summary

	PCA	PPCA	Factor Analysis
<i>Decoder</i>	$\mathbf{x}_n = \mathbf{W}\mathbf{z}_n$ (deterministic)	$\mathbf{x} = \mathbf{W}\mathbf{z} + \boldsymbol{\mu} + \boldsymbol{\epsilon}$, $\boldsymbol{\epsilon} \sim \mathcal{N}(\mathbf{0}, \sigma^2 \mathbf{I})$	$\mathbf{x} = \mathbf{W}\mathbf{z} + \boldsymbol{\mu} + \boldsymbol{\epsilon}$, $\boldsymbol{\epsilon} \sim \mathcal{N}(\mathbf{0}, \boldsymbol{\Psi})$
<i>Noise structure</i>	none	isotropic: same variance σ^2 in every dimension	diagonal: different variance ψ_i per dimension
<i>Prior on $\mathbf{z}$</i>	none	$p(\mathbf{z}) = \mathcal{N}(\mathbf{0}, \mathbf{I})$	$p(\mathbf{z}) = \mathcal{N}(\mathbf{0}, \mathbf{I})$
<i>Constraint on $\mathbf{W}$</i>	orthonormal columns	orthonormal columns	none
<i>Encoder (inference)</i>	deterministic projection: $\mathbf{z}_n = \mathbf{W}^T \mathbf{x}_n$	posterior mean: $\mathbb{E}[\mathbf{z} \mathbf{x}] = \mathbf{M}^{-1} \mathbf{W}^T (\mathbf{x} - \boldsymbol{\mu})$, $\mathbf{M} = \mathbf{W}^T \mathbf{W} + \sigma^2 \mathbf{I}$	posterior mean: $\mathbb{E}[\mathbf{z} \mathbf{x}] = \mathbf{W}^T \boldsymbol{\Sigma}_x^{-1} (\mathbf{x} - \boldsymbol{\mu})$, using full $\boldsymbol{\Sigma}_x = \mathbf{W} \mathbf{W}^T + \boldsymbol{\Psi}$
<i>Encoder uncertainty</i>	none	yes: $\text{Cov}(\mathbf{z} \mathbf{x}) = \sigma^2 \mathbf{M}^{-1}$	yes: full posterior covariance, generally non-diagonal
<i>Fitting method</i>	eigendecomposition of $\mathbf{S}$ (direct, no likelihood)	closed-form MLE via eigendecomposition of $\mathbf{S}$ (Tipping & Bishop)	no closed form — requires EM algorithm
<i>Marginal $p(\mathbf{x})$</i>	none	$\mathcal{N}(\boldsymbol{\mu}, \mathbf{W} \mathbf{W}^T + \sigma^2 \mathbf{I})$	$\mathcal{N}(\boldsymbol{\mu}, \mathbf{W} \mathbf{W}^T + \boldsymbol{\Psi})$
<i>Generative?</i>	No	Yes	Yes

Table 1: Comparison of PCA, PPCA, and Factor Analysis in encoder/decoder terms.

4 Autoencoders

Recall that PCA and FA as learning a linear mapping from $\mathbf{x} \rightarrow \mathbf{z}$, called the **encoder**, f_e , and learning another (linear) mapping $\mathbf{z} \rightarrow \mathbf{x}$, called the **decoder**, f_d . The overall reconstruction function has the form $r(\mathbf{x}) = f_d(f_e(\mathbf{x}))$. The model is trained to minimize $\mathcal{L}(\boldsymbol{\theta}) = \|\mathbf{r}(\mathbf{x}) - \mathbf{x}\|_2^2$, if isotropic generative model's noise variance is assumed. More generally, $\mathcal{L}(\boldsymbol{\theta}) = -\log p(\mathbf{x} | r(\mathbf{x}))$.

Autoencoders uses neural networks to implement non-linear mappings in the encoder and decoder.




Figure 1: An autoencoder with one hidden layer MLP

The hidden units in the middle is a low-dimensional bottleneck between the input and its reconstruction.

If the hidden layer is wide enough, the model may learn the identity function. Thus, a narrow bottleneck layer with $L \ll D$ can be used, called **undercomplete representation**; a wider bottleneck layer with $L \gg D$ can also be used but to impose other kind of regularization, called **overcomplete representation**.

Regularization techniques to avoid trivial memorization,

1. **Bottleneck autoencoders:** This is a special case of a **linear autoencoder**, in which there is one hidden layer, the hidden units are computed using $\mathbf{z} = \mathbf{W}_1 \mathbf{x}$, and the output is reconstructed using $\hat{\mathbf{x}} = \mathbf{W}_2 \mathbf{z}$, where $\mathbf{W}_1$ is a $L \times D$ matrix, $\mathbf{W}_2$ is a $D \times L$ matrix, and $L < D$ (restricting model capacity). Hence $\hat{\mathbf{x}} = \mathbf{W}_2 \mathbf{W}_1 \mathbf{x} = \mathbf{W} \mathbf{x}$ is the output of the model. If we train this model to minimize the squared reconstruction error, $\mathcal{L}(\mathbf{W}) = \sum_{n=1}^N \|\mathbf{x}_n - \mathbf{W} \mathbf{x}_n\|_2^2$, literature shows that $\hat{\mathbf{W}}$ is an orthogonal projection onto the first L eigenvectors of the empirical covariance matrix of the data, equivalent to PCA. If non-linearities are introduced, such as replacing MLP with CNN, the model is strictly more powerful than PCA without significantly changes the model size and training time.
2. **Denoising autoencoders:** Adding noises such as Gaussian noise to the input, the model is able to recover details that are missing in the input, since it has seen similar images before, and can store this information in the parameters of the model. Suppose we train a DAE using Gaussian corruption and squared error reconstruction, i.e., we use $p_c(\tilde{\mathbf{x}} | \mathbf{x}) = \mathcal{N}(\tilde{\mathbf{x}} | \mathbf{x}, \sigma^2 \mathbf{I})$ and $\ell(\mathbf{x}, r(\tilde{\mathbf{x}})) = \|\mathbf{r}(\tilde{\mathbf{x}}) - \mathbf{x}\|_2^2$ for sample $\mathbf{x}$. DAE can learn a vector field, corresponding to

the gradient of the log data density $e(\mathbf{x}) = r(\tilde{\mathbf{x}}) - \mathbf{x} \approx \nabla_{\mathbf{x}} \log p(\mathbf{x})$. Thus points that are close to the data manifold will be projected onto it via the sampling process.

3. **Contractive autoencoders:** Adding a penalty term to the reconstruction loss, $\lambda \left\| \frac{\partial f_e(\mathbf{x})}{\partial \mathbf{x}} \right\|_F^2 = \lambda \sum_k \|\nabla_{\mathbf{x}} h_k(\mathbf{x})\|_2^2$ (i.e. penalize the Frobenius norm of the encoder's Jacobian), forcing the encoding to be insensitive to small input perturbations.
4. **Sparse autoencoders:** Adding a penalty term to the reconstruction loss, $\lambda \|\mathbf{z}\|_1$, forcing some embeddings to be small/zero. Thus, even under the overcomplete representation ($L > D$), it restricts how much of the model capacity gets used at any one time, which can actually give a richer, more interpretable representation

4.1 Variational autoencoders (VAE)

	Deterministic	Probabilistic
<i>Linear</i>	PCA	PPCA, FA
<i>Nonlinear</i>	Bottleneck/Denoising/Contractive/Sparse AE	VAE

Table 2: Comparison of dimensionality reduction algorithms.