Quiz 8

1

1/1 point (graded)

In statistics,

- (•) we use the sample statistics to learn about the distribution parameters, ✔
- ( ) we use the distribution parameters to learn about the sample statistics.

Submit

Answers are displayed within the problem

2

1/1 point (graded)

If an estimator is unbiased, then

- [ ] its value is always the value of the parameter,
- [x] its expected value is always the value of the parameter, ✔
- [ ] it variance is the same as the variance of the parameter.

✔

Submit

Answers are displayed within the problem

3

1/1 point (graded)

The unbiased estimator for the standard deviation

- exists,
- doesn't exist. ✔

Submit

Answers are displayed within the problem

4

1/1 point (graded)

Two estimators s_1 and s_2 have the same MSE. If s_1 is unbiased and s_2 is biased, then

- s_1 has the smaller variance,
- s_2 has the smaller variance, ✔
- they have the same variance,
- none of the above always holds.

Submit

Answers are displayed within the problem

5

1/1 point (graded)

To correct the raw sample variance to make it unbiased, we multiply it by

- $\frac{n-1}{n}$
- $\frac{n}{n-1}$ ✔
- $\frac{n}{n+1}$
- $\frac{n+1}{n}$

Submit

Answers are displayed within the problem

6

1/1 point (graded)

If all the observations in a sample increase by 5

- [x] the sample mean increases by 5, ✔
- [ ] the sample mean stays the same,

- [ ] the sample variance increases by 5,
- [x] the sample variance stays the same. ✔

✔

Submit

Answers are displayed within the problem

7

1/1 point (graded)

As the sample size n grows, the effect of the Bessel correction

- [ ] becomes larger,
- [x] becomes smaller, ✔
- [ ] stays the same.

Submit

Answers are displayed within the problem

8

1/1 point (graded)

As the sample size n grows, the sample mean estimates the distribution mean better. Because

- [ ] its bias decreases,

- [x] its variance decreases, ✔
- [x] its mean square error decreases, ✔
- [ ] none of the above.

✔

Submit

Answers are displayed within the problem