

[Course](#) > [Week 1...](#) > [10.7 Pr...](#) > [Proble...](#)

Problem Set 10

1

2.0/2.0 points (graded)

Which of the following will **increase** the length of the confidence interval?

☒ Increase confidence level ✓

☐ Decrease confidence level

☐ Increase sample size

☒ Decrease sample size ✓



Submit

You have used 3 of 3 attempts

i Answers are displayed within the problem

2

0/1 point (graded)

The standard deviation of the diameter of rivet heads manufactured by a factory is estimated to be 0.15mm. Given a sample with size 50 and sample mean 18.45mm, what is the upper limit of the confidence interval of the distribution mean with confidence level 98%?

✖ Answer: 18.499349290713994

Submit

You have used 4 of 4 attempts

❏ Answers are displayed within the problem

3

0/2 points (graded)

One- and two-sided tests

We know the male students' height is approximately normal, and has standard deviation 4 inches. In a sample of 10 male students, the mean height is 68 inches. Calculate the p value corresponding to the following null hypotheses.

- H_0 : The average height of male students in this college is 70 inches.

H_1 : The average height of male students in this college is **not** 70 inches.

✖ Answer: 0.11384629800665805

- H_0 : The average height of male students in this college is **at least** 70 inches.

H_1 : The average height of male students in this college **less than** 70 inches.

✖ Answer: 0.056923149003329024

Submit

You have used 4 of 4 attempts

❏ Answers are displayed within the problem

4

0/1 point (graded)

The null hypothesis says that a sprinter's reaction time follows a normal distribution with mean **at most** 0.150 seconds. Six measurements of a sprinter's reaction time show 0.152, 0.154, 0.166, 0.147, 0.161, and 0.159 seconds. What is the p value?

1

✗ Answer: 0.033611332848360709

1

Submit

You have used 4 of 4 attempts

i Answers are displayed within the problem

5

0/1 point (graded)

A psychologist estimates the standard deviation of a driver's reaction time to be 0.05 seconds. How large a sample of measurements must be taken to derive a confidence interval for the mean with margin of error at most 0.01 second, and confidence level 95%?

1

✗ Answer: 97

1

Submit

You have used 4 of 4 attempts

i Answers are displayed within the problem