

Problem 8.1:

- a. ii, iii, iv and v.

For v., consider the Markov blanket of D .

Remember that a node in the Bayesian network is conditionally independent from all other nodes given its parents, children and children's parents (this evidence is the **Markov blanket** of the node).

- b. $\mathbf{P}(D, TP, CC, UP, PP, W) = \mathbf{P}(D) \mathbf{P}(TP) \mathbf{P}(CC) \mathbf{P}(UP|D, TP) \mathbf{P}(PP|CC, UP) \mathbf{P}(W|CC, UP, PP)$.

- c. To calculate these values we can directly use the equation derived in **b.**. In fact, we have to evaluate it using the requested values of the random variables:

$$P(\neg d, tp, cc, \neg up, pp, w) = P(\neg d) P(tp) P(cc) P(\neg up|\neg d, tp) P(pp|cc, \neg up) P(w|cc, \neg up, pp) = 0.75 \cdot 0.35 \cdot 0.3 \cdot 0.9 \cdot 0.9 \cdot 0.9 = 0.05740875.$$

$$P(\neg d, tp, cc, \neg up, \neg pp, w) = P(\neg d) P(tp) P(cc) P(\neg up|\neg d, tp) P(\neg pp|cc, \neg up) P(w|cc, \neg up, \neg pp) = 0.75 \cdot 0.35 \cdot 0.3 \cdot 0.9 \cdot 0.1 \cdot 0.5 = 0.00354375.$$

- d. The required probability is $P(w|tp, \neg d, pp)$.

To calculate this we can use enumeration:

$$\mathbf{P}(W|tp, \neg d, pp) = \alpha P(\neg d) P(tp) \sum_{UP} \mathbf{P}(UP|\neg d, tp) \sum_{CC} \mathbf{P}(CC) \mathbf{P}(pp|CC, UP) \mathbf{P}(W|CC, UP, pp).$$

$$\begin{aligned} P(w|tp, \neg d, pp) &= \alpha P(\neg d) P(tp) \{P(up|\neg d, tp) [P(cc) P(pp|cc, up) P(w|cc, up, pp)] \\ &\quad + P(\neg cc) P(pp|\neg cc, up) P(w|\neg cc, up, pp)] \\ &\quad + P(\neg up|\neg d, tp) [P(cc) P(pp|cc, \neg up) P(w|cc, \neg up, pp)] \\ &\quad + P(\neg cc) P(pp|\neg cc, \neg up) P(w|\neg cc, \neg up, pp)]\} \\ &= \alpha \cdot 0.75 \cdot 0.35 \{0.1[0.3 \cdot 0.5 \cdot 0.4 + 0.7 \cdot 0.05 \cdot 0.05] + 0.9[0.3 \cdot 0.9 \cdot 0.9 + 0.7 \cdot 0.1 \cdot 0.15]\} \\ &\approx \alpha \cdot 0.0615, \end{aligned}$$


$$\begin{aligned} P(\neg w|tp, \neg d, pp) &= \alpha P(\neg d) P(tp) \{P(up|\neg d, tp) [P(cc) P(pp|cc, up) P(\neg w|cc, up, pp)] \\ &\quad + P(\neg cc) P(pp|\neg cc, up) P(\neg w|\neg cc, up, pp)] \\ &\quad + P(\neg up|\neg d, tp) [P(cc) P(pp|cc, \neg up) P(\neg w|cc, \neg up, pp)] \\ &\quad + P(\neg cc) P(pp|\neg cc, \neg up) P(\neg w|\neg cc, \neg up, pp)]\} \\ &= \alpha \cdot 0.75 \cdot 0.35 \{0.1[0.3 \cdot 0.5 \cdot 0.6 + 0.7 \cdot 0.05 \cdot 0.95] + 0.9[0.3 \cdot 0.9 \cdot 0.1 + 0.7 \cdot 0.1 \cdot 0.85]\} \\ &\approx \alpha \cdot 0.0237, \end{aligned}$$

$$\alpha = \frac{1}{0.0615 + 0.0237} \approx 11.7371, \\ P(w|tp, \neg d, pp) \approx 11.7371 \cdot 0.0615 \approx 0.7218.$$

- e. The random variable EF has a direct influence on the race. We can reasonably assume that when there is an engine failure during the race, the pilot does not win. Another reasonable assumption is that a competitive car is also more reliable with a direct influence on the random variable EF . In conclusion, we introduce the node EF as a parent of W and child of CC .

Problem 8.2:

a. The corresponding network is the following:



b. Using enumeration:

$$\mathbf{P}(CD|\neg a, gf) = \alpha \mathbf{P}(CD) \sum_{DF} \mathbf{P}(gf|\neg a, DF) \mathbf{P}(\neg a|DF) \sum_H \mathbf{P}(DF|CD, H) \mathbf{P}(H).$$

$$\begin{aligned} P(cd|\neg a, gf) &= \alpha P(cd) \{P(gf|df, \neg a) P(\neg a|df) [P(df|cd, h) P(h) + P(df|cd, \neg h) P(\neg h)] \\ &\quad + P(gf|\neg df, \neg a) P(\neg a|\neg df) [P(\neg df|cd, h) P(h) + P(\neg df|cd, \neg h) P(\neg h)]\} \\ &= \alpha \cdot 0.6 \{0.4 \cdot 0.3 [0.15 \cdot 0.5 + 0.01 \cdot 0.5] + 0.05 \cdot 0.75 [0.85 \cdot 0.5 + 0.99 \cdot 0.5]\} \\ &\approx \alpha \cdot 0.0265, \end{aligned}$$

$$\begin{aligned} P(\neg cd|\neg a, gf) &= \alpha P(\neg cd) \{P(gf|df, \neg a) P(\neg a|df) [P(df|\neg cd, h) P(h) + P(df|\neg cd, \neg h) P(\neg h)] \\ &\quad + P(gf|\neg df, \neg a) P(\neg a|\neg df) [P(\neg df|\neg cd, h) P(h) + P(\neg df|\neg cd, \neg h) P(\neg h)]\} \\ &= \alpha \cdot 0.4 \{0.4 \cdot 0.3 [0.99 \cdot 0.5 + 0.1 \cdot 0.5] + 0.05 \cdot 0.75 [0.01 \cdot 0.5 + 0.9 \cdot 0.5]\} \\ &\approx \alpha \cdot 0.0330, \end{aligned}$$

$$\alpha = \frac{1}{0.0265 + 0.0330} \approx 16.8067,$$

$$\mathbf{P}(CD|\neg a, gf) \approx 16.8067 \cdot \begin{bmatrix} 0.0265 \\ 0.0330 \end{bmatrix} \approx \begin{bmatrix} 0.4454 \\ 0.5546 \end{bmatrix}.$$

c. Using variable elimination:

$$\mathbf{P}(CD|\neg a, gf) = \underbrace{\alpha \mathbf{P}(CD)}_{f_1(CD)} \sum_{DF} \underbrace{P(gf|\neg a, DF)}_{f_2(DF)} \underbrace{P(\neg a|DF)}_{f_3(DF)} \sum_H \underbrace{P(DF|CD, H)}_{f_4(DF, CD, H)} \underbrace{P(H)}_{f_5(H)}.$$

In the following notation we represent a $2 \times 2 \times 2$ matrix as $\{[2 \times 2] [2 \times 2]\}$.

$$\begin{aligned} \mathbf{f}_4(DF, CD, H) &= \begin{bmatrix} \mathbf{P}(DF|CD, h) & \mathbf{P}(DF|CD, \neg h) \end{bmatrix} \\ &= \left\{ \begin{bmatrix} P(df|cd, h) & P(df|\neg cd, h) \\ P(\neg df|cd, h) & P(\neg df|\neg cd, h) \end{bmatrix} \begin{bmatrix} P(df|cd, \neg h) & P(df|\neg cd, \neg h) \\ P(\neg df|cd, \neg h) & P(\neg df|\neg cd, \neg h) \end{bmatrix} \right\} \\ &= \left\{ \begin{bmatrix} 0.15 & 0.99 \\ 0.85 & 0.01 \end{bmatrix} \begin{bmatrix} 0.01 & 0.1 \\ 0.99 & 0.9 \end{bmatrix} \right\}, \end{aligned}$$

$$\mathbf{f}_1(CD) = \begin{bmatrix} P(cd) & P(\neg cd) \end{bmatrix} = \begin{bmatrix} 0.6 & 0.4 \end{bmatrix}, \quad \mathbf{f}_2(DF) = \begin{bmatrix} P(gf|\neg a, df) \\ P(gf|\neg a, \neg df) \end{bmatrix} = \begin{bmatrix} 0.4 \\ 0.05 \end{bmatrix},$$

$$\mathbf{f}_3(DF) = \begin{bmatrix} P(\neg a|df) \\ P(\neg a|\neg df) \end{bmatrix} = \begin{bmatrix} 0.3 \\ 0.75 \end{bmatrix}, \quad \mathbf{f}_5(H) = \begin{bmatrix} P(h) & P(\neg h) \end{bmatrix} = \begin{bmatrix} 0.5 & 0.5 \end{bmatrix}.$$

After defining the factors, we can write:

$$\mathbf{P}(CD|\neg a, gf) = \alpha \mathbf{f}_1(CD) \times \sum_{DF} \mathbf{f}_2(DF) \times \mathbf{f}_3(DF) \times \sum_H \mathbf{f}_4(DF, CD, H) \times \mathbf{f}_5(H).$$

In the following we use the pointwise product and we sum out variables.

$$\mathbf{f}_4(DF, CD, H) \times \mathbf{f}_5(H) = \left\{ \begin{bmatrix} 0.075 & 0.495 \\ 0.425 & 0.005 \end{bmatrix} \begin{bmatrix} 0.005 & 0.050 \\ 0.495 & 0.450 \end{bmatrix} \right\},$$

$$\mathbf{f}_6(DF, CD) = \sum_H \mathbf{f}_4(DF, CD, H) \times \mathbf{f}_5(H) = \sum_H \left\{ \begin{bmatrix} 0.075 & 0.495 \\ 0.425 & 0.005 \end{bmatrix} \begin{bmatrix} 0.005 & 0.050 \\ 0.495 & 0.450 \end{bmatrix} \right\} = \begin{bmatrix} 0.080 & 0.545 \\ 0.920 & 0.455 \end{bmatrix},$$

$$\mathbf{f}_2(DF) \times \mathbf{f}_3(DF) \times \mathbf{f}_6(DF, CD) = \begin{bmatrix} 0.0096 & 0.0654 \\ 0.0345 & 0.0171 \end{bmatrix},$$

$$\mathbf{f}_7(CD) = \sum_{DF} \mathbf{f}_2(DF) \times \mathbf{f}_3(DF) \times \mathbf{f}_6(DF, CD) = \sum_{DF} \begin{bmatrix} 0.0096 & 0.0654 \\ 0.0345 & 0.0171 \end{bmatrix} = \begin{bmatrix} 0.0441 & 0.0825 \end{bmatrix}.$$

And finally we get:

$$\mathbf{P}(CD|\neg a, gf) = \alpha \mathbf{f}_1(CD) \times \mathbf{f}_7(CD) \approx \alpha \begin{bmatrix} 0.0265 & 0.0330 \end{bmatrix},$$

$$\alpha = \frac{1}{0.0265+0.0330} \approx 16.8067,$$

$$\mathbf{P}(CD|\neg a, gf) \approx 16.8067 \cdot \begin{bmatrix} 0.0265 & 0.0330 \end{bmatrix} \approx \begin{bmatrix} 0.4454 & 0.5546 \end{bmatrix}.$$


d. Besides the last calculation to complete the normalization that both methods require, we have:

	Enumeration	Variable Elimination
additions	6	6
multiplications	18	16

In this case we do not have a significant improvement using variable elimination. The improvement becomes more significant for larger Bayesian networks.

Problem 8.3:

- a. The network is composed of four random variables: The coin that has been drawn DC with domain $\langle a, b, c \rangle$ and the results of the three flips $X1$, $X2$ and $X3$ with domains $\langle heads, tails \rangle$.



- b. The required probability is $\mathbf{P}(DC|X1 = heads, X2 = heads, X3 = tails)$.

We first write the full joint probability:

$$\mathbf{P}(DC, X1, X2, X3) = \mathbf{P}(X1|DC)\mathbf{P}(X2|DC)\mathbf{P}(X3|DC)\mathbf{P}(DC).$$

The result is directly obtained introducing the evidence and using normalization:

$$\begin{aligned} \mathbf{P}(DC|X1 = heads, X2 = heads, X3 = tails) &= \\ &= \alpha \mathbf{P}(X1 = heads|DC)\mathbf{P}(X2 = heads|DC)\mathbf{P}(X3 = tails|DC)\mathbf{P}(DC) = \\ &= \alpha \begin{bmatrix} 0.2 \\ 0.6 \\ 0.8 \end{bmatrix} \times \begin{bmatrix} 0.2 \\ 0.6 \\ 0.8 \end{bmatrix} \times \begin{bmatrix} 0.8 \\ 0.4 \\ 0.2 \end{bmatrix} \times \begin{bmatrix} \frac{1}{3} \\ \frac{1}{3} \\ \frac{1}{3} \end{bmatrix} \approx \alpha \begin{bmatrix} 0.0107 \\ 0.0480 \\ 0.0427 \end{bmatrix}, \end{aligned}$$

$$\alpha = \frac{1}{(0.0107+0.0480+0.0427)} \approx 9.8619329,$$

$$\mathbf{P}(DC|X1 = heads, X2 = heads, X3 = tails) \approx 9.8619329 \begin{bmatrix} 0.0107 \\ 0.0480 \\ 0.0427 \end{bmatrix} = \begin{bmatrix} 0.1055 \\ 0.4734 \\ 0.4211 \end{bmatrix}.$$

The coin that is most likely to have been drawn is b.