

# **P9185 - Project 3: Protocol of a Cluster-randomized trial for Asthma-PASS**

Amy Pitts, Yuqi Miao

2022-04-25

# Background

Our interest is in persistent asthma in minority children.


- Comprehensive school-based interventions in collaboration with communities to reduce asthma morbidity and promote physical activity in urban areas.
- A pilot cluster RCT was conducted exploring this intervention in Bronx elementary schools
  - **Goal:** whether Children in schools receiving Asthma-PASS intervention may experience a greater improvement in the number of SFD at 6 **or** 12 months follow up than the children in the comparison group.
  - 4 Bronx elementary schools were recruited into the pilot study.
  - A total of 108 children recruited including ages 4-11 years with physician-diagnosed persistent or uncontrolled asthma attending kindergarten to 5th grades

- ① Data Overview
- ② Exploration into Pilot Study data
  - Model Specifications
  - Results
- ③ Phase III Proposal
  - Model Specifications
  - Sample Size Suggestions

Variable	Definition
ID	Participant's ID
Time	Follow up time (Baseline, 6 months, 12 months)
Group	Intervention group (control or Intervention)
SFD	Symptom free days in the past two weeks
School	School recruited for the study

*Table 1: Data Descriptions*

# Data Exploration



## **Current outcome:** SFD (Count data)


- Due to the skewed distribution towards higher values a poisson model will not fit our data well
- Outcome does not seem linear over the time observations.

*Interested in the change from baseline to observation times.*


## **New outcome:** Change in SFD (Continuous Data)

- Transform the SFD by calculated:
  - 6 month observation - baseline
  - 12 month observation - baseline
- Baseline with become covariate

# New Continuous Outcome



# Exploring variation between school and within school






# Data Description

Characteristic	Overall, N = 108 <sup>1</sup>	Group Control		Group Intervention		p-value <sup>2</sup>
		School 1, N = 28 <sup>1</sup>	School 3, N = 31 <sup>1</sup>	School 2, N = 21 <sup>1</sup>	School 4, N = 28 <sup>1</sup>	
baseline	11.0 (7.0, 13.0)	12.0 (7.5, 13.0)	9.0 (6.2, 11.8)	11.0 (9.0, 14.0)	11.0 (7.5, 12.5)	0.3
Unknown	3	1	1	0	1	
Change_6months	2 (0, 5)	2 (0, 3)	3 (0, 8)	1 (0, 2)	0 (0, 4)	0.3
Unknown	10	3	4	2	1	
Change_12months	1.0 (0.0, 4.0)	1.0 (0.0, 3.5)	2.0 (0.5, 5.5)	0.5 (-0.2, 3.0)	0.0 (0.0, 3.0)	0.2
Unknown	15	5	4	1	5	

<sup>1</sup> Median (IQR)

<sup>2</sup> Kruskal-Wallis rank sum test

# Missing Data



# Model Specifications

To model change in SFD let  $i$  for school,  $j$  for subjects,  $k$  for measures.

We will use mixed effect model.

$$\begin{aligned} Y_{ijk} = & \beta_0 + \beta_1 \text{Baseline}_{ij} \\ & + \beta_2 \text{Group}_{ij} + \beta_3 \text{Compare}_{ijk} \\ & + \beta_4 \text{Group}_{ij} \times \text{Compare}_{ijk} \\ & + \alpha_{0i} + \alpha_{0j} + \epsilon_{ijk} \end{aligned}$$

where  $\alpha_{0i} \sim N(0, \sigma_w^2)$ ,  $\alpha_{0j} \sim N(0, \sigma_b^2)$ , and  $\epsilon_{ijk} \sim N(0, \sigma^2)$ .

# Missing Data Assumptions

We will be assuming data is missing at random (MAR)<sup>[1]</sup>.

- MAR assumption:  $R \perp\!\!\!\perp Y_{mis} | X, Y_{obs}$
- Separable parameter assumption
- Ignorability condition

$$\begin{aligned} L_i^O(\theta, \psi) &\propto f_{\theta, \psi}(Y_{obs,i}, R_i, X_i) \\ &= f_{\psi}(R_i | Y_{obs,i}, X_i) f_{\theta}(Y_{obs,i} | X_i) \end{aligned}$$

# Model Result

## Fixed Effects Estimates:

Characteristic	Beta	95% CI <sup>†</sup>	p-value
baseline	-0.82	-1.0, -0.69	<0.001
group			
0	—	—	
1	1.1	-0.46, 2.6	0.2
compare			
m6_m0	—	—	
m12_m0	-0.05	-1.4, 1.3	>0.9
group * compare			
1 * m12_m0	-0.27	-2.2, 1.6	0.8

<sup>†</sup> CI = Confidence Interval

## Random Effects Estimates:

group	Std.Dev	Variance
id	2.055	4.221
school	0.000	0.000
Residual	3.250	10.560


## Model Interpretations:

- 1 interpretations
- 2 interpretations
- 3 interpretations


*Note:* This model is singular

# Model Quality

Residual plot



Normal Q-Q Plot



# Model without Baseline

## Fixed Effects Estimates:

Characteristic	Beta	95% CI <sup>†</sup>	p-value
group			
0	—	—	
1	1.7	-2.5, 5.8	0.3
compare			
m6_m0	—	—	
m12_m0	-0.08	-1.5, 1.3	>0.9
group * compare			
1 * m12_m0	-0.26	-2.2, 1.7	0.8

<sup>†</sup> CI = Confidence Interval

## Random Effects Estimates:

group	Std.Dev	Variance
id	4.384	19.215
school	0.597	0.357
Residual	3.275	10.726


## Model Interpretations:

- 1 interpretations
- 2 interpretations
- 3 interpretations




# Model Quality

Residual plot



Normal Q-Q Plot





# Cluster RCT design

- The investigators wish to propose a **cluster-randomized clinical trial (RCT)** in 30 Bronx schools to evaluate the effectiveness of their intervention program.
- Primary hypothesis: compared to the control group, children in schools randomized to intervention group will experience a greater improvement in the number of SFD at **any of the 3, 6, 9, and 12 months** assessment.
- The investigators would like to have at **80% probability** to declare the trial is successful if the true effect size in **improvement of SFD over time** is **at least 1/3 standard deviation**.

# Study design proposal:

## 3 level structure<sup>[2]</sup>:

$$y_{ijk} = \beta_0 + \delta_{(3)}X_{ijk} + \mu_i + \mu_j + e_{ijk}$$

- $i$  for school,  $j$  for subjects,  $k$  for measures
- $\mu_i \sim N(0, \sigma_b^2)$  random intercept between schools
- $\mu_j \sim N(0, \sigma_w^2)$  random intercept for subjects within schools
- $e_{ijk} \sim N(0, \sigma^2)$  random error term

## Intraclass Correlation Coefficient

$$\rho_1 = \frac{\sigma_b^2 + \sigma_w^2}{\sigma_b^2 + \sigma_w^2 + \sigma^2}$$

$$\rho_2 = \frac{\sigma_b^2}{\sigma_b^2 + \sigma_w^2 + \sigma^2}$$

# Hypothesis Set Up

**Hypothesis:**  $H_0 : \delta_{(3)} = 0$ ,  $H_1 : \delta_{(3)} \neq 0$

- *Clinical Interest:*  $H_1 : \delta_{(3)} > 0$ ?
- Calculate  $N$  based on normal distribution
- Interested in when  $\beta = 0.20$ ,  $\alpha^* = 0.05/2$  and  $\Delta = 1/3$

## Three levels of Sampling

- $N_1$ : Number of Observation looks (4 in our case)
- $N_2$ : Number of Individuals in each School (*What we want to estimate*)
- $N_3$ : Number of Schools for one treatment arm (15 in our case)

# Sample Size Calculation

## Sample Size formula

$$N_2 = \frac{2(1 + (\rho_1 - \rho_2)N_1 - \rho_1)z_{\alpha^*,\beta}^2}{N_1 N_3 \Delta^2 - 2\rho_2 N_1 z_{\alpha^*,\beta}^2}$$

Where  $z$  is calculated based on the normal distribution.

$$z_{\alpha^*,\beta}^2 = (z_{\alpha^*/2} + z_{\beta})^2 = [\Phi^{-1}(1 - \alpha^*/2) + \Phi^{-1}(1 - \beta)^{-1}]^2$$

# Intra Class Correlation in our Models

## Model 1 Random Effects:

group	Std.Dev	Variance
id	2.055	4.221
school	0.000	0.000
Residual	3.250	10.560

①  $\rho_1 = 0.286$

②  $\rho_2 = 0.000$

## Model 2 Random Effects:

group	Std.Dev	Variance
id	4.384	19.215
school	0.597	0.357
Residual	3.275	10.726

①  $\rho_1 = 0.646$

②  $\rho_2 = 0.012$

# Sample Size Suggested

label	rho1	rho2	class_size	group_size	total_samp
Est w/ Baseline	0.286	0.000	4.372	65.580	131.160
Average	0.466	0.006	5.918	88.766	177.532
Est w/o Baseline	0.646	0.012	7.656	114.846	229.692

[1] Hogan, J. W., Roy, J., & Korkontzelou, C. (2004). Handling drop-out in longitudinal studies. *Statistics in Medicine*, 23(9), 1455–1497.

<https://doi.org/10.1002/sim.1728>

[2] Ahn, C., Heo, M., & Zhang, S. (2014). *Sample size calculations for clustered and longitudinal outcomes in clinical research*. CRC Press.