

Bayesian dynamic pricing in discrete time

Yuqiong Wang

(joint work with Erik Ekström)

**Department of Mathematics
Uppsala University**

12th Bachelier Finance Society Congress, Rio de Janeiro, Brazil

July 8, 2024

Outline

- 1 **Introduction**
 - Incomplete information
 - Applications
- 2 **Problem formulation and results**
 - Formulation
 - Construct the value function
 - Consequence of convexity
- 3 **Some remarks**

Stopping with incomplete information

- Optimal stopping problems often concern:


$$V = \sup_{\tau} \mathbb{E}[G(X_{\tau})].$$

Stopping with incomplete information

- Optimal stopping problems often concern:

$$V = \sup_{\tau} \mathbb{E}[G(X_{\tau})].$$

- Usually uncertainty is associated with the assumptions:




Stopping with incomplete information

- Optimal stopping problems often concern:

$$V = \sup_{\tau} \mathbb{E}[G(X_{\tau})].$$

- Usually uncertainty is associated with the assumptions:




– Is X a Gaussian with mean 0 or mean 1?

Stopping with incomplete information

- Optimal stopping problems often concern:

$$V = \sup_{\tau} \mathbb{E}[G(X_{\tau})].$$

- Usually uncertainty is associated with the assumptions:



– Is X a Gaussian with mean 0 or mean 1?


- Combine optimal stopping and filtering: optimising while **learning**.

Stopping with incomplete information

- Optimal stopping problems often concern:

$$V = \sup_{\tau} \mathbb{E}[G(X_{\tau})].$$

- Usually uncertainty is associated with the assumptions:



– Is X a Gaussian with mean 0 or mean 1?

- Combine optimal stopping and filtering: optimising while **learning**.
- Applications: e.g. **statistics**, and **option pricing**.

In statistics: sequential testing

Classical problem: Testing the unknown drift of a BM.

- Observe the trajectory of a BM with unknown drift:

$$X_t = \theta t + W_t.$$

where $\mathbb{P}(\theta = 1) = \pi = 1 - \mathbb{P}(\theta = 0)$, $\pi \in (0, 1)$.

In statistics: sequential testing

Classical problem: Testing the unknown drift of a BM.

- Observe the trajectory of a BM with unknown drift:

$$X_t = \theta t + W_t.$$

where $\mathbb{P}(\theta = 1) = \pi = 1 - \mathbb{P}(\theta = 0)$, $\pi \in (0, 1)$.

- Want to test: $H_1 : \theta = 1$, $H_0 : \theta = 0$, as accurately as possible.

In statistics: sequential testing

Classical problem: Testing the unknown drift of a BM.

- Observe the trajectory of a BM with unknown drift:

$$X_t = \theta t + W_t.$$

where $\mathbb{P}(\theta = 1) = \pi = 1 - \mathbb{P}(\theta = 0)$, $\pi \in (0, 1)$.

- Want to test: $H_1 : \theta = 1$, $H_0 : \theta = 0$, as **accurately** as possible.
- Observation is not free: $c > 0$ per unit time of observation.

In statistics: sequential testing

Classical problem: Testing the unknown drift of a BM.

- Observe the trajectory of a BM with unknown drift:

$$X_t = \theta t + W_t.$$

where $\mathbb{P}(\theta = 1) = \pi = 1 - \mathbb{P}(\theta = 0)$, $\pi \in (0, 1)$.

- Want to test: $H_1 : \theta = 1$, $H_0 : \theta = 0$, as **accurately** as possible.
- Observation is not free: $c > 0$ per unit time of observation.
- Need to test as **fast** as possible.

In statistics: sequential testing

Classical problem: Testing the unknown drift of a BM.

- Observe the trajectory of a BM with unknown drift:

$$X_t = \theta t + W_t.$$

where $\mathbb{P}(\theta = 1) = \pi = 1 - \mathbb{P}(\theta = 0)$, $\pi \in (0, 1)$.

- Want to test: $H_1 : \theta = 1$, $H_0 : \theta = 0$, as **accurately** as possible.
- Observation is not free: $c > 0$ per unit time of observation.
- Need to test as **fast** as possible.
- The time to stop observing is part of the decision.

In statistics: sequential testing

- The **minimised cost** V :

$$V = \inf_{\tau, d} \{ \mathbb{P}(d = 0, \theta = 1) + \mathbb{P}(d = 1, \theta = 0) + c\mathbb{E}[\tau] \}. \quad (1)$$

In statistics: sequential testing

- The **minimised cost** V :

$$V = \inf_{\tau, d} \{ \mathbb{P}(d = 0, \theta = 1) + \mathbb{P}(d = 1, \theta = 0) + c\mathbb{E}[\tau] \}. \quad (1)$$

Solution: embed in another coordinate, solves a free-boundary problem.

- Defining the **posterior probability process**

$$\Pi_t := \mathbb{P}_\pi(\theta = 1 | \mathcal{F}_t^X),$$

In statistics: sequential testing

- The **minimised cost** V :

$$V = \inf_{\tau, d} \{ \mathbb{P}(d = 0, \theta = 1) + \mathbb{P}(d = 1, \theta = 0) + c\mathbb{E}[\tau] \}. \quad (1)$$

Solution: embed in another coordinate, solves a free-boundary problem.

- Defining the **posterior probability process**

$$\Pi_t := \mathbb{P}_\pi(\theta = 1 | \mathcal{F}_t^X),$$

- Problem (1) can be written as

$$V(\pi) = \inf_{\tau} \mathbb{E}_\pi [c\tau + \Pi_\tau \wedge (1 - \Pi_\tau)]$$

where

$$d\Pi_t = \Pi_t(1 - \Pi_t)d\tilde{W}_t,$$

In statistics: sequential testing

- The **minimised cost** V :

$$V = \inf_{\tau, d} \{ \mathbb{P}(d = 0, \theta = 1) + \mathbb{P}(d = 1, \theta = 0) + c\mathbb{E}[\tau] \}. \quad (1)$$

Solution: embed in another coordinate, solves a free-boundary problem.

- Defining the **posterior probability process**

$$\Pi_t := \mathbb{P}_\pi(\theta = 1 | \mathcal{F}_t^X),$$

- Problem (1) can be written as

$$V(\pi) = \inf_{\tau} \mathbb{E}_\pi [c\tau + \Pi_\tau \wedge (1 - \Pi_\tau)]$$

where

$$d\Pi_t = \Pi_t(1 - \Pi_t)d\tilde{W}_t,$$

- Standard method applies: explicit solution. [c.f. Shiryaev \(1969\)](#).

In finance: American option pricing

- Consider the stock of a company with:

$$dX_t = \theta X_t dt + \sigma X_t dW_t, \quad X_0 = x,$$

In finance: American option pricing

- Consider the stock of a company with:

$$dX_t = \theta X_t dt + \sigma X_t dW_t, \quad X_0 = x,$$

- where the company can be either in a **good** or **bad** state:

$$\mathbb{P}(\theta = 1) = \pi = 1 - \mathbb{P}(\theta = 0), \quad \pi \in (0, 1)$$

.

In finance: American option pricing

- Consider the stock of a company with:

$$dX_t = \theta X_t dt + \sigma X_t dW_t, \quad X_0 = x,$$

- where the company can be either in a **good** or **bad** state:

$$\mathbb{P}(\theta = 1) = \pi = 1 - \mathbb{P}(\theta = 0), \quad \pi \in (0, 1)$$

- The **buyer** of the option needs to estimate θ .
- The value of an American put:

$$V = \sup_{\tau \leq T} \mathbb{E}_{t,x} [e^{-r\tau} (K - X_\tau)^+]$$

In finance: American option pricing

- Consider the stock of a company with:

$$dX_t = \theta X_t dt + \sigma X_t dW_t, \quad X_0 = x,$$

- where the company can be either in a **good** or **bad** state:

$$\mathbb{P}(\theta = 1) = \pi = 1 - \mathbb{P}(\theta = 0), \quad \pi \in (0, 1)$$

- The **buyer** of the option needs to estimate θ .
- The value of an American put:

$$V = \sup_{\tau \leq T} \mathbb{E}_{t,x} [e^{-r\tau} (K - X_\tau)^+]$$


- Solution: rely on the Π process again:

$$V = \sup_{\tau} \mathbb{E}_{t,x,\pi} [e^{-r\tau} (K - X_\tau)^+].$$

c.f. Decamps et al. (2005), Gapeev (2012).

In finance: American option pricing

- Can reduce to a one (spatial)-dimensional problem.



- The behavior of boundaries depends on the parameter.
c.f. Ekström et al. (2019)

What about control?

- Can we formulate a similar **control** problem?

What about control?

- Can we formulate a similar **control** problem?
- Why not start with a **Bernoulli prior** again?

What about control?

- Can we formulate a similar **control** problem?
- Why not start with a **Bernoulli prior** again?
- We consider a **learning-and-earning problem** in discrete time.

Outline

1 Introduction

- Incomplete information
- Applications

2 Problem formulation and results

- Formulation
- Construct the value function
- Consequence of convexity

3 Some remarks

Problem formulation

- Consider a **seller** who offers a product for sale.

Problem formulation

- Consider a **seller** who offers a product for sale.
- The potential **buyers** arrive in a sequential fashion.

Problem formulation

- Consider a **seller** who offers a product for sale.
- The potential **buyers** arrive in a sequential fashion.
- At time n , the seller offers a price p_n .

Problem formulation

- Consider a **seller** who offers a product for sale.
- The potential **buyers** arrive in a sequential fashion.
- At time n , the seller offers a price p_n .
- Let θ be the unknown state of the world, and priorly,

$$\mathbb{P}(\theta = 1) = \pi = 1 - \mathbb{P}(\theta = 0), \quad \pi \in (0, 1).$$

Problem formulation

- Consider a **seller** who offers a product for sale.
- The potential **buyers** arrive in a sequential fashion.
- At time n , the seller offers a price p_n .
- Let θ be the unknown state of the world, and priorly,

$$\mathbb{P}(\theta = 1) = \pi = 1 - \mathbb{P}(\theta = 0), \quad \pi \in (0, 1).$$

- At time n , with a proposed price p , we observe the amount of product sold, $X_n(\theta, p)$.

Problem formulation

- Consider a **seller** who offers a product for sale.
- The potential **buyers** arrive in a sequential fashion.
- At time n , the seller offers a price p_n .
- Let θ be the unknown state of the world, and priorly,

$$\mathbb{P}(\theta = 1) = \pi = 1 - \mathbb{P}(\theta = 0), \quad \pi \in (0, 1).$$

- At time n , with a proposed price p , we observe the amount of product sold, $X_n(\theta, p)$.
- X_i 's are independent conditioning on θ .

Problem formulation

- Consider a **seller** who offers a product for sale.
- The potential **buyers** arrive in a sequential fashion.
- At time n , the seller offers a price p_n .
- Let θ be the unknown state of the world, and priorly,

$$\mathbb{P}(\theta = 1) = \pi = 1 - \mathbb{P}(\theta = 0), \quad \pi \in (0, 1).$$

- At time n , with a proposed price p , we observe the amount of product sold, $X_n(\theta, p)$.
- X_i 's are independent conditioning on θ .
- We know this conditional distribution:

$$F_\theta(x, p) = \mathbb{P}(X(\theta, p) \leq x)$$

Problem formulation

- The seller seeks to maximise the discounted profit:

$$V = \sup_{\{p_n\}_{n \geq 0}} \mathbb{E} \left[\sum_{n=0}^{\infty} a^n p_n X_n^{p_n} \right].$$

where $0 < a < 1$.

Problem formulation

- The seller seeks to maximise the discounted profit:

$$V = \sup_{\{p_n\}_{n \geq 0}} \mathbb{E} \left[\sum_{n=0}^{\infty} a^n p_n X_n^{p_n} \right].$$

where $0 < a < 1$.

- The case where $X(\theta, p)$ being **Bernoulli** is studied in Economic & operations research literatures, c.f. [McLennan 1984](#), [Harrison 2012](#).

Problem formulation

- The seller seeks to maximise the discounted profit:

$$V = \sup_{\{p_n\}_{n \geq 0}} \mathbb{E} \left[\sum_{n=0}^{\infty} a^n p_n X_n^{p_n} \right].$$

where $0 < a < 1$.

- The case where $X(\theta, p)$ being **Bernoulli** is studied in Economic & operations research literatures, c.f. [McLennan 1984](#), [Harrison 2012](#).
- Observing one customer at a time.

Problem formulation

- The seller seeks to maximise the discounted profit:

$$V = \sup_{\{p_n\}_{n \geq 0}} \mathbb{E} \left[\sum_{n=0}^{\infty} a^n p_n X_n^{p_n} \right].$$

where $0 < a < 1$.

- The case where $X(\theta, p)$ being **Bernoulli** is studied in Economic & operations research literatures, c.f. [McLennan 1984](#), [Harrison 2012](#).
- Observing one customer at a time.
- “incomplete learning”, myopic strategies, examples (linear).

Problem formulation

In a Bayesian setting, the problem can be embedded in a Markovian framework.

Problem formulation

In a Bayesian setting, the problem can be embedded in a Markovian framework.

- Define the usual posterior probability process

$$\Pi_n^p := \mathbb{P}(\theta = 1 | \mathcal{F}_n^{X^p}).$$

Problem formulation

In a Bayesian setting, the problem can be embedded in a Markovian framework.

- Define the usual posterior probability process

$$\Pi_n^p := \mathbb{P}(\theta = 1 | \mathcal{F}_n^{X^p}).$$

- By conditioning, the value function can be written as

$$V(\pi) = \sup_{\{p_k\}_{k=0}^{\infty}} \mathbb{E}_{\pi} \left[\sum_{k=0}^{\infty} a^k p_k \left(\Pi_k^{p_k} \int_{\mathbb{R}} x F_1(dx, p_k) + (1 - \Pi_k^{p_k}) \int_{\mathbb{R}} x F_0(dx, p_k) \right) \right]$$

- Only π dependent! Time-homogeneous.

Properties of the value function

- We can find **convex** upper and lower bounds for V

Properties of the value function

- We can find **convex** upper and lower bounds for V
- By DPP, V satisfies

$$V(\pi) = \sup_p \left\{ a \mathbb{E}_\pi [V(\Pi_1^p)] + p \left(\pi \int_{\mathbb{R}} x F_1(dx, p) + (1 - \pi) \int_{\mathbb{R}} x F_0(dx, p) \right) \right\}.$$

Properties of the value function

- We can find **convex** upper and lower bounds for V
- By DPP, V satisfies

$$V(\pi) = \sup_p \left\{ a\mathbb{E}_\pi [V(\Pi_1^p)] + p \left(\pi \int_{\mathbb{R}} xF_1(dx, p) + (1 - \pi) \int_{\mathbb{R}} xF_0(dx, p) \right) \right\}.$$

Definition

Define an operator T :

$$(Tu)(\pi) = \sup_p \left\{ a\mathbb{E}_\pi [u(\Pi_1^p)] + p \left(\pi \int_{\mathbb{R}} xF_1(dx, p) + (1 - \pi) \int_{\mathbb{R}} xF_0(dx, p) \right) \right\}.$$

- Define a sequence of functions $\{u_n\}_{n \geq 0}$ by letting

$$u_0 = 0, \quad u_{n+1} = Tu_n.$$

Properties of the value function

We are able to show that

- T preserves monotonicity and convexity,

Properties of the value function

We are able to show that

- T preserves monotonicity and convexity,
- T is a contraction,

Properties of the value function

We are able to show that

- T preserves monotonicity and convexity,
- T is a contraction,
- $u_\infty := \lim u_n$ exists and is the unique fixed point of T .

Properties of the value function

We are able to show that

- T preserves monotonicity and convexity,
- T is a contraction,
- $u_\infty := \lim u_n$ exists and is the unique fixed point of T .
- $V = u_\infty$.

Properties of the value function

We are able to show that

- T preserves monotonicity and convexity,
- T is a contraction,
- $u_\infty := \lim u_n$ exists and is the unique fixed point of T .
- $V = u_\infty$.

Theorem

The value function V is convex and continuous on $[0, 1]$.

Consequence of convexity: how to choose a model

How to find a model where V is larger? Convexity gives us the answer.



Theorem

For any $(\pi, p) \in [0, 1] \times \mathbb{R}$, take $\Pi_0 = \tilde{\Pi}_0 = \pi$. Assume the following two conditions hold, then $V_n(\pi) \geq \tilde{V}_n(\pi)$ for all $n \geq 0$ and $\pi \in [0, 1]$.

- 1 (Earning) $\int_{\mathbb{R}} x F_1(dx, p) \geq \int_{\mathbb{R}} x \tilde{F}_1(dx, p)$, and
 $\int_{\mathbb{R}} x F_0(dx, p) \geq \int_{\mathbb{R}} x \tilde{F}_0(dx, p)$.
- 2 (Learning) Π_1^p dominates $\tilde{\Pi}_1^p$ in convex order.



Properties of the optimal strategy

Example: Bernoulli observations



Properties of the optimal strategy

Example: Bernoulli observations



Other results

- Conditions for monotonicity of V .
- Conditions for monotonicity of the optimal strategy.

Outline

- 1 **Introduction**
 - Incomplete information
 - Applications
- 2 **Problem formulation and results**
 - Formulation
 - Construct the value function
 - Consequence of convexity
- 3 **Some remarks**

Some remarks

Incomplete learning?

Some remarks

Incomplete learning?

- It's possible, and fine.

Some remarks

Incomplete learning?

- It's possible, and fine.

Myopic strategy?

Some remarks

Incomplete learning?

- It's possible, and fine.

Myopic strategy?

- Not always optimal (obviously).

Some remarks

Incomplete learning?

- It's possible, and fine.

Myopic strategy?

- Not always optimal (obviously).

Generalisations?

Some remarks

Incomplete learning?

- It's possible, and fine.

Myopic strategy?

- Not always optimal (obviously).

Generalisations?

- Can relax the prior to an **arbitrary** distribution, and let X be in the **exponential family**.

Some remarks

Incomplete learning?

- It's possible, and fine.

Myopic strategy?

- Not always optimal (obviously).

Generalisations?

- Can relax the prior to an **arbitrary** distribution, and let X be in the **exponential family**.
- But the problems becomes time-dependent and very difficult.

Some remarks

Continuous time version?

Some remarks

Continuous time version?

- Consider observing the process

$$dX_t = D(p_t)dt + dW_t, \quad X_0 = x.$$

Some remarks

Continuous time version?

- Consider observing the process

$$dX_t = D(p_t)dt + dW_t, \quad X_0 = x.$$

- Dynamic pricing formulation:

$$V = \sup_p \mathbb{E} \left[\int_0^\infty e^{-rt} p_t D(p_t) dt \right].$$

Some remarks

Continuous time version?

- Consider observing the process

$$dX_t = D(p_t)dt + dW_t, \quad X_0 = x.$$

- Dynamic pricing formulation:

$$V = \sup_p \mathbb{E} \left[\int_0^\infty e^{-rt} p_t D(p_t) dt \right].$$

- Can write down HJB, characterise solutions.

Some remarks

Continuous time version?

- Consider observing the process

$$dX_t = D(p_t)dt + dW_t, \quad X_0 = x.$$

- Dynamic pricing formulation:

$$V = \sup_p \mathbb{E} \left[\int_0^\infty e^{-rt} p_t D(p_t) dt \right].$$

- Can write down HJB, characterise solutions.
- Explicit examples? In working progress...

Some remarks

Continuous time version?

- Consider observing the process

$$dX_t = D(p_t)dt + dW_t, \quad X_0 = x.$$

- Dynamic pricing formulation:

$$V = \sup_p \mathbb{E} \left[\int_0^\infty e^{-rt} p_t D(p_t) dt \right].$$

- Can write down HJB, characterise solutions.
- Explicit examples? In working progress...
- Arbitrary prior?

Thank you for your attention!

Contact: yuqiong.wang@math.uu.se