

1 ^{1,†} **and** ^{1,†}

2 1
3 2

4 *corresponding author(s):

5 †these authors contributed equally to this work

6 **ABSTRACT**

7

8 **Introduction**

9 GAP: No presynaptic model has taken into account temperature differences in release probability only calcium¹.

10 **Model**

11 **Transitions between the presynaptic vesicle pools**

12 Three pools of vesicles:

- 13 • Recycling pool
- 14 • Docked pool (i.e. readily releasable pool RRP)
- 15 • Fused pool

Calcium-dependent release probability (from docked to fused):

$$p_{rel} =$$

16 **Calcium dynamics**

$$\begin{aligned}\frac{dCa_{pre}}{dt} &= -\frac{Ca_{pre}}{\tau_{pre}} \\ \frac{dCa_{jump}}{dt} &= \frac{1 - Ca_{jump}}{\tau_{rec}} - \delta_{Ca} \cdot Ca_{jump} \cdot Ca_{pre} \\ Ca_{pre} &\leftarrow Ca_{pre} + Ca_{jump} \text{ upon spike arrival}\end{aligned}$$

Table 1. Model variables.

Symbol	Description	Unit
R	Number of vesicles in the recycling pool	-
D	Number of docked vesicles	-
F	Number of fused vesicles	-
Ca_{pre}	Intracellular presynaptic calcium concentration	mM
Ca_{jump}	Jump in Ca_{pre} upon spike arrival	mM

Table 2. Model parameters.

Symbol	Description	Unit	Reference values
D_0	Docked pool capacity	-	5-20 vesicles per bouton in cultured rat hipp. neurons ^{2,3}
R_0	Recycling pool capacity	-	15-40 vesicles per bouton in cultured rat hipp. neurons ($\sim 3\times$ larger than the docked pool capacity) ^{2,3}
$\tau_{R \rightarrow D}$	Rate from recycling to docked		
$\tau_{D \rightarrow R}$	Rate from docked to recycling (mixing)		
$\tau_{F \rightarrow R}$	Rate from fused to recycling		
τ_{pre}	Ca_{pre} attenuation recovery rate	s	
τ_{rec}	Calcium jump recovery rate	s	
δ_{Ca}			
Sigmoid parameters			

Table 3. Datasets.

Reference	Stim. conditions	Tissue	Age	Method	aCSF
Fernández-Alfonso T. et al. 2008 ⁴	300AP at 2, 5, 10, 20, 30 Hz 1800AP at 30 Hz (Baf, 0.5 μ M) (for both 25C and 35C)	Rat cult. hipp. (autaptic)	P1-3	spHluorin (pH imaging)	119 NaCl, 2.5 KCl, 2 CaCl₂ , 2 MgCl ₂ , 30 glucose 25 HEPES (buf.pH 7.4) 10 μ M CNQX and 50 μ M AP5
delvendahl2016fast ⁵	temperature 25, 30 and 35	Mice cult. hipp. (autaptic)			

References

- Wang, B. & Dudko, O. K. A theory of synaptic transmission. *Elife* **10**, e73585 (2021).
- Rizzoli, S. O. & Betz, W. J. Synaptic vesicle pools. *Nat. Rev. Neurosci.* **6**, 57–69, [10.1038/nrn1583](https://doi.org/10.1038/nrn1583) (2005).
- Schikorski, T. & Stevens, C. F. Morphological correlates of functionally defined synaptic vesicle populations. *Nat. Neurosci.* **4**, 391–395, [10.1038/86042](https://doi.org/10.1038/86042) (2001).
- Fernández-Alfonso, T. & Ryan, T. A. The Kinetics of Synaptic Vesicle Pool Depletion at CNS Synaptic Terminals. *Neuron* **41**, 943–953, [10.1016/S0896-6273\(04\)00113-8](https://doi.org/10.1016/S0896-6273(04)00113-8) (2004).
- Delvendahl, I., Vyleta, N. P., von Gersdorff, H. & Hallermann, S. Fast, temperature-sensitive and clathrin-independent endocytosis at central synapses. *Neuron* **90**, 492–498 (2016).