

1,† and 1,†

1

2

3

4 *corresponding author(s):

5 †these authors contributed equally to this work

6 ABSTRACT

7

8 Introduction

9 Model

Table 1. Model parameters.

Symbol	Description	Unit	Experimental litterature
R_0	Recycling pool capacity	-	
D_0	Docked pool capacity	-	5-20 vesicles per bouton in cultured rat hippocampal neurons ^{1,2}
$\tau_{R \rightarrow D}$	Vesicle transition rate from recycling to docked		
$\tau_{D \rightarrow R}$	Vesicle transition rate from docked to recycling (mixing)		
$\tau_{F \rightarrow R}$	Vesicle transition rate from fused to recycling		
$[Ca^{2+}]_o$	Extracellular calcium concentration	mM	
τ_{pre}	$[Ca^{2+}]_i$ attenuation recovery rate	s	
τ_{rec}	Calcium jump recovery rate	s	
δ_{Ca}			

10 References

1. Rizzoli, S. O. & Betz, W. J. Synaptic vesicle pools. *Nat. Rev. Neurosci.* **6**, 57–69, [10.1038/nrn1583](https://doi.org/10.1038/nrn1583) (2005).
2. Schikorski, T. & Stevens, C. F. Morphological correlates of functionally defined synaptic vesicle populations. *Nat. Neurosci.* **4**, 391–395, [10.1038/86042](https://doi.org/10.1038/86042) (2001).