

**NANYANG
TECHNOLOGICAL
UNIVERSITY**
SINGAPORE

Experiment 1: Parametric Curves

CZ2003 Computer Graphics and Visualization

SS3

Name	Matric Number
Pang Yu Shao	U17216 <u>80</u> D

SCHOOL OF COMPUTER SCIENCE AND ENGINEERING
NANYANG TECHNOLOGICAL UNIVERSITY
SINGAPORE
2nd Febraury 2021

CONTENTS

Contents	1
1 Defining Surfaces Parametrically	2
1.1 Plane Passing Through Three Defined Points	2
1.2 Triangular Polygon with Three Defined Vertices	2
1.3 Origin-Centered Ellipsoid with Defined Semi-axes	3

1 DEFINING SURFACES PARAMETRICALLY

1.1 Plane Passing Through Three Defined Points

To define the plane parametrically, we can use the following formula: $P = P1 + u(P2 - P1) + v(P3 - P1)$

Therefore, with the 3 points $(N, M, 0)$, $(0, M, N)$, $(N, 0, M)$, we get:

$$x(u, v) = N - Nu = 8 - 8u$$

$$y(u, v) = M + Mv = 10 + 10v$$

$$z(u, v) = Nu + Mv = 8u + 10v$$

$$u, v \in [0, 1]$$




Fig. 1. Plots of the plane defined in "1a.wrl" with differing resolutions

As seen in Fig. 1 above, a sampling resolution of **1** for both u and v is sufficient for drawing the plane as it has no curvature and having a higher resolution would produce the exact same drawing.

1.2 Triangular Polygon with Three Defined Vertices

To define the Triangular Polygon, we use the formula for defining Bilinear Surface Parametrically, and we set two of the points to be the same point, essentially resulting in a Triangular polygon.

$$P = P1 + u(P2 - P1) + v(P3 - P1 + u(P4 - P3 - (P2 - P1)))$$

Let $P4 = P3$, we get:

$$P = P1 + u(P2 - P1) + v(P3 - P1) + uv(P1 - P2)$$

Therefore, with the 3 points $(N, M, 0)$, $(0, M, N)$, $(N, 0, M)$, we get:

$$x(u, v) = N - Nu + Nu v = 8 - 8u + 8uv$$

$$y(u, v) = M - Mv = 10 - 10v$$

$$z(u, v) = Nu + Mv - Nu v = 8u + 10v - 8uv$$

$$u, v \in [0, 1]$$




Fig. 2. Plots of the Triangular Polygon defined in "1b.wrl" with differing resolutions

As seen in Fig. 2 above, a sampling resolution of **1** for both u and v is sufficient for drawing the triangular polygon as it has no curvature and having a higher resolution would produce the exact same drawing.

1.3 Origin-Centered Ellipsoid with Defined Semi-axes

An ellipsoid can be parametrically defined using the following functions:

$$x = a \cos(u) \sin(v)$$

$$y = b \sin(u)$$

$$z = c \cos(u) \cos(v)$$

Where $u \in [-\pi/2, \pi/2]$ and $v \in [-\pi, \pi]$. Therefore, we can get:

$$x = N \cos(-\pi/2 + \pi u) \sin(-\pi + 2\pi v) = 8 \cos(-\pi/2 + \pi u) \sin(-\pi + 2\pi v)$$

$$y = M \sin(-\pi/2 + \pi u) = 10 \sin(-\pi/2 + \pi u)$$

$$z = (N + M/2) \cos(-\pi/2 + \pi u) \cos(-\pi + 2\pi v) = 9 \cos(-\pi/2 + \pi u) \cos(-\pi + 2\pi v)$$

$$u, v \in [0, 1]$$




Fig. 3. Plots of the Triangular Polygon defined in "1c.wrl" with differing resolutions

The sampling resolution for v is chosen to be 2 times larger than that of u as the coefficient of v is 2 times larger. As seen in Fig. 3 above, the best resolution obtained was 50, 100 for u and v respectively. By decreasing the resolution to 25 and 50, the interpolations could be seen when zoomed in. When the resolution was increased instead to 100 and 200, no visible difference could be seen.