

11.6 Differential Derivatives and the Gradient Vector

$D_{\vec{u}}f(x_0, y_0)$, \vec{u} is a unit vector

Def

The directional derivative of f at (x_0, y_0) is the direction of a unit vector $\vec{u} = \langle a, b \rangle$ is

$$D_{\vec{u}}f(x_0, y_0) = \lim_{h \rightarrow 0} \frac{f(x_0 + h_0, y_0 + h_0) - f(x_0, y_0)}{h}$$

if this limit exists.

Theorem

If f is a differentiable function of x & y , then

$$D_{\vec{u}}f(x, y) = f_x(x, y)a + f_y(x, y)b \quad \vec{u} = \langle a, b \rangle \quad \& \quad \vec{u} = \text{unit vector}$$

Note

If \vec{u} makes angle θ with the positive x -axis, then $\vec{u} = \cos \theta, \sin \theta$.

Ex 1

Find $D_{\vec{u}}f(x, y)$ if $f(x, y) = x^3 - 3xy + 4y^2$ & \vec{u} is the unit vector given by angle $\theta = \frac{\pi}{6}$. What is $D_{\vec{u}}f(1, 2)$?

$$a = \cos \frac{\pi}{6} = \frac{\sqrt{3}}{2} \quad b = \sin \frac{\pi}{6} = \frac{1}{2}$$

$$f_x(x, y) = 3x^2 - 3y \quad f_y(x, y) = -3x + 8y$$

$$D_{\vec{u}}f(x, y) = (3x^2 - 3y)\frac{\sqrt{3}}{2} + (-3x + 8y)\frac{1}{2}$$

$$D_{\vec{u}}f(x, y) = (3(1)^2 - 3(2))\frac{\sqrt{3}}{2} + (-3(1) + 8(2))\frac{1}{2}$$

$$D_{\vec{u}}f(x, y) = -\frac{3\sqrt{3}}{2} + \frac{13}{2}$$

$$D_{\vec{u}}f(x, y) = \frac{-3\sqrt{3} + 13}{2}$$

The Gradient Vector

$$D_{\vec{u}}f(x, y) = f_x(x, y)a + f_y(x, y)b$$

$$\langle f_x(x, y), f_y(x, y) \rangle \cdot \langle a, b \rangle$$

$$D_{\vec{u}}f(x, y) = \nabla f(x, y) \cdot \vec{u} \quad \nabla f(x, y) = \text{gradient vector of } f$$

Ex 2

Find the directional derivative of the function $f(x, y) = x^2y^3 - 4y$ of the point $(2, -1)$ in the direction of the vector $\vec{v} = 2\vec{i} + 5\vec{j}$

$$\nabla f(x, y) = \langle 2xy^3, 3x^2y^2 - 4 \rangle$$

$$\nabla f(2, -1) = \langle 2(2)(-1)^3, 3(2)^2(-1)^2 - 4 \rangle$$

$$\nabla f(2, -1) = \langle -4, 8 \rangle$$

$$\vec{u} = \frac{\vec{v}}{|\vec{v}|} = \frac{\langle 2, 5 \rangle}{\sqrt{29}}$$

$$D_{\vec{u}}f(2, -1) = \nabla f(x, y) \cdot \vec{u}$$

$$D_{\vec{u}}f(2, -1) = \langle -4, 8 \rangle \cdot \left\langle \frac{2}{\sqrt{29}}, \frac{5}{\sqrt{29}} \right\rangle$$

$$D_{\vec{u}}f(2, -1) = \frac{32}{\sqrt{29}}$$

Functions of 3 Variables

For a function f of 3 variables, the gradient vector ∇f is

$$\nabla f(x, y, z) = \langle f_x(x, y, z), f_y(x, y, z), f_z(x, y, z) \rangle$$

The directional derivative of f at (x_0, y_0, z_0) in the direction of a unit vector $\vec{u} = \langle a, b, c \rangle$ is

$$D_{\vec{u}}f(x_0, y_0, z_0) = \nabla f(x_0, y_0, z_0) \cdot \vec{u}$$

Ex 3A

If $f(x, y, z) = x \sin(yz)$, find ∇f

$$f_x(x, y, z) = \sin yz \quad f_y(x, y, z) = x \cdot \cos yz \cdot z = xz \cos yz \quad f_z(x, y, z) = x \cdot \cos yz \cdot y = xy \cos(yz)$$

$$\nabla f = \langle \sin(yz), xz \cos(yz), xy \cos yz \rangle$$

Ex 3B

Find the directional derivative of f at $(1, 3, 0)$ in the direction of $\vec{v} = \langle 1, 2, -1 \rangle$

$$\vec{u} = \frac{\vec{v}}{|\vec{v}|} = \frac{\langle 1, 2, -1 \rangle}{\sqrt{6}} = \left\langle \frac{1}{\sqrt{6}}, \frac{2}{\sqrt{6}}, -\frac{1}{\sqrt{6}} \right\rangle$$

$$D_{\vec{u}}f(x, y, z) = \nabla f(x, y, z) \cdot \vec{u}$$

$$D_{\vec{u}}f(x, y, z) = \frac{\sin(yz)}{\sqrt{6}} + \frac{2xz \cos(yz)}{\sqrt{6}} - \frac{xy \cos(yz)}{\sqrt{6}}$$

$$D_{\vec{u}}f(1, 3, 0) = \frac{\sin 0}{\sqrt{6}} + 0 - \frac{3 \cos 0}{\sqrt{6}}$$

$$\boxed{D_{\vec{u}}f(1, 3, 0) = -\frac{3}{\sqrt{6}}}$$