Rebuttal for ICML'25 Submission #3922

Table 1: Corresponding to Q4,Q6 of Reviewer HcSt, Q3 of Reviewer Wxrf. The RMSE result on correlation prediction of $|\psi_{\rm HB}\rangle$ with varied N and $n_{\rm sft}$. M is fixed to 64. ResMLP(CNN)-a-b-cxd represents neural network MLP (CNN) that composed of d+2 in the order a-,b-, d layers of c-width fully connected (convolutional) layers with residual connection. The best results are highlighted in boldface while the second-best results are distinguished in underlined.

Methods	N = 48				N = 63			N = 100		N = 127		
Methods	$n_{\rm sft} = 20$	$n_{\rm sft} = 60$	$n_{\rm sft} = 100$	$n_{\rm sft} = 20$	$n_{\rm sft} = 60$	$n_{\rm sft} = 100$	$n_{\rm sft} = 20$	$n_{\rm sft} = 60$	$n_{\rm sft} = 100$	$n_{\rm sft} = 20$	$n_{\rm sft} = 60$	$n_{\rm sft} = 100$
CS	0.21113			0.21257			0.21399			0.21447		
ResMLP-128x2	0.08282	0.07752	0.06616	0.12055	0.08776	0.07086	0.10848	0.08158	0.07405	0.10091	0.10083	0.08245
ResMLP-128x3	0.06214	0.04853	0.04494	0.07256	0.05506	0.04467	0.07740	0.06496	0.07098	0.08535	0.08280	0.08691
ResMLP-128x4	0.05428	0.03825	0.03524	0.06463	0.04435	0.03833	0.07532	0.05952	0.06010	0.07971	0.09173	0.08608
ResMLP-128x5	0.07228	0.04721	0.03764	0.07308	0.05957	0.05091	0.08046	0.07146	0.07174	0.08408	0.08650	0.08458
ResCNN-32-64	0.07160	0.04723	0.03795	0.07176	0.04066	0.03042	0.06549	0.04566	0.03464	0.06468	0.03189	0.07404
ResCNN-32-64-128	0.08089	0.03422	0.03435	0.09003	0.03401	0.03159	0.07603	0.03245	0.03295	0.08420	0.03179	0.03025
ResCNN-32-64-128x2	0.06484	0.04899	0.03456	0.06621	0.03608	0.03100	0.06436	0.03425	0.02808	0.07441	0.03196	0.05221
ResCNN-32-64-128x2-64	0.17049	0.15600	0.16109	0.18302	0.13459	0.14636	0.13584	0.14809	0.11732	0.17961	0.09261	0.12525
LLM4QPE-T	0.05189	0.03368	0.03197	0.06111	0.03364	0.02863	0.05050	0.03227	0.02726	0.05079	0.03184	0.02634
RBFK	0.05452	0.04176	0.04101	0.04726	0.03829	0.03922	0.04096	0.03299	0.03282	0.03850	0.03115	0.03086
Lasso	0.04221	0.02636	0.02489	0.04856	0.02791	0.02326	0.04219	0.02602	0.02646	0.04137	0.03292	0.02083
Ridge	0.04247	0.02884	0.02475	0.04216	0.02816	0.02402	0.04191	0.02711	0.02251	0.04110	0.02620	0.02161

Table 2: Corresponding to Q4,Q6 of Reviewer HcSt, Q3 of Reviewer Wxrf. The RMSE result on correlation prediction of $|\psi_{\text{TFIM}}\rangle$ with varied N and n_{sft} . M is fixed to 64. ResXXX-a-b-cxd represents neural network MLP (CNN) that composed of d+2 in the order a-,b-, d layers of c-width fully connected (convolutional) layers with residual connection. The best results are highlighted in **boldface** while the second-best results are distinguished in <u>underlined</u>.

Methods		N = 48			N = 63			N=100			N=127		
Methods	$n_{\rm sft} = 20$	$n_{\rm sft} = 60$	$n_{\rm sft} = 100$	$n_{sft} = 20$	$n_{\rm sft} = 60$	$n_{\rm sft} = 100$	$n_{sft} = 20$	$n_{\rm sft} = 60$	$n_{\rm sft} = 100$	$n_{\rm sft} = 20$	$n_{\rm sft} = 60$	$n_{\rm sft} = 100$	
CS	0.20924				0.20990			0.21092			0.21180		
ResMLP-128x2	0.07899	0.06371	0.05524	0.07986	0.05279	0.04283	0.08293	0.05303	0.04630	0.07908	0.05006	0.04333	
ResMLP-128x3	0.06080	0.05664	0.06074	0.06514	0.06928	0.06914	0.06301	0.06358	0.07317	0.06324	0.06510	0.07327	
ResMLP-128x4	0.05912	0.05794	0.05980	0.05899	0.05705	0.06163	0.05678	0.05628	0.06977	0.05535	0.06496	0.07197	
ResMLP-128x5	0.07422	0.06545	0.05739	0.07341	0.06921	0.069215	0.06648	0.06556	0.07044	0.06941	0.07222	0.06867	
ResCNN-32-64	0.12845	0.15039	0.08935	0.12227	0.16686	0.10315	0.10084	0.08879	0.05177	0.10495	0.08535	0.04647	
ResCNN-32-64-128	0.13545	0.17135	0.12004	0.12545	0.17026	0.11778	0.11433	0.11267	0.05027	0.13312	0.03562	0.05347	
ResCNN-32-64-128x2	0.13624	0.17178	0.12015	0.12608	0.17103	0.13809	0.12221	0.11046	0.06586	0.13757	0.10498	0.05556	
ResCNN-32-64-128x2-64	0.13719	0.16750	0.14122	0.13184	0.16034	0.14158	0.12356	0.14996	0.11282	0.11823	0.03601	0.03785	
LLM4QPE-T	0.05088	0.03493	0.03006	0.05252	0.03566	0.03082	0.05217	0.03476	0.03012	0.05259	0.03641	0.03084	
Lasso	0.04624	0.03219	0.02812	0.04633	0.03930	0.02859	0.04073	0.03256	0.02899	0.04583	0.03283	0.02932	
Ridge	0.04473	0.03173	0.02807	0.04561	0.03226	0.02839	0.04598	0.03277	0.02883	0.04570	0.03285	0.02911	

Table 3: Corresponding to Q2,Q5 of Reviewer HcSt, Q3 of Reviewer 3fBm. The RMSE results on correlation prediction of $|\psi_{\rm HB}\rangle$ with varied N. Training set and testing set are both have 10^4 samples, with noise-free labels $(M \to \infty)$. The best results are highlighted in **boldface**.

$M o \infty$	N=8	N = 10	N = 12	N = 16	N = 25	N = 31
Ridge	0.00367	0.00444	0.00566	0.00636	0.00599	0.00579
ResMLP-128x4	0.03961	0.03677	0.03460	0.03129	0.02769	0.02625
ResCNN-32-64-128x2	0.02056	0.03710	0.03432	0.03050	0.02582	0.02381
LLM4QPE-F	0.04666	0.04385	0.03969	0.03728	0.03083	0.02951

Table 4: Corresponding to Q2,Q5 of Reviewer HcSt, Q3 of Reviewer 3fBm. The RMSE results on predicting correlation of $|\psi_{\rm HB}\rangle$ with varied training size n. System size N=8. The number of testing set is fix to 2×10^4 . Labels are noise-free $(M\to\infty)$. The best results are highlighted in **boldface**.

$M o \infty$	# Params.	$n = 10^2$	$n = 10^{3}$	$n = 10^4$	$n = 10^5$
Ridge	< 0.01M	0.00780	0.00528	0.00367	0.00660
ResMLP-128x4	0.09M	0.04219	0.04172	0.03961	0.03956
ResCNN-32-64-128x2	1.14M	0.01987	0.02078	0.02056	0.02054
LLM4QPE-F	9.89M	0.03966	0.04304	0.04916	0.04659

Table 5: Corresponding to Q2,Q5 of Reviewer HcSt, Q3 of Reviewer 3fBm. The RMSE results on predicting entanglement entropy of $|\psi_{\rm HB}\rangle$ with varied training size n. System size N=8. The number of testing set is fix to 2×10^4 . Labels are noise-free $(M\to\infty)$. The best results are highlighted in **boldface**.

$M \to \infty$	# Params.	$n = 10^2$	$n = 10^3$	$n = 10^4$	$n = 10^5$
Ridge	< 0.01M	0.01563	0.00947	0.00753	0.00851
ResMLP-128x4	0.09M	0.10817	0.09142	0.05398	0.05302
ResCNN-32-64-128x2	1.14M	0.04334	0.02410	0.03520	0.02073
LLM4QPE-F	9.89M	0.10648	0.11171	0.10895	0.10826

Table 6: Corresponding to Q1 of Reviewer m6FX. The RMSE results of LLM4QPE-F on correlation prediction of N-qubit $|\psi_{\rm HB}\rangle$, with embedding $M_{\rm emb}$ random measurement outcomes. Training set and testing set are both have 10⁴ samples, with noise-free labels $(M \to \infty)$. $M_{\rm emb}$ is the actual number of embedded measurement outcomes.

	N=8	N = 10	N = 12	N = 16	N = 25	N = 31
$M_{ m emb} = 1$ $M_{ m emb} = 8$ $M_{ m emb} = 64$	0.04666	0.04385	0.04126	0.03728	0.03083	0.03125
$M_{\rm emb} = 8$	0.04746	0.04926	0.03969	0.03984	0.03408	0.02951
$M_{\rm emb} = 64$	0.04795	0.04791	0.04785	0.04043	0.03637	0.03524
$M_{\rm emb} = 512$	0.04913	0.04521	0.04506	0.03905	0.03406	0.03268

Table 7: Corresponding to Q1 of Reviewer m6FX, Q1 of Reviewer 3fBm. The RMSE results of LLM4QPE-F on correlation prediction of N-qubit $|\psi_{\rm HB}\rangle$, with embedding $M_{\rm emb}$ real measurement outcomes over the finetuning phase. testing size is set to 200. M is fixed to 512. $M_{\rm emb} \leq M$ is the actual number of embedded measurement outcomes. $n_{\rm sft}$ is the training size over the finetuning phase.

		N = 63			N = 100		N = 127			
	$n_{\rm sft} = 20$	$n_{\rm sft} = 60$	$n_{\rm sft} = 100$	$n_{\rm sft} = 20$	$n_{\rm sft} = 60$	$n_{\rm sft} = 100$	$n_{\rm sft} = 20$	$n_{\rm sft} = 60$	$n_{\rm sft} = 100$	
$M_{\rm emb} = 1$	0.02555	0.02104	0.02019	0.02307	0.01872	0.01760	0.02239	0.01739	0.01635	
$M_{ m emb} = 8$	0.02556	0.02106	0.02019	0.02309	0.01873	0.01760	0.02242	0.01739	0.01635	
$M_{\rm emb} = 64$	0.02556	0.02104	0.02019	0.02309	0.01872	0.01759	0.02239	0.01739	0.01636	
$M_{\rm emb} = 512$	0.02560	0.02104	0.02019	0.02309	0.01872	0.01759	0.02240	0.01740	0.01635	

Table 8: Corresponding to Q1 of Reviewer 3fBm. The RMSE results of predicting correlation of N-qubit $|\psi_{\rm HB}\rangle$, with MLP, Lasso and Ridge as learning models. Measurement outcomes are embedded as input features of MLP by two ways: raw tensor directly characterizing, or averaging (Avg.) with M for each qubit $(M \times N \to 1 \times N)$. $N \in \{63, 100, 127\}$. Training size $n \in \{20, 80, 100\}$. Measurement shots $M \in \{64, 128, 256, 512\}$.

			N = 63			N = 100			N = 127	
		n=20	n = 60	n = 100	n=20	n = 60	n = 100	n=20	n = 60	n = 100
	Raw	0.08964	0.05522	0.04872	0.08666	0.04949	0.04055	0.08878	0.05068	0.04076
M = 64	Avg.	0.05572	0.03522	0.02984	0.05525	0.03972	0.02801	0.05505	0.03951	0.03242
WI-04	Lasso	0.04856	0.02791	0.02326	0.04219	0.02602	0.02646	0.04137	0.03292	0.02083
	Ridge	0.04216	0.02816	0.02402	0.04191	0.02711	0.02251	0.04110	0.02620	0.02161
M=128	Raw	0.10921	0.05905	0.04835	0.10966	0.06137	0.04485	0.10408	0.06359	0.04554
	Avg.	0.04403	0.03034	0.02552	0.04699	0.03561	0.02603	0.04435	0.03421	0.03007
M-120	Lasso	0.03168	0.02171	0.01905	0.03127	0.02045	0.01735	0.03041	0.01980	0.01647
	Ridge	0.03169	0.02178	0.01921	0.03069	0.02067	0.01786	0.03053	0.02087	0.01726
	Raw	0.14085	0.08316	0.06045	0.12558	0.08648	0.05983	0.11720	0.08232	0.06089
M = 256	Avg.	0.03581	0.02673	0.02272	0.04022	0.02966	0.02168	0.03883	0.03188	0.02893
M-250	Lasso	0.02556	0.01749	0.12125	0.02406	0.01747	0.01467	0.02283	0.01542	0.01324
	Ridge	0.02556	0.01751	0.01572	0.02408	0.01697	0.01494	0.02286	0.01576	0.01377
	Raw	0.15943	0.11187	0.08246	0.13586	0.10826	0.08329	0.12608	0.10324	0.08330
M=512	Avg.	0.03020	0.02475	0.02211	0.03713	0.02864	0.02272	0.03644	0.02962	0.02618
W131Z	Lasso	0.02037	0.01586	0.11038	0.01892	0.01403	0.01263	0.01702	0.01257	0.01117
	Ridge	0.02036	0.01583	0.01436	0.01891	0.01404	0.01271	0.01798	0.01285	0.01186

Table 9: Corresponding to Q4 of Reviewer 3fBm. The RMSE results of Ridge on predicting correlation of N-qubit $|\psi_{\rm HB}\rangle$ and $|\psi_{\rm TFIM}\rangle$. The input dimension d is both fixed to 20. Regularization of Ridge is set to $\lambda=1$.

Dataset	N = 63					N = 100				N = 127					
Dataset	n = 20	n = 40	n = 60	n = 80	n = 100	n = 20	n = 40	n = 60	n = 80	n = 100	n = 20	n = 40	n = 60	n = 80	n = 100
HB	0.09998	0.10555	0.09941	0.09322	0.08782	0.10015	0.10395	0.09867	0.09278	0.08692	0.09964	0.10491	0.09898	0.09241	0.08680
TFIM	0.10185	0.10333	0.09845	0.09189	0.08565	0.10093	0.10436	0.09847	0.09193	0.08824	0.10148	0.10372	0.10106	0.09426	0.08716