


Varicose Veins



Definition

- Dilated, usually tortuous, subcutaneous veins ≥ 3 mm in diameter measured in the upright position with demonstrable reflux.



Epidemiology

- Adult prevalence of visible varicose veins is 25–30 per cent in women and 15 per cent in men.
- Common in females.
- Incidence increases with age.



Risk factors

- Strong family history
- Age
- Obesity
- Pregnancy
- Smoking
- Constipation
- Prolonged standing

Inconclusive
evidence



Causes

Primary

- Cause not known.
- Often familial.
- Very rarely, congenital absence of the valves.

Secondary

- **Obstruction to venous outflow**
 - Pregnancy
 - Fibroids/ovarian cyst
 - Abdominal lymphadenopathy
 - Pelvic cancer (cervix, uterus, ovary, rectum)
 - Ascites
 - Iliac vein thrombosis
 - Retroperitoneal fibrosis
- **Valve destruction**
 - Deep vein thrombosis
- **High flow and pressure**
 - Arteriovenous fistula



CEAP classification

(clinical – etiology – anatomy – pathophysiology)

Clinical

- C0: No signs of venous disease
- C1: Telangectasia or reticular veins
- C2: Varicose veins
- C3: Oedema
- C4a: Pigmentation or eczema
- C4b: Lipodermatosclerosis or atrophie blanche
- C5: Healed venous ulcer
- C6: Active venous ulcer

Etiologic

- Ec: Congenital
- Ep: Primary
- Es: Secondary (post-thrombotic)
- En: No venous cause identified



Anatomical

- As: Superficial veins
- Ap: Perforator veins
- Ad: Deep veins
- An: No venous location identified

Pathophysiological

- Pr: Reflux
- Po: Obstruction
- Pr,o: Reflux and obstruction
- Pn: No venous pathophysiology identifiable



Clinical features

1. Symptoms

Due to varicose veins

- Aching or heaviness
 - ☐ Increases throughout the day or with prolonged standing.
 - ☐ Relieved by elevation or compression hosiery.
- Ankle swelling
- Itching
- Cosmetic concerns

Due to complications


- Bleeding
- Superficial thrombophlebitis
- Eczema
- Lipodermatosclerosis
- Ulceration



2. Signs

- Tortuous dilated subcutaneous veins
 - ❑ Long saphenous incompetence - medial thigh and calf varicosities
 - ❑ Short saphenous incompetence - posterolateral calf varicosities
 - ❑ Isolated incompetence of the proximal anterolateral long saphenous tributary - anterolateral thigh and calf varicosities
- Telangectasia
- Reticular veins
- Saphena varix
- Atrophie blanche
- Corona phlebectasia
- Lipodermatosclerosis
- Ulceration
- Pigmentation
- Eczema







Reticular veins





Eczema



Severe chronic venous disease



Localised chronic inflammation and fibrosis of the skin and subcutaneous tissues of the leg.



Lipodermatosclerosis and Lead Champagne bottle appearance.



RISHACADEMY

educate yourself to empower yourself

www.rishacademy.com

Leakage of RBC
from engorged
vessels



Degradation of
haemoglobin and
deposition of
haemosiderin



Pigmentation



RISHACADEMY

educate yourself to empower yourself


www.rishacademy.com


Tourniquet tests

1. Procedure

- The patient should lie on a couch.
- The limb to be examined is then elevated to empty the veins.
- A tourniquet made from a long length of 1 cm diameter soft rubber tubing is then pulled tight around the upper thigh and held in place by strong artery forceps.
- The patient is then asked to stand up quickly, and the legs are observed for 10–15 seconds.







2. Interpretation

Saphenofemoral junction incompetence

- Veins above the tourniquet will rapidly fill, but those below it will remain collapsed when patient stands up.
- Confirmed by suddenly releasing the tourniquet and watching the veins below the site of the tourniquet rapidly distend from above.

Other sites of superficial to deep incompetence

- Veins below the tourniquet fill immediately when patient stands up.
- Test can be repeated with the tourniquet moved progressively down the whole length of the leg to try to define all the sites of superficial to deep vein incompetence.
- Apply it once below the knee to exclude short saphenous incompetence.



Investigation

- **Venous duplex**


- ☐ Shows superficial and deep veins
- ☐ Can identify the site of incompetence
- ☐ Recommended for all the patients who undergoing intervention

- **Contrast venography**

- ☐ Conventional
 - If duplex is inconclusive
- ☐ MRV/CTV
 - Assess intra-abdominal and pelvic veins



Management



Compression hosiery

- It relies on graduated external pressure to improve deep venous return.
- Compression hosiery can be knee length or thigh length.
- Compression hosiery is classified according to the pressure exerted,
 - class 1 exert 14–17 mmHg
 - class 2 exert 18–24 mmHg
 - class 3 exert 25–35 mmHg
- Recommended Stockings are - Class 2 and Class 3.
- Significantly improves varicose vein symptoms.



Ultrasound-guided foam sclerotherapy

- Sclerosants
 - ☐ **Chemical irritants** - glycerine
 - ☐ **Osmotic** - hypertonic saline
 - ☐ **Detergent** - sodium tetradecyl sulphate (**STD**), Polidocanol
- Injection of detergent directly into the superficial veins.
- Only 1 or 2 mL of foam should be injected at a time and the distribution of the foam should be monitored and massaged with the ultrasound probe.
- Compression bandaging or hosiery is then applied and left in situ for 7–10 days.
- Sclerotherapy improves varicose vein-related symptoms, but recurrence rates and the need for reintervention are relatively high.
- Complications - phlebitis, pigmentation, headache, visual disturbance, chest tightness, cough.



Endovenous laser ablation(EVLA)

- Insertion of a laser fibre into the lumen of an incompetent truncal vein, with subsequent thermal ablation of the vein.
- The tumescent anaesthesia,
 - Dilute lidocaine with adrenaline and bicarbonate warmed to body temperature.
 - Provides analgesia and compresses the vein.
 - Increasing the contact area between the vein wall and laser fibre.
 - Protects adjacent structures (skin, nerves).
- EVLA treats only junctional and truncal incompetence.
- Compression and analgesia are usually recommended in the early postoperative period.



Radiofrequency ablation(RFA)

- It is a minimally invasive endovascular therapy that uses a bipolar catheter to generate thermal energy to ablate the vein.
- Preoperative planning and consent, patient positioning, tumescent anaesthesia administration and postoperative management are identical to EVLA.
- The main differences include vein cannulation and method of ablation.



- Local 'stab' avulsions.
 - Deals with varicosities.
- Saphenofemoral or saphenopopliteal ligation.
- Long saphenous vein stripping.
 - Effectively avulses all incompetent thigh perforators
 - Not usually done below the knee due to risk of saphenous nerve injury.



Post operative care

- Graduated compression stockings or bandages are worn day and night for 7-10 days. Thereafter only during day for 1 month.
- Oral analgesics
- Leg elevation



Complications of surgery

- Bleeding or bruising
- Healing with fibrosis- contracture
- Wound infections
- Neuropraxia or complete damage
 - Saphenous nerve accompanies LSV
 - Sural nerve accompanies SSV
 - Sapheno popliteal ligation may lead to sural nerve and common peroneal nerve damage.
- Recurrence
- DVT



Non-Thermal Non –Tumescent methods

- Mechano-chemical endovenous ablation(MOCA).
- Venaseal closure system.

