


# Liver Abscess




**RISH**ACADEMY


educate yourself to empower yourself

[www.rishacademy.com](http://www.rishacademy.com)

# LIVER ABSCESS



Pyogenic liver  
abscess



Amoebic liver  
abscess



# PYOGENIC LIVER ABSCESS

- Common causes include biliary stone disease and other causes of intra-abdominal sepsis, including appendicitis and diverticular disease.
- Usually found in the right lobe
- Occurs secondary to other sources of bacterial sepsis



- The diagnosis is suggested by the finding of a multiloculated cystic mass on ultrasound or CT scan and is confirmed by aspiration for culture and sensitivity.



# Causative bacteria

- Escherichia coli- commonest
- Streptococcus milleri
- Klebsiella
- Proteus vulgaris
- Pseudomonas
- Clostridia



# Clinical features

- Fever (high) with chills and rigors
- Right upper quadrant pain
- Weight loss
- Jaundice- occasionally
- Intercostal tenderness
- Tender, soft liver



# Investigations

Blood



- Leukocytosis
- Increased alkaline phosphatase

Chest X Ray



- Elevation of right hemidiaphragm
- R/S pleural effusion?
- Soft tissue shadow

CT scan



# Treatment

Antibiotics



Penicillin  
Aminoglycoside  
Metronidazole  
Cephalosporin  
Metronidazole

ultrasound-guided  
aspiration



Repeated  
aspirations may be  
necessary





# AMOEBIIC LIVER ABSCESS

- Caused by *Entamoeba histolytica*
- 7-10 times more frequent in adult men. More common in alcoholic and cirrhotic patients
- spread by the fecal–oral route.



Entamoeba histolytica is endemic in many parts of the world. It exists in vegetative form outside the body and is The most common presentation is with dysentery, but it may also present with an amoebic abscess, the common sites being paracaecal and in the liver




Amoebic  
cyst



Ingested and develops  
into the trophozoite  
form in the colon



liver



through the  
bowel wall  
and to the  
liver via the



passes  
through the  
bowel wall



# Clinical features

- Persistent fever
- Right upper quadrant pain
- Hepatomegaly- soft, tender, smooth
- Point tenderness over the liver- Intercostal tenderness
- Right pleural effusion- Due to hyperaemia
- Peritonitis due to rupture of abscess
- Empyema
- Retroperitoneal abscess
- Pericardial tamponade



# Investigations

- USS
- CT scan
- Chest X ray- for pleural effusion
- Isolation of the parasite from the liver lesion or the stool



# Treatment

- Oral or IV Metronidazole (400– 800 mg tds for 7–10 days)
- Oral or IV Ciprofloxacin- for control of 2ry infection
- Needle aspiration
- Surgery- Rarely needed

