

TYPE 2 DIABETES

Age-group effects, organ-specific physiology, ADMET-style profiling, and what changes when medication dose is increased or decreased

Based on the ADMET dashboard concept in the reference image: absorption, distribution, metabolism, excretion, and toxicity are used here as a way to organize medication behavior. For diabetes medicines, these are combined with target-organ physiology, age-related considerations, and dose-response patterns.

Important terminology: Type 2 diabetes is generally not classified clinically into “Phase 1” and “Phase 2.” This document interprets your “phase 2” wording as **established Type 2 diabetes** requiring ongoing glucose-lowering management.

Scope: metformin, sulfonylureas, pioglitazone, DPP-4 inhibitors, SGLT2 inhibitors, GLP-1 receptor agonists, tirzepatide, and basal insulin concepts. Dose examples are reference-level ranges/levels from labeling or guideline tables; they are **not instructions to change a patient’s dose**.

Prepared as a research/education reference - September 2026

1. Executive overview

Type 2 diabetes is a chronic disorder in which insulin resistance, progressive beta-cell dysfunction, abnormal hepatic glucose production, altered incretin signaling, and changes in renal glucose handling contribute to elevated blood glucose. The clinical picture varies considerably by age and comorbidity.

Key idea for an ML/ADMET system: dose is only one input. A better representation combines medication + dose + age + kidney function + baseline HbA1c/glucose + weight/BMI + comorbidities + co-medications + duration of treatment + observed response.

Domain	What to capture	Typical signal in Type 2 diabetes
Glucose	Fasting glucose, post-meal glucose, HbA1c	Improves as effective therapy lowers glucose; insufficient therapy leaves hyperglycemia.
Pancreas	Beta-cell function and insulin secretion	Often progressively impaired over years; secretagogues increase insulin release but can cause hypoglycemia.
Liver	Hepatic glucose production	Excess hepatic glucose output contributes to fasting hyperglycemia; metformin targets this pathway.
Kidney	eGFR, albuminuria, volume status	Kidney disease changes medication choice/dosing and also represents a major complication pathway.
Heart/vessels	ASCVD, heart failure, BP, lipids	Cardiovascular risk is a central treatment consideration; some agents provide cardiorenal benefit beyond A1C lowering.
Adipose/weight	Body weight, appetite, waist/BMI	Weight gain is common with some secretagogues and insulin; GLP-1/GIP therapies tend to reduce appetite and weight.
Nervous system	Hypoglycemia awareness, cognition, function	Older adults can have less obvious hypoglycemia symptoms and greater consequences from falls/confusion.

The 2026 ADA Standards emphasize person-centered medication selection based on comorbidities, adverse effects, hypoglycemia risk, treatment burden, and patient goals. They also emphasize cardiorenal-protective therapies when relevant.

2. Effects across age groups

Age changes the risk-benefit balance of treatment. The important distinction is not that the same medication suddenly becomes “toxic” at a certain birthday; rather, kidney function, frailty, nutrition, cognition, polypharmacy, and cardiovascular disease become more influential with aging.

Age group	Typical concerns in established T2D	Medication-response considerations	Monitoring emphasis
Children/adolescents (12-17)	Weight/obesity, insulin resistance, puberty-related variability, adherence, family support.	Therapies with pediatric indications are more limited than adult options; dosing must follow pediatric labeling and specialist guidance.	Glucose/HbA1c, growth, weight/BMI, adherence, psychosocial factors.
Younger adults (18-39)	Long lifetime exposure to hyperglycemia; obesity, fatty liver, reproductive considerations may matter.	Long-term weight, cardiovascular and kidney effects matter; treatment may be intensified when glucose is not at goal.	HbA1c, glucose patterns, weight, BP, lipids, kidney and eye screening.
Middle-aged adults (40-64)	Accumulating ASCVD, CKD, hypertension, dyslipidemia, neuropathy and retinopathy risk.	Medication selection increasingly depends on cardiovascular/kidney status as well as glucose lowering.	HbA1c/glucose, eGFR/UACR, BP, lipids, cardiovascular risk, eye/foot assessments.
Older adults (65-74)	Frailty may begin to emerge; polypharmacy and hypoglycemia consequences are more important.	Avoid overtreatment; simplify regimens when appropriate while maintaining individualized goals.	Hypoglycemia, renal function, cognition, falls, nutrition, functional status, treatment burden.
Older adults (75+)	Higher likelihood of CKD, frailty, cognitive/functional impairment, dehydration, and medication interactions.	The most aggressive glucose lowering is not automatically the safest. ADA highlights individualized goals and deintensification when burdens/risks outweigh benefits.	Renal function, dehydration/volume status, cognition, meals, falls, hypoglycemia, caregiver support.

ADA 2026 explicitly addresses age as a life-span issue and notes that intensive regimens using insulin and/or sulfonylureas can constitute overtreatment in some older adults, particularly when health status is complex. Deintensification can include lowering doses or discontinuing medicines while maintaining individualized goals.

3. What established Type 2 diabetes can do to the body

Persistent hyperglycemia contributes to vascular and metabolic injury. The following is a system-level map for a dataset; it does not mean every person develops every complication.

Organ/system	Main diabetes-related effect	Useful dataset variables
Pancreas	Progressive beta-cell dysfunction; reduced capacity to compensate for insulin resistance.	Fasting C-peptide (if available), insulin use, duration of diabetes, HbA1c.
Liver	Excess hepatic glucose production; metabolic fatty liver can coexist with insulin resistance.	ALT/AST, liver fat diagnosis, fasting glucose, BMI/waist.
Kidneys	Diabetic kidney disease can cause albuminuria and declining eGFR; kidney function also changes medication exposure/eligibility.	eGFR, UACR, creatinine, potassium, hydration/volume status.
Eyes	Retinal microvascular injury can progress to diabetic retinopathy and vision loss.	Retinopathy stage, eye exam timing, HbA1c duration.
Nerves	Peripheral and autonomic neuropathy can cause pain, numbness, altered sensation, GI or orthostatic symptoms.	Neuropathy status, foot findings, pain score, orthostatic symptoms.
Heart and vessels	Higher risk of atherosclerotic cardiovascular disease and heart failure.	ASCVD history, HF status, BP, lipids, smoking, kidney disease.
Feet/skin	Loss of sensation plus vascular disease can increase ulcer/infection risk.	Foot exam, pulses, neuropathy, prior ulcer/amputation.
Brain/function	Hypoglycemia can cause confusion, seizure, injury; chronic vascular disease can affect cognitive health.	Hypoglycemia events, cognition, falls, frailty, glucose variability.

4. ADMET-style profiling for Type 2 diabetes medicines

The reference image shows five axes: Absorption, Distribution, Metabolism, Excretion, and Toxicity. For diabetes pharmacotherapy, these can be extended with target organ/system and clinical response.

Dimension	What it means	Diabetes-specific examples
Absorption	How drug enters systemic circulation and how food/formulation changes exposure.	Oral vs injectable route; food effects; delayed gastric emptying matters for some GLP-1 therapies.
Distribution	How drug moves through body tissues/fluids.	Protein binding, tissue penetration, and body composition can influence exposure; clinically relevant details depend on the specific drug.
Metabolism	Biotransformation, often involving liver enzymes or other pathways.	Some drugs have hepatic metabolism; others are more dependent on unchanged renal excretion.
Excretion	How drug leaves the body.	Renal function is central for several diabetes drugs; eGFR can therefore be a critical model feature.
Toxicity/safety	Adverse effects and organ-specific hazards at therapeutic exposure.	Hypoglycemia, GI intolerance, dehydration, genital infections, edema/heart failure risk, and other class-specific effects.
Target organ/system	The principal physiological site/pathway affected.	Liver for metformin; pancreatic beta cells for sulfonylureas; kidney tubules for SGLT2 inhibitors; pancreas/stomach/brain appetite pathways for GLP-1/GIP agents.

For a predictive system, “affected organ” should mean **target/physiological organ**, not automatically “damaged organ.” A second variable such as **adverse-event organ** can separately record where toxicity presents.

5. Medication dose-response map

The table below describes the direction of change expected when exposure is moved upward or downward within a medicine's labeled range. It is intentionally phrased as a physiological pattern rather than a patient-specific dose adjustment instruction.

Class/example	Main target organ/system	Dose increased within labeled range - typical direction	Dose decreased - typical direction	Key dose-related concerns
Metformin	Liver	More suppression of hepatic glucose production; GI intolerance may increase.	Less glucose-lowering exposure; GI burden may fall.	GI effects; renal contraindication/limitations; over time.
Glimepiride / glipizide	Pancreatic beta cells	More insulin secretion -> lower glucose, but hypoglycemia and weight gain become more likely.	Less insulin-secretagogue effect -> glucose may rise; hypoglycemia risk tends to fall.	Hypoglycemia, especially with missed meals; older adults; higher-consequence group
Pioglitazone	Adipose, muscle, liver	Greater insulin sensitization over time; edema/weight gain may increase.	Less insulin-sensitizing effect; weight/fluid effects may lessen.	Edema, weight gain, heart-failure risk considerations
Sitagliptin	Pancreatic incretin pathway	Typically modest additional glucose lowering; low hypoglycemia risk alone.	Less glucose-lowering effect; renal dosing may dictate the level.	Dose is renal-adjusted rather than freely titrated across all patients.
Empagliflozin / dapagliflozin	Kidney proximal tubule	More glucosuria/diuresis within the labeled range; glucose-lowering may be limited by kidney function.	Less glucosuria and often less diuretic effect; glycemia may worsen.	Volume depletion, genital infections, rare euglycemic DKA; kidney function affects use/effect.
Semaglutide / dulaglutide	Pancreas, stomach, appetite pathways	Usually stronger average glucose and weight effects, but GI adverse effects may increase during escalation.	Usually less glucose/weight effect; GI burden may improve.	GI intolerance and dehydration; dose escalation is gradual to improve tolerability.
Tirzepatide	Pancreas, stomach, appetite pathways	Higher exposure can further lower glucose and body weight on average, while GI burden may rise.	Less incretin effect and generally less glucose/weight reduction.	GI effects, dehydration, and product-specific contraindications/precautions
Basal insulin (concept)	Whole-body glucose regulation	Stronger glucose lowering, including overnight/fasting glucose; hypoglycemia and weight gain risk rise if exposure exceeds needs.	Less insulin effect; persistent hyperglycemia becomes more likely if insufficient.	Hypoglycemia is the major dose-related risk; careful titration and monitoring are essential.

6. Reference dose levels for selected medicines

These are representative labeled strengths/levels useful for a research dataset. They are not a recommended regimen for an individual. Exact maximums and renal restrictions differ by formulation, indication, country, and product label.

Medication	Representative levels	Primary target	Why the dose is moved up/down
Metformin	500, 1000, 1500, 2000 mg/day are common reference levels; some formulations/labels differ.	Liver	Up for additional glycemic effect if tolerated; down/stop if contraindications, intolerance, or clinical reasons require.
Glimepiride	1, 2, 4, 8 mg/day reference sequence.	Pancreatic beta cells	Up for glycemic effect only when appropriate; down when hypoglycemia or overtreatment risk emerges.
Glipizide IR	5, 10, 20 mg and higher labeled daily exposure depending on formulation; IR and ER differ.	Pancreatic beta cells	Dose/formulation specific; reduction may be needed when hypoglycemia or renal/food-intake concerns exist.
Pioglitazone	15, 30, 45 mg/day reference levels.	Adipose, muscle, liver	Slow onset; dose constrained by edema/weight and heart-failure considerations.
Sitagliptin	100 mg/day standard; 50 or 25 mg/day are renal-adjusted levels.	Pancreatic incretin pathway	Renal function, not "stronger glycemic need," determines lower levels.
Empagliflozin	10 and 25 mg/day labeled strengths in relevant T2D use.	Kidney proximal tubule	Higher dose may add glycemic effect when appropriate; kidney function and indication matter.
Semaglutide	0.25, 0.5, 1, 2 mg weekly reference sequence for Ozempic-type glycemic use.	Pancreas, stomach, appetite pathways	0.25 mg is an initiation/tolerability step; maintenance escalation is gradual.
Dulaglutide	0.75, 1.5, 3, 4.5 mg weekly reference sequence.	Pancreas, stomach, appetite pathways	Escalation can increase glycemic effect but also GI burden; changes are spaced for tolerability.
Tirzepatide	2.5, 5, 7.5, 10, 12.5, 15 mg weekly reference sequence.	GIP/GLP-1 pathways	2.5 mg is an initiation dose; higher levels are reached gradually based on response/tolerability.

7. What happens when the dose is too low or too high?

Scenario	Glucose effects	Possible body effects	Dataset label
Dose too low / insufficient exposure	Fasting/post-meal glucose may remain elevated; HbA1c may remain above individualized target.	Persistent hyperglycemia over time increases risk of dehydration, infections, and microvascular/macrovacular complications.	under-treatment / insufficient response
Appropriate effective exposure	Glucose moves toward the individualized goal without unacceptable adverse effects.	Lower hyperglycemic burden; other class-specific benefits may occur (e.g., weight or cardiorenal effects).	adequate response
Dose too high / excessive exposure	Glucose may fall too far for agents that can cause hypoglycemia; for many non-hypoglycemic drugs, other toxicity rather than low glucose is the limiting problem.	Examples: hypoglycemia, GI intolerance, dehydration/volume depletion, edema/fluid retention, infection risk, or other class-specific adverse effects.	overtreatment / adverse-effect limited

For ML use, separate **pharmacologic effect** from **adverse effect**. A drug can produce the desired glucose effect while simultaneously producing an unwanted organ-specific effect. This is exactly why an ADME-style multi-axis score is more informative than a single “drug score.”

8. High-priority safety signals for a dataset

Signal	Classes where it matters	Interpretation
Hypoglycemia	Insulin; sulfonylureas; some combinations	Can cause sweating, tremor, confusion, seizure, loss of consciousness; consequences be especially serious in older adults.
Volume depletion	SGLT2 inhibitors; diuretic combinations	Watch for dizziness, hypotension and dehydration; risk can increase during illness/poor intake.
Genital mycotic infection	SGLT2 inhibitors	Increased glucosuria can predispose to fungal genital infection.
GI intolerance	Metformin; GLP-1 RAs; tirzepatide	Nausea, diarrhea, vomiting, abdominal discomfort can limit dose escalation and can contribute to dehydration.
Edema/heart-failure concern	Pioglitazone	Fluid retention and weight gain are important dose-limiting safety considerations.
DKA risk signal	SGLT2 inhibitors	DKA is uncommon in T2D but can occur, sometimes without very high glucose; risk rises with fasting, dehydration and other triggers.

9. Recommended dataset schema for your ADMET/diabetes application

A useful model should represent the patient state, drug exposure, target organ, and observed outcome separately.

Field	Type	Example	Why it matters
age_group	categorical	65-74	Captures age-related changes in safety, frailty, and treatment burden.
age_years	numeric	68	More granular than age group.
baseline_hba1c	numeric	8.9%	Represents pre-treatment glycemic burden.
baseline_glucose	numeric	210 mg/dL	Useful for short-term response modeling.
bmi	numeric	31.2	Captures obesity/insulin-resistance context and weight-response potential.
egfr	numeric	52 mL/min/1.73m ²	Critical for kidney disease staging and medication exposure/eligibility.
medication_class	categorical	SGLT2 inhibitor	Supports class-level analysis.
medication	categorical	Empagliflozin	Supports drug-specific effects.
dose	numeric	10 mg/day	Core exposure variable.
dose_change	categorical	increase / decrease / unchanged	Allows longitudinal dose-response analysis.
target_organ_system	categorical	Kidney proximal tubule	Links drug action to physiology.
organ_response	text/category	Increased urinary glucose excretion	Captures mechanism-level response.
admet_absorption	score/text	oral; food/formulation dependent	Drug disposition axis.
admet_distribution	score/text	drug-specific	Distribution axis.
admet_metabolism	score/text	hepatic/other	Metabolic clearance axis.
admet_excretion	score/text	renal/other	Elimination and kidney dependence.
toxicity_score	numeric	0-100	Composite safety feature; should be derived transparently rather than guessed.
adverse_event	categorical	nausea / hypoglycemia / edema	Outcome classification.
outcome_hba1c	numeric	7.4%	Primary response target.
weight_change	numeric	-4.2 kg	Important whole-body outcome.

10. Sources and evidence notes

The clinical framing in this report uses current 2026 ADA Standards of Care and FDA labeling where relevant. Because labels can change, the product-specific prescribing information should always be checked before clinical use.

1. American Diabetes Association Professional Practice Committee. "9. Pharmacologic Approaches to Glycemic Treatment: Standards of Care in Diabetes-2026." Diabetes Care 49(Suppl 1):S183-S215.
https://diabetesjournals.org/care/article/49/Supplement_1/S183/163934/9-Pharmacologic-Approaches-to-Glycemic-Treatment
2. American Diabetes Association Professional Practice Committee. "13. Older Adults: Standards of Care in Diabetes-2026." Diabetes Care 49(Suppl 1):S277-S296.
https://diabetesjournals.org/care/article/49/Supplement_1/S277/163921/13-Older-Adults-Standards-of-Care-in-Diabetes-2026
3. American Diabetes Association. "6. Glycemic Goals, Hypoglycemia, and Hyperglycemic Crises: Standards of Care in Diabetes-2026." https://diabetesjournals.org/care/article/49/Supplement_1/S132/163927/6-Glycemic-Goals-Hypoglycemia-and-Hyperglycemic
4. FDA. Ozempic (semaglutide) prescribing information, 2025 label.
https://www.accessdata.fda.gov/drugsatfda_docs/label/2025/209637s025lbl.pdf
5. FDA. Trulicity (dulaglutide) prescribing information, 2025 label.
https://www.accessdata.fda.gov/drugsatfda_docs/label/2025/125469s056lbl.pdf
6. FDA. Mounjaro (tirzepatide) prescribing information, 2025 label.
https://www.accessdata.fda.gov/drugsatfda_docs/label/2025/217806s000lbl.pdf
7. FDA Drug Labeling reference materials for metformin-containing products and pioglitazone. Product labels vary by formulation and indication.
<https://www.accessdata.fda.gov/drugsatfda/>

Clinical safety note: Never use this report to start, stop, increase, decrease, or substitute diabetes medicines for a specific person. Medication changes require individualized clinical assessment, especially when insulin or sulfonylureas are involved, or when kidney disease, heart failure, dehydration, severe illness, or hypoglycemia is present.