


Christian B. Mendl, Irene López Gutiérrez, Keefe Huang

Exercise 2.1 (Bloch sphere and single qubit rotation gates)

Recall from the lecture that an arbitrary single qubit quantum state can be parametrized as

$$|\psi\rangle = e^{i\gamma} \left(\cos \frac{\theta}{2} |0\rangle + e^{i\varphi} \sin \frac{\theta}{2} |1\rangle \right)$$

where θ , φ and γ are real numbers, which can be chosen such that $0 \leq \theta \leq \pi$ and $0 \leq \varphi \leq 2\pi$. The angles θ and φ define the Bloch sphere representation of $|\psi\rangle$, as shown on the right.



https://commons.wikimedia.org/wiki/File:Bloch_sphere.svg

- (a) Find the Bloch angles θ and φ of $|\psi\rangle = \frac{i}{2}|0\rangle - \frac{\sqrt{3}}{2}|1\rangle$, and the corresponding Bloch vector

$$\vec{r} = (\cos(\varphi) \sin(\theta), \sin(\varphi) \sin(\theta), \cos(\theta)).$$

For a real unit vector $\vec{v} \in \mathbb{R}^3$, the rotation by an angle ω about the \vec{v} axis is defined as

$$R_{\vec{v}}(\omega) = \exp(-i\omega \vec{v} \cdot \vec{\sigma}/2) = \cos(\omega/2)I - i \sin(\omega/2)(\vec{v} \cdot \vec{\sigma}),$$

where $\vec{\sigma} = (\sigma_1, \sigma_2, \sigma_3)$ is the Pauli vector. The rotations R_x , R_y , R_z about the standard axes correspond to the special cases $\vec{v} = (1, 0, 0)$, $\vec{v} = (0, 1, 0)$ and $\vec{v} = (0, 0, 1)$, respectively.

- (b) Compute $R_x(\frac{2\pi}{3})|\psi\rangle$ for the state $|\psi\rangle$ defined in (a), and visualize this operation on the Bloch sphere.

Hint: $\cos(\frac{\pi}{3}) = \frac{1}{2}$ and $\sin(\frac{\pi}{3}) = \frac{\sqrt{3}}{2}$.

- (c) The Z-Y decomposition theorem states the following: given any unitary 2×2 matrix U , there exist real numbers $\alpha, \beta, \gamma, \delta \in \mathbb{R}$ such that

$$U = e^{i\alpha} R_z(\beta) R_y(\gamma) R_z(\delta).$$

Find the Z-Y decomposition of the Hadamard gate $H = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix}$.

Hint: There exists a solution with $\beta = 0$.

Solution hints

- (a) Absorb the prefactor i of $|\psi\rangle$ into $e^{i\gamma}$ by setting $\gamma = \frac{\pi}{2}$. The last entry of the Bloch vector \vec{r} is $-\frac{1}{2}$.
- (b) You should obtain $R_x(\frac{2\pi}{3})|\psi\rangle = i|0\rangle$. R_x rotates $|\psi\rangle$ within the y - z -plane to the north pole. (The prefactor i in $i|0\rangle$ does not affect the Bloch vector representation.)
- (c) Set $\gamma = \frac{\pi}{2}$ and $\delta = \pi$, and absorb an overall prefactor in $e^{i\alpha}$.