

Visual Data Analytics Data Reconstruction and Interpolation

Dr. Johannes Kehrer

Disclaimer


These lecture slides and materials are for personal use by students and are intended for educational purposes only. Distribution or reproduction of the material is prohibited.

The views and opinions expressed by any individual during the course of these lectures are their own and do not necessarily represent the views, opinions, or positions of Siemens or the Technical University of Munich.


Although reasonable efforts have been made to ensure the information in these lecture slides is current and accurate, we do not assume any responsibility for the content of the posted material.

These slides also contain links to websites of third parties. As the content of these websites is not under our control, we cannot assume any liability for such external content.


Sampling & Reconstruction



Continuous signal




Sampled signal



Reconstructed signal



Data reconstruction

- From discrete samples to a continuous representation



Overview

- Scattered data interpolation
 - Continuous interpolation functions
 - Piecewise interpolation via triangulation




- Interpolation on grids (next lecture)

Data reconstruction


- Initial data often given at a discrete set of **scattered points (samples)** in the domain

Precipitation stations
in the territory of Moravia and Silesia





Data reconstruction

- We want to derive a **continuous** representation from data given at scattered points
 - Better communication of spatial data distribution




Data reconstruction

- Some analysis techniques require a continuous representation



Data distribution





Isocontours –
curves on which
all points have a
certain value

Data reconstruction



- Isocontours/lines in continuous data fields

Geographic map with isolines




Data reconstruction

- Data reconstruction from scattered points




data interpolation



piece-wise constant
interpolation


Assumes some similarity
between data values
inversely proportional to
distance



continuous interpolation

Data reconstruction


- Data reconstruction from scattered points



- Strategies to obtain values in-between data points...
 1. from a **continuous function** which interpolates the given values and varies smoothly in-between
 2. from a grid which is constructed from the given points, i.e., a **triangulation**

Continuous representation

- Given a set of **scattered points** p_i in a 2D parameter domain with **scalar values** f_i
 - The principles are applicable to arbitrary parameter domain dimensions (1D/2D/3D)
- **Goal:** Construct a **continuous function** f from given set of p_i, f_i which approximates (“follows”) the given values



- Radial Basis Functions


- Independent of dimension of parameter domain (1D/2D/3D)
- Function f represented as **weighted sum** of N radial functions φ

$$f(\mathbf{x}) = \sum_{i=1}^N f_i \varphi(\|\mathbf{p}_i - \mathbf{x}\|)$$


- Each (\mathbf{p}_i, f_i) influences $f(\mathbf{x})$ based on Euclidean distance

$$r = \|\mathbf{p}_i - \mathbf{x}\|$$

- Nearby points have higher influence than far-away points




Continuous representation




- Radial Basis Functions

- Each radial function $\varphi(r)$ is centered around a data point p_i
- Value of $\varphi(r)$ decreases quickly with increasing distance $r = \|p_i - \mathbf{x}\|$ to the function's center p_i




Continuous representation




- Radial Basis Functions

- Each radial function $\varphi(r)$ is centered around a data point p_i
- Value of $\varphi(r)$ decreases quickly with increasing distance $r = \|p_i - \mathbf{x}\|$ to the function's center p_i



Continuous representation




- Radial Basis Functions

- Each radial function $\varphi(r)$ is centered around a data point p_i
- Value of $\varphi(r)$ decreases quickly with increasing distance $r = \|p_i - \mathbf{x}\|$ to the function's center p_i
- Function f represented as **weighted sum** of N radial functions φ


$$f(\mathbf{x}) = \sum_{i=1}^N f_i \varphi(\|p_i - \mathbf{x}\|)$$

where $\|\cdot\|$ is the Euclidean distance (length of a vector)



- Radial Basis Functions
 - Instead of the **black** curve we want the **red** one, i.e., a curve which is **going through the initial data points**
 - This is called an **interpolation**
 - **Question:** How do we have to select the **weights w_i** so that the red curve is obtained?


$$f(x) = \sum_i w_i \varphi(\|p_i - x\|)$$



Continuous representation

- Example:
 - Data points: $p_1 = 1, p_2 = 3, p_3 = 4$
 - Data values: $f_1 = 1, f_2 = 3/5, f_3 = 0$
- Find the weights w_i such that f interpolates all points

$$f(x) = \sum_{i=1}^N w_i \varphi(\|p_i - x\|) \quad \text{where} \quad \varphi(r) = e^{-r^2}$$



| r | 0 | 0.5 | 1 | 1.5 | 2 | 2.5 | 3 |
|--------------|---|-------|-------|--------|---|-----|---|
| $\varphi(r)$ | 1 | $4/5$ | $2/5$ | $1/10$ | 0 | 0 | 0 |

- Radial Basis Functions – finding the weights w_i
 - For $j = 1, \dots, N$,
specify w_i such that $f(p_j)$ interpolates the value f_j

$$f(p_j) = \sum_{i=1}^N w_i \varphi(\|p_i - p_j\|) = f_j$$

Continuous representation

- Radial Basis Functions – finding the weights w_i
 - For $j = 1, \dots, N$,
specify w_i such that $f(p_j)$ interpolates the value f_j
$$f(p_j) = \sum_{i=1}^N w_i \varphi(\|p_i - p_j\|) = f_j$$
 - Yields a system of linear equations (per point) to be solved for w_i

$$\begin{bmatrix} \varphi(\|p_1 - p_1\|) & \varphi(\|p_2 - p_1\|) & \cdots & \varphi(\|p_N - p_1\|) \\ \varphi(\|p_1 - p_2\|) & \varphi(\|p_2 - p_2\|) & \cdots & \varphi(\|p_N - p_2\|) \\ \vdots & \vdots & \ddots & \vdots \\ \varphi(\|p_1 - p_N\|) & \varphi(\|p_2 - p_N\|) & \cdots & \varphi(\|p_N - p_N\|) \end{bmatrix} \begin{bmatrix} w_1 \\ w_2 \\ \vdots \\ w_N \end{bmatrix} = \begin{bmatrix} f_1 \\ f_2 \\ \vdots \\ f_N \end{bmatrix}$$

N equations in N unknowns

Continuous representation


- Radial Basis Functions – finding the weights w_i
 - For $j = 1, \dots, N$,
specify w_i such that $f(p_j)$ interpolates the value f_j
$$f(p_j) = \sum_{i=1}^N w_i \varphi(\|p_i - p_j\|) = f_j$$
 - Yields a system of linear equations (per point) to be solved for w_i

$$\begin{bmatrix} \varphi(0) & \varphi(\|p_2 - p_1\|) & \cdots & \varphi(\|p_N - p_1\|) \\ \varphi(\|p_1 - p_2\|) & \varphi(0) & \cdots & \varphi(\|p_N - p_2\|) \\ \vdots & \vdots & \ddots & \vdots \\ \varphi(\|p_1 - p_N\|) & \varphi(\|p_2 - p_N\|) & \cdots & \varphi(0) \end{bmatrix} \begin{bmatrix} w_1 \\ w_2 \\ \vdots \\ w_N \end{bmatrix} = \begin{bmatrix} f_1 \\ f_2 \\ \vdots \\ f_N \end{bmatrix}$$

Continuous representation

- Example (cont.)

- Data points: $p_1 = 1, p_2 = 3, p_3 = 4$
- Data values: $f_1 = 1, f_2 = 3/5, f_3 = 0$
- Radial function: $\varphi(r) = e^{-r^2}$



$$\Rightarrow \underbrace{\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 2/5 \\ 0 & 2/5 & 1 \end{bmatrix}}_{\mathbf{R}} \underbrace{\begin{bmatrix} w_1 \\ w_2 \\ w_3 \end{bmatrix}}_{\mathbf{w}} = \underbrace{\begin{bmatrix} 1 \\ 3/5 \\ 0 \end{bmatrix}}_{\mathbf{f}}$$

$$\begin{aligned} 1 &= w_1 \\ 3/5 &= w_2 + \frac{2}{5}w_3 \\ 0 &= \frac{2}{5}w_2 + w_3 \end{aligned}$$


If matrix \mathbf{R} is invertible,

$$\mathbf{w} = \mathbf{R}^{-1}\mathbf{f} \text{ with solution: } w_1 = 1, w_2 = 5/7, w_3 = -2/7$$

Continuous representation

- Example

$$f(x) = \varphi(\|1 - x\|) + \frac{5}{7} \varphi(\|3 - x\|) - \frac{2}{7} \varphi(\|4 - x\|)$$





Continuous representation

- 2D Example

- Data points: $p_1 = (1, 1)^T$, $p_2 = (2, 3)^T$, $p_3 = (4, 2)^T$

- Data values: $f_1 = 0.5$, $f_2 = 0.8$, $f_3 = 0.4$

$$f(\mathbf{x}) = 0.49 \varphi(\|(1, 1)^T - \mathbf{x}\|) + 0.79 \varphi(\|(2, 3)^T - \mathbf{x}\|) + 0.39 \varphi(\|(4, 2)^T - \mathbf{x}\|)$$




- Drawbacks of radial basis functions
 - Every sample point has influence on whole domain
 - Adding a new sample requires re-solving the equation system
 - Computationally expensive (solving a system of linear equations)
- What can we do?
 - Find a different radial function
 - Give up finding a smooth reconstruction
 - Try finding a piecewise (local) reconstruction function

- Radial Basis Functions

- Instead of the **black** curve we want the **red** one, i.e., the curve which is going through the initial data points
- This is called an interpolation
- **Question:** How do we have to select the **radial function φ** so that the red curve is obtained?

$$f(x) = \sum_i f_i \varphi(\|p_i - x\|)$$



Continuous representation


- Inverse distance weighting

- Sample positions p_i and values f_i
- Assumption: Nearby points are more similar than those further away \rightarrow they have more influence

$$f(\mathbf{x}) = \sum_i f_i \varphi(\|p_i - \mathbf{x}\|)$$

$$d_i = \|p_i - \mathbf{x}\|, \quad \varphi(r) = \frac{1}{r^2} / \sum_{i=1}^N \frac{1}{d_i^2}$$

Attention for $d_i < \epsilon$




Continuous representation

- Inverse distance weighting
 - Sample positions p_i and values f_i
 - Assumption: Nearby points are more similar than those further away \rightarrow they have more influence

$$f(\mathbf{x}) = \sum_i f_i \varphi(\|p_i - \mathbf{x}\|) =$$


$$= \sum_{i=1}^N \frac{f_i}{\|p_i - \mathbf{x}\|^2} / \sum_{i=1}^N \frac{1}{\|p_i - \mathbf{x}\|^2}$$

$$f(\mathbf{x}) = \frac{\frac{22}{4^2} + \frac{34}{1^2} + \frac{27}{2.5^2} + \frac{30}{3^2} + \frac{33}{2^2}}{\frac{1}{4^2} + \frac{1}{1^2} + \frac{1}{2.5^2} + \frac{1}{3^2} + \frac{1}{2^2}} = 32.38$$



Continuous representation


- Inverse distance weighting
 - Sample positions p_i and values f_i
 - Assumption: Nearby points are more similar than those further away



function is zero at p_i , $i \neq 1$

Continuous representation

- Inverse distance weighting
 - Sample positions p_i and values f_i
 - Assumption: Nearby points are more similar than those further away




function is zero at p_i , $i \neq 2$

Continuous representation

- Inverse distance weighting
 - Sample positions p_i and values f_i
 - Assumption: Nearby points are more similar than those further away

$$f(x) = \frac{\frac{f_1}{\|p_1 - x\|^2} + \frac{f_2}{\|p_2 - x\|^2} + \frac{f_3}{\|p_3 - x\|^2} + \frac{f_4}{\|p_4 - x\|^2} + \frac{f_5}{\|p_5 - x\|^2}}{\frac{1}{\|p_1 - x\|^2} + \frac{1}{\|p_2 - x\|^2} + \frac{1}{\|p_3 - x\|^2} + \frac{1}{\|p_4 - x\|^2} + \frac{1}{\|p_5 - x\|^2}}$$





Continuous representation

- Inverse distance weighting
 - We no longer have to solve a system of linear equations to find the weights
 - However, every sample point still has global influence
- What can we do?
 - Give up smooth reconstruction by constructing a grid from the given points, i.e., a **triangulation**

Generating a triangulation


- Try finding a piecewise (local) reconstruction function
 - Connect the points so that a **triangulation** is obtained
 - Interpolate locally within the triangles

Value obtained by
only considering
values at triangle
corners




Generating a triangulation

- Once a triangulation is given
 - Let the scalar values at vertices be interpolated across the triangles
 - I.e., **piecewise linear** interpolation of values at interior points





Generating a triangulation

- Given **irregularly distributed** positions without connectivity information
- For a set of points **many** triangulations exist
- The challenge is to find the **connectivity** so that a “good” triangulation is generated






Generating a triangulation

- What is a good triangulation?





Generating a triangulation

- What is a good triangulation?
 - A measure for the quality of a triangulation is the **aspect ratio** of the so-defined triangles
 - Avoid long, thin triangles
 - Make triangles as “round” as possible



Generating a triangulation

- An “optimal” triangulation
 - Makes triangles as “round” as possible
 - Maximizes the **minimum angle** in the triangulation
 - Maximizes $\frac{\text{radius of in-circle}}{\text{radius of circumcircle}}$




- A **Delaunay triangulation** is an optimal triangulation

Delaunay triangulation

- Delaunay triangulation


- The **circumcircle** of any triangle does not contain another point of the set

- Maximizes the minimum angle in the triangulation
- Such a triangulation is **unique** (independent of the order of samples) for all but trivial cases




Delaunay triangulation

- How to build a Delaunay triangulation from an initial, non-optimal triangulation?
 - Can be performed by successively improving the initial triangulation via local operations



Delaunay triangulation

- Edge flip operation




- An edge is **local Delaunay** if there exists an empty circumcircle
- If an edge shared by two triangles is illegal, a **flip operation** generates a new edge that is legal!


Delaunay triangulation

Edge flip algorithm:


```
put all edges in stack and mark them
while stack is non-empty do
  pop edge  $uv$  from stack and unmark it
  if  $uv$  is illegal then
    substitute  $pq$  for  $uv$     //edge flip
    for  $ab \in \{up, pv, vq, qu\}$  do
      if  $ab$  is unmarked then
        push  $ab$  on the stack and mark it
      endif
    endfor
  endif
endwhile
```










Edge flip - example



Initial triangulation



Illegal edge → flip




End result

Illegal edge → flip

Edge flip - example

- **Result:** Delaunay triangulation




- All edges are local Delaunay
- The circumcircle of any triangle does not contain another point of the set

<http://multivis.net/lecture/delaunay.html>


Delaunay triangulation

- Local vs. global optimality
 - Edge is locally Delaunay ... but not globally




Delaunay triangulation

- Local vs. global optimality
 - If a triangulation is locally Delaunay everywhere
→ globally Delaunay




Voronoi diagram

- **Problem:** Looking for nearest neighbor




Voronoi diagram

- Partitions domain into **Voronoi regions**
 - Each Voronoi region contains one initial sample – the **Voronoi samples**




Voronoi diagram

- Partitions domain into **Voronoi regions**
 - Each Voronoi region contains one initial sample – the **Voronoi samples**
 - Points in Voronoi region are closer to respective sample than to any other sample




Voronoi diagram

- Centers of circumcircles of Delauney triangulation




Voronoi diagram

- The **geometric dual** (topologically equal) of Delaunay triangulation
 - Voronoi samples are vertices in Delaunay triangulation



Voronoi diagram


- Construction
 - Points in a Voronoi region are closer to the respective sample than to any other sample




Voronoi diagram

- Construction

- Points in a Voronoi region are closer to the respective sample than to any other sample



Contact information



Dr. Johannes Kehrer

Siemens Technology
T RDA BAM IBI-DE
Otto-Hahn-Ring 6
81739 München, Deutschland

E-mail:
kehrer.johannes@siemens.com

Internet
siemens.com/innovation

Intranet
intranet.ct.siemens.com