

The Geometry of Quantization

Yusuf Berk AKÇAY

Chapter 1

The Harmonic Oscillator

1.1 Classical Setting

Classically, the harmonic oscillator is described by a 2nd order ODE, called Newton's Second Law

$$m\ddot{x} = -kx$$

in the 'configuration space' $\mathbb{R} = \{x \mid x \in \mathbb{R}\}$. This 2nd order ODE can be turned into a system of 1st order ODEs as follows: First, define the canonical variables as

$$q := x \quad p := m\dot{x}$$

Then, we obtain

$$\dot{q} = p/m \quad \dot{p} = -kx$$

These equations define a flow in the "phase space"

$$\mathbb{R}^2 = \{(q, p) \mid q, p \in \mathbb{R}\}$$

If we define the Hamiltonian (energy) function $H : \mathbb{R}^2 \rightarrow \mathbb{R}$ as

$$H(q, p) = \frac{p^2}{2m} + \frac{kq^2}{2}$$

we get

$$\dot{q} = \frac{\partial H}{\partial p} \quad \dot{p} = -\frac{\partial H}{\partial q}$$

the so-called Hamilton's equations.

- L is an n -dimensional submanifold of $H^{-1}(E)$.
- The pullback to L of the form

$$\alpha_n = \sum_{j=1}^n p_j dq_j$$

is exact, i.e., we have

$$\alpha_n|_L = \sum_j \frac{\partial S}{\partial q_j} dq_j = dS$$

- The canonical projection $\pi : T^*\mathbb{R}^n \rightarrow \mathbb{R}^n$ induces a diffeomorphism $L \simeq \mathbb{R}^n$. Thus, L is projectable.

Of course, these assertions hold when S is globally defined. If it is not, we have the weakened version:

- L is an n -dimensional submanifold of $H^{-1}(E)$.
- The pullback to L of the form

$$\alpha_n = \sum_{j=1}^n p_j dq_j$$

is closed, i.e., we have

$$d\alpha_n|_L := \omega_n|_L = 0$$

In the latter case, we say L is a Lagrangian submanifold of $\mathbb{R}^{2n}$. If L satisfies the former strong assertions, we say L is an exact, projectable Lagrangian submanifold. Example: Consider the harmonic oscillator with $L = H^{-1}(E)$, i.e.,

an ellipse.

- The pullback to L of $\alpha = pdq$ is closed since $\dim L = 1$, but not exact since

$$\int_L pdq = \int dp \wedge dq = \text{Area of ellipse} \neq 0$$

- L fails to project diffeomorphically to $\mathbb{R}$ since the former is compact while the latter is not.

1.5 Hamilton-Jacobi Theorem

Consider the Hamiltonian vector field

$$X_H = \sum_j \left(\frac{\partial H}{\partial p_j} \frac{\partial}{\partial q_j} - \frac{\partial H}{\partial q_j} \frac{\partial}{\partial p_j} \right)$$

satisfying

$$X_H \lrcorner d\alpha_n = -dH \quad \text{i.e.,} \quad -d\alpha_n(X_H, -) = dH(-)$$

If the Lagrangian submanifold L lies in a level set of H , then TL lies in the kernel of dH since H is constant on L . Thus, the two-form $d\alpha_n$ vanishes on the subspace

$$\langle T_p L, X_H(p) \rangle \leq T_p \mathbb{R}^{2n}$$

for each $p \in L$. Since $d\alpha_n$ defines a nondegenerate, skew-symmetric bilinear form, by linear algebra, a subspace on which $d\alpha_n$ vanishes can be at most n -dimensional. Thus, we have

$$X_H(p) \in T_p L$$

In other words, X_H is tangent to L , or the restriction of X_H defines a vector field on L . Conversely, assume X_H is tangent to the Lagrangian submanifold L . Then, we have

$$dH|_L(T_p L) = -d\alpha_n|_L(X_H(p), T_p L) = 0 \Rightarrow dH|_L = 0$$

Thus, H is locally constant on L .

A function $H : \mathbb{R}^{2n} \rightarrow \mathbb{R}$ is locally constant on a Lagrangian submanifold L of $\mathbb{R}^{2n}$ iff X_H is tangent to L .

1.6 The WKB Approximation

Recall that if $S : \mathbb{R}^n \rightarrow \mathbb{R}$ is an admissible function, then $\varphi = e^{iS/\hbar}$ satisfies

$$(\hat{H} - E)\varphi = O(\hbar)$$

To obtain a better approximation, consider

$$\varphi(x) = a(x)e^{iS(x)/\hbar}$$

This gives

$$(\hat{H} - E)\varphi = -\frac{1}{2m} \left[i\hbar \left(a\Delta S + 2 \sum_j \frac{\partial a}{\partial x_j} \frac{\partial S}{\partial x_j} \right) + \hbar^2 \Delta a \right] e^{iS/\hbar}$$

Thus, as a second approximation we have

$$a\Delta S + 2 \sum_j \frac{\partial a}{\partial x_j} \frac{\partial S}{\partial x_j} = 0 \rightarrow \text{Transport Equation}$$

Thus, we obtain a solution satisfying

$$(\hat{H} - E)\varphi = O(\hbar^2)$$

Example: In 1-d, we have

$$aS'' + 2a'S' = 0 \Rightarrow a = \frac{c}{\sqrt{S'}} = \frac{c}{[2m(E - V)]^{1/4}}$$

Since a is independent of S , we can use this expression for non-projectable Lagrangian submanifolds as well. However, a is not well-defined. For instance, if $V(x) = x^2$ we have

$$a = \begin{cases} |x| < \sqrt{E} \rightarrow \text{Well-defined} \\ |x| = \sqrt{E} \rightarrow \text{Singularity} \\ |x| > \sqrt{E} \rightarrow \mathbb{C} \end{cases}$$

Remarks:

- Notice that the semiclassical states of the H.O. are not singular.
- Our construction does not depend on S , thus it makes sense for arbitrary Lagrangian submanifolds.

As a further step, we identify the quantum states with the Hilbert space

$$\mathcal{H}_{\mathbb{R}^n} = \left\{ \sigma \text{ half-density on } \mathbb{R}^n \mid \langle \sigma, \tau \rangle = \int_{\mathbb{R}^n} \sigma \bar{\tau} \right\}$$

and define the Hamiltonian operator as

$$\hat{H}(a\sqrt{|dx|}) = (\hat{H}a)\sqrt{|dx|}$$

Then, the second-order approximate solution to the eigenvalue problem $\hat{H}I_{\hbar} = EI_{\hbar}$ is given by

$$I_{\hbar} = [(dS)^*a] e^{iS/\hbar} \in \mathcal{H}_{\mathbb{R}^n}$$

Chapter 2

Symplectic Geometry

2.1 Symplectic Vector Spaces

Let V be an m -dimensional vector space over $\mathbb{R}$ endowed with a bilinear form

$$\omega : V \times V \rightarrow \mathbb{R}$$

This gives a map $\tilde{\omega}$ defined by contraction:

$$\tilde{\omega} : V \rightarrow V^* \quad x \mapsto \omega(x, -)$$

The ω -orthogonal of a subspace $W \leq V$ is defined by

$$W^\perp = \{x \in V \mid \omega(x, W) = 0\}$$

The bilinear form ω is called a symplectic form if

- ω is nondegenerate $\iff \tilde{\omega}$ is an isomorphism $\iff V^\perp = 0$
- $\omega(x, y) = -\omega(y, x)$

(Notice that these two properties imply $m = 2n$) In that case, we say (V, ω) is a symplectic vector space. We can also define the symmetries of a symplectic vector space, i.e., symplectomorphisms, as follows:

$$Sp(V) = \{A \in GL(V) \mid \omega(Ax, Ay) = \omega(x, y)\}$$

Canonical Example: Let E be a real n -dimensional vector space and $V = E \oplus E^*$. Define $\omega : V \times V \rightarrow \mathbb{R}$ as

$$\omega((x, \lambda), (x', \lambda')) = \lambda'(x) - \lambda(x')$$

2.2 Lagrangian Subspaces

The ω -orthogonal of a subspace W of a symplectic vector space (V, ω) is called the symplectic orthogonal of W . We have

- $W^{\perp\perp} = W$
- $\dim W^\perp = \dim V - \dim W$
- $A \leq B \Rightarrow B^\perp \leq A^\perp$
- $(A + B)^\perp = A^\perp \cap B^\perp$
- $(A \cap B)^\perp = A^\perp + B^\perp$

A subspace $L \leq V$ is called a Lagrangian subspace provided

- $\omega|_L = 0$
- $\dim L = \frac{1}{2} \dim V$

Let me sketch some results without proofs:

Theorem 2.2.1. *Every symplectic vector space has a Lagrangian subspace.*

Theorem 2.2.2. *Every $2n$ -dimensional symplectic vector space is symplectomorphic to $(\mathbb{R}^{2n}, \omega_n)$ where*

$$\omega_n = \begin{pmatrix} 0 & I_n \\ -I_n & 0 \end{pmatrix}$$

2.3 The Lagrangian Grassmannian

Let V be a $2n$ -dimensional symplectic vector space. The Lagrangian Grassmannian $\mathcal{L}(V)$ is defined to be

$$\mathcal{L}(V) = \{\text{Lagrangian subspaces of } V\}$$

This set has a natural manifold structure, and in fact, we have

$$\mathcal{L}(V) \simeq U(n)/O(n)$$

In particular, the fundamental group of $\mathcal{L}(V)$ (i.e., the group of closed loops (up to homotopy) in $\mathcal{L}(V)$) is isomorphic to that of S^1 :

$$\pi_1(\mathcal{L}(V)) \simeq \pi_1(S^1) \simeq \mathbb{Z}$$

Thus, we can define the 'winding number' of a closed loop in $\mathcal{L}(V)$, giving the cohomology element

$$\mu_V \in H^1(\mathcal{L}(V), \mathbb{Z})$$

Example: Consider $\mathbb{R}^2$ and the closed loop

$$L(t) = \langle \cos t, \sin t \rangle \in \mathcal{L}(\mathbb{R}^2) \quad t \in [0, \pi]$$

This loop performs one rotation, so the corresponding winding number is 1.

2.4 Symplectic Manifolds

To motivate the definition of a symplectic manifold, let's recall the $\mathbb{R}^{2n}$ case. First, define the differential forms

$$\alpha_n = \sum_{i=1}^n p_i dq_i \quad \omega_n = -d\alpha_n = \sum_{i=1}^n dq_i \wedge dp_i$$

The 2-form ω_n defines a symplectic vector space structure on each tangent space of $\mathbb{R}^{2n}$. The dynamics of the system are determined by the Hamiltonian vector field

$$X_H = \sum_{i=1}^n \left(\frac{\partial H}{\partial p_i} \frac{\partial}{\partial q_i} - \frac{\partial H}{\partial q_i} \frac{\partial}{\partial p_i} \right)$$

This vector field is uniquely determined by the relation

$$i_{X_H} \omega_n = dH \Rightarrow X_H = \tilde{\omega}_n^{-1}(dH)$$

where we used the non-degeneracy of ω_n . Thus, the symplectic form ω_n completely determines Hamilton's equations. This description leads us to the following:

- Energy Conservation

$$\mathcal{L}_{X_H} H = X_H H = dH(X_H) = \omega_n(X_H, X_H) = 0$$

where we used the 2-form (antisymmetric) nature of ω_n .

- Preservation of Symplectic Area

$$\mathcal{L}_{X_H} \omega_n = d(i_{X_H} \omega_n) + i_{X_H} d\omega_n = 0$$

where we used the closedness of ω_n .

So, let's make the following definition: A symplectic structure on a manifold M is a closed, nondegenerate 2-form ω on M . In this case, we say (M, ω) is a symplectic manifold.

Example: $(\mathbb{R}^{2n}, \omega_n = \sum_{i=1}^n dq_i \wedge dp_i)$

2.5 Maslov Index

A symplectic vector bundle is a vector bundle $E \rightarrow B$ where each fiber is a symplectic vector space. The Lagrangian subbundle is defined similarly. Now, let's define the famous Maslov index: Suppose we are given the manifold $\mathbb{R}^{2n} \simeq T^*\mathbb{R}^n$ and a Lagrangian submanifold L . Let $\gamma(t)$ be a closed loop in L . By means of the Gauss map

$$G : L \rightarrow \mathcal{L}(\mathbb{R}^{2n}) \quad p \mapsto T_p L$$

we can get a closed loop in $\mathcal{L}(\mathbb{R}^{2n})$. Now, define the Maslov index $\mu_L(\gamma(t))$ of the closed loop $\gamma(t)$ to be the winding number of this closed loop in $\mathcal{L}(\mathbb{R}^{2n})$. Example: In $\mathbb{R}^2$ consider $L = S^1$ and

$$\gamma(t) = (\cos t, \sin t) \quad t \in [0, 2\pi]$$

When t moves from 0 to 2π , the Lagrangian subspace $T_{\gamma(t)}L$ performs two rotations, so

$$\mu_L(\gamma(t)) = 2$$

2.6 Cotangent Bundles

The most important examples of symplectic manifolds are cotangent bundles. Let M be a smooth manifold. Recall that T_p^*M denotes the vector space of covectors at $p \in M$. Now, the cotangent bundle T^*M is defined to be

$$T^*M = \{(x, b) \mid x \in M, b \in T_x^*M\}$$

We have the natural projection map

$$\pi : T^*M \rightarrow M \quad \pi(x, b) = x$$

Locally, T^*M can be parametrized by $2n$ coordinates $(q_1, \dots, q_n; p_1, \dots, p_n)$ where the q 's stand for coordinates of M . Now, define the 1-form α_M on T^*M , called the tautological 1-form as follows:

$$\alpha_M|_{(x,b)}(v) = b(\pi_* v)$$

Locally, this form can be expressed as

$$\alpha_M = \sum_{i=1}^n p_i dq_i$$

The exterior derivative of the tautological 1-form gives us a symplectic form

$$-d\alpha_M = \omega_M = \sum_{i=1}^n dq_i \wedge dp_i$$

Chapter 3

Quantization of Cotangent Bundles

3.1 Mechanics on Manifolds

Let's extend our discussion of classical/quantum mechanics to manifolds. Let M be a configuration space of a physical system and T^*M be the corresponding phase space. The Hamiltonian $H : M \rightarrow \mathbb{R}$ is defined to be $H(q, p) = T(q, p) + V(q)$ where

$$T(q, p) = \frac{1}{2} \sum_{ij} g^{ij}(q, p) p_i p_j$$

is the kinetic energy and

$$V \rightarrow \mathbb{R}$$

is the potential energy. The case $V = 0$ is especially important in geometry, which gives us the geodesics of M . In quantum mechanics, one defines the Hamiltonian operator $\hat{H} : C^\infty(M) \rightarrow C^\infty(M)$ as

$$\hat{H} = -\frac{\hbar^2}{2m} \Delta + V$$

where Δ is the Laplace-Beltrami operator on M . One can also define the corresponding Hamiltonian operator $\hat{H} : \mathcal{H}_M \rightarrow \mathcal{H}_M$ as usual. Now, the Schrödinger equation

$$\hat{H}\psi = E\psi$$

has the second-order WKB approximation

$$\psi = e^{iS/\hbar} a$$

where

$$H \circ \iota_{dS} = E \quad a\Delta S + 2\mathcal{L}_{\nabla S}a = 0$$

Now, let's talk about the geometry of this WKB solution. If $L \subseteq H^{-1}(E)$ is an exact Lagrangian submanifold, i.e.,

- $\pi_L := \pi|_L$ is a diffeomorphism
- $\alpha_M|_L$ is an exact 1-form on L

Let $\phi \rightarrow \mathbb{R}$ be a primitive of $\alpha_M|_L$ and a be a half-form on L invariant under the action of $X_H|_L$ (whose existence is guaranteed by the Hamilton-Jacobi theorem), then a semiclassical state on M is given by

$$(\pi_L^{-1})^* e^{i\phi/\hbar} a$$

Thus, a semiclassical state is determined by:

- A Lagrangian submanifold $L \subseteq T^*M$ satisfying $L \subseteq H^{-1}(E)$
- A half-form a satisfying

$$\mathcal{L}_{X_H|_L} a = 0$$

The question we must answer is the following: What will we do if L is not exact (no ϕ) and not projectable (no π_L^{-1})?

3.2 Handling Non-Exactness

Assume L is not an exact Lagrangian, i.e., the 1-form $\alpha_M|_L$ is not exact. It is still closed, however, and thus locally exact:

$$L = \bigcup_j L_j \quad \exists \phi_j : L_j \rightarrow \mathbb{R} \text{ such that } d\phi_j = \alpha_M|_{L_j}$$

Then, we can quantize each L_j as

$$(\pi_{L_j}^{-1})^* e^{i\phi_j/\hbar} a$$

To quantize L , we must combine all of these. To get consistent results at the intersections of L_j 's, we must have

$$\phi_j - \phi_k \in 2\pi\hbar\mathbb{Z} \quad \text{on } L_j \cap L_k$$

This condition is equivalent to the so-called **quantization condition**

$$\lambda_L = 2\pi\hbar \times \text{Integral Class}$$

i.e., for all closed loops in L

$$\int \alpha_M \in 2\pi\hbar\mathbb{Z}$$

Example: For the Hamiltonian $H = (p^2 + \omega^2 q^2)/2$, this condition gives

$$E = n\hbar\omega$$

3.3 Handling Non-Projectability

Assume L is non-projectable. The strategy in this case is as follows: Decompose L as $L = \sqcup_j L_j$ where each L_j is projectable, and sum the corresponding contributions:

$$\sum_j (\pi_{L_j}^{-1})^* e^{i\phi_j/\hbar} a$$

Example: In $T^*\mathbb{R} = \{(q, p)\}$ consider the Hamiltonian $H = (p^2 - q)/2$ and $E = 0$. The Lagrangian submanifold is given as

$$L = H^{-1}(0) = \{(x^2, x) \mid x \in \mathbb{R}\} \simeq \mathbb{R}$$

This Lagrangian is clearly non-projectable, but it consists of projectable components

$$L_{\pm} = \{(x^2, x) \mid x \in \mathbb{R}^{\pm}\}$$

Let me first determine what $a(x)$ should be. By direct computation, $X_H|_L \sim \partial_x$, so we have $a(x) = A \in \mathbb{R}$. Now, $\alpha_M|_L$ can be evaluated as

$$\alpha_M|_L = 2x^2 dx \Rightarrow \phi = \frac{2x^3}{3}$$

Summing the contributions from $L_{\pm}$ gives

$$\psi(x) \sim x^{1/4} \cos(2x^{3/2}/3\hbar)$$

3.4 The Maslov Correction

We start by considering a Lagrangian submanifold $L \subseteq T^*\mathbb{R} \simeq \mathbb{R}^2$. Consider the projection of L to p -axis:

$$\pi_p : \mathbb{R}^2 \rightarrow p\text{-axis}$$

If the map π_p is a diffeomorphism, then L is said to be p -projectable.

For a p -projectable Lagrangian, we define the function $\tau : L \rightarrow \mathbb{R}$ such that its differential yields the symplectic potential in the p -representation:

$$\tau : L \rightarrow \mathbb{R} \quad \text{with} \quad d\tau = -qdp|_L$$

Example: Let's consider a simple example where the position is constant. Let

$$L = \{(q_0, x) \mid x \in \mathbb{R}\}$$

In this case, the position is fixed at $q = q_0$ while momentum $p = x$ varies. For a constant half-density a defined on L , we physically expect the wave function in the position representation to be highly localized around q_0 , yielding a Dirac delta function $\delta(q - q_0)$.

Now, by direct computation we have

$$\tau = -q_0x$$

Pulling this back via the inverse projection, we get

$$(\pi_p^{-1})^* e^{i\tau/\hbar} a = \underbrace{e^{-iq_0p/\hbar} a |dp|^{1/2}}_{\text{Fourier transform of } \delta(q - q_0)}$$

This precisely matches the expected momentum representation of a delta function localized at q_0 .

Now, let's generalize this. For any p -projectable $L \subseteq T^*\mathbb{R}$ with $d\tau = -qdp|_L$, we define

$$B|dp|^{1/2} = (\pi_p^{-1})^* e^{i\tau/\hbar} a$$

The quantum state associated with this Lagrangian submanifold is then given by the inverse semiclassical Fourier transform of B :

$$J_\hbar(L, \iota, \tau, a) := \mathcal{F}_\hbar^{-1}(B) |dq|^{1/2}$$

Let's compare this p -projectable approach with our earlier method that relied on q -projectability.

Example: Consider $L = \{(x, kx) \mid x \in \mathbb{R}\}$. Assume a is a constant half-density on L .

Using the q -representation:

$$pdq \rightarrow \frac{kx^2}{2}$$

$$(\pi_L^{-1})^* a = A|dq|^{1/2}$$

This leads to the state:

$$I_{\hbar} = e^{ikq^2/2\hbar} A|dq|^{1/2}$$

Using the p -representation:

$$-qdp \rightarrow -\frac{kx^2}{2}$$

$$(\pi_p^{-1})^* a = |k|^{-1/2} A|dp|^{1/2}$$

We evaluate the generating function in the p -coordinates:

$$\pi_p^* \left(-\frac{kx^2}{2} \right) = -\frac{p^2}{2k}$$

Applying the semiclassical Fourier transform, we find the relation between the two states:

$$J_{\hbar} = e^{-i\pi \operatorname{sgn}(k)/4} I_{\hbar}$$

Notice the emergence of the phase factor $e^{-i\pi \operatorname{sgn}(k)/4}$, which is intimately related to the Maslov index.

3.5 Arbitrary Biprojectable Lagrangian Embeddings

Now, consider an arbitrary Lagrangian submanifold $L \subseteq T^*\mathbb{R}$ that is biprojectable, meaning it can be projected onto both the q and p axes locally.

We relate the corresponding functions for the two projections. Let ϕ be the generating function for the q -projection and τ for the p -projection:

$$\begin{aligned}\phi &= \tau + \iota^*(qp) \\ S &= \phi \circ \pi_L^{-1} \quad T = \tau \circ \pi_p^{-1}\end{aligned}$$

From this Legendre transformation structure, we can express $S(q)$ in terms of $T(p)$:

$$S(q) = -p(q)T'(p(q)) + T(p(q))$$

where the coordinate transformations are governed by:

$$p(q) = S'(q) \longleftrightarrow q = -T'(p)$$

Differentiating the relation $q = -T'(p(q))$ with respect to q , we obtain the relation between the second derivatives:

$$\begin{aligned}1 &= -T''(p(q)) \cdot p'(q) = -T''(p(q)) \cdot S''(q) \\ T''(p(q)) &= -(S''(q))^{-1}\end{aligned}$$

We have two pullbacks for the half-density a :

$$\begin{aligned}(\pi_L^{-1})^* a &= A|dq|^{1/2} \\ (\pi_p^{-1})^* a &= B|dp|^{1/2}\end{aligned}$$

The amplitudes are related by the Jacobian of the transformation between p and q :

$$A(q) = \left| \frac{dp}{dq} \right|^{1/2} B(p(q)) = |S''(q)|^{1/2} B(p(q))$$

The state $J_\hbar$ is given by the integral:

$$J_\hbar = (2\pi\hbar)^{-1/2} \int_{\mathbb{R}} e^{i(pq+T(p))/\hbar} B(p) dp |dq|^{1/2}$$

We evaluate this integral using the stationary phase approximation. The phase is $\Phi(p) = pq + T(p)$, and the stationary points occur where $\Phi'(p) = q + T'(p) = 0$, which is consistent with our coordinate relation $q = -T'(p)$. The second derivative of the phase at the stationary point is $T''(p(q))$.

Applying the stationary phase formula yields:

$$J_\hbar = e^{-i\pi \operatorname{sgn}(T''(p(q)))/4} e^{iS(q)/\hbar} |T''(p(q))|^{-1/2} B(p(q)) + O(\hbar)$$

Recognizing the leading term as our previously defined state $I_{\hbar}$, we establish the connection:

$$J_{\hbar} = e^{-i\pi \operatorname{sgn}(T''(p(q)))/4} I_{\hbar} + O(\hbar)$$

Example: Consider $L = \{(x^2, x)\}$ as before.

First, we find

$$d\tau = \iota^*(-qdp) = -(x^2)dx \implies \tau(x) = -\frac{x^3}{3}$$

$$d\phi = \iota^*(pdq) = x(2xdx) = 2x^2dx \implies \phi(x) = \frac{2x^3}{3}$$

The function $T(p)$ is straightforward since $p = x$:

$$T(p) = -\frac{p^3}{3}$$

The relation between q and p is $q = -T'(p) = p^2$. Notice that for $q > 0$, there are two branches of momentum: $p = \pm q^{1/2}$.

The second derivative is:

$$T''(p(q)) = -2p = -2\operatorname{sgn}(p)q^{1/2}$$

Because there are two stationary points (for $p > 0$ and $p < 0$) contributing to a given $q > 0$, the integral expression for $J_{\hbar}$ becomes a sum of two terms:

$$J_{\hbar} = \left(\underbrace{e^{-i\pi/4} e^{2iq^{3/2}/3\hbar} B(-q^{1/2})}_{p<0} + \underbrace{e^{i\pi/4} e^{-2iq^{3/2}/3\hbar} B(q^{1/2})}_{p>0} \right) q^{-1/4} |dq|^{1/2}$$

It is crucial to note that while the semiclassical expression diverges at the caustic point $q = 0$ (due to the $q^{-1/4}$ factor), the exact state $J_{\hbar}$, defined as an integral expression, remains smooth around $q = 0$.

$J_{\hbar} \rightarrow$ As an integral expression, smooth around $q = 0$

With this new method of Maslov, our quantization condition becomes

$$\int_{\gamma} \alpha_M = 2\pi\hbar(n + \mu_L(\gamma)/4)$$