

The Dirichlet Pólya Problem for Spherical Shells in General Dimension: Exact Dimension Lifting and the Shifted-Tail Bottleneck

Working manuscript

18 July 2026

Abstract

For a spherical shell in dimension $d \geq 3$, the separated radial problem has Bessel orders $\ell + (d-2)/2$ and spherical-harmonic multiplicities $\kappa_{d,\ell}$. We give an exact branching decomposition of the strict shell-action proxy into positive integer- and half-integer shifted planar tails. We also prove a signed primitive comparison and retain its full positive remainder. This yields the exact identity

$$W_d - P_d^< = \mathcal{B}_d(A) + \sum_{m \geq 0} \binom{m+d-3}{d-3} D_A \left(m + \frac{d-2}{2} \right).$$

Consequently, one dimension-free positive shifted-tail inequality implies the sharp Dirichlet Pólya inequality for every spherical shell and every $d \geq 3$. The dimension lift is proved here; the inner shifted-tail inequality remains the central open lemma. We prove the outer and terminal tail regions, close the first interface-cell milestone by exact rational wall estimates, and prove all shifted tails above an explicit fixed-ratio high-energy threshold. A wall-safe first-shelf reduction compresses every remaining many-cell tail to one scalar coupled to its terminal one-interface reserve. In the thin sector it also improves the uniform threshold to $K \geq 125/[8(1-\rho)^2]$. A scale-free shelf-curvature bound and a fractional-top inverse-action estimate give an explicit exact-angle lower bound for the terminal reserve, including the previously unused zero-level triangle. We also prove an automatic ultra-thin high-floor sector and exclude every noncompact escape in both the first-floor and high-floor first-drop branches. A directed-rounding certificate proves all 19602 finite first-floor chambers with $r \geq 3/2$, $s \in \{0,1\}$, $x < 100$. A second certificate handles all 398 finite small-start chambers, while an analytic comparison closes their infinite rays. Together with the continuous large-ray certificate, this proves the entire first-floor sector $s \in \{0,1\}$. In the no-drop branch we prove every $n = 1$ case and reduce the remaining data to finitely many discrete chambers, with the explicit bound $n < 380$ in the two low-floor rows. These results do not prove every remaining compact chamber, and no assertion in this manuscript treats the full residual terminal-shelf scalar as proved.

1 Statement and status

For $0 < \rho < 1$ let

$$A_{\rho,1}^{(d)} = \{x \in \mathbb{R}^d : \rho < |x| < 1\}, \quad d \geq 3,$$

and use the strict counting convention

$$N_D(A_{\rho,1}^{(d)}, K^2) := \#\{j : \lambda_j^D(A_{\rho,1}^{(d)}) < K^2\}.$$

Write

$$\omega_d = \frac{\pi^{d/2}}{\Gamma(d/2 + 1)}, \quad L_d = \frac{\omega_d}{(2\pi)^d}.$$

The desired estimate is

$$N_D(A_{\rho,1}^{(d)}, K^2) \leq L_d |A_{\rho,1}^{(d)}| K^d = \frac{1 - \rho^d}{2^d \Gamma(d/2 + 1)^2} K^d. \quad (1)$$

For $a > 0$, define the zero-extended ball action

$$G_a(z) = \begin{cases} \frac{1}{\pi} \left(\sqrt{a^2 - z^2} - z \arccos \frac{z}{a} \right), & 0 \leq z \leq a, \\ 0, & z > a, \end{cases} \quad (2)$$

put $\mu = \rho K$, and set

$$A(z) = A_{\rho,K}(z) := G_K(z) - G_\mu(z). \quad (3)$$

For $x > 0$, let

$$[x]_< := \#\{n \in \mathbb{N} : n < x\} = \max\{0, [x] - 1\}.$$

In particular, $[n]_< = n - 1$ at a positive integral wall.

The only unproved statement on the primary path is the following.

Conjecture 1.1 (Positive shifted shell tails). *For $0 < \rho < 1$, $K > 0$, and*

$$r \in \tfrac{1}{2}\mathbb{N},$$

one has

$$T_A(r) := [A(r) + \tfrac{1}{4}]_< + 2 \sum_{j \geq 1} [A(r + j) + \tfrac{1}{4}]_< \leq 2 \int_r^K A(z) \, dz. \quad (4)$$

The zero extension makes the sum finite and makes the assertion trivial when $r \geq K$.

Sections 2–5 prove that Conjecture 1.1 implies (1) in every dimension $d \geq 3$. Section 6 records the proved tail regions, the completed first interface-cell theorem, fixed-ratio and thin high-energy theorems, the terminal reserve, and the current finite reductions. It also states explicitly which finite wall problems remain.

2 General-dimensional spectral proxy

Put

$$a_d = \frac{d-2}{2}, \quad \nu_{\ell,d} = \ell + a_d.$$

The degree- ℓ spherical harmonics on $\mathbb{S}^{d-1}$ have multiplicity

$$\kappa_{d,\ell} = \binom{\ell + d - 1}{d - 1} - \binom{\ell + d - 3}{d - 1} = \frac{(2\ell + d - 2)(\ell + d - 3)!}{\ell!(d - 2)!}. \quad (5)$$

Here and below a binomial coefficient with lower index larger than its nonnegative upper index is zero.

Under the unitary radial substitution

$$u(r) = r^{(d-1)/2} R(r),$$

the degree- ℓ block becomes

$$L_{\ell,d} = -\frac{d^2}{dr^2} + \frac{\nu_{\ell,d}^2 - \frac{1}{4}}{r^2} \quad \text{on } (\rho, 1), \quad u(\rho) = u(1) = 0. \quad (6)$$

Its positive eigenfrequencies are the zeros of

$$D_{\nu,\rho}(k) = J_\nu(\rho k)Y_\nu(k) - Y_\nu(\rho k)J_\nu(k), \quad \nu = \nu_{\ell,d}.$$

With the standard continuous Bessel phase, let

$$\gamma_{\rho,K}(\nu) = \frac{\theta_\nu(K) - \theta_\nu(\rho K)}{\pi}.$$

The determinant factorization and phase monotonicity give the exact strict count

$$N_D(A_{\rho,1}^{(d)}, K^2) = \sum_{\ell \geq 0} \kappa_{d,\ell} [\gamma_{\rho,K}(\nu_{\ell,d})]_{<}. \quad (7)$$

The real-order phase-difference enclosure of [2, 3] gives, for $0 \leq \nu \leq K$,

$$\gamma_{\rho,K}(\nu) < A_{\rho,K}(\nu) + \frac{1}{4}. \quad (8)$$

For $\nu \geq K$, both the exact count and the action proxy vanish. Since $x < y$ implies $[x]_{<} \leq [y]_{<}$, including when y is integral, (7)–(8) imply the wall-safe proxy bound

$$N_D(A_{\rho,1}^{(d)}, K^2) \leq P_d^<(\rho, K), \quad P_d^< := \sum_{\ell \geq 0} \kappa_{d,\ell} q_\ell, \quad (9)$$

where

$$q_\ell = [A(\ell + a_d) + \frac{1}{4}]_{<}. \quad (10)$$

The argument above is dimension-independent after the substitutions $\nu = \ell + a_d$ and (5). In particular, the published phase enclosure is already a real-order result: no interpolation is introduced when d changes parity.

3 Exact harmonic branching

Let $p = d - 2$, so $a_d = p/2$, and define

$$r_m = m + \frac{p}{2}, \quad c_{d,m} = \binom{m+p-1}{p-1} = \binom{m+d-3}{d-3}. \quad (11)$$

Proposition 3.1 (Exact shifted-tail decomposition). *For every finitely supported sequence $(q_\ell)_{\ell \geq 0}$,*

$$\sum_{\ell \geq 0} \kappa_{d,\ell} q_\ell = \sum_{m \geq 0} c_{d,m} \left(q_m + 2 \sum_{j \geq 1} q_{m+j} \right). \quad (12)$$

Consequently, for (10),

$$\boxed{P_d^< = \sum_{m \geq 0} c_{d,m} T_A(r_m)}. \quad (13)$$

This identity is exact at every strict action wall.

Proof. The generating functions are

$$\sum_{\ell \geq 0} \kappa_{d,\ell} s^\ell = \frac{1+s}{(1-s)^{d-1}}, \quad \sum_{m \geq 0} c_{d,m} s^m = \frac{1}{(1-s)^{d-2}},$$

and

$$1 + 2 \sum_{j \geq 1} s^j = \frac{1+s}{1-s}.$$

Coefficient comparison gives

$$\kappa_{d,\ell} = c_{d,\ell} + 2 \sum_{m=0}^{\ell-1} c_{d,m}.$$

Substitution and finite reindexing prove (12). For (10), one has $q_{m+j} = [A(r_m + j) + 1/4]_<$, which proves (13). No floor has been approximated or moved. $\square$

The shifts are

$$r_m \in \begin{cases} \mathbb{N}_0 + \frac{1}{2}, & d \text{ odd}, \\ \mathbb{N}, & d \text{ even}. \end{cases}$$

Thus Conjecture 1.1 is asked only on the two grids that actually occur.

4 The signed Weyl primitive

Define the right-continuous branching staircase

$$S_d(z) = \sum_{\substack{m \geq 0 \\ r_m \leq z}} c_{d,m} \tag{14}$$

and its discrepancy primitive

$$\Delta_d(z) = \int_0^z \left(S_d(u) - \frac{u^p}{p!} \right) du. \tag{15}$$

Proposition 4.1 (Signed branching primitive). *For every $d \geq 3$ and $z \geq 0$,*

$$\boxed{\Delta_d(z) \leq 0}. \tag{16}$$

For $d = 3$, equality at positive z occurs precisely at the positive integers. For $d \geq 4$, the inequality is strict for $z > 0$.

Proof. For $0 \leq z < r_0 = p/2$, $S_d(z) = 0$, so

$$\Delta_d(z) = -\frac{z^{p+1}}{(p+1)!} \leq 0.$$

Let $M \geq 0$ and $r_M = M + p/2$. The hockey-stick identity gives

$$\int_0^{r_M} S_d(u) du = \sum_{m=0}^{M-1} (r_M - r_m) c_{d,m} = \binom{M+p}{p+1}.$$

Therefore

$$(p+1)! \Delta_d(r_M) = \prod_{j=0}^p (M+j) - \left(M + \frac{p}{2} \right)^{p+1} \leq 0, \tag{17}$$

because the arithmetic mean of $M, M+1, \dots, M+p$ is $M+p/2$.

On the half-open cell $[r_M, r_{M+1})$,

$$S_d(z) = \binom{M+p}{p}, \quad \Delta'_d(z) = \binom{M+p}{p} - \frac{z^p}{p!}.$$

If the critical point lies in this cell, it is

$$z_M^* = \left(\prod_{j=1}^p (M+j) \right)^{1/p}.$$

Direct substitution into the cell formula gives

$$(p+1)! \Delta_d(z_M^*) = p(z_M^*)^p \left(z_M^* - M - \frac{p+1}{2} \right) \leq 0, \quad (18)$$

since z_M^* is the geometric mean of $M+1, \dots, M+p$ and their arithmetic mean is $M+(p+1)/2$. If the critical point is outside the cell, the cell maximum is at an endpoint, already covered by (17). This proves (16).

For $p=1$, the cell maximum is $z=M+1$ and has value zero. For $p \geq 2$, the consecutive factors in both AM–GM applications are unequal, which gives the stated strictness. $\square$

5 Exact defect identity and conditional closure

Define

$$D_A(r) = 2 \int_r^K A(z) dz - T_A(r) \quad (19)$$

and the continuous payment

$$W_d = \frac{2}{p!} \int_0^K z^p A(z) dz. \quad (20)$$

Lemma 5.1 (Exact Weyl normalization). *For $p = d-2$,*

$$\int_0^a z^p G_a(z) dz = \frac{\Gamma((p+1)/2)}{4\sqrt{\pi} (p+2)\Gamma((p+4)/2)} a^{p+2}. \quad (21)$$

Consequently,

$$W_d = \frac{1 - \rho^d}{2^d \Gamma(d/2 + 1)^2} K^d = L_d |A_{\rho,1}^{(d)}| K^d. \quad (22)$$

Proof. Scale $z = ax$ in (2), integrate the term containing $\arccos x$ by parts, and evaluate the resulting beta integrals. This gives (21). Multiplying the difference of (21) at $a = K$ and $a = \rho K$ by $2/p!$, the duplication formula for Γ reduces the coefficient to

$$\frac{1}{2^{p+2} \Gamma((p+4)/2)^2} = \frac{1}{2^d \Gamma(d/2 + 1)^2}.$$

The last equality in (22) follows from $|A_{\rho,1}^{(d)}| = \omega_d(1 - \rho^d)$. $\square$

Proposition 5.2 (Branching-defect identity). *Let*

$$\mathcal{B}_d(A) = 2 \int_0^K (-A'(z))(-\Delta_d(z)) dz. \quad (23)$$

Then $\mathcal{B}_d(A) \geq 0$ and

$$\boxed{W_d - P_d^< = \mathcal{B}_d(A) + \sum_{m \geq 0} c_{d,m} D_A(r_m)}. \quad (24)$$

Proof. The shell action is absolutely continuous, decreasing, and vanishes at K . Fubini gives

$$\sum_{m \geq 0} c_{d,m} 2 \int_{r_m}^K A(z) dz = 2 \int_0^K S_d(z) A(z) dz.$$

Since $\Delta'_d = S_d - z^p/p!$ almost everywhere and both endpoint terms vanish, integration by parts gives

$$2 \int_0^K S_d A = W_d + 2 \int_0^K A \Delta'_d = W_d - 2 \int_0^K A' \Delta_d = W_d - \mathcal{B}_d(A).$$

Both A' and Δ_d are nonpositive, so $\mathcal{B}_d(A) \geq 0$. Subtract (13) and use (19). $\square$

Theorem 5.3 (Conditional all-dimensional shell theorem). *If Conjecture 1.1 holds, then (1) holds for every $d \geq 3$, $0 < \rho < 1$, and $K \geq 0$. It then also holds for every shell $A_{r,R}^{(d)} = \{x \in \mathbb{R}^d : r < |x| < R\}$ and every spectral parameter.*

Proof. The case $K = 0$ is trivial, so assume $K > 0$. Every r_m in (11) lies in $\frac{1}{2}\mathbb{N}$, so the conjecture gives $D_A(r_m) \geq 0$. Equations (9), (24), and (22) yield

$$N_D(A_{\rho,1}^{(d)}, K^2) \leq P_d^< \leq W_d = L_d |A_{\rho,1}^{(d)}| K^d.$$

For a shell with outer radius R , take $\rho = r/R$ and $K = R\sqrt{\Lambda}$ and use Dirichlet dilation. The non-strict counting form follows by applying the strict estimate at $\Lambda + \varepsilon$ and letting $\varepsilon \downarrow 0$. $\square$

6 Proved tail regions and the first interface cell

The action derivative is

$$-A'(z) = \frac{1}{\pi} \begin{cases} \arccos \frac{z}{K} - \arccos \frac{z}{\mu}, & 0 < z < \mu, \\ \arccos \frac{z}{K}, & \mu \leq z < K. \end{cases} \quad (25)$$

Thus A is decreasing and concave on $(0, \mu)$, decreasing and convex on (μ, K) , and A, A' match at μ .

Proposition 6.1 (Outer and terminal tails). *Conjecture 1.1 holds in either of the following cases:*

1. $r \geq \mu$;
2. $G_K(r) \leq 3/4$.

Proof. If $r \geq \mu$, then $A = G_K$ on $[r, K]$; the claim is trivial when $r \geq K$. Otherwise set $B = \lceil K - r \rceil$ and zero-extend

$$g(s) = G_K(r + s) \quad (0 \leq s \leq K - r)$$

to the integer interval $[0, B]$. Since $G_K(K) = G'_K(K) = 0$, this extension is nonnegative, decreasing, convex, $1/2$ -Lipschitz, and vanishes at B . The convex shifted-floor theorem of [1] therefore bounds its ordinary-floor tail by $2 \int_0^B g$. At every sample,

$$[g(j) + 1/4]_< \leq \lfloor g(j) + 1/4 \rfloor.$$

The strict tail is consequently bounded by $2 \int_0^B g = 2 \int_r^K G_K$.

If $G_K(r) \leq 3/4$, monotonicity and $A \leq G_K$ imply $A(r + j) + 1/4 \leq 1$ for every $j \geq 0$. Every strict bracket is therefore zero. Notice that equality at 1 is harmless precisely because the bracket is strict. $\square$

We next sharpen the prescribed first milestone. For the ball action set

$$T_K(r) = [G_K(r) + \tfrac{1}{4}]_< + 2 \sum_{j \geq 1} [G_K(r + j) + \tfrac{1}{4}]_<, \quad D_K(r) = 2 \int_r^K G_K(z) dz - T_K(r). \quad (26)$$

The wall-safe translation in the preceding proof gives $D_K(r) \geq 0$.

Lemma 6.2 (A low-action $1/3$ -Lipschitz reserve). *Let g be strictly decreasing and convex on $[0, b]$, with $g(b) = 0$ and $\text{Lip}(g) \leq 1/3$. If*

$$\frac{3}{4} < g(0) < 1,$$

then

$$2 \int_0^b g(z) dz - \left([g(0) + \tfrac{1}{4}]_< + 2 \sum_{j \geq 1} [g(j) + \tfrac{1}{4}]_< \right) > \frac{11}{16}. \quad (27)$$

The function is understood to be zero-extended after b .

Proof. Let $h = g(0)$ and let $x \in (0, b)$ be defined by $g(x) = 3/4$. Since $g(j) < 1$, a sample contributes exactly one if and only if $j < x$. Therefore the bracketed count is

$$2\lceil x \rceil - 1.$$

The Lipschitz bound and $g(b) = 0$ give

$$g(z) \geq \max\{h - z/3, 0\}, \quad 2 \int_0^b g(z) dz \geq 3h^2 > \frac{27}{16}.$$

If $x \leq 1$, the count is one and (27) follows.

Suppose $x > 1$ and put $a = (h - 3/4)/x$. Convexity implies that, for $z \geq x$,

$$g(z) \geq \max\left\{\frac{3}{4} - a(z - x), 0\right\}.$$

Hence

$$\int_0^b g(z) dz \geq \frac{3}{4}x + \frac{(3/4)^2}{2a} = x \left(\frac{3}{4} + \frac{9}{32(h - 3/4)} \right) > \frac{15}{8}x.$$

Since $2\lceil x \rceil - 1 < 2x + 1$, the defect is larger than

$$\frac{15}{4}x - (2x + 1) = \frac{7}{4}x - 1 > \frac{3}{4} > \frac{11}{16}.$$

$\square$

Proposition 6.3 (Exact one-interface reduction). *Assume*

$$0 < \mu - r < 1, \quad r \in \frac{1}{2}\mathbb{N}.$$

Define

$$n_0 = [G_K(r) + \frac{1}{4}]_{<} - [G_K(r) - G_\mu(r) + \frac{1}{4}]_{<} \in \mathbb{N}_0 \quad (28)$$

and

$$\delta_\mu(r) = \int_r^\mu G_\mu(z) \, dz.$$

Then

$$\boxed{D_A(r) = D_K(r) + n_0 - 2\delta_\mu(r).} \quad (29)$$

Moreover,

$$2\delta_\mu(r) < \frac{4(\mu - r)^{5/2}}{15\sqrt{\mu}} < \frac{4\sqrt{2}}{15} < 1. \quad (30)$$

Consequently, the shifted-tail inequality holds if any of the following is true:

1. $n_0 \geq 1$;
2. $G_K(\max\{r, K/2\}) \geq 1$;
3. $r \geq K/2$.

The only remaining one-interface configurations satisfy

$$\boxed{r < \frac{K}{2}, \quad K < \frac{1}{\omega_0}, \quad n_0 = 0, \quad \omega_0 := G_1(1/2) = \frac{\sqrt{3}}{2\pi} - \frac{1}{6}.} \quad (31)$$

In particular,

$$r \in \left\{ \frac{1}{2}, 1, \frac{3}{2}, 2, \frac{5}{2}, 3, \frac{7}{2}, 4, \frac{9}{2} \right\}.$$

Proof. Since $r + j > \mu$ for every $j \geq 1$, all samples after the first agree with the ball samples. Thus $T_A(r) = T_K(r) - n_0$. Subtracting $2 \int_r^\mu G_\mu$ from the ball integral proves (29).

The turning estimate

$$G_\mu(z) < \frac{(\mu - z)^{3/2}}{3\sqrt{\mu}} \quad (0 < z < \mu)$$

integrates to (30), using $\mu > r \geq 1/2$ and $\mu - r < 1$. If $n_0 \geq 1$, then $D_K(r) \geq 0$ and (29) is positive.

For $r < K$, use the same integer endpoint $B = \lceil K - r \rceil$ and zero-extended function $g(s) = G_K(r + s)$. Put

$$s_* = \max\{0, K/2 - r\}.$$

Then g is $1/3$ -Lipschitz on $[s_*, B]$ and $g(s_*) = G_K(\max\{r, K/2\})$. The refined ordinary-floor theorem of [3, Theorem 3.4], followed by $[\cdot]_{<} \leq \lfloor \cdot \rfloor$, gives

$$D_K(r) \geq \frac{1}{2} \lfloor G_K(\max\{r, K/2\}) \rfloor. \quad (32)$$

For $r \geq K$ the same inequality is trivial. Hence the second condition and (30) prove the claim.

If $r \geq K/2$ and $G_K(r) < 1$, the translated ball action is globally $1/3$ -Lipschitz. Its tail is either zero or Lemma 6.2 applies. In the latter case $D_K(r) > 11/16 > 4\sqrt{2}/15$, so (29) is again positive. This proves the third condition.

If none of the three conditions holds, then $r < K/2$ and $G_K(K/2) = \omega_0 K < 1$, giving (31). The elementary rational bounds

$$\frac{265}{153} < \sqrt{3} < \frac{26}{15}, \quad \frac{333}{106} < \pi < \frac{355}{113}$$

(the square-root bounds follow by squaring) give $1/10 < \omega_0 < 1/9$. Consequently

$$\frac{9}{2} < \frac{1}{2\omega_0} < 5.$$

Since $r < K/2 < 1/(2\omega_0)$ and $r \in \frac{1}{2}\mathbb{N}$, only the nine displayed shifts remain. $\square$

Remark 6.4 (The residual first-interface family). Proposition 6.3 replaces the only unresolved part of the first-interface problem by a bounded nine-shift family: $2r < K < 1/\omega_0$, $r < \mu < \min\{r+1, K\}$, and $n_0 = 0$. The inequality to be checked there is exactly

$$D_K(r) \geq 2\delta_\mu(r).$$

Proposition 6.5 and Lemma 6.6 below close this family.

Proposition 6.5 (Finite ball-wall reduction). *In the residual family of Proposition 6.3, put*

$$B = G_K(r) + \frac{1}{4}, \quad q = [B]_<, \quad e = B - q \in (0, 1],$$

and set

$$E_r(K) = D_K(r) - e.$$

Then

$$2\delta_\mu(r) < e. \tag{33}$$

Consequently, the residual one-interface inequality follows from $E_r(K) \geq 0$.

On every open K -interval containing no ball-sample wall, E_r is strictly increasing. At a wall of the first sample $G_K(r) + 1/4$, the two unit drops in $D_K(r)$ and e cancel, so E_r is continuous. At a wall of a later sample $G_K(r+j) + 1/4$, $j \geq 1$, E_r drops by two for each crossed level. It is therefore enough to check the right-hand limits at the finitely many later-sample walls, together with the first positive-count wall and any lower endpoint $K = 2r$ at which the count is already positive.

More explicitly, all walls that can occur have the form

$$G_K(r+j) = n - \frac{1}{4}, \quad j \geq 0, \quad n \in \{1, 2, 3\}, \quad r+j < 10. \tag{34}$$

Proof. Let $h = G_\mu(r)$. If $q = 0$, strict increase of $G_a(r)$ in the radius a gives $h < G_K(r) < B = e$. If $q \geq 1$, write $B = q + e$ and use the strict bracket convention; then $[B-h]_< = [B]_<$ holds exactly when $h < e$. Thus, in both cases,

$$n_0 = 0 \iff [B-h]_< = [B]_< \iff h < e.$$

When $q \geq 1$, equality $h = e$ lands on the integral wall q and causes a strict-count drop. The function G_μ is strictly convex on $[r, \mu]$, with endpoint values h and 0. It lies strictly below its endpoint chord, and therefore

$$2\delta_\mu(r) < h(\mu - r) < e,$$

which proves (33). If $T_K(r) = 0$, then $T_A(r) = 0$ and the desired inequality is immediate. Otherwise it is enough, by (29), to prove $D_K(r) \geq e$.

Away from sample walls the integer-valued tail and q are constant. Writing $x = r/K$, differentiation of the ball action and its tail integral gives

$$\pi E'_r(K) = K \left(\arccos x - x\sqrt{1-x^2} \right) - \sqrt{1-x^2}, \quad (35)$$

and

$$\pi E''_r(K) = \arccos x + x\sqrt{1-x^2} - \frac{x^2}{K\sqrt{1-x^2}}. \quad (36)$$

Here $x < 1/2$ and $K > 1$. Hence

$$\pi E''_r(K) > \frac{\pi}{3} - \frac{1}{2\sqrt{3}} > 0.$$

For $r \geq 1$, evaluation at $K = 2r$ gives

$$\pi E'_r(2r) = \frac{2\pi r}{3} - \frac{\sqrt{3}}{2}(r+1) > 0.$$

For $r = 1/2$, positivity of the tail count implies $G_K(r) > 3/4$. Since G_K is $1/2$ -Lipschitz and vanishes at K , this forces $K > 2$. Formula (35) is already positive at $(r, K) = (1/2, 2)$, since

$$\pi E'_{1/2}(2) = 2 \arccos \frac{1}{4} - \frac{3\sqrt{15}}{8} > \frac{2\pi}{3} - \frac{3}{2} > 0.$$

Equation (36) then keeps it positive. Thus E_r is strictly increasing between walls.

The stated jump rules follow directly from the weights $1, 2, 2, \dots$ in $T_K(r)$. At a first-sample wall, e equals 1 and its right-hand limit is 0, exactly matching the unit right-hand drop of D_K . Simultaneous later-sample crossings are combined at the same wall abscissa. Finally, $K < 1/\omega_0 < 10$ and $G_K(z) < K/\pi < 10/3$. Thus only $n = 1, 2, 3$ and samples $r + j < 10$ can occur, and

$$\partial_K G_K(z) = \frac{1}{\pi} \sqrt{1 - \frac{z^2}{K^2}} > 0 \quad (K > z)$$

shows that every pair (j, n) produces at most one wall. This proves the finite reduction. $\square$

Lemma 6.6 (Exact residual wall certificate). *In the residual family of Proposition 6.5, if $T_K(r) > 0$, then*

$$E_r(K) > 0. \quad (37)$$

More precisely, $E_r(K) > 3/20$ for $r \leq 4$, while for $r = 9/2$ one has

$$E_{9/2}(K) > \frac{1253}{555}.$$

Proof. Put

$$H_r(K) = 2 \int_r^K G_K(z) dz - G_K(r) - \frac{1}{4}, \quad M_r(K) = \sum_{j \geq 1} [G_K(r+j) + \frac{1}{4}]_<.$$

The first-sample bracket cancels from the definition of E_r , including at its strict walls, and hence

$$E_r(K) = H_r(K) - 2M_r(K). \quad (38)$$

For $x = r/K$, direct integration gives

$$H_r(K) = \frac{A}{2} - \frac{A \arcsin x + B\sqrt{1-x^2}}{\pi} - \frac{1}{4}, \quad A = \frac{K^2}{2} + r^2 + r, \quad B = K \left(\frac{3r}{2} + 1 \right). \quad (39)$$

We record rational envelopes sufficient to check every wall. For $0 \leq x \leq 13/20$, let

$$a_m = \frac{\binom{2m}{m}}{4^m(2m+1)}, \quad b_m = \frac{\binom{2m}{m}}{4^m(2m-1)},$$

and set

$$L(x) = \sum_{m=0}^6 a_m x^{2m+1}, \quad U(x) = L(x) + \frac{x^{15}}{1-x^2},$$

$$V(x) = 1 - \sum_{m=1}^6 b_m x^{2m}, \quad W(x) = V(x) - \frac{x^{14}}{1-x^2}.$$

The power series, whose omitted positive coefficients are smaller than one, give

$$L \leq \arcsin x \leq U, \quad W \leq \sqrt{1-x^2} \leq V.$$

Together with $333/106 < \pi < 355/113$, they imply

$$\begin{aligned} \underline{G}(K, z) &= K \left[\frac{113}{355} (W + xL) - \frac{x}{2} \right] \leq G_K(z) \\ &\leq K \left[\frac{106}{333} (V + xU) - \frac{x}{2} \right] = \overline{G}(K, z), \end{aligned} \quad (40)$$

where now $x = z/K$, and

$$\frac{A}{2} - \frac{106}{333} (AU + BV) - \frac{1}{4} \leq H_r(K) \leq \frac{A}{2} - \frac{113}{355} (AL + BW) - \frac{1}{4}. \quad (41)$$

Thus every comparison below is a rational-number comparison.

Let $k_n(z)$ denote the unique solution of $G_K(z) = n-1/4$. Substitution in (40) gives the following pairwise disjoint brackets:

wall	bracket	wall	bracket
$k_1(3/2)$	$(22/5, 9/2)$	$k_1(2)$	$(5, 26/5)$
$k_1(5/2)$	$(57/10, 29/5)$	$k_1(3)$	$(63/10, 32/5)$
$k_1(7/2)$	$(69/10, 7)$	$k_1(4)$	$(15/2, 38/5)$
$k_2(3/2)$	$(77/10, 39/5)$	$k_1(9/2)$	$(81/10, 41/5)$
$k_2(2)$	$(83/10, 17/2)$	$k_1(5)$	$(87/10, 44/5)$
$k_2(5/2)$	$(9, 227/25)$		

(40a)

These are all later-sample walls. Indeed, the same rational bounds give

$$\frac{227}{25} < \frac{1}{\omega_0} < \frac{459}{50}$$

and, at $K = 459/50$,

$$\overline{G}(K, 11/2) < \frac{18}{25} < \frac{3}{4}, \quad \overline{G}(K, 3) < \frac{8}{5} < \frac{7}{4}, \quad \overline{G}(K, 3/2) < \frac{9}{4} < \frac{11}{4}.$$

Monotonicity in K and z excludes, respectively, every omitted level-3/4, level-7/4, and level-11/4 or higher wall.

At a right-hand wall limit, the value of M_r is the event number in the following exact order:

r	$(z, n; M_r)$
$\frac{1}{2}$	$(\frac{3}{2}, 1; 1), (\frac{5}{2}, 1; 2), (\frac{7}{2}, 1; 3), (\frac{3}{2}, 2; 4), (\frac{9}{2}, 1; 5), (\frac{5}{2}, 2; 6)$
1	$(2, 1; 1), (3, 1; 2), (4, 1; 3), (2, 2; 4), (5, 1; 5)$
$\frac{3}{2}$	$(\frac{5}{2}, 1; 1), (\frac{7}{2}, 1; 2), (\frac{9}{2}, 1; 3), (\frac{5}{2}, 2; 4)$
2	$(3, 1; 1), (4, 1; 2), (5, 1; 3)$
$\frac{5}{2}$	$(\frac{7}{2}, 1; 1), (\frac{9}{2}, 1; 2)$
3	$(4, 1; 1), (5, 1; 2)$
$\frac{7}{2}$	$(\frac{9}{2}, 1; 1)$
4	$(5, 1; 1)$
$\frac{9}{2}$	$\emptyset.$

Equations (39)–(41) then give the strict rational bounds

r	first-wall lower	first-wall upper	activation or $K = 2r$	other-wall lower
$\frac{1}{2}$	$> 3/20$	$< 8/25$	$> 11/25$	$> 13/20$
1	$> 17/100$	$< 1/2$	$> 14/25$	$> 7/10$
$\frac{3}{2}$	$> 9/25$	$< 53/100$	$> 69/100$	$> 37/50$
2	$> 2/5$	$< 29/50$	$> 37/50$	$> 39/50$
$\frac{5}{2}$	$> 23/50$	$< 63/100$	$> 22/25$	$> 81/100$
3	$> 13/25$	$< 69/100$	$> 47/50$	$> 17/20$
$\frac{7}{2}$	$> 29/50$	$< 19/25$	$> 109/100$	—
4	$> 33/50$	$< 83/100$	$> 163/100$	— .

For each $r \leq 4$, the two first-wall columns concern the right-hand value at $G_K(r+1) = 3/4$. Its displayed upper bound is smaller than both competing lower bounds, so this is the controlling wall. Formula (35) shows that H_r , and hence E_r between its later-sample walls, is increasing throughout the positive-count residual region. For $r \leq 3$ the tail is zero before its first-sample activation; for $r = 7/2, 4$ the displayed $K = 2r$ value owns the lower endpoint. Finally, for $r = 9/2$ there is no later wall and

$$E_{9/2}(9) = \frac{43}{2} - \frac{279\sqrt{3}}{8\pi} > \frac{1253}{555}.$$

This proves (37). A dependency-free replay of all 628 rational comparisons is supplied in `scripts/general_d_finite_wall_fraction_verify.py`; it is an audit of, not a numerical premise for, the displayed estimates. $\square$

Theorem 6.7 (One-interface shifted tails). *Conjecture 1.1 holds whenever*

$$0 < \mu - r < 1.$$

It also holds at the endpoint $r = \mu$ by Proposition 6.1.

Proof. The three regions in Proposition 6.3 are already proved there. In its residual family, if $T_K(r) = 0$, then $T_A(r) = 0$ and the claim is immediate. Otherwise Lemma 6.6 and Proposition 6.5 give

$$D_K(r) = E_r(K) + e > e > 2\delta_\mu(r).$$

The exact identity (29) now gives $D_A(r) > 0$. $\square$

Proposition 6.8 (Fixed-ratio high-energy tails). *For $0 < \rho < 1$, put*

$$c_\rho = \frac{\arccos \rho}{\pi}, \quad d_\rho = 1 - 2c_\rho = \frac{2 \arcsin \rho}{\pi},$$

$$\eta_\rho = G_1 \left(\max \left\{ \rho, \frac{1}{2} \right\} \right), \quad a_\rho = \frac{2\pi\rho}{1-\rho}, \quad C_* = \frac{1}{2} + \frac{8\sqrt{2}}{15},$$

and

$$K_0(\rho) = \frac{a_\rho + 2\eta_\rho C_* + \sqrt{a_\rho^2 + 4a_\rho \eta_\rho C_* + \eta_\rho^2}}{2\eta_\rho^2}. \quad (42)$$

If $K \geq K_0(\rho)$, then Conjecture 1.1 holds for every $r \in \frac{1}{2}\mathbb{N}$.

Proof. Only $r < \mu$ requires proof. Set

$$n = \lfloor \mu - r \rfloor, \quad q = r + n, \quad B = \lceil K - r \rceil.$$

Thus $q \leq \mu < q + 1$. For integers $a < b$, write

$$\mathbf{T}(f; a, b) = \frac{1}{2} \lfloor f(a) \rfloor + \sum_{j=a+1}^{b-1} \lfloor f(j) \rfloor + \frac{1}{2} \lfloor f(b) \rfloor.$$

The strict brackets in T_A are bounded above by ordinary floors. Additivity at the split point and $A \leq G_K$ therefore give, when $n \geq 1$,

$$\frac{T_A(r)}{2} \leq \mathbf{T} \left(A(r + \cdot) + \frac{1}{4}; 0, n \right) + \mathbf{T} \left(G_K(r + \cdot) + \frac{1}{4}; n, B \right). \quad (43)$$

For $n = 0$, the first term is absent and the same inequality follows by direct pointwise majorization.

Assume first $n \geq 1$ and put

$$h(t) = A(r + t) + \frac{1}{4}, \quad m_j = \lfloor h(j) \rfloor.$$

If the nonincreasing sequence $(m_j)_{j=0}^n$ leaves its first shelf, let p be its last index on that shelf; otherwise put $p = n$. On $[0, n]$, h is decreasing, concave, and c_ρ -Lipschitz. The quantitative concave trapezoidal-floor theorem [3, Theorem 1.4], with the basic concavity bound in the no-drop case, yields

$$\mathbf{T}(h; 0, n) \leq \int_0^n A(r + t) dt + \frac{n}{4} - \frac{1 - c_\rho}{2}(n - p). \quad (44)$$

For $n = 0$, take $p = 0$ and read this as an absent identity.

Let $\sigma = -A'$. On $0 < z < \mu$,

$$\sigma'(z) = \frac{1}{\pi\sqrt{\mu^2 - z^2}} - \frac{1}{\pi\sqrt{K^2 - z^2}} \geq \kappa := \frac{1 - \rho}{\pi\rho K}.$$

Indeed, after subtracting the value at $z = 0$, the remaining bracket is decreasing in its radius parameter. Since $\sigma(0) = 0$,

$$A(x) - A(y) = \int_x^y \sigma(z) dz \geq \frac{\kappa}{2}(y^2 - x^2) \geq \frac{\kappa}{2}(y - x)^2.$$

The samples at r and $r + p$ lie on the same floor shelf, so their action drop is smaller than one. Keeping the r -term in the preceding estimate and using $r \geq 1/2$ gives

$$p < \sqrt{r^2 + a_\rho K} - r \leq \sqrt{a_\rho K + \frac{1}{4}} - \frac{1}{2}. \quad (45)$$

For the convex suffix, set

$$s = \max \left\{ q, \frac{K}{2} \right\}, \quad t_* = s - r.$$

Then $t_* \in [n, B]$, the translated ball action is $1/2$ -Lipschitz on $[n, B]$ and $1/3$ -Lipschitz on $[t_*, B]$, and

$$G_K(s) \geq K\eta_\rho.$$

The quantitative convex theorem [3, Theorem 3.4] gives

$$\mathbf{T} \left(G_K(r + \cdot) + \frac{1}{4}; n, B \right) \leq \int_n^B G_K(r + t) dt - \frac{1}{4} \lfloor K\eta_\rho \rfloor. \quad (46)$$

The only mismatch between the two integrals on the right of (43) and the shell integral is

$$\delta = \int_q^\mu G_\mu(z) dz.$$

The turning bound from [3, Lemma 6.2] and $0 \leq \mu - q < 1$, $\mu > 1/2$, give

$$0 \leq \delta \leq \frac{2(\mu - q)^{5/2}}{15\sqrt{\mu}} < \frac{2\sqrt{2}}{15}. \quad (47)$$

Combining (43), (44), and (46) gives

$$\frac{T_A(r)}{2} \leq \int_r^K A(z) dz + \delta - \frac{M}{4}, \quad M = \lfloor K\eta_\rho \rfloor + d_\rho(n - p) - p. \quad (48)$$

By (45),

$$M > K\eta_\rho - \sqrt{a_\rho K + \frac{1}{4}} - \frac{1}{2}.$$

The definition (42) is exactly the positive-root condition

$$K\eta_\rho - \sqrt{a_\rho K + \frac{1}{4}} \geq C_*.$$

Thus $M > 8\sqrt{2}/15 > 4\delta$, and (48) proves the result. All inequalities use ordinary floors on the majorizing side, so strict action walls require no limiting argument. $\square$

The fixed-ratio threshold is deliberately uniform in the shift, but it is not sharp as $\rho \uparrow 1$. Retaining the location of the first shelf improves the thin-shell scale by two powers of $1 - \rho$.

Lemma 6.9 (Optimized thin first-shelf loss). *Let*

$$\rho = 1 - \varepsilon, \quad 0 < \varepsilon \leq \frac{1}{100}, \quad K\varepsilon^2 \geq \frac{125}{8}.$$

Use n, p from the proof of Proposition 6.8, put $m = n - p$, and set

$$d = d_\rho, \quad \mathcal{R} = p - dm.$$

If $\mathcal{R} > 0$, then

$$\mathcal{R} < \frac{361}{80\sqrt{\varepsilon}}. \quad (49)$$

Proof. This is a scale-free consequence of the exact local slope and the same-floor condition. Put

$$y = \sqrt{\varepsilon}, \quad \varkappa = K\varepsilon^2, \quad P = py, \quad \mathfrak{r} = \mathcal{R}y, \quad S = (p + m)y.$$

Elementary trigonometric bounds give

$$d > \frac{9}{10}, \quad S = P + \frac{P - \mathfrak{r}}{d} < \frac{19}{9}P. \quad (50)$$

Write

$$U = \mu - r, \quad t = \frac{r}{\mu}, \quad w = \frac{U}{\mu} = 1 - t,$$

and define

$$\mathcal{Q} = 1 + \frac{y^2}{\varkappa} + \frac{yS}{\varkappa}, \quad \hat{q} = \frac{y^2}{\rho}(\mathcal{Q} - 1).$$

The unconditional first-shelf bound $p^2 < 2\pi\rho K/\varepsilon$ and (50) give

$$w < \frac{1 + p + m}{\rho K} = \frac{y^4 + y^3 S}{\varkappa \rho} = \hat{q} < \frac{y^4}{\varkappa \rho} + \frac{19}{9}y\sqrt{\frac{2\pi}{\varkappa \rho}}.$$

Here the first term is at most $8/1237500 < 1/1000$, while the square of the factor following $19/9$ is at most $352/86625 < 1/225$. Hence

$$w < \hat{q} < \frac{1}{1000} + \frac{19}{135} < \frac{1}{7}, \quad t > 1 - \hat{q} > \frac{6}{7}. \quad (51)$$

Using

$$-A'(z) = \frac{1}{\pi} \left(\arccos \frac{z}{K} - \arccos \frac{z}{\mu} \right)$$

in the same-floor inequality $A(r) - A(r + p) < 1$ gives, after scaling,

$$P^2 < \mathcal{H}(P, \mathfrak{r}) := \frac{2\pi^2 \mathcal{Q}}{(1 - \hat{q})^2}. \quad (52)$$

On the preceding domain, $\mathcal{H}$ decreases in $\mathfrak{r}$ and

$$\partial_P \mathcal{H} < \frac{673816}{1366875} < \frac{1}{2}.$$

Let $B_* = 361/80$. If $\mathfrak{r} \geq B_*$, then $P \geq \mathfrak{r}$ and $P^2 - \mathcal{H}(P, B_*)$ is increasing for $P \geq B_*$. Indeed, every intermediate $P' \in [B_*, P]$ retains the unconditional shelf ceiling and satisfies $S(P', B_*) < 19P'/9$, so (51) and the displayed derivative bound hold throughout the comparison interval. Its minimum over $y \leq 1/10$ and $\varkappa \geq 125/8$ is at $P = B_*$, $y = 1/10$, and $\varkappa = 125/8$. There

$$\mathcal{Q} = \frac{12869}{12500}, \quad \hat{q} = \frac{41}{137500},$$

and, using $\pi < 22/7$,

$$B_*^2 - 2 \left(\frac{22}{7} \right)^2 \frac{12869/12500}{(1 - 41/137500)^2} = \frac{72581986185449}{5925464687161600} > 0.$$

This contradicts (52) and proves (49). All displayed comparisons are rational once the elementary bound for π is inserted; the dependency-free script `computations/round9_verify_thin_plateau_constants.py` replays the finite arithmetic. $\square$

Proposition 6.10 (Thin high-energy shifted tails). *Let $\rho = 1 - \varepsilon$ with $0 < \varepsilon \leq 1/100$. If*

$$K \geq \frac{125}{8\varepsilon^2}, \quad (53)$$

then Conjecture 1.1 holds for every $r \in \frac{1}{2}\mathbb{N}$.

Proof. The outer tails are covered by Proposition 6.1. For an inner tail, retain the notation of Proposition 6.8. Before its final estimate, (48) gives

$$\frac{T_A(r)}{2} \leq \int_r^K A(z) dz + \delta - \frac{M}{4}, \quad M = \lfloor KG_1(\rho) \rfloor - \mathcal{R},$$

with $4\delta < 8\sqrt{2}/15$. If $\mathcal{R} \leq 0$, the convex gain already gives $M > 1 > 4\delta$. If $\mathcal{R} > 0$, Lemma 6.9,

$$G_1(1 - \varepsilon) \geq \frac{2\sqrt{2}}{3\pi}\varepsilon^{3/2}, \quad \sqrt{2} > \frac{140}{99}, \quad \sqrt{2} < \frac{99}{70}, \quad \pi < \frac{1571}{500},$$

and $\lfloor x \rfloor > x - 1$ give

$$M > \frac{2829397}{3732696} > \frac{132}{175} > \frac{8\sqrt{2}}{15} > 4\delta.$$

Thus (48) proves the claim. The proof uses only a positive shift and a unit-spaced sampling grid; it is unchanged on the integer and half-integer grids, and every threshold equality is owned by the strict-floor majorant. $\square$

Corollary 6.11 (A uniform small-hole sector). *If*

$$0 < \rho < \omega_0, \quad K(\omega_0 - \rho) \geq \frac{1}{2} + \frac{8\sqrt{2}}{15},$$

then Conjecture 1.1 holds for every $r \in \frac{1}{2}\mathbb{N}$.

Proof. For $r < \mu$, use the split (43), basic concavity on the head, and (46) with $s = K/2$. This gives

$$\frac{T_A(r)}{2} \leq \int_r^K A(z) dz + \delta - \frac{\lfloor \omega_0 K \rfloor - n}{4}.$$

Since $r \geq 1/2$,

$$\lfloor \omega_0 K \rfloor - n > K(\omega_0 - \rho) - \frac{1}{2} \geq \frac{8\sqrt{2}}{15} > 4\delta.$$

The outer tails were proved in Proposition 6.1. $\square$

7 Cell recurrence and the corrected global target

For $x_j = r + j$, set

$$Q_j = [A(x_j) + \frac{1}{4}]_<, \quad D_j = 2 \int_{x_j}^K A(z) dz - \left(Q_j + 2 \sum_{k>j} Q_k \right).$$

Then

$$D_j - D_{j+1} = 2 \int_{x_j}^{x_{j+1}} A(z) dz - Q_j - Q_{j+1}. \quad (54)$$

If

$$e_j = A(x_j) + \frac{1}{4} - Q_j \in (0, 1]$$

and

$$C_j = 2 \int_{x_j}^{x_j+1} A(z) dz - A(x_j) - A(x_j + 1),$$

then

$$D_j - D_{j+1} = C_j - \frac{1}{2} + e_j + e_{j+1}. \quad (55)$$

On a fully inner cell,

$$C_j = - \int_0^1 u(1-u) A''(x_j + u) du \geq 0.$$

Individual increments in (55) can nevertheless be negative, and a negative increment need not be paid by the next floor-drop cell. Thus the viable block formulation must retain the terminal convex reserve. If J is the first index with $x_J \geq \mu$, the exact target is

$$\boxed{D_0 = \sum_{j=0}^{J-1} (D_j - D_{j+1}) + D_J \geq 0,} \quad (56)$$

where $D_J \geq 0$ is the translated convex ball-tail defect. A proof of (56) needs a quantitative, not merely nonnegative, lower bound for D_J or an equivalent matched shelf reserve.

The many-cell floor dynamics can in fact be compressed to one initial shelf and one terminal one-interface tail. This is the form used below.

Proposition 7.1 (Wall-safe first-shelf reduction). *Suppose $r < \mu$ and put*

$$n = \lfloor \mu - r \rfloor, \quad q = r + n, \quad d_\rho = \frac{2 \arcsin \rho}{\pi}.$$

For $0 \leq j \leq n$, let

$$F_j = \left\lfloor A(r + j) + \frac{1}{4} \right\rfloor.$$

If this sequence leaves its first shelf, let p be the last index on that shelf; otherwise put $p = n$. Define

$$R_p = 2 \int_r^{r+p} A(z) dz - 2pF_0. \quad (57)$$

Then, at ordinary and strict floor walls alike,

$$\boxed{D_A(r) \geq D_A(q) + R_p + \frac{d_\rho}{2}(n - p).} \quad (58)$$

Moreover, if

$$\varepsilon_j = A(r + j) + \frac{1}{4} - F_j \in [0, 1), \quad \sigma = -A',$$

then

$$R_p = C_p + p \left(\varepsilon_0 + \varepsilon_p - \frac{1}{2} \right), \quad (59)$$

$$C_p := - \int_0^p u(p - u) A''(r + u) du \geq 0, \quad (60)$$

and

$$R_p \geq -\frac{p}{2} + 2 \int_0^p u \sigma(r+u) \, du. \quad (61)$$

In particular, with

$$\kappa = \frac{1-\rho}{\pi \rho K}, \quad a_\rho = \frac{2\pi \rho}{1-\rho},$$

one has

$$p < \sqrt{r^2 + a_\rho K} - r. \quad (62)$$

Proof. Put $h(t) = A(r+t) + 1/4$. At half-tail scale, the strict trapezoidal sum

$$\mathbf{T}_<(h; a, b) := \frac{1}{2}[h(a)]_< + \sum_{j=a+1}^{b-1} [h(j)]_< + \frac{1}{2}[h(b)]_<$$

on $[0, p]$ is at most pF_0 . If $p < n$, the ordinary floor of h drops immediately after the left endpoint of $[p, n]$. The quantitative concave trapezoidal-floor theorem [3, Theorem 1.4] gives

$$\mathbf{T}_<(h; p, n) \leq \int_p^n A(r+t) \, dt - \frac{d_\rho}{4}(n-p).$$

If $p = n$, this block is absent. Additivity at p and n leaves exactly one half of the strict tail beginning at q , and hence

$$\frac{T_A(r)}{2} \leq pF_0 + \int_{r+p}^q A(z) \, dz - \frac{d_\rho}{4}(n-p) + \frac{T_A(q)}{2}.$$

Subtracting this from $\int_r^K A$ proves (58). At an ordinary wall the strict bracket is smaller than the ordinary floor used in the majorant, so no limiting argument is needed.

Since $F_0 = F_p$, the trapezoidal identity gives (59)–(60). Also $F_0 \leq A(r+p) + 1/4$, and therefore

$$R_p \geq -\frac{p}{2} + 2 \int_0^p (A(r+t) - A(r+p)) \, dt,$$

which is (61) after reversing the order of integration. Finally, on the inner branch

$$\sigma'(z) = \frac{1}{\pi \sqrt{\mu^2 - z^2}} - \frac{1}{\pi \sqrt{K^2 - z^2}} \geq \kappa, \quad \sigma(0) = 0.$$

Equality of the two endpoint floors implies $A(r) - A(r+p) < 1$. Consequently

$$1 > \int_r^{r+p} \sigma(z) \, dz \geq \frac{\kappa}{2}((r+p)^2 - r^2),$$

which is exactly (62). $\square$

The proposition identifies the sharp scalar which remains. The point q is a one-interface start, so Theorem 6.7 gives $D_A(q) \geq 0$ and, more precisely,

$$D_A(q) = D_K(q) + \lambda_q - 2 \int_q^\mu G_\mu(z) \, dz, \quad \lambda_q = [G_K(q) + \tfrac{1}{4}]_< - [A(q) + \tfrac{1}{4}]_<. \quad (63)$$

Thus every tail satisfying

$$d_\rho(n-p) \geq p \quad (64)$$

is already proved by (58) and $R_p \geq -p/2$. Any residual tail must satisfy the explicit interface-band restriction

$$n < \frac{1 + d_\rho}{d_\rho} \left(\sqrt{r^2 + a_\rho K} - r \right), \quad (65)$$

and the sole remaining inequality is

$$\boxed{D_A(q) + R_p + \frac{d_\rho}{2}(n - p) \geq 0.} \quad (66)$$

Unlike a local bad-cell pairing, this scalar retains the terminal convex reserve forced by the counterexamples above.

Lemma 7.2 (Scale-free shelf curvature). *In the notation of Proposition 7.1, suppose $p \geq 1$ and put*

$$\Delta = A(r) - A(r + p).$$

Then

$$C_p \geq \frac{p^2}{3(2r + p)} \Delta. \quad (67)$$

Consequently, since $\Delta = \varepsilon_0 - \varepsilon_p$ on the first ordinary-floor shelf,

$$R_p \geq \frac{p^2}{3(2r + p)}(\varepsilon_0 - \varepsilon_p) + p \left(\varepsilon_0 + \varepsilon_p - \frac{1}{2} \right). \quad (68)$$

Moreover,

$$A(0) - A(r) \leq \frac{r\Delta}{2p}. \quad (69)$$

Proof. On the inner branch $\sigma = -A'$ is nonnegative, increasing, and convex, with $\sigma(0) = 0$. Indeed,

$$\sigma'(z) = \frac{1}{\pi\sqrt{\mu^2 - z^2}} - \frac{1}{\pi\sqrt{K^2 - z^2}},$$

and differentiating once more preserves the sign. Use the positive hinge representation

$$\sigma(z) = az + \int_{[0, \infty)} (z - s)_+ d\nu(s), \quad a \geq 0, \quad \nu \geq 0.$$

Both sides of (67) are linear in σ . For the generator $\sigma(z) = z$ their ratio is

$$\frac{p^3/6}{p(r + p/2)} = \frac{p^2}{3(2r + p)}.$$

For a hinge with $s \leq r$, the same ratio is

$$\frac{p^3/6}{p(r - s + p/2)} \geq \frac{p^2}{3(2r + p)}.$$

If $r < s < r + p$ and $L = r + p - s$, the ratio is $p - 2L/3 \geq p/3$, which is again at least the required constant. Hinges to the right contribute zero. This proves (67), and (68) follows from (59).

Finally, convexity and $\sigma(0) = 0$ imply that $\sigma(z)/z$ is nondecreasing. Hence

$$\int_0^r \sigma(z) dz \leq \frac{r}{2} \sigma(r), \quad \Delta \geq p\sigma(r),$$

which proves (69). □

Lemma 7.3 (Fractional-top convex reserve). *Let g be nonnegative, strictly decreasing, convex, and C^1 on $[0, L]$, with $g(L) = 0$, and extend it by zero. Set*

$$v = g(0), \quad c_v = -g'(0) > 0,$$

and

$$D_g = 2 \int_0^L g(s) ds - \left([g(0) + \tfrac{1}{4}]_< + 2 \sum_{j \geq 1} [g(j) + \tfrac{1}{4}]_< \right), \quad B = [g(0) + \tfrac{1}{4}]_<.$$

For $1 \leq k \leq B$, let y_k be the unique point satisfying $g(y_k) = k - 1/4$, and put $c_k = -g'(y_k)$. Then

$$D_g \geq \sum_{k=1}^B \left(\frac{1}{2c_k} - 1 \right) + \frac{(v - B)_+^2}{c_v}. \quad (70)$$

Proof. Let $y(t)$ be the decreasing inverse of g , extended by zero for $t \geq g(0)$. Convexity of g makes y convex; the zero extension remains convex because its right derivative is zero. Layer cake gives

$$2 \int_0^L g = 2 \int_0^{g(0)} y(t) dt.$$

The level $k - 1/4$ contributes exactly $2\lceil y_k \rceil - 1 \leq 2y_k + 1$ to the strict tail, including when y_k is integral. The tangent to y at $k - 1/4$ gives

$$\int_{k-1}^k y(t) dt \geq y_k - \frac{1}{4} y'(k - \tfrac{1}{4}) = y_k + \frac{1}{4c_k}.$$

These disjoint unit intervals cover $[0, B]$. If $v > B$, the supporting tangent at the top endpoint is

$$y(t) \geq \frac{v - t}{c_v} \quad (B \leq t \leq v),$$

and therefore the unused top interval contributes

$$2 \int_B^v y(t) dt \geq \frac{(v - B)^2}{c_v}.$$

If $v \leq B$ this interval is empty. Adding this contribution proves (70). At a top action wall the equality level is absent from B , exactly as required by the strict convention. $\square$

Corollary 7.4 (Exact-angle terminal reserve). *At the one-interface point q , put*

$$B = [G_K(q) + \tfrac{1}{4}]_<, \quad Q = [A(q) + \tfrac{1}{4}]_<, \quad v_q = G_K(q), \quad \beta_q = \frac{\arccos(q/K)}{\pi}.$$

For $1 \leq k \leq B$, let $\theta_k \in (0, \pi/2)$ be the unique solution of

$$\frac{K}{\pi} (\sin \theta_k - \theta_k \cos \theta_k) = k - \frac{1}{4}.$$

Then

$$D_A(q) \geq \max \left\{ 0, \frac{\pi}{2} \sum_{k=1}^B \frac{1}{\theta_k} - Q - 2 \int_q^\mu G_\mu(z) dz + \frac{(v_q - B)_+^2}{\beta_q} \right\}. \quad (71)$$

Proof. Apply Lemma 7.3 to $g(s) = G_K(q + s)$. At the level $k - 1/4$ the ball slope is $c_k = \theta_k/\pi$, while the top slope is β_q , so

$$D_K(q) \geq \sum_{k=1}^B \left(\frac{\pi}{2\theta_k} - 1 \right) + \frac{(v_q - B)_+^2}{\beta_q}.$$

Insert this in (63); the $-B$ from the sum cancels the $+B$ in $\lambda_q = B - Q$. The other entry in the maximum is the completed one-interface theorem. $\square$

Corollary 7.5 (Zero-level terminal triangle). *Suppose $q \leq \mu < q + 1$ and $B = [G_K(q) + 1/4]_< = 0$. Put*

$$v = G_K(\mu), \quad c = \frac{\arccos \rho}{\pi}.$$

Then the shifted shell tail beginning at q has zero discrete count and

$$D_A(q) = 2 \int_q^K A(z) dz \geq 2 \int_\mu^K G_K(z) dz \geq \frac{v^2}{c}. \quad (72)$$

Consequently the scalar (66) is nonnegative whenever

$$\frac{v^2}{c} \geq \frac{p - d_\rho(n - p)}{2}. \quad (73)$$

Proof. The condition $B = 0$ is $G_K(q) \leq 3/4$. Since $A \leq G_K$ and both actions decrease, every strict bracket in the shell tail is zero, including at the equality wall. On $[\mu, K]$ one has $A = G_K$. Moreover G_K is convex there and lies above its tangent at μ ,

$$G_K(z) \geq v - c(z - \mu).$$

The tangent reaches zero no later than K , because $G_K(K) = 0$; integrating its positive triangle gives $2 \int_\mu^K G_K \geq v^2/c$. Finally use $R_p \geq -p/2$ in (58). $\square$

7.1 A certified first-floor large ray

We record a computer-assisted result for the ordinary first-floor branch $F_0 = 1$, $p < n$. Put

$$x = r + p, \quad y = x + 1, \quad s = n - p - 1, \quad q = x + s + 1,$$

so that $q \leq \mu < q + 1$. At the first-shelf wall $A(x) = 3/4$, set

$$v = A(y), \quad \lambda = \frac{K - \mu}{\pi\mu}, \quad c = \frac{1}{\pi} \arccos \frac{\mu}{K}, \quad S_z = \sqrt{\mu^2 - z^2},$$

and define the terminal envelope

$$\mathcal{E}(y, v) = 2 \int_y^\mu [v - \lambda(S_y - S_z)]_+ dz + \frac{(v - \lambda S_y)_+^2}{c}. \quad (74)$$

Before specializing to the wall, concavity on the first shelf and its first drop cell proves (66) whenever

$$p(A(r) + A(x)) + A(x) + A(y) + \mathcal{E}(y, A(y)) - (1 + 2p) \geq 0.$$

At $A(x) = 3/4$ this becomes the weakened wall certificate

$$\mathcal{W} := p \left(A(r) - \frac{5}{4} \right) + v - \frac{1}{4} + \mathcal{E}(y, v) \quad (75)$$

For fixed (μ, r, p) , every term before subtracting the constant in the unspecialized certificate is strictly increasing in K . Indeed, the sampled actions increase, while v/λ and λ increase in the integral in (74); on the positive outer branch, $(v - \lambda S_y)/\arccos(\mu/K)$ also increases. Thus the minimum in a fixed first-floor chamber is its post-crossing value at the unique wall $A(x) = 3/4$.

Theorem 7.6 (First-floor large-ray certificate). *On the implicit wall $A(x) = 3/4$, one has*

$$\boxed{\mathcal{W} > 0} \quad \text{if} \quad r \geq \frac{3}{2}, \quad s \in \{0, 1\}, \quad x \geq 100. \quad (76)$$

The assertion holds on the continuous relaxation of r, p and $\alpha = \mu - q \in [0, 1]$, and therefore on the required half-integral and integral chamber parameters.

Proof. Write

$$\delta = K - \mu, \quad t = x^{-1/3}, \quad \kappa = t\delta, \quad P = tp, \quad \beta = \mu - x \in [s + 1, s + 2].$$

Changing variables $R = x + u$ in the wall equation gives the exact identity

$$\frac{3\pi}{4} = \int_{\beta}^{\beta+\delta} \frac{\sqrt{u(2x+u)}}{x+u} du. \quad (77)$$

The elementary bounds

$$\sqrt{\frac{u}{2x}} \leq \frac{\sqrt{u(2x+u)}}{x+u} \leq \sqrt{\frac{2u}{x}}$$

on the wall interval, with a preliminary contradiction if that interval reaches $u = x$, give

$$\frac{22}{25} < \kappa < \frac{731}{250}. \quad (78)$$

Moreover, a possible failure must have $0 < A(r) - 3/4 < 1/2$. Comparing the radius derivatives at r and x then gives

$$A(r) - \frac{3}{4} \geq \frac{(22/25)P}{2\pi(583/500)\sqrt{(1083/500)(3571/1000 + P)}}.$$

At $P = 41$ the right side is strictly larger than $1/2$; hence every possible failure lies in $0 \leq P < 41$.

It remains to certify a compact four-variable box while retaining the implicit wall. For $0 \leq \xi \leq 1$ put $w_{\xi} = \beta t + \xi \kappa$ and define

$$\begin{aligned} \Phi_x(\xi) &= \frac{\sqrt{w_{\xi}(2 + t^2 w_{\xi})}}{1 + t^2 w_{\xi}}, \\ \Phi_r(\xi) &= \frac{\sqrt{(w_{\xi} + P)(2 + t^2(w_{\xi} - P))}}{1 + t^2 w_{\xi}}, \\ \Phi_y(\xi) &= \frac{\sqrt{((\beta - 1)t + \xi \kappa)(2 + t^2((\beta + 1)t + \xi \kappa))}}{1 + t^2(\beta t + \xi \kappa)}. \end{aligned}$$

At $R = \mu + \xi\delta$ these are the exact scaled radius profiles for x, r, y . For a profile F , put

$$T_2[F] = \frac{F(0) + 2F(1/2) + F(1)}{4}, \quad M_2[F] = \frac{F(1/4) + F(3/4)}{2}.$$

The radius profile is increasing and strictly concave, so two-panel trapezoid and midpoint quadrature enclose every wall root by

$$\frac{\kappa}{\pi} T_2[\Phi_x] \leq \frac{3}{4} \leq \frac{\kappa}{\pi} M_2[\Phi_x]. \quad (79)$$

The same lower quadrature gives

$$\underline{A}_r = \frac{\kappa}{\pi} T_2[\Phi_r], \quad \underline{v} = \frac{\kappa}{\pi} T_2[\Phi_y].$$

Put $L = \beta - 1$ and

$$H = \frac{3}{4} - \frac{\kappa\sqrt{t}\sqrt{\beta(2+\beta t^3)}}{\pi(1+\beta t^3)}, \quad \vartheta = \frac{\kappa}{1+\beta t^3}, \quad \mathcal{Q} = \frac{\sqrt{\vartheta(2+t^2\vartheta)}}{\pi}.$$

The anchored slope envelope gives $v - \lambda S_y \geq H$. When $H > 0$, (74) and $c/t \leq \mathcal{Q}$ therefore yield the stable lower certificate

$$t\mathcal{W} \geq \mathcal{C}_* := P\left(\underline{A}_r - \frac{5}{4}\right) + t\left(\underline{v} - \frac{1}{4} + 2LH\right) + \frac{H^2}{\mathcal{Q}}. \quad (80)$$

This expression has a finite continuous limit at $t = 0$.

The directed-rounding subdivision covers

$$0 \leq t \leq \frac{53861}{250000}, \quad s+1 \leq \beta \leq s+2, \quad 0 \leq P \leq 41,$$

encloses every compatible κ through (79) and (78), and retains the exact domain condition

$$1 - Pt^2 - \frac{3}{2}t^3 \geq 0$$

equivalent to $r \geq 3/2$. Each terminal box is proved either outside this domain, automatic from $\underline{A}_r \geq 5/4$, or strictly positive from $H > 0$ and (80). At 384-bit Arb precision the two runs process, respectively,

$$17697 \quad (s=0), \quad 25367 \quad (s=1)$$

boxes, to maximum depths 21 and 22. The identical certificate passes at 512 bits, and a 768-bit replay with tighter wall-root bisection also passes. The verifier is `scripts/general_d_first_floor_ray_verify.py`. All acceptance tests use directed interval signs; floating point only orders the subdivision queue. This proves (76). $\square$

The stronger certificate needed on the compact part of the wall keeps the actual interface geometry instead of the relaxed terminal envelope. Put

$$q = x + s + 1, \quad \alpha = \mu - q \in [0, 1],$$

and, on $A(x) = 3/4$, write

$$a_r = A(r), \quad u = A(q), \quad h = G_K(\mu), \quad c = \frac{1}{\pi} \arccos \frac{\mu}{K}.$$

Concavity of A on $[r, \mu]$ and the outer tangent triangle give

$$\boxed{\mathcal{C}_{\text{ff}} := p \left(a_r + \frac{3}{4} \right) + (s+1) \left(\frac{3}{4} + u \right) + \alpha(u+h) + \frac{h^2}{c} - (1+2p) \leq D_A(r)}. \quad (81)$$

The right-hand side is interpreted as the post-crossing value. At the wall itself the strict bracket can only lower the count.

Theorem 7.7 (Finite first-floor wall certificate). *On the implicit wall $A(x) = 3/4$,*

$$\boxed{\mathcal{C}_{\text{ff}} > 0} \quad \text{if} \quad r \geq \frac{3}{2}, \quad s \in \{0, 1\}, \quad x < 100. \quad (82)$$

Consequently Theorem 7.6 and (82) close the complete sector $r \geq 3/2$, $s \in \{0, 1\}$.

Proof. For fixed x , the wall has the radius-average form

$$\int_{\mu}^{K(\mu)} \sqrt{1 - \frac{x^2}{R^2}} dR = \frac{3\pi}{4},$$

and hence

$$K'(\mu) = \frac{\sqrt{1 - x^2/\mu^2}}{\sqrt{1 - x^2/K^2}} \in (0, 1).$$

Along this wall $A(r)$ is nonincreasing, $A(q)$ is increasing, and both h and c are decreasing. These signs follow by differentiating the radius average; for example

$$\frac{d}{d\mu} A(z) = \frac{\sqrt{1 - x^2/\mu^2}}{\pi} \left(\frac{\sqrt{1 - z^2/K^2}}{\sqrt{1 - x^2/K^2}} - \frac{\sqrt{1 - z^2/\mu^2}}{\sqrt{1 - x^2/\mu^2}} \right),$$

and the squared ratio in parentheses has derivative with the sign of $z^2 - x^2$. Also $K' < 1$ makes μ/K increase, while direct differentiation gives $h' < 0$.

Thus, on an exact rational panel $\alpha_0 \leq \alpha \leq \alpha_1$, strict rational wall brackets $K_i^- < K(q + \alpha_i) < K_i^+$ give a one-sided lower bound for every term in (81): evaluate a_r, h at the upper endpoint with K_1^- , evaluate u at the lower endpoint with K_0^- , and bound c from above at the lower endpoint with K_0^+ . The wall roots are isolated by exact rational bisection, and all transcendental signs are then decided with directed-rounding Arb arithmetic.

The exact enumeration is

$$r = \frac{m}{2}, \quad 3 \leq m \leq 199, \quad p \geq 0, \quad r + p < 100, \quad s \in \{0, 1\}.$$

It contains 9801 chambers for each s , hence 19602 in total. At both 384 and 512 bits the verifier accepts 19606 panels after processing 19610 panels and isolating 39208 endpoint roots. Only $(r, p, s) = (3/2, 4, 0), \dots, (3/2, 7, 0)$ require one bisection in α ; no deeper split occurs. A 768-bit replay with 160 root bisections per endpoint gives the identical counts. The checked script is `scripts/general_d_first_floor_finite_verify.py`. Since every leaf has a strictly positive Arb lower endpoint, this proves (82). $\square$

Theorem 7.8 (Complete small-start certificate). *On the implicit wall $A(x) = 3/4$, one has*

$$\boxed{\mathcal{C}_{\text{ff}} > 0} \quad \text{if} \quad r \in \left\{ \frac{1}{2}, 1 \right\}, \quad s \in \{0, 1\}, \quad p \geq 0. \quad (83)$$

Together with Theorems 7.6 and 7.7, this closes the whole ordinary first-floor sector $s \in \{0, 1\}$.

Proof. For $x < 100$, the correlated wall-panel construction in the proof of Theorem 7.7 applies without change. The exact enumeration

$$r \in \left\{ \frac{1}{2}, 1 \right\}, \quad s \in \{0, 1\}, \quad p \geq 0, \quad r + p < 100$$

contains 398 chambers. Directed-rounding replays at 384 and 512 bits process 420 panels, accept 409 leaves, and isolate 807 endpoint roots, with maximum depth one. A 1024-bit replay with 200 root steps has identical counts. The verifier is `scripts/general_d_first_floor_small_starts_verify.py`.

It remains to treat $x \geq 100$. Put $\beta = \mu - x = s + 1 + \alpha \in [1, 3]$ and $\delta = K - \mu$. The wall equation is

$$\frac{3\pi}{4} = \int_{\beta}^{\beta+\delta} \frac{\sqrt{v(2x+v)}}{x+v} dv. \quad (84)$$

It first implies $\delta < x - \beta$: otherwise the integration interval contains $[x/2, x]$, where the integrand is at least $\sqrt{5}/3$, already a contradiction at $x = 100$. Hence

$$\delta \leq \left(\frac{9\pi\sqrt{2}}{8} \right)^{2/3} x^{1/3} < \frac{731}{250} x^{1/3}.$$

With $T_* = 53861/250000 > 100^{-1/3}$ this gives, exactly,

$$\frac{K}{x} < 1 + 3T_*^3 + \frac{731}{250} T_*^2 < \frac{6}{5}.$$

For $f_z(R) = \sqrt{1 - z^2/R^2}$, the ratio f_r/f_x decreases with R . Since $r \leq 1$, $x \geq 100$, and $K < 6x/5$,

$$\left(\frac{f_r(R)}{f_x(R)} \right)^2 \geq \frac{K^2 - r^2}{K^2 - x^2} > \frac{36x^2 - 25}{11x^2} \geq \frac{14399}{4400} > \frac{25}{9}.$$

Integration over $R \in [\mu, K]$ therefore yields

$$A(r) > \frac{5}{3} A(x) = \frac{5}{4}.$$

Finally, if $v = A(x + 1)$, the one-cell action loss is smaller than $\arccos(\mu/K)/\pi < 1/2$, so $v > 1/4$. The weaker wall certificate

$$p \left(A(r) - \frac{5}{4} \right) + v - \frac{1}{4} + \mathcal{E}$$

is thus strictly positive, since $\mathcal{E} \geq 0$. This proves the analytic ray and completes (83). $\square$

The last possible first-floor escape requires a separate limiting estimate, because no complete terminal level is present below the first action wall.

Lemma 7.9 (Uniform critical first-floor gap). *Let*

$$H_{\kappa}(X) = \frac{c_0}{\sqrt{\kappa}} ((\kappa + X)^{3/2} - X^{3/2}), \quad w = H_{\kappa}(0) = c_0 \kappa,$$

where $c_0 = 2\sqrt{2}/(3\pi)$. For $0 < w < 3/4$, let $H_{\kappa}(M) = 3/4$ and $H_{\kappa}(N) = 1$, and set

$$\mathcal{L}(w) = -2 \int_M^N (1 - H_{\kappa}(X)) dX + \frac{M}{2}, \quad \mathcal{Z}(w) = \frac{\pi}{\sqrt{2}} w^2.$$

Then

$$\boxed{\mathcal{L}(w) + \mathcal{Z}(w) > \frac{\pi}{8\sqrt{2}}.} \quad (85)$$

At $w = 3/4$ the same conclusion holds with the smaller wall-safe terminal value $\pi/(2\sqrt{2})$.

Proof. The function $1 - H_\kappa$ is convex between M and N , so its endpoint chord gives $\mathcal{L} \geq (3M - N)/4$. Write

$$X = \kappa u(t), \quad u(t) = \frac{(1 - t^2)^2}{4t^2}, \quad \frac{H_\kappa(X)}{w} = \frac{3/t + t^3}{4}.$$

If ξ, η are the parameters at M, N , then

$$w = \frac{3\xi}{3 + \xi^4} = \frac{4\eta}{3 + \eta^4}.$$

Using that u decreases, the ranges $0 < w \leq 1/2$, $1/2 \leq w \leq 2/3$, and $2/3 \leq w < 3/4$ give, respectively,

$$\frac{\sqrt{2}}{\pi}(\mathcal{L} + \mathcal{Z}) \geq \frac{1}{8} + \frac{85469}{51518208}, \quad \frac{1}{8} + \frac{41923}{1281424}, \quad \frac{1}{8} + \frac{743}{4608}.$$

Here one uses $\xi \leq (259/243)w$ in the first range, $\xi \leq (283/256)w$ in the second, and $\eta \geq 1/2$ in the third. At the wall the one-sided strict convention can lower the normalized terminal payment by only $1/16$, leaving $1/8 + 455/4608$. This proves (85). $\square$

Theorem 7.10 (Global exhaustion of the ordinary first floor). *In the universal active region $K > \pi/(1 - \rho)$, every negative ordinary first-floor first-drop scalar*

$$F_0 = 1, \quad p < n, \quad D_A(q) + R_p + \frac{d_\rho}{2}(n - p) < 0$$

belongs to a relatively compact subset of the exact shell parameter space. Consequently the possible discrete labels (r, n, p, Q, B) are bounded. The case $p = 0$ is automatic, and the conclusion includes the whole cone $s = n - p - 1 \geq 2$.

Proof. Put $\tau = \sqrt{1 - \rho}$, $a = K - \mu$, and $\kappa = \tau a$. The fixed-ratio, compact-optical, and thin high-energy results already proved above show that an escaping negative sequence must have

$$\tau \rightarrow 0, \quad a \rightarrow \infty, \quad \tau\kappa = (1 - \rho)a < \frac{125}{8},$$

while the first-drop interface bound gives $c_0\kappa < 3/4$. The exact noncancelling turning profile is

$$\mathcal{H}_{\tau, \kappa}(X) = \frac{1}{\pi} \int_0^\kappa \frac{\sqrt{(\kappa + X - u)[2\kappa - (\kappa + X + u)\tau^2]}}{\kappa - u\tau^2} du. \quad (86)$$

On the complete possible negative head its ratio to $H_\kappa(X)$ tends to one uniformly.

After passage to a subsequence, $\kappa \rightarrow \kappa_0 \in [0, 3/(4c_0)]$. If $\kappa_0 > 0$, exact level convergence, the first-shelf reduction, and the wall-safe terminal estimate give

$$\liminf \tau \left(D_A(q) + R_p + \frac{d_\rho}{2}(n - p) \right) \geq \mathcal{Z}(c_0\kappa_0) + \mathcal{L}(c_0\kappa_0) > 0$$

by Lemma 7.9.

If $\kappa_0 = 0$, put $Y = \kappa X$. Uniformly on the physical level window,

$$\mathcal{H}_{\tau,\kappa}(Y/\kappa) \longrightarrow H_0(Y) := \frac{\sqrt{2}}{\pi} \sqrt{Y}.$$

The $3/4$ - and 1 -levels converge to $Y_{3/4} = 9\pi^2/32$ and $Y_1 = \pi^2/2$. The exact rescaled prefix then satisfies

$$\begin{aligned} \liminf \kappa\tau \left(R_p + \frac{d_p}{2}(n-p) \right) &\geq -2 \int_{Y_{3/4}}^{Y_1} (1 - H_0(Y)) dY + \frac{Y_{3/4}}{2} \\ &= \frac{17\pi^2}{192} > 0. \end{aligned}$$

The nonnegative one-interface reserve is enough on this ray. Both possible limits contradict negativity, so no negative sequence escapes. The two shift grids have $r \geq 1/2$, and $r < \rho K$; the fixed-ratio bound therefore also prevents an escape through $\rho \downarrow 0$. Relative compactness follows. It bounds the discrete labels, although it does not assert finiteness of the connected components of the remaining real-analytic wall set. $\square$

Remark 7.11 (Remaining first-floor wall sectors). Theorems 7.6 and 7.7 close $r \geq 3/2$, $s \in \{0, 1\}$, while Theorem 7.8 closes the remaining two starts. Thus the whole sector $s \in \{0, 1\}$ is proved. Theorem 7.10 removes every noncompact part of the remaining cone. Only the compact residue with $s \geq 2$ is not yet certified.

Proposition 7.12 (Exact no-drop identities). *Suppose $p = n \geq 1$, and write $F_0 = \dots = F_n = f$. Put*

$$\varepsilon_q = A(q) + \frac{1}{4} - f \in [0, 1), \quad \chi_q = \mathbf{1}_{\{A(q)+1/4=f\}}.$$

Then

$$D_A(r) = D_A(q) + R_n + \chi_q, \tag{87}$$

$$R_n = -\frac{n}{2} + 2n\varepsilon_q + 2 \int_0^n u \sigma(r+u) du. \tag{88}$$

Thus the no-drop branch is automatic when $f = 0$, or $\varepsilon_q \geq 1/4$, or

$$2 \int_0^n u \sigma(r+u) du \geq n \left(\frac{1}{2} - 2\varepsilon_q \right).$$

Away from the endpoint wall, the exact shifted-tail condition is the quantitative payment

$$D_A(q) \geq \frac{n}{2} - 2n\varepsilon_q - 2 \int_0^n u \sigma(r+u) du > 0. \tag{89}$$

At the endpoint wall the right side is reduced by the favorable unit $\chi_q = 1$ in (87); proving the wall-safe scalar (66) directly asks for the stronger displayed payment.

Proof. Strict decrease shows that a lower ordinary-floor wall cannot occur at an inner sample before q without ending the shelf. Hence the strict brackets at $r, \dots, q$ equal f , except that the bracket at q is $f - 1$ when $\chi_q = 1$. Therefore

$$T_A(r) - T_A(q) = 2nf - \chi_q,$$

which proves (87). Since $f = A(q) + 1/4 - \varepsilon_q$,

$$\begin{aligned} R_n &= 2 \int_0^n (A(r+t) - A(q)) dt + 2n \left(\varepsilon_q - \frac{1}{4} \right) \\ &= 2 \int_0^n \int_t^n \sigma(r+u) du dt + 2n \left(\varepsilon_q - \frac{1}{4} \right), \end{aligned}$$

and reversing the order proves (88). The remaining statements follow from $D_A(q) \geq 0$. $\square$

7.2 High floors: an automatic thin sector and the critical ray

Assume that the first ordinary floor is $F_0 = f \geq 2$, and write

$$\varepsilon = 1 - \rho, \quad a = \varepsilon K = K - \mu, \quad h = f - \frac{1}{4}, \quad m = n - p. \quad (90)$$

Since $A(0) = a/\pi > A(r) \geq h$, every active high-floor configuration satisfies

$$a > \pi h. \quad (91)$$

The reduced scalar is

$$\mathcal{S} = D_A(q) + R_p + \frac{d_\rho}{2}m.$$

For $b > 0$, put $B_b(y) = \pi^{-1} \sqrt{b^2 - y^2}$. The radius-average formula for the shell action gives, for $0 \leq y \leq a\rho$,

$$\frac{1}{2}B_a(y) + \frac{1}{2\rho}B_{a\rho}(y) \leq \mathcal{A}_{\varepsilon,a}(y) := A_{1-\varepsilon,a/\varepsilon}(y/\varepsilon) \leq B_a(y). \quad (92)$$

Lemma 7.13 (Quantitative product-shelf margin). *Let $f \geq 3$, $h = f - 1/4$, $a > \pi h$, and $y_b = \sqrt{a^2 - \pi^2 h^2}$. If $0 \leq y_0 \leq s \leq y_b$, then*

$$\boxed{2 \int_{y_0}^s B_a(y) dy - 2f(s - y_0) + \frac{a-s}{2} \geq \frac{f^2}{4a}.} \quad (93)$$

Proof. For fixed y_0 the left side is concave in s , so it is enough to use $s = y_0$ and $s = y_b$. The first value is at least $(a - y_b)/2 \geq \pi^2 h^2/(4a) > f^2/(4a)$. At $s = y_b$, put $R = a/\pi$. Differentiation with respect to y_0 gives $2(f - B_a(y_0))$; hence the minimum occurs where $B_a(y_0) = f$ if $R \geq f$, and at $y_0 = 0$ if $h \leq R < f$.

In the first case set $u = \sqrt{R^2 - h^2}$ and $v = \sqrt{R^2 - f^2}$. The chord estimate gives

$$\frac{\mathcal{L}}{\pi} > \frac{2R - 3u + v}{4} \geq \frac{3h^2/2 - f^2}{4R} > \frac{f^2}{4\pi^2 R}.$$

The last comparison follows from $\pi^2 > 9$ and $112f^2 - 216f + 27 > 0$. In the second case write $R = h + t$, $0 \leq t < 1/4$. Rationalizing the same chord estimate gives

$$\frac{\mathcal{L}}{\pi} \geq \frac{P(h, t)}{6R}, \quad P(h, t) = (h + t)^2 - 4(1 - t)^2 t (2h + t) \geq h^2 - 2h - \frac{1}{4}.$$

Here $18(h^2 - 2h - 1/4) - 3f^2 = 15f^2 - 45f + 45/8 > 0$ for $f \geq 3$. Again $\mathcal{L}/\pi > f^2/(4\pi^2 R)$, which is (93). $\square$

Proposition 7.14 (An automatic finite-shell sector). *In the high-floor notation above, suppose*

$$f \geq 3, \quad \varepsilon a^4 \leq \frac{f^3}{144}. \quad (94)$$

Then

$$\boxed{\varepsilon \left(R_p + \frac{d_\rho}{2} m \right) \geq \frac{f^2}{8a} > 0.} \quad (95)$$

In particular, (66) holds without using a terminal reserve.

Proof. Put $y_0 = \varepsilon r$, $s = \varepsilon(r + p)$, and $y_q = \varepsilon q$. Since $A(r + p) \geq h$ and the upper half of (92) holds, $s \leq y_b$. Conditions (91)–(94) imply $\rho > 99/100$ and $y_b \leq a\rho$: indeed

$$1 - \rho^2 < 2\varepsilon \leq \frac{f^3}{72a^4} < \frac{\pi^2 h^2}{a^2}.$$

Thus the lower half of (92) applies on the whole shelf. With

$$E = \int_0^{y_b} (B_a - \rho^{-1} B_{a\rho}) \, dy,$$

direct rationalization, followed by $h \geq 11f/12$, gives

$$E \leq \frac{a^3(\rho^{-2} - 1)}{3\pi^2 h} < \frac{\varepsilon a^3}{11f} \leq \frac{f^2}{1584a}. \quad (96)$$

Moreover $y_q \geq a\rho - \varepsilon$, and hence

$$d_\rho(y_q - s) - (a - s) \geq -(1 - d_\rho)a - \varepsilon(a + 1). \quad (97)$$

Writing $\rho = \cos \theta$ gives $1 - d_\rho = 2\theta/\pi \leq \sqrt{2\varepsilon}$; condition (94) then yields

$$\frac{(1 - d_\rho)a}{2} \leq \frac{f^2}{24a}, \quad \frac{\varepsilon(a + 1)}{2} \leq \frac{f^2}{24a}.$$

Finally, the exact scaled identity is

$$\varepsilon \left(R_p + \frac{d_\rho}{2} m \right) = 2 \int_{y_0}^s \mathcal{A}_{\varepsilon,a}(y) \, dy - 2f(s - y_0) + \frac{d_\rho}{2}(y_q - s).$$

Combining the last four displays with Lemma 7.13 gives

$$\varepsilon \left(R_p + \frac{d_\rho}{2} m \right) \geq \frac{f^2}{4a} - \frac{f^2}{1584a} - \frac{f^2}{12a} > \frac{f^2}{8a}.$$

Ordinary shelf walls and strict terminal walls only improve these bounds. $\square$

The complementary unbounded scaling is the grazing ray

$$\varepsilon \downarrow 0, \quad K\varepsilon^{3/2} \longrightarrow \kappa \in (0, \infty), \quad n\sqrt{\varepsilon} = O(1). \quad (98)$$

Set $c_0 = 2\sqrt{2}/(3\pi)$. At the coordinate $X = (\mu - z)\sqrt{\varepsilon}$, the ball formula gives, locally uniformly,

$$A(z) \longrightarrow H_\kappa(X) := \frac{c_0}{\sqrt{\kappa}} \left((\kappa + X)^{3/2} - X^{3/2} \right), \quad H_\kappa(0) = w := c_0\kappa. \quad (99)$$

If $B = [w + 1/4]_<$, the exact-angle reserve gives the wall-safe limit

$$\liminf_{\varepsilon \downarrow 0} \sqrt{\varepsilon} D_A(q) \geq \mathcal{T}(w) := \frac{\pi}{2\sqrt{2}} w^{1/3} \sum_{k=1}^B (k - \tfrac{1}{4})^{-1/3}. \quad (100)$$

Indeed, at the level $\tau_k = k - 1/4$,

$$\sqrt{\varepsilon} \frac{\pi}{2\theta_k} \longrightarrow \frac{\pi}{2\sqrt{2}} w^{1/3} \tau_k^{-1/3},$$

whereas the finite Q -term and the interface cap vanish after multiplication by $\sqrt{\varepsilon}$. The fractional-top term in (71) is nonnegative and is not needed below. At a limiting action wall, the strict value of B is the smaller one-sided complete-level sum and is therefore the safe convention.

Theorem 7.15 (No negative leading obstruction on the critical ray). *For every fixed limiting first-floor level $f \geq 2$, each no-drop or first-drop configuration on (98) has a nonnegative leading scalar after (100) is included. If its limiting shelf/head prefix is negative, the terminal reserve exceeds the magnitude of that prefix strictly.*

Proof. Write $X = \kappa u$ and $t = \sqrt{1+u} - \sqrt{u} \in (0, 1]$. Direct algebra gives

$$\frac{H_\kappa(X)}{w} = F(t) := \frac{3/t + t^3}{4}, \quad u = \frac{(1-t^2)^2}{4t^2}, \quad H'_\kappa(X) = \frac{3c_0}{2}t. \quad (101)$$

Let M be the point $H_\kappa(M) = h = f - 1/4$ and $N_f \geq M$ the point $H_\kappa(N_f) = f$. Scaling the first-shelf reduction and minimizing over its starting point shows that the worst first-drop prefix is

$$\mathcal{L} = -2 \int_M^{N_f} (f - H_\kappa(X)) dX + \frac{M}{2}. \quad (102)$$

Indeed, the first shelf gives the displayed integral, the remaining post-drop length gives $M/2$ because $d_\rho \rightarrow 1$, and moving the starting point beyond the f -level adds only nonnegative shelf area. Since $0 \leq f - H_\kappa \leq 1/4$ on this interval, $\mathcal{L} < 0$ forces $N_f > 2M$.

This last inequality forces

$$w > \frac{3}{4}, \quad t_f > \frac{1}{4}, \quad (103)$$

where t_f is the parameter at N_f . To see this without an asymptotic estimate, use $3/(4t) \leq F(t) \leq 1/t$ and $h/f \geq 7/8$. If $w \leq 3/4$, then $t_f < 3/10$ and

$$\frac{u_f}{u_h} \leq \left(\frac{7}{6}\right)^2 \left(\frac{400}{351}\right)^2 < 2,$$

contrary to $N_f > 2M$. If $t_f \leq 1/4$, the same computation gives

$$\frac{u_f}{u_h} < \left(\frac{6}{5}\right)^2 \left(\frac{100}{91}\right)^2 < 2.$$

Thus (103) holds. Equations (101)–(102) now give

$$-\mathcal{L} < \frac{N_f - M}{2} < \frac{1}{3c_0} = \frac{\pi}{2\sqrt{2}}.$$

On the other hand, $B \geq 1$ and the first term of (100) gives

$$\mathcal{T}(w) \geq \frac{\pi}{2\sqrt{2}} w^{1/3} (3/4)^{-1/3} > \frac{\pi}{2\sqrt{2}}.$$

This strictly pays a negative first-drop prefix. In the no-drop branch $M = 0$ and $w \geq h$. If $w \geq f$ its prefix is already nonnegative; otherwise $F(t_f) = f/w \leq f/h \leq 8/7$, so $t_f > 1/4$ and the same comparison applies. $\square$

Corollary 7.16 (Uniform critical no-drop gap for $f = 2, 3$). *For a no-drop sequence on (98) with fixed $f \in \{2, 3\}$,*

$$\liminf_{\varepsilon \downarrow 0} \sqrt{\varepsilon} \mathcal{S} \geq \gamma_0 := \frac{\pi}{2\sqrt{2}} \left[\left(\frac{7}{3} \right)^{1/3} - 1 \right] > \frac{1}{3}. \quad (104)$$

Proof. No-drop gives $w \geq f - 1/4 \geq 7/4$. If the prefix is negative, Theorem 7.15 bounds its magnitude by $\pi/(2\sqrt{2})$, while the first complete terminal level is at least $\pi(7/3)^{1/3}/(2\sqrt{2})$. If the prefix is nonnegative the estimate is stronger. Finally, $\pi/(2\sqrt{2}) > 15/14$ and $(7/3)^{1/3} > 59/45$. $\square$

Theorem 7.17 (Global exhaustion of high-floor first drops). *The set of negative high-floor first-drop configurations*

$$F_0 = f \geq 2, \quad p < n, \quad D_A(q) + R_p + \frac{d_\rho}{2}(n - p) < 0$$

is relatively compact in the exact shell parameter space. In particular, the possible discrete labels (f, n, p, Q, B) are bounded, although the bounds furnished by this argument are not numerical.

Proof. Write $\epsilon = 1 - \rho$, $s = \sqrt{\epsilon}$, $a = K - \mu$, $\kappa = sa$, $m = n - p$, $x = r + p$, and $h = f - 1/4$. If $H(t) = A(\mu - t)$, the first drop gives

$$W := H(0) = G_K(\mu) < h, \quad c_0 \kappa < h.$$

Let $H(M) = h$ and let T be the first point at which $H(T) = f$, or put $T = \mu$ if that level is absent. The shelf inequalities and the bound $0 \leq f - H \leq 1/4$ on its negative part give

$$R_p + \frac{d_\rho}{2}m \geq \frac{(1 + d_\rho)M - T - d_\rho}{2}. \quad (105)$$

Thus negativity forces $T > (1 + d_\rho)M - d_\rho$.

The required uniform interface control follows from the shell-increment elasticity inequality

$$\frac{H(t_2) - W}{H(t_1) - W} \geq \sqrt{\frac{t_2(2\mu - t_2)}{t_1(2\mu - t_1)}} \quad (0 < t_1 < t_2 \leq \mu). \quad (106)$$

Indeed, for each $R \in [\mu, K]$ the quotient of

$$\sqrt{1 - \mu^2/R^2 + t(2\mu - t)/R^2} - \sqrt{1 - \mu^2/R^2}$$

by $\sqrt{t(2\mu - t)}$ is nondecreasing; integration in R proves (106). Applying it at M, T along a negative sequence with $\rho \rightarrow 1$ yields

$$f - W \leq 1 + \frac{\sqrt{3}}{2} + o(1). \quad (107)$$

As in the first-floor exhaustion, the fixed-ratio, compact-optical, and thin high-energy theorems leave only

$$s \rightarrow 0, \quad a \rightarrow \infty, \quad s\kappa < \frac{125}{8}.$$

The turning estimate $W = c_0\kappa + o(1)$ and (107) then pin

$$f - 1 - \frac{\sqrt{3}}{2} - o(1) \leq c_0\kappa \leq f - \frac{1}{4} + o(1). \quad (108)$$

They also confine the dangerous level band to a bounded turning window.

If $f \rightarrow \infty$, the root-free exact-angle reserve contains $B \geq f - O(1)$ complete levels. Since $sK^{1/3} = \kappa^{1/3}$ and $\sum_{k \leq B} (k - 1/4)^{-1/3} \gg B^{2/3}$, it gives $sD_A(q) \rightarrow +\infty$, whereas (105) and the bounded turning window give $s(R_p + d_\rho m/2) \geq -O(1)$. Hence an escaping negative sequence has bounded f .

Pass finally to a subsequence with fixed f and $\kappa \rightarrow \kappa_0 > 0$. Exact profile and level convergence give

$$\liminf s \left(R_p + \frac{d_\rho}{2} m \right) \geq \mathcal{L}_f := -2 \int_{M_0}^{N_0} (f - H_{\kappa_0}(X)) dX + \frac{M_0}{2}.$$

The wall-safe complete-level reserve has limit

$$\mathcal{T}(w) = \frac{\pi}{2\sqrt{2}} w^{1/3} \sum_{k=1}^{[w+1/4]<} (k - \frac{1}{4})^{-1/3}, \quad w = c_0\kappa_0.$$

If $\mathcal{L}_f < 0$, the exact prefix geometry forces $N_0 > 2M_0$. The t -parameterization in (101) then gives, uniformly for $f \geq 2$,

$$-\mathcal{L}_f < \frac{3\pi}{8\sqrt{2}}, \quad \mathcal{T}(w) > \frac{\pi}{2\sqrt{2}},$$

and hence

$$\mathcal{L}_f + \mathcal{T}(w) > \frac{\pi}{8\sqrt{2}}. \quad (109)$$

If $\mathcal{L}_f \geq 0$, the same comparison forces a complete terminal level, so the sum is again positive. This excludes the last escape. Relative compactness follows, and compactness bounds the discrete labels without asserting a finite number of analytic wall components. $\square$

7.3 Global finite exhaustion of the no-drop branch

We first record the root-free consequence of the terminal estimate that drives the exhaustion. For $J \geq 1$, put

$$C_0^3 = \frac{\pi}{12}, \quad c_* = \frac{4\sqrt{2}}{15}, \quad S_J = \sum_{k=1}^J (k - \frac{1}{4})^{-1/3}. \quad (110)$$

Proposition 7.18 (Root-free no-drop cutoff). *In the no-drop branch of Proposition 7.12, let*

$$Q = [A(q) + \frac{1}{4}]< \in \{f - 1, f\}, \quad \mathcal{S}_n = D_A(q) + R_n.$$

Then

$$D_A(q) \geq \max\{0, C_0 K^{1/3} S_Q - Q - c_*\}. \quad (111)$$

Consequently the no-drop scalar is nonnegative whenever

$$K \geq K_{\text{nd}}(n, Q) := \left(\frac{Q + n/2 + c_*}{C_0 S_Q} \right)^3. \quad (112)$$

Every branch below this cutoff is confined, for fixed (n, f, Q) , to a compact strip with finitely many action walls.

Proof. For $0 \leq \theta \leq \pi/2$,

$$\sin \theta - \theta \cos \theta = \int_0^\theta t \sin t \, dt \geq \frac{2\theta^3}{3\pi}.$$

Thus the angles in Corollary 7.4 satisfy

$$\frac{\pi}{2\theta_k} \geq C_0 K^{1/3} (k - \tfrac{1}{4})^{-1/3}.$$

The cap bound (30) is smaller than c_* , and the fractional-top term is nonnegative. Since the ball level count B is at least Q , this proves (111). Under (112) it gives $D_A(q) \geq n/2$, whereas (88) gives $R_n \geq -n/2$.

Finally, $K - \mu > \pi(f - 1/4)$ and $\mu \geq r + n$, so

$$K > \pi(f - \tfrac{1}{4}) + r + n.$$

Together with $K < K_{\text{nd}}(n, Q)$ this bounds r, K, μ , every terminal level, and every sample index. On each open chamber the exact scalar has

$$\partial_K \mathcal{S}_n = \frac{2}{\pi} \int_r^K \sqrt{1 - z^2/K^2} \, dz > 0, \quad \partial_\mu \mathcal{S}_n = -\frac{2}{\pi} \int_r^\mu \sqrt{1 - z^2/\mu^2} \, dz < 0.$$

It therefore reduces to the lower- K , upper- μ , and finitely many one-sided sample walls; the strict equality face is retained rather than passed through by continuity. $\square$

Proposition 7.19 (Uniform half-unit terminal reserve). *Let $q \leq \mu < q + 1$, $q \geq 3/2$, and $A(q) \geq 7/4$. Then, including at the lower ordinary-floor wall at q ,*

$$\boxed{D_A(q) > \frac{1}{2}}. \quad (113)$$

Proof. Put

$$v = G_K(q), \quad h = G_\mu(q), \quad f = \left\lfloor A(q) + \frac{1}{4} \right\rfloor \geq 2, \quad Q = [A(q) + \tfrac{1}{4}]_<, \quad B = [v + \tfrac{1}{4}]_<.$$

If $q = \mu$, the inner cap vanishes. If $q < \mu$, strict convexity of G_μ on $[q, \mu]$ and $\mu - q < 1$ give

$$2 \int_q^\mu G_\mu(z) \, dz < h(\mu - q) < h.$$

Moreover $h \leq G_{q+1}(q)$, and $G_{q+1}(q)$ decreases with q . Outward arithmetic therefore gives the uniform cap bound

$$2 \int_q^\mu G_\mu(z) \, dz < G_{5/2}(3/2) < \frac{1}{5}. \quad (114)$$

For $1 \leq k \leq B$, let θ_k be the exact ball angle at level $k - 1/4$. Corollary 7.4, with its nonnegative fractional-top term omitted, gives

$$D_A(q) \geq \frac{\pi}{2} \sum_{k=1}^B \frac{1}{\theta_k} - Q - 2 \int_q^\mu G_\mu(z) dz. \quad (115)$$

If $q \geq K/2$, every retained level crosses strictly to the right of q , so $\theta_k < \arccos(q/K) \leq \pi/3$. Away from the lower wall, $Q = f$ and $B \geq f$, which gives more than $f/2 - 1/5$. At the lower wall with $q < \mu$, one has $Q = f - 1$ but $B \geq f$; with $q = \mu$, the cap is zero and $B = Q = f - 1$, so pairing all $f - 1$ count units with their angle terms gives $D_A(q) > (f - 1)/2 \geq 1/2$.

If $q < K/2$, let K_* be the unique solution of

$$G_{K_*}(3/2) = \frac{7}{4}.$$

Monotonicity gives $K \geq K_*$. At that radius, a directed 512-bit Arb calculation proves

$$\theta_1 < \frac{503}{500}, \quad \theta_2 < \frac{11}{8},$$

and hence

$$\frac{\pi}{2} \left(\frac{1}{\theta_1} + \frac{1}{\theta_2} \right) - 2 > \frac{7}{10}, \quad \frac{\pi}{2\theta_1} - 1 > \frac{1}{2}. \quad (116)$$

The angles decrease as K increases, and every later net level is positive. Away from the lower wall, the first inequality in (116) and (114) give more than $1/2$. At the lower wall the same two cases $q < \mu$ and $q = \mu$ are handled, respectively, by the extra ball level and the second inequality in (116). The computation is reproduced by `scripts/general_d_no_drop_round5_verify.py` and passes unchanged at 1024 bits. $\square$

Corollary 7.20 (Closure of the one-cell no-drop branch). *Every no-drop branch with $f \geq 2$ and $n = 1$ satisfies*

$$\boxed{\mathcal{S}_1 = D_A(q) + R_1 > 0.} \quad (117)$$

Proof. Concavity of A on $[r, q]$ gives

$$2 \int_r^q A(z) dz \geq A(r) + A(q) \geq 2f - \frac{1}{2}.$$

Thus $R_1 \geq -1/2$, while Proposition 7.19 is strict. $\square$

Corollary 7.21 (A finite-chamber discard sector). *For a no-drop branch with $f \geq 2$, put*

$$\varepsilon = A(q) + \frac{1}{4} - f, \quad \Delta = A(r) - A(q).$$

If

$$2\varepsilon + \Delta + \frac{n\Delta}{3(2r+n)} \geq \frac{1}{2} - \frac{1}{2n}, \quad (118)$$

then $\mathcal{S}_n > 0$.

Proof. The scale-free shelf-curvature bound gives

$$R_n \geq \frac{n^2\Delta}{3(2r+n)} + n \left(2\varepsilon + \Delta - \frac{1}{2} \right) \geq -\frac{1}{2}.$$

Apply Proposition 7.19. $\square$

Theorem 7.22 (Quantitative no-drop transfer). *Let $f \in \{2, 3\}$, $b = f - 1/4$, and*

$$c_f = \frac{\sqrt{1+b^2}-1}{2}, \quad C_K(f) = \left(\frac{f+1/2+c_*}{C_0 S_{f-1}} \right)^3, \quad C_\delta(f) = \left(8\pi^2 f^2 C_K(f) \right)^{1/3}.$$

Suppose a residual no-drop candidate satisfies, with $\delta = K - \mu$,

$$\frac{c_f}{2}n \leq \delta \leq C_\delta(f)n, \quad \frac{c_f^2}{8\pi^2(C_\delta(f)+2)}n^3 < K < C_K(f)n^3. \quad (119)$$

Put

$$s = \sqrt{\delta/K}, \quad \kappa = s\delta = s^3 K, \quad N = sn.$$

If $0 < s \leq 1/10$, then, including at every endpoint wall,

$$\boxed{s\mathcal{S}_n = s(D_A(q) + R_n) > \frac{1}{10}.} \quad (120)$$

Proof. Put $\alpha = \mu - q \in [0, 1)$ and $X_q = s\alpha$. The residual endpoint floors give

$$b \leq A(q) < f \leq 3, \quad \frac{21}{8} \leq \kappa. \quad (121)$$

The last inequality follows from the radius representation and $A(q) \leq 2\kappa/\pi$.

Use the exact head coordinate $X = s(\mu - z)$ and set

$$\mathcal{H}_{s,\kappa}(X) = A(\mu - X/s), \quad H_\kappa(X) = \frac{c_0}{\sqrt{\kappa}} \left((\kappa + X)^{3/2} - X^{3/2} \right), \quad c_0 = \frac{2\sqrt{2}}{3\pi}.$$

Unlike a two-ball subtraction, the shell has the non-cancelling exact representation

$$\mathcal{H}_{s,\kappa}(X) = \frac{1}{\pi} \int_0^\kappa \frac{\sqrt{(\kappa + X - u)[2\kappa - (\kappa + X + u)s^2]}}{\kappa - us^2} du. \quad (122)$$

The ratio of its integrand to the integrand defining H_κ is

$$\mathcal{R}(u, X) = \frac{\sqrt{1 - a(u, X)s^2}}{1 - b_0(u)s^2}, \quad a(u, X) = \frac{\kappa + X + u}{2\kappa}, \quad b_0(u) = \frac{u}{\kappa}.$$

For $0 \leq X \leq 2$ at every physical point,

$$0 \leq b_0 \leq 1, \quad 0 \leq a \leq 1 + \frac{1}{\kappa} \leq \frac{29}{21}.$$

The inequalities

$$1 - y \leq \sqrt{1 - y} \leq 1, \quad \frac{1}{1 - s^2} - 1 \leq \frac{100}{99}s^2$$

therefore give $|\mathcal{R} - 1| \leq (29/21)s^2$. At $X_q < s$, the sharper bound $a < 107/105$ gives

$$H_\kappa(X_q) \leq \frac{\mathcal{H}_{s,\kappa}(X_q)}{1 - (107/105)s^2} < \frac{31}{10}.$$

Since

$$0 < H'_\kappa(X) = \frac{3c_0}{2\sqrt{\kappa}} (\sqrt{\kappa + X} - \sqrt{X}) < \frac{1}{2},$$

one has $H_\kappa < 41/10$ on $[0, 2]$. Integrating the ratio bound against the positive limiting integrand proves

$$|\mathcal{H}_{s,\kappa}(X) - H_\kappa(X)| < 6s^2 \quad (0 \leq X \leq 2). \quad (123)$$

Put

$$\zeta = \frac{s}{2} + 6s^2 \leq \frac{11}{100}, \quad w = H_\kappa(0) = c_0\kappa.$$

The endpoint estimate, (123), and $H'_\kappa < 1/2$ give

$$w > b - \zeta. \quad (124)$$

For $X = \kappa u$, set

$$t = \sqrt{1+u} - \sqrt{u}, \quad F(t) = \frac{3/t + t^3}{4}.$$

Then

$$\frac{H_\kappa(X)}{w} = F(t), \quad H'_\kappa(X) = \frac{3c_0}{2}t.$$

Let X_* be the point at which $H_\kappa(X_*) = f + \zeta$. This point is positive: otherwise (123) at X_q would contradict $A(q) < f$. Furthermore

$$\frac{f + \zeta}{w} < \frac{f + \zeta}{f - 1/4 - \zeta} \leq \frac{211}{164} < \frac{163}{125} = F(3/5).$$

Because F decreases, $t_* > 3/5$, and hence $H'_\kappa > 9c_0/10$ on $[0, X_*]$. With $d = f + \zeta - w$, this gives

$$0 < d < \frac{1}{4} + 2\zeta \leq \frac{47}{100}, \quad X_* < \frac{10d}{9c_0} < \frac{19}{10}.$$

Thus $X_q + X_* < 2$, so the error estimate applies on the whole possible negative head. Since $f + \zeta - H_\kappa$ is convex with endpoint values d and 0, its integral is at most $dX_*/2$. The exact increasing profile and the no-drop head identity therefore give

$$sR_n \geq -2 \int_0^{X_*} (f + \zeta - H_\kappa(X)) dX \geq -\frac{10d^2}{9c_0}. \quad (125)$$

The first complete terminal level $\tau = 3/4$ is present because $G_K(q) \geq A(q) \geq 7/4$, including at the strict wall. Its angle lies in $(0, \pi/2)$. If

$$x_\tau = \left(\frac{3\pi\tau}{K} \right)^{1/3},$$

then $x_\tau^3 = 3(\sin \theta - \theta \cos \theta)$. The chord bound for sine and (121) imply $\theta < \sqrt{3}s$, while

$$\sin \theta - \theta \cos \theta \geq \frac{\theta^3}{3} - \frac{\theta^5}{30}$$

gives $x_\tau/\theta > 1 - 3s^2/10$. The exact-angle reserve, the cap bound, and $Q \leq 3$ now give

$$sD_A(q) \geq \frac{\pi}{2\sqrt{2}} \left(\frac{b - \zeta}{3/4} \right)^{1/3} \left(1 - \frac{3s^2}{10} \right) - \frac{17}{5}s. \quad (126)$$

Here all further exact-angle and fractional-top terms were discarded in the favorable direction; the count and cap cost is less than $(3 + 2/5)s = 17s/5$.

Finally, all constants close rationally:

$$\frac{\pi}{2\sqrt{2}} > 1, \quad c_0 > \frac{49}{165}, \quad \frac{b-\zeta}{3/4} \geq \frac{164}{75} > \left(\frac{129}{100}\right)^3.$$

Adding (125) and (126) yields

$$s\mathcal{S}_n > \frac{129}{100} \frac{997}{1000} - \frac{17}{50} - \frac{10}{9} \frac{165}{49} \left(\frac{47}{100}\right)^2 = \frac{1758611}{14700000} > \frac{1}{10}.$$

At the lower ordinary-floor wall the separate identity (87) supplies one additional favorable unit for the original defect. The outward-rounded constants used above are replayed by `scripts/general_d_no_drop_round5_verify.py`; the certificate passes at both 512 and 1024 bits. $\square$

Theorem 7.23 (Global finite exhaustion of no-drop data). *Suppose $f \geq 2$, $n \geq 1$, and $\mathcal{S}_n < 0$. Then Proposition 7.18 must fail, and the following properties hold:*

1. *if $r \leq K/2$, necessarily*

$$2 \leq f \leq 49, \quad 2 \leq n \leq 48; \quad (127)$$

2. *if $r > K/2$ and $f \geq 4$, necessarily*

$$4 \leq f \leq 8, \quad 2 \leq n \leq 39; \quad (128)$$

3. *in the two exceptional rows $f = 2, 3$, necessarily $2 \leq n \leq 379$. Before the quantitative transfer, an unbounded sequence could occur only in these rows, and then*

$$K - \mu \asymp n, \quad K \asymp n^3, \quad r/K \longrightarrow 1, \quad \rho \longrightarrow 1. \quad (129)$$

Theorem 7.22 gives the explicit conclusion $n < 380$ in both rows. Thus the residual triples (n, f, Q) form a completely explicit finite set.

The certified central list has 647 necessary pairs after the two $n = 1$ pairs are removed; the outer $f \geq 4$ list has 60 necessary pairs and is unchanged. These counts precede certification of the continuous action chambers. This is a finite reduction, not a proof of the residual finite wall chambers.

Proof. Put

$$\delta = K - \mu, \quad \alpha = \mu - q \in [0, 1), \quad b = f - \frac{1}{4}, \quad A(q) = b + \xi, \quad A(r) = b + \xi_0, \quad \Delta = \xi_0 - \xi.$$

Only $R_n < 0$ can remain. The exact shelf-curvature identity then gives

$$\xi_0 + \xi < \frac{1}{2}, \quad 0 \leq \xi < \frac{1}{4}, \quad 0 \leq \Delta < \frac{1}{2}, \quad A(q) < f, \quad A(r) < f + \frac{1}{4}. \quad (130)$$

In addition to $K < K_{\text{nd}}(n, Q)$, three elementary action estimates are needed. The radius derivative and $K - q = \delta + \alpha < \delta + 1$ give

$$K \leq U_f(\delta) := \frac{2\delta^2(\delta + 1)}{\pi^2 b^2}. \quad (131)$$

The slope representation

$$\sigma(z) = \frac{1}{\pi} \int_{\mu}^K \frac{z}{a\sqrt{a^2 - z^2}} da \geq \frac{\delta z}{\pi K^2}$$

and (130) imply

$$bn(2r+n) < K^2, \quad bn(n+1) < K^2. \quad (132)$$

Finally, (69) with $p = n$ gives

$$\frac{\delta}{\pi} < f + \frac{1}{4} + \frac{r}{4n}. \quad (133)$$

If $r \leq K/2$, then $r < \delta + n + 1$, and hence

$$\delta < D_{n,f} := \frac{f + 1/2 + 1/(4n)}{1/\pi - 1/(4n)}.$$

Therefore every residual central pair satisfies

$$bn(n+1) < U_f(D_{n,f})^2. \quad (134)$$

The left side increases with n and the right side decreases. Also

$$S_Q \geq \frac{3}{2} \left((Q + \frac{3}{4})^{2/3} - (\frac{3}{4})^{2/3} \right) \geq \frac{5}{4} Q^{2/3} \quad (Q \geq 7).$$

For $f \geq 16$ this first forces $n > f/6$; then $D_{n,f} < 5f$, $U_f(D_{n,f}) < 30f$, $n < 31\sqrt{f}$, and thus $f < 34596$. The remaining monotone endpoint comparisons are finite. Outward Arb arithmetic verifies, for both $Q = f - 1, f$, that every $50 \leq f \leq 34595$ is excluded and leaves exactly 649 necessary pairs before the half-unit reserve is used. Its two $n = 1$ pairs are $(f, n) = (2, 1), (3, 1)$; Corollary 7.20 removes them and leaves 647 necessary pairs inside (127).

If $r > K/2$, the inequality $R_n < 0$ and monotonicity of σ give $\sigma(r) < 1/(2n)$. The radius representation also gives

$$\sigma(r) > \frac{\delta}{2\pi\sqrt{2K(\delta+n+1)}},$$

and therefore

$$K > \frac{\delta^2 n^2}{2\pi^2(\delta+n+1)}, \quad \delta+1 > c_f n, \quad c_f = \frac{\sqrt{1+b^2}-1}{2}. \quad (135)$$

With $\delta_{n,f} = \max\{\pi b, c_f n - 1\}$, every residual outer pair must satisfy

$$\frac{\delta_{n,f}^2 n^2}{2\pi^2(\delta_{n,f} + n + 1)} < K_{\text{nd}}(n, Q). \quad (136)$$

The same sum bound gives $f \leq 66$. Once $c_f n - 1$ is active, the left side of the normalized comparison increases with n , while its right side decreases. Its limiting coefficients are

$$\gamma_f = \frac{c_f^2}{2\pi^2(c_f + 1)}, \quad \beta_J = \frac{3}{2\pi S_J^3}.$$

The certified signs $\gamma_2 < \beta_2$, $\gamma_3 < \beta_3$, and $\gamma_4 > \beta_3$ leave only $f = 2, 3$ as possible unbounded rows. For $4 \leq f \leq 66$, the finite monotone replay leaves exactly 60 necessary pairs in (128); none has $n = 1$.

For reproducibility, all outward-rounded comparisons just invoked are in `scripts/general_d_no_drop_exhaustion_verify.py`. At 256-bit precision it performs 138,184 central endpoint comparisons and 980 outer comparisons, and reports the pre-reserve counts 649 and 60 above. The same certificate was independently replayed through 1024 bits; the Round 5 replay confirms the updated counts 647 and 60.

It remains to rule out escape in $f = 2, 3$. On the upper half of $[\mu, K]$,

$$A(q) \geq \frac{\delta}{2\pi} \sqrt{\frac{\delta}{2K}}.$$

Together with $A(q) < f$, (112), and (135), this proves $\delta \asymp n$ and $K \asymp n^3$. Set

$$\eta = \frac{\delta}{K}, \quad \kappa = K\eta^{3/2}, \quad N = n\sqrt{\eta}.$$

This is exactly the critical scaling (98). For an explicit transfer, define $C_K(f)$ and $C_\delta(f)$ as in Theorem 7.22. If $n \geq \lceil 2/c_f \rceil$, then (135), the failed cutoff, and $\delta^3 < 8\pi^2 f^2 K$ give precisely

$$\frac{c_f}{2}n \leq \delta \leq C_\delta(f)n, \quad \frac{c_f^2}{8\pi^2(C_\delta(f) + 2)}n^3 < K < C_K(f)n^3.$$

The endpoint-action bound $A(q) \geq 2\kappa/(3\pi)$ and $A(q) < f$ give $\kappa < 3\pi f/2$. Since the cone also gives $\delta \geq c_f n/2$,

$$s = \frac{\kappa}{\delta} < \frac{3\pi f}{c_f n} < \frac{38}{n} \quad (f = 2, 3). \quad (137)$$

Indeed, the two coefficients are $48\pi/(\sqrt{65} - 4)$ and $72\pi/(\sqrt{137} - 4)$, both strictly smaller than 38. Thus Theorem 7.22 excludes every residual candidate with $n \geq 380$, since then $s < 1/10$. Candidates below the preliminary threshold $\lceil 2/c_f \rceil$ already satisfy this bound. The two low-floor rows therefore satisfy the explicit bound $n < 380$. $\square$

8 A dimension-dependent no-mode region

The radial form (6), the one-dimensional Dirichlet Poincaré inequality, and $r^{-2} \geq 1$ on $(\rho, 1)$ give

$$\lambda_{\ell,n} \geq \frac{n^2\pi^2}{(1-\rho)^2} + \nu_{\ell,d}^2 - \frac{1}{4}.$$

Since $\nu_{0,d} = (d-2)/2$, one obtains the strengthened wall

$$N_D(A_{\rho,1}^{(d)}, K^2) = 0 \quad \text{if} \quad K^2 \leq \frac{\pi^2}{(1-\rho)^2} + \frac{(d-2)^2 - 1}{4}. \quad (138)$$

The equality face belongs to the no-mode region because the counting function is strict.

In particular, for the general-dimensional Pólya theorem it is enough to prove the shifted-tail estimate in the universal active region

$$K > \frac{\pi}{1-\rho}. \quad (139)$$

Indeed, the complementary region has no eigenvalues already in dimension three, and the additional centrifugal term only enlarges it for $d > 3$. This observation does not prove the stronger standalone Conjecture 1.1 below the wall, but it removes that region from the spectral critical path.

9 Remaining proof obligations

The first interface-cell milestone is complete, and Proposition 6.8 proves that for each fixed ρ only a bounded frequency interval remains; in the thin sector Proposition 6.10 improves the sufficient threshold to $K \geq 125/[8(1 - \rho)^2]$. Proposition 7.1 further replaces the many-cell dynamics by the single scalar (66). For the spectral theorem this scalar is needed only in (139). Lemma 7.2 sharpens its head term, while Corollary 7.4 replaces qualitative terminal nonnegativity by a fractional-top positive sum, and Corollary 7.5 covers the empty-level case. Proposition 7.14 closes an explicit high-floor finite-shell sector, while Theorem 7.15 rules out a negative leading obstruction on the grazing ray. For $p = n$, Proposition 7.12 identifies the exact endpoint phase and Theorem 7.23 proves a global finite reduction. It does not certify every member of that finite family. Theorems 7.10 and 7.17 remove every noncompact first-drop escape. The primary route now has the following precise steps.

1. In the no-drop branch, certify the residual one-sided walls in the explicit central and outer boxes (127)–(128), together with the two rows $f = 2, 3$, $2 \leq n \leq 379$ from Theorem 7.22. The $n = 1$ branch is already closed by Corollary 7.20; in the remaining chambers the discard sector (118) and the favorable endpoint unit in (87) should be retained.
2. For $p < n$, certify the compact residual wall sets left by Theorems 7.10 and 7.17. In the first-floor branch, Theorems 7.6 and 7.7 already close the full sector $r \geq 3/2$, $s \in \{0, 1\}$, and Theorem 7.8 closes the two remaining starts. Thus only the compact $s \geq 2$ residue remains on the first floor. In the high-floor branch the integer data are bounded qualitatively, but explicit bounds and the resulting directed-rounding wall certificates are still required.
3. Insert the resulting all-frequency shifted-tail theorem into Theorem 5.3. No dimension-dependent angular or optical theorem is then needed.

If the first step fails only for an explicitly controlled family of tails, the exact identity (24) is the appropriate aggregate fallback: the weighted negative part must be compared directly with $\mathcal{B}_d(A)$.

References

- [1] N. Filonov, M. Levitin, I. Polterovich, and D. A. Sher, Pólya’s conjecture for Euclidean balls, *Invent. Math.* **234** (2023), 129–169. <https://doi.org/10.1007/s00222-023-01198-1>.
- [2] N. Filonov, M. Levitin, I. Polterovich, and D. A. Sher, Uniform enclosures for the phase and zeros of Bessel functions and their derivatives, *SIAM J. Math. Anal.* **56** (2024), no. 6, 7644–7682. <https://doi.org/10.1137/24M1642032>.
- [3] N. Filonov, M. Levitin, I. Polterovich, and D. A. Sher, Pólya’s conjecture for Dirichlet eigenvalues of annuli, *J. Lond. Math. Soc.* **113** (2026), no. 2, e70425. <https://doi.org/10.1112/jlms.70425>.