

Purely Analytic Supplement for A Dirichlet Pólya Inequality on Three-Dimensional Spherical Shells

Anonymous for peer review

17 July 2026

Abstract

This support volume contains the detailed hand proofs and exact-rational tables used in the spherical-shell theorem. The live finite argument in Parts I–VI includes the 38-state closed post-event domination theorem in Part III; Parts VII–IX prove the retained-remainder theorem above the final staircase, and Part X assembles the residual closure. The global 611-row allocation also records superseded rows and regression checks and is not a second live high-event premise; individual inequalities cited by the live modules remain proof premises. The former aggregate-tail argument, programs, and interval certificates remain available only as optional independent checks and are not premises of this analytic proof.

Reading convention. The volume is modular so that each bottleneck lemma can be reviewed and built independently. Every included part is generated from its corresponding LaTeX source in the `manuscript/analytic` directory. The repeated title pages deliberately mark the trust boundary between exhibits.

Two review paths. A conceptual review should read the phase and layer-cake reductions in the main paper, then Parts VII–IX here, followed by the assembly in Part X. A finite audit should read the zero/index and event material in Parts I–III, the live owner statements in Part IV, and Parts V–VI and X. The twenty-two seam rows in Part IV, together with any other rows classified as retired by the dependency table and manifest, are provenance or regression material and may be skipped when reconstructing the live implication chain.

Minimal dependency graph

result	premises	external theorem	owner	status		
phase and lattice proxy	separation, determinant, exact strict phase count	FLPS phase; annuli Theorem 5.1 and Lemma 5.2	main paper	used	in	final proof
weighted scaffold and primitive owners	multiplicity identity, action integral, aggregate clause	balls Theorem 5.1; annuli Proposition 3.1 and Theorems 1.4, 3.4	main paper	used	in	final proof
zero/index registry	recurrences, rational signs, fixed-channel comparison	DLMF (10.21.2), Parts I–III (10.47.3); Lorch abstract inequalities (i), (ii)		used	in	final proof
finite lower/high staircases	inventories, caps, closed post-event domination	none beyond the preceding row	Parts I–IV	used	in	final proof
optical and owner subtraction	product/radial deficits and exact regional logic	none	Parts V–VI	used	in	final proof
all-frequency middle theorem	retained remainder, angular block, scalar positivity	none	Parts VII–IX	used	in	final proof
final residual closure	all preceding used rows	none	Part X	used	in	final proof
retired seam, $K = 200$, aggregate, executable and interval replays	independent regression contours	Arb only in retired interval replay	Part IV and archive/legacy	optional	cross-check	

The source/output manifest and numerical build order are given in `manuscript/analytic/MANIFEST.md`. The master deliberately includes the ten component PDFs so that each source can also be compiled and reviewed as a standalone document.

Provenance census of the exact ledger

This census is included for traceability, not as an additional theorem or dependency. The canonical registry has 553 core rows and 58 supplemental rows. The following inclusive ranges give a one-to-one allocation; the part numbers refer to this volume. The family codes denote, in order, rational- π checks (PI), thresholds (TH), strict-floor conventions (SF), zero specializations (Z), lower inventories (LI), checkpoint inventories (KI), cap payments (CP), localizations (LOC), the upper owner (U), optical screens (OPT), exact subtraction (SUB), and supplemental seam/payment rows (SUP).

canonical family	row range	unique owner
PI	PI1–PI4	II
TH	TH1–TH7, TH31–TH34 / TH8–TH30	I / II
SF	SF1–SF35	II
Z	Z1–Z48	I
LI	LI1–LI8, LI10–LI44 / LI9	I / II
KI	KI1–KI117	II
CP	CP1–CP43, CP56, CP60–CP77 / remaining rows	II / III
LOC	LOC1–LOC47	III
U	U1–U8	V
OPT	OPT1–OPT14 / OPT15–OPT50	V / II
SUB	SUB1–SUB68	VI
SUP	SUP1–SUP57 / SUP58	IV / I

Here “remaining rows” in CP means CP44–CP55, CP57–CP59, and CP78–CP98. Thus the six disjoint owner totals are

$$103 + 278 + 83 + 57 + 22 + 68 = 611.$$

In particular, the ten lower- d identities SUP36–SUP45 are assigned to Part IV although Part II also proves their general odd-sum identity; SUP58 is assigned to Part I; and OPT15–OPT50 are assigned to Part II although Part V reuses their consequences. These are proof redundancies, not row overlaps. The displayed ranges exhaust every canonical row and have empty pairwise intersections.

Contents

Minimal dependency graph	2
Provenance census of the exact ledger	2
I Remaining hand-visible ledger rows	4
II Universal arithmetic and checkpoint inventories	13
III Connected high-event registry	22
IV Seam and two-sided payments	34
V Upper owner and optical screens	40
VI Exact owner subtraction	45
VII Global retained-remainder identity	48
VIII Global middle-ratio angular bound	54
IX Upward-closed scalar positivity	59
X Final all-frequency residual closure	67

Hand-visible analytic derivations for 103 finite-ledger predicates

1 Scope and logical convention

This note exposes, in hand-checkable form, the predicates not assigned to the other five ledger packets. They are precisely

11 threshold predicates + 48 zero predicates + 43 lower-closure predicates + 1 cap-74 predicate = 103.

The omitted lower-closure predicate is the single six-cap tuple identity; it belongs to the cap-inventory packet. Every inequality below is proved from a displayed analytic inequality, integer arithmetic, rational cross multiplication, or positive-side squaring. An executable ledger may check the same arithmetic, but no executable output is used as a premise.

We use

$$\omega_0 = \frac{\sqrt{3}}{2\pi} - \frac{1}{6}, \quad C_0 = 1 + \frac{8\sqrt{2}}{15}, \quad \rho_* = \frac{\omega_0}{2C_0}, \quad \rho_0 = \frac{1}{\sqrt{337}}, \quad \sigma = \frac{3\omega_0}{4}, \quad \rho_c = \frac{1}{1+2\pi}, \quad d = \frac{\sqrt{337}}{2}.$$

The elementary enclosure

$$\frac{333}{106} < \pi < \frac{355}{113} < \frac{22}{7} \tag{1}$$

is analytic: the lower and middle bounds follow from alternating truncations of Machin's identity $\pi = 16 \arctan(1/5) - 4 \arctan(1/239)$, and the last one is immediate by cross multiplication. For reference, 355/113 exceeds the upper Machin truncation by the positive rational numerator 45167474711 over denominator 189820334689453125. The classical integral

$$0 < \int_0^1 \frac{x^4(1-x)^4}{1+x^2} dx = \frac{22}{7} - \pi$$

also gives the final upper bound directly.

2 The eleven threshold predicates

First we obtain a rational two-sided enclosure for ω_0 . The exact integer residuals

$$3 \, 1101^2 113^2 - 607^2 355^2 = 1998482 > 0, \quad 25 \, 333^2 - 243 \, 106^2 = 41877 > 0$$

and (1) imply

$$\sqrt{3} > \frac{607 \, 355}{1101 \, 113} > \frac{607\pi}{1101}, \quad 9\sqrt{3} < 5\pi.$$

Consequently

$$\frac{40}{367} < \omega_0 < \frac{1}{9}. \tag{2}$$

Indeed, $607/2202 - 1/6 = 40/367$, while $5/18 - 1/6 = 1/9$.

Since $2 \cdot 32^2 - 45^2 = 23 > 0$, one has $\sqrt{2} > 45/32$, hence $C_0 > 7/4$ and $\rho_* < 2/63$. The following chain consists entirely of strict rational comparisons, with the indicated radical comparisons justified by squaring:

$$\rho_* < \frac{2}{63} < \frac{1}{\sqrt{337}} < \frac{1}{18} < \frac{3}{40} < \frac{3\omega_0}{4} < \frac{1}{12} < \frac{7}{51} < \frac{1}{1+2\pi} < \frac{1}{7} < \frac{39}{50} < \frac{7}{8}.$$

Here

$$4 \cdot 337 < 63^2, \quad 18^2 < 337, \quad 367 < 400, \quad 51 < 84, \quad 367 < 420,$$

and $7/51 < \rho_c < 1/7$ follows respectively from $\pi < 22/7$ and $\pi > 3$. This proves the five ratio-order rows. It also proves

$$\omega_0 < \frac{1}{9} < \frac{7}{51} < \rho_c.$$

Moreover

$$\omega_0 d > \frac{20\sqrt{337}}{367} > 1, \quad 400 \cdot 337 - 367^2 = 111 > 0.$$

The four terminal comparisons follow from the same bounds:

$$\begin{aligned} \sqrt{114} < 18 < \frac{2}{\omega_0}, & \quad \frac{2}{\omega_0} < \frac{367}{20}, \\ \rho_c < \frac{1}{7} < \frac{60}{367} < \frac{3\omega_0}{2}, & \quad \omega_0 \sqrt{114} > \frac{40\sqrt{114}}{367} > 1. \end{aligned}$$

The last positive-side squaring has residual

$$1600 \cdot 114 - 367^2 = 47711 > 0.$$

Thus the one-to-one threshold registry is

ID	source row	proved predicate
T1–T5	1–5	$\rho_* < \rho_0 < \sigma < \rho_c < 39/50 < 7/8$, one adjacent pair per row
T6	6	$\omega_0 < \rho_c$
T7	7	$\omega_0 d > 1$
T8	31	$\sqrt{114} < 2/\omega_0$
T9	32	$2/\omega_0 < 367/20$
T10	33	$\rho_c < 3\omega_0/2$
T11	34	$\omega_0 \sqrt{114} > 1$

3 The forty-eight zero specializations

3.1 A recurrence and elementary tangent bounds

Put

$$j_\ell(x) = \sqrt{\frac{\pi}{2x}} J_{\ell+1/2}(x), \quad P_\ell(x) = x^{\ell+1} j_\ell(x).$$

Then

$$P_{\ell+1} = (2\ell+1)P_\ell - x^2 P_{\ell-1}, \quad P_0 = \sin x, \quad P_1 = \sin x - x \cos x. \quad (3)$$

Write $P_\ell = A_\ell(t) \sin x + x C_\ell(t) \cos x$, $t = x^2$. The recurrence gives

$$\begin{aligned} A_2 &= 3 - t, & C_2 &= -3, \\ A_3 &= 15 - 6t, & C_3 &= t - 15, \\ A_5 &= 945 - 420t + 15t^2, & C_5 &= -945 + 105t - t^2, \\ A_6 &= 10395 - 4725t + 210t^2 - t^3, & C_6 &= -10395 + 1260t - 21t^2, \\ A_7 &= 135135 - 62370t + 3150t^2 - 28t^3, & C_7 &= -135135 + 17325t - 378t^2 + t^3. \end{aligned}$$

For $0 < s < \pi/2$, differentiation and integration give

$$\tan s > s, \quad \tan s > s + \frac{s^3}{3}. \quad (4)$$

For $0 < s < 1$, $\sin s < s$ and $\cos s > 1 - s^2/2$ give

$$\tan s < \frac{s}{1 - s^2/2}. \quad (5)$$

3.2 Sixteen half-period locations

For each internal face, the first displayed margin below is $w - (h_L/2)(22/7)$; the second is $(h_R/2)(333/106) - w$. Positivity therefore proves both strict half-period location rows without a decimal approximation.

(ℓ, n)	w	(h_L, h_R)	left margin	right margin
(1, 2)	77/10	(4, 5)	99/70	163/1060
(2, 2)	9	(5, 6)	8/7	45/106
(3, 2)	103/10	(6, 7)	61/70	737/1060
(5, 2)	129/10	(8, 9)	23/70	1311/1060
(1, 3)	21/2	(6, 7)	15/14	105/212
(2, 3)	61/5	(7, 8)	6/5	97/265
(6, 1)	10	(6, 7)	4/7	211/212
(7, 1)	23/2	(7, 8)	1/2	113/106

3.3 Sixteen half-period signs

At an integral or half-integral multiple of π , exactly one of $\sin x, \cos x$ vanishes. Direct substitution in the displayed polynomials gives the following complete sign table.

(ℓ, n)	h_L	left sign	(ℓ, n)	h_R	right sign
(1, 2)	4	—	(1, 2)	5	+
(2, 2)	5	—	(2, 2)	6	+
(3, 2)	6	—	(3, 2)	7	+
(5, 2)	8	—	(5, 2)	9	+
(1, 3)	6	+	(1, 3)	7	—
(2, 3)	7	+	(2, 3)	8	—
(6, 1)	6	+	(6, 1)	7	—
(7, 1)	7	+	(7, 1)	8	—

Here are explicit polynomial witnesses for the only signs not immediate from $A_1 = 1, C_1 = -1, A_2 = 3 - t, C_2 = -3, A_3 = 15 - 6t, C_3 = t - 15$. From (1), at $x = 4\pi$ one has $144 < t < 159$, at $x = 9\pi/2$ one

has $t > 182$, and at $x = 7\pi/2$ one has $110 < t < 121$. Monotonicity on these rational intervals and direct evaluation give

$$\begin{aligned} C_5(t) &< C_5(144) = -6561, & A_5(t) &> A_5(182) = 421365, \\ C_6(t) &< C_6(81) = -46116, & A_6(t) &> A_6(110) = 700645, \\ A_7(t) &< A_7(110) = -5878565, & C_7(t) &< C_7(144) = -2492559. \end{aligned}$$

For example, $A'_6 = -4725 + 420t - 3t^2$ is positive on $[110, 121]$, while A'_7 is negative there; the other asserted monotonicities follow at once from their displayed derivatives. Combining these signs with the quadrant signs of sine and cosine proves all sixteen entries.

3.4 Eight rational-wall signs

Substitution in (3) reduces each rational-wall sign to one of the following exact tangent comparisons. The shifted angle is always in $(0, \pi/2)$, as already proved by the location table.

(ℓ, n)	wall	hand certificate	$\text{sgn } j_\ell(w)$
(1, 2)	77/10	For $u = 5\pi/2 - w$, $\tan u > 3/20 > 10/77 = 1/w$; hence $\tan(w - 2\pi) = \cot u < w$.	−
(2, 2)	9	For $u = 3\pi - w$, $\tan u > 21/50 > 9/26 = 3w/(w^2 - 3)$.	−
(3, 2)	103/10	For $u = 7\pi/2 - w$, $\tan u > 69/100 > 621540/938227 = (6w^2 - 15)/(w(w^2 - 15))$.	−
(5, 2)	129/10	For $y = w - 4\pi$, $0 < y < 17/50$, so $\tan y < 1700/4711 < 177800829/427700150 = -wC_5/A_5$.	−
(1, 3)	21/2	For $u = 7\pi/2 - w$, $\tan u > 49/100 > 2/21 = 1/w$, hence $\tan(w - 3\pi) < w$.	+
(2, 3)	61/5	For $u = 4\pi - w$, $\tan u > 9/25 > 915/3646 = 3w/(w^2 - 3)$.	+
(6, 1)	10	For $y = w - 3\pi$, $4/7 < y < 29/50$, so $\tan y < 2900/4159 < 188790/127579 = -wC_6/A_6$.	+
(7, 1)	23/2	For $u = 4\pi - w$, $\tan u > u + u^3/3 > 546377/375000 > 3153151489/2276518664 = wC_7/A_7$.	+

For complete cross-multiplication visibility, the four less immediate positive gaps in that table are

$$\begin{aligned} \frac{69}{100} - \frac{621540}{938227} &= \frac{2583663}{93822700}, \\ \frac{177800829}{427700150} - \frac{1700}{4711} &= \frac{110529450419}{2014895406650}, \\ \frac{188790}{127579} - \frac{2900}{4159} &= \frac{415198510}{530601061}, \\ \frac{546377}{375000} - \frac{3153151489}{2276518664} &= \frac{3837851856583}{53355906187500}. \end{aligned}$$

The four small gaps are respectively $31/1540$, $24/325$, $829/2100$, $9939/91150$.

3.5 Intermediate rational-wall signs used by the index induction

The coefficient signs make all unmentioned entries in the sign vectors immediate. The remaining intermediate entries require the following exact thresholds. At $w = 10$, with $y = 10 - 3\pi$,

$$\tan y < \frac{2900}{4159} < \frac{170}{117} < 10, \quad \tan y > y > \frac{4}{7} > \frac{890}{21789}.$$

Substitution in P_1, P_3, P_5 gives $\text{sgn}(j_1, j_3, j_5) = (+, -, -)$, and $P_4 = 7P_3 - w^2P_2 < 0$. At $w = 103/10$, with $u = 7\pi/2 - w$,

$$\tan u > \frac{69}{100} > \frac{10}{103},$$

which gives $j_1(w) > 0$. At $w = 23/2$, with $u = 4\pi - w$,

$$\tan u > \frac{546377}{375000} > \frac{3153151489}{2276518664} > \frac{224020}{186301} > \frac{138}{517}.$$

Besides the target j_7 sign, the last two thresholds give $j_4(w) < 0$ and $j_2(w) > 0$; the recurrence gives the intervening signs. Finally, at $w = 129/10$, with $y = w - 4\pi$,

$$\tan y < \frac{1700}{4711} < \frac{177800829}{427700150} < \frac{651063}{327820} < \frac{129}{10}.$$

Besides the target j_5 sign, this proves $j_3(w) > 0$ and $j_1(w) < 0$, and the recurrence gives the remaining signs. Every sign used below is therefore supported by a displayed rational comparison.

3.6 Exact zero indices from adjacent-order interlacing

For $x > 0$, define the strict positive-zero count

$$Z_\ell(x) := \#\{n \in \mathbb{N} : j_{\ell+1/2,n} < x\}.$$

Lemma 3.1 (Interlacing-to-index principle). *For every $\ell \in \mathbb{N}_0$ and $n \in \mathbb{N}$,*

$$j_{\ell+1/2,n} < j_{\ell+3/2,n} < j_{\ell+1/2,n+1}. \quad (6)$$

If x is not a zero of $j_0, \dots, j_L$, then, for $0 \leq \ell < L$,

$$Z_{\ell+1}(x) \in \{\max\{Z_\ell(x) - 1, 0\}, Z_\ell(x)\}, \quad \text{sgn } j_{\ell+1}(x) = (-1)^{Z_{\ell+1}(x)}. \quad (7)$$

Consequently $Z_0(x)$, together with the signs of $j_1(x), \dots, j_L(x)$, determines all the counts $Z_1(x), \dots, Z_L(x)$ uniquely.

Proof. DLMF 10.21.2 states, for $\nu \geq 0$,

$$j_{\nu,1} < j_{\nu+1,1} < j_{\nu,2} < j_{\nu+1,2} < j_{\nu,3} < \dots.$$

Taking $\nu = \ell + 1/2$ proves (6). Put $q = Z_\ell(x)$. The first inequality in (6) shows that at most q zeros of order $\ell + 3/2$ lie below x . If $q \geq 1$, the second inequality shows that its first $q - 1$ zeros lie below $j_{\ell+1/2,q} < x$. Thus

$$\max\{q - 1, 0\} \leq Z_{\ell+1}(x) \leq q,$$

which is the first assertion in (7).

The spherical Bessel function j_ℓ is positive for all sufficiently small positive arguments. Its positive zeros are simple by uniqueness for the Bessel equation, so its sign changes at every positive zero. This proves the parity identity in (7). When $q \geq 1$, the two possible counts have opposite parity; when $q = 0$, only zero is possible. The sign therefore selects the count uniquely. $\square$

Because $j_0(x) = \sin x/x$, one has $j_{1/2,n} = n\pi$. The strict locations in the first table therefore give $Z_0(w)$ exactly. Applying Lemma 3.1 successively to the nonzero sign vectors obtained from (3) and the exact tangent comparisons above gives:

wall w	target (ℓ, n)	location of w	$Z_0(w)$	$\text{sgn}(\mathbf{j}_0, \dots, \mathbf{j}_\ell)(w)$	$(Z_0, \dots, Z_\ell)(w)$
77/10	(1, 2)	$(2\pi, 5\pi/2)$	2	(+, -)	(2, 1)
9	(2, 2)	$(5\pi/2, 3\pi)$	2	(+, +, -)	(2, 2, 1)
10	(6, 1)	$(3\pi, 7\pi/2)$	3	(-, +, +, -, -, -, +)	(3, 2, 2, 1, 1, 1, 0)
103/10	(3, 2)	$(3\pi, 7\pi/2)$	3	(-, +, +, -)	(3, 2, 2, 1)
21/2	(1, 3)	$(3\pi, 7\pi/2)$	3	(-, +)	(3, 2)
23/2	(7, 1)	$(7\pi/2, 4\pi)$	3	(-, -, +, +, -, -, -, +)	(3, 3, 2, 2, 1, 1, 1, 0)
61/5	(2, 3)	$(7\pi/2, 4\pi)$	3	(-, -, +)	(3, 3, 2)
129/10	(5, 2)	$(4\pi, 9\pi/2)$	4	(+, -, -, +, +, -)	(4, 3, 3, 2, 2, 1)

In every row the target component is $Z_\ell(w) = n - 1$, and the displayed sign is nonzero. Hence, with the convention $j_{\nu,0} = 0$,

$$j_{\ell+1/2, n-1} < w < j_{\ell+1/2, n}.$$

Thus the unique next zero is the claimed n -th zero, and the eight rows prove

$$\begin{aligned} j_{3/2,2} &> 77/10, & j_{5/2,2} &> 9, & j_{7/2,2} &> 103/10, & j_{11/2,2} &> 129/10, \\ j_{3/2,3} &> 21/2, & j_{5/2,3} &> 61/5, & j_{13/2,1} &> 10, & j_{15/2,1} &> 23/2. \end{aligned} \tag{8}$$

3.7 Six Lorch rows and two wall-order rows

For $\nu = \ell + 1/2$, the second strict inequality, item (ii), displayed in the published abstract of Lorch's article *Some inequalities for the first positive zeros of Bessel functions*, SIAM J. Math. Anal. **24** (1993), 814–823, doi:10.1137/0524050, is valid for $-1 < \nu < \infty$ and states

$$j_{\nu,1}^2 > \frac{24(\nu+1)^2}{1-2\nu+\sqrt{(2\nu+3)(2\nu+11)}} - 2(\nu^2-1) =: R_\ell.$$

After rationalization,

$$R_\ell = \frac{3(2\ell+3)\sqrt{(\ell+2)(\ell+6)} - 2\ell^2 + \ell + 6}{4}.$$

All sides being squared are positive, and the following integer residuals prove the three required strict specializations:

ℓ	wall c	positive squared residual proving $R_\ell > c^2$
8	71/6	$1026^2 \cdot 35 - 6067^2 = 35171$
9	64/5	$1575^2 \cdot 165 - 20059^2 = 6939644$
10	69/5	$3450^2 \cdot 3 - 5911^2 = 767579$

Since $j_\ell(x) > 0$ for sufficiently small $x > 0$, and its first positive zero is strictly to the right of c , these same three rows imply $j_\ell(c) > 0$. Thus the three executable sign rows are analytic corollaries, not independent numerical assumptions. Finally,

$$\frac{103}{10} - \frac{81}{8} = \frac{7}{40} > 0, \quad \frac{61}{5} - \frac{21}{2} = \frac{17}{10} > 0.$$

The one-to-one zero registry is as follows. Number the eight internal faces in the order of the location table. For face r , IDs $Z_{5r-4}, \dots, Z_{5r}$ are, in order: right of the left half-period, left of the right half-period, left half-period sign, right half-period sign, and rational-wall sign. This bijects IDs $Z1$ – $Z40$ with the first forty source rows. The remaining bijection is

ID	source row	predicate
Z41	41	Lorch specialization $\ell = 8, c = 71/6$
Z42	42	positive sign at $\ell = 8, c = 71/6$
Z43	43	Lorch specialization $\ell = 9, c = 64/5$
Z44	44	positive sign at $\ell = 9, c = 64/5$
Z45	45	Lorch specialization $\ell = 10, c = 69/5$
Z46	46	positive sign at $\ell = 10, c = 69/5$
Z47	47	$103/10 > 81/8$
Z48	48	$61/5 > 21/2$

4 The forty-three lower-closure predicates

4.1 Eight activation and exclusion rows

The complete arithmetic is

$$\begin{aligned}
(23/2)^2 - 114 &= 73/4 > 0, & 10^2 &= 100, \\
(129/10)^2 - 114 &= 5241/100 > 0, & 103/10 - 81/8 &= 7/40 > 0, \\
(81/8)^2 + 8 &= 7073/64, & (81/8)^2 + 18 - 114 &= 417/64 > 0, \\
(21/2)^2 + 4 - 114 &= 1/4 > 0, & 16\pi^2 &> 16 \cdot 3^2 > 114.
\end{aligned}$$

These are exactly lower IDs L1–L8, corresponding to source rows 1–8.

4.2 Five Weyl payments

For $0 < \rho < \rho_c < 1/7$, (1) yields the uniform analytic lower bound

$$W(\rho, K) = \frac{2}{9\pi}(1 - \rho^3)K^3 > \frac{38}{539}K^3. \quad (9)$$

Using $d > 9$ and $\sqrt{7073}/8 > 21/2$, the five payments are the following exact positive reserves:

ID	source row	threshold/cap	reserve from (9)
L9	10	$K > d, 46$	$(38/539)9^3 - 46 = 2908/539$
L10	11	$K > 10, 59$	$(38/539)10^3 - 59 = 6199/539$
L11	12	$K > 81/8, 66$	$(38/539)(81/8)^3 - 66 = 990435/137984$
L12	13	$K > 21/2, 69$	$(38/539)(21/2)^3 - 69 = 555/44$
L13	14	$K > \sqrt{7073}/8, 78$	$(38/539)(21/2)^3 - 78 = 159/44$

4.3 The cap–395 cell: eighteen rows

Put $R = 367/20$, $z_\ell = \ell + 1/2$. The analytic turning estimate

$$G_K(z) < \frac{(K - z)^{3/2}}{3\sqrt{K}}$$

is increasing in K , so on $K < R$ a strict cap c_ℓ follows from

$$\frac{(R - z_\ell)^3}{9R} = \frac{(357 - 20\ell)^3}{1321200} < \left(c_\ell + \frac{3}{4}\right)^2. \quad (10)$$

The following table gives all fifteen caps and the exact positive numerator $82575(4c_\ell + 3)^2 - (357 - 20\ell)^3$ obtained by clearing the positive denominators in (10).

ℓ	0	1	2	3	4	5	6	7
c_ℓ	6	5	5	4	4	3	3	3
margin	14697882	5409422	11827162	3611502	8555642	1604782	5267322	8361062

ℓ	8	9	10	11	12	13	14
c_ℓ	2	2	1	1	1	1	0
margin	2346202	4446342	176282	1474822	2444562	3133502	286642

Thus L15–L29 correspond bijectively to source rows 16–30, with L(15 + ℓ) the ℓ -th table column. Since $G_R(z)$ is strictly decreasing in z , the $c_{14} = 0$ row removes every $\ell \geq 14$ tail; this is L30/source row 31.

The weighted cap itself, L14/source row 15, is visible from the contributions

$$6, 15, 25, 28, 36, 33, 39, 45, 34, 38, 21, 23, 25, 27, \quad \sum_{\ell=0}^{13} (2\ell + 1)c_\ell = 395.$$

Finally, on the exceptional cell $\rho K \geq 5/2$, with $\rho < 11/80$ and $1/\pi > 113/355$,

$$\begin{aligned} W(\rho, K) &> \frac{2}{9} \frac{113}{355} \left(\frac{5}{2}\right)^3 \left[\left(\frac{80}{11}\right)^3 - 1 \right] \\ &= \frac{160293325}{378004} = 395 + \frac{10981745}{378004} > 395. \end{aligned}$$

This is L31/source row 32.

4.4 Twelve floor and boundary-owner rows

The nine finite arithmetic rows are instances of one identity:

$$\frac{3}{2}(p+1) \leq 2p + \frac{1}{2} \quad (p \geq 2), \quad \left(2p + \frac{1}{2}\right) - \frac{3}{2}(p+1) = \frac{p-2}{2} \geq 0. \quad (11)$$

At $p = 2$, $(3/2)3 = 9/2$; this is L32/source row 33. For $p = 2, \dots, 9$, formula (11) is respectively L33–L40/source rows 34–41.

The half-integer face is exact: $5/2 = 2 + 1/2$. Assigning it to the high side makes the interface remainder vanish and leaves the exceptional cell closed; this is L41/source row 42. At $K = 2/\omega_0$,

$$p = \lfloor \omega_0 K \rfloor = 2, \quad \rho K \leq \frac{2\rho_c}{\omega_0} < 3$$

by T10. This proves L42/source row 43. Moreover

$$\rho K - \frac{1}{2} < \frac{5}{2}, \quad m = \left\lfloor \rho K - \frac{1}{2} \right\rfloor \leq 2 \leq 2p - 1 = 3,$$

and the strict exceptional condition $K < 2/\omega_0$ fails at equality. Thus the complementary average-floor branch owns the equality face, proving L43/source row 44. More generally, if $K \geq 2/\omega_0$, then $p \geq 2$ and

$$\rho K < \frac{3}{2}(p+1) \leq 2p + \frac{1}{2}, \quad m \leq 2p - 1,$$

so the finite instances conceal no unproved large- p case.

5 The supplemental cap–74 incompatibility

The propagated ball wall requires $z^2 > 2409/100$, while the product wall requires $z^2 < 70/3$. They are incompatible because

$$\frac{2409}{100} - \frac{70}{3} = \frac{7227 - 7000}{300} = \frac{227}{300} > 0.$$

This is supplemental ID S1 and is a hand proof, not merely a reported certificate value.

6 Census and source-visibility verdict

family	count	IDs	hand-visible status before this note
threshold remainder	11	T1–T11	printed, with a few implications compressed
zero specializations	48	Z1–Z48	tables printed; several recurrence steps compressed
lower remainder	43	L1–L43	payments printed; floor instances generated implicitly
supplemental cap–74	1	S1	printed exactly
total	103		

No predicate in this 103-row allocation is an unsupported executable-only mathematical premise. Before this note, the principal visibility gaps were the one-to-one expansion of the forty zero sign/location rows, the fact that the three Lorch-wall signs are logical corollaries of the strict first-zero bounds, and the expansion of the twelve floor/boundary rows. The displayed recurrences, rational margins, general floor identity, and exact ownership arguments close those presentation gaps.

Exact-ledger decomputerization: Packets A and B

16 July 2026

Abstract

This note replaces two groups of executable exact-arithmetic checks in the three-dimensional spherical-shell Pólya proof. Packet A proves once and for all the strict-bracket convention, odd-multiplicity block sums, the optical product-deficit identity, and monotonicity of propagated zero and channel cuts. Packet B gives exact hand-checkable tables for the five moving $3z$ -crossings and for the seven first-excluded checkpoint channels. It then proves, without a finite search, that the five printed checkpoint inventories are exhaustive. Every numerical comparison below is an equality of rational numbers, a cross multiplication, or a positive-side squaring. No program output, decimal approximation, or interval calculation is a premise.

Contents

1	Scope and notation	1
2	A rational enclosure for π	2
3	Packet A: universal exact-arithmetic lemmas	2
3.1	Strict floors and equality owners	2
3.2	Odd multiplicities and block caps	3
3.3	The symbolic optical product deficit	4
3.4	Monotonic propagated cuts	5
4	Packet B: moving-$3z$ ordering	5
5	Packet B: checkpoint boundary table and exhaustiveness	7
6	Executable-row coverage contract	8

1 Scope and notation

Throughout $0 < \rho < 1$. Put

$$z = z_\rho := \frac{\pi}{1 - \rho}, \quad k_m(\rho)^2 := z_\rho^2 + m(m + 1),$$

and define the two ratio endpoints

$$\rho_0 := \frac{1}{\sqrt{337}}, \quad \rho_c := \frac{1}{1 + 2\pi}.$$

The high-checkpoint argument uses the strict proxy

$$b_{\ell,n}(z)^2 := \max\{n^2 z^2 + \ell(\ell + 1), \beta_{\ell,n}^2\}. \quad (1)$$

A channel may be counted at K only when $b_{\ell,n}(z) < K$. The constants $\beta_{\ell,n}$ are strict lower bounds for the corresponding ball zeros. The zero bounds used in the checkpoint table are displayed explicitly in Section 5.

The purpose of this note is deliberately limited. It replaces the general exact-arithmetic and checkpoint-ordering rows of the old ledger; it does not replace the zero-sign proofs, high-event payment registry, seam arithmetic, optical endpoint screens, or formal owner subtraction.

2 A rational enclosure for π

Lemma 2.1 (Hand enclosure for π). *One has*

$$\frac{333}{106} < \pi < \frac{22}{7}, \quad \frac{1}{\pi} < \frac{106}{333}. \quad (2)$$

Proof. For $0 < x \leq 1$, the alternating series for $\arctan x$ gives a strict lower bound after a negative term and a strict upper bound after a positive term. Hence

$$\arctan \frac{1}{5} > \frac{1}{5} - \frac{1}{3 \cdot 5^3} + \frac{1}{5 \cdot 5^5} - \frac{1}{7 \cdot 5^7} = \frac{323852}{1640625}, \quad \arctan \frac{1}{239} < \frac{1}{239}.$$

Machin's identity therefore yields

$$\pi = 16 \arctan \frac{1}{5} - 4 \arctan \frac{1}{239} > \frac{1231847548}{392109375},$$

and direct subtraction gives the positive margin

$$\frac{1231847548}{392109375} - \frac{333}{106} = \frac{3418213}{41563593750} > 0.$$

For the upper bound, polynomial division and elementary integration give

$$0 < \int_0^1 \frac{x^4(1-x)^4}{1+x^2} dx = \frac{22}{7} - \pi.$$

Finally, reciprocation of the positive lower bound for π gives the last inequality. Notice also that $22/7 - 333/106 = 1/742 > 0$, so the enclosure is nonempty. $\square$

3 Packet A: universal exact-arithmetic lemmas

3.1 Strict floors and equality owners

Definition 3.1. For $x \in \mathbb{R}$, let

$$[x]_{<} := \#\{q \in \mathbb{Z}_{\geq 1} : q < x\}.$$

Thus the bracket counts positive integers under a *strict* cutoff.

Lemma 3.2 (Strict-bracket identity). *For every real x ,*

$$[x]_{<} = \max\{0, [x] - 1\}. \quad (3)$$

In particular, for every integer $m \geq 1$,

$$[m]_{<} = m - 1, \quad [m + \tfrac{1}{2}]_{<} = m.$$

Consequently an integer equality wall is owned by the excluded side.

Proof. If $x \leq 1$, both sides of (3) vanish. If $x > 1$, the positive integers below x are exactly $1, \dots, [x] - 1$. This also proves the two specializations. $\square$

The same lemma handles the two equality conventions used later. If $A + 1/4 = n \in \mathbb{Z}_{\geq 1}$, then $[A + 1/4]_{<} = n - 1$. If $a = m\pi$ in the optical radial decomposition, the strict radial layers are represented by $(N, t) = (m - 1, 1)$, not by $(m, 0)$.

3.2 Odd multiplicities and block caps

Lemma 3.3 (Odd-sum and block-cap identity). *For every integer $h \geq 0$,*

$$\sum_{\ell=0}^h (2\ell + 1) = (h + 1)^2, \quad \sum_{\ell=0}^{h-1} (2\ell + 1) = h^2. \quad (4)$$

More generally, if radial layer n contains a complete angular block $0 \leq \ell \leq h_n$, with $h_n = -1$ denoting an empty block, then its total multiplicity is

$$\sum_n (h_n + 1)^2. \quad (5)$$

If $x^2 = h(h + 1)$ is a strict product equality wall, the equality channel $\ell = h$ is absent and the contribution is h^2 .

Proof. The first identity follows either by induction or by $(\ell + 1)^2 - \ell^2 = 2\ell + 1$. Summing over independent radial layers gives (5). At $x^2 = h(h + 1)$, $\sqrt{x^2 + 1/4} - 1/2 = h$; Lemma 3.2 excludes $\ell = h$, leaving $0 \leq \ell < h$ and hence h^2 . $\square$

For completeness, the cap arithmetic collapsed by this lemma is printed in compact form below. A dash means that another analytic argument has declared the cell impossible; the lemma asserts only the arithmetic of the non-dash entries.

$k_8 : H \setminus h$	$\emptyset$	0	1	2	3
7	64	65	68	73	80
6	49	50	53	—	—
5	36	37	40	45	52
4	25	26	29	34	41

$k_9 : H \setminus h$	$\emptyset$	0	1	2	3	4
8	81	—	—	—	—	—
7	64	65	68	73	—	—
6	49	50	53	58	65	74
5	36	37	40	45	52	61
4	25	26	29	34	41	50

Indeed every entry is $(H+1)^2$ plus 0 or $(h+1)^2$. At k_7 , the first blocks $H = 4, 5, 6$ give 25, 36, 49, and second channels 0, 1 contribute $1 + 3 = 4$. The remaining printed caps decompose as

$$\begin{aligned}
k_{10} : & \quad 10^2; \quad 9^2, 9^2 + 4^2; \\
& \quad 8^2, 8^2 + 4^2, 8^2 + 5^2, 8^2 + 6^2; \\
& \quad 7^2 + 5^2, 7^2 + 4^2 + 2^2, 7^2 + 5^2 + 2^2, 7^2 + 6^2, 7^2 + 6^2 + 2^2 \\
& \quad = 100; \quad 81, 97; \quad 64, 80, 89, 100; \quad 74, 69, 78, 85, 89, \\
k_{11} : & \quad 11^2; \quad 10^2, 10^2 + 5^2; \quad 9^2 + 5^2, 9^2 + 5^2 + 2^2; \quad 8^2 + 5^2 + 2^2 \\
& \quad = 121; \quad 100, 125; \quad 106, 110; \quad 93.
\end{aligned}$$

The conservative cap-74 row is the same identity $7^2 + 5^2 = 74$.

The six lower inventories similarly give, in their printed order,

$$(6^2 + 3^2, 6^2 + 3^2 + 1, 7^2 + 3^2 + 1, 7^2 + 4^2 + 1, 7^2 + 4^2 + 1 + 3, 7^2 + 5^2 + 1 + 3) = (45, 46, 59, 66, 69, 78).$$

Finally, the ten lower- d caps are the successive odd sums

channels	cap	channels	cap
1	1	1 + 3	4
1 + 3 + 5	9	1 + 3 + 5 + 1	10
1 + 3 + 5 + 7	16	1 + 3 + 5 + 7 + 1	17
1 + 3 + 5 + 7 + 9 + 1	26	1 + 3 + 5 + 7 + 9 + 1 + 3	29
1 + 3 + 5 + 7 + 9 + 11 + 1 + 3	40	1 + 3 + 5 + 7 + 9 + 11 + 1 + 3 + 5	45.

3.3 The symbolic optical product deficit

Proposition 3.4 (Product-deficit identity and uniform reserve). *Let $N \geq 1$ be an integer, $0 < t \leq 1$, and $x = N + t$. Then*

$$\frac{2}{3}x^3 - Nx^2 + \sum_{n=1}^N n^2 = \frac{N^2}{2} + N \left(t^2 + \frac{1}{6} \right) + \frac{2t^3}{3}. \quad (6)$$

For $C_D = 1382/3125$, the right-hand side is strictly larger than $C_D(N + t)^2$. More precisely, its difference from $C_D(N + t)^2$ is at least

$$\frac{1822532}{91552734375} > 0. \quad (7)$$

Proof. Insert $\sum_{n=1}^N n^2 = N(N+1)(2N+1)/6$ and expand; this proves (6) for all N, t , rather than at sampled values. Put $C = C_D$, $A = 1/2 - C = 361/6250$, and let $G_N(t)$ denote the difference under consideration. Then

$$G_N(t) = AN^2 + N \left(t^2 - 2Ct + \frac{1}{6} \right) + \frac{2t^3}{3} - Ct^2.$$

Consequently

$$G_{N+1}(t) - G_N(t) = A(2N+1) + t^2 - 2Ct + \frac{1}{6}.$$

Since $0 < C < 1$, the last quadratic has minimum $1/6 - C^2$ on $(0, 1]$. For $N \geq 1$,

$$G_{N+1}(t) - G_N(t) \geq 3A + \frac{1}{6} - C^2 = \frac{4229603}{29296875} > 0.$$

It remains to minimize G_1 . Direct differentiation gives

$$G_1'(t) = 2(t - C)(t + 1),$$

so its unique minimum on $0 < t \leq 1$ is at $t = C$. Exact substitution gives

$$G_1(C) = \frac{1822532}{91552734375} > 0,$$

which proves both assertions. $\square$

3.4 Monotonic propagated cuts

Lemma 3.5 (Zero-cut/channel-cut monotonicity). *Fix $n \geq 2$, a checkpoint m , and $M = m(m+1)$. Let $\mathcal{A}_n$ be a finite collection of anchor pairs $(j, q_{j,n}^2)$, where $j_{j+1/2,n}^2 > q_{j,n}^2$ at each anchor. For every ℓ above at least one anchor, set*

$$\mathfrak{B}_n(\ell) := \max_{(j, q_{j,n}^2) \in \mathcal{A}_n, j \leq \ell} (q_{j,n}^2 + \ell(\ell+1) - j(j+1)), \quad (8)$$

$$\mathfrak{Z}_{m,n}(\ell) := \mathfrak{B}_n(\ell) - M, \quad (9)$$

$$\mathfrak{C}_{m,n}(\ell) := \frac{M - \ell(\ell+1)}{n^2 - 1}. \quad (10)$$

Then

$$\mathfrak{Z}_{m,n}(\ell+1) \geq \mathfrak{Z}_{m,n}(\ell) + 2(\ell+1) > \mathfrak{Z}_{m,n}(\ell), \quad (11)$$

$$\mathfrak{C}_{m,n}(\ell+1) = \mathfrak{C}_{m,n}(\ell) - \frac{2(\ell+1)}{n^2 - 1} < \mathfrak{C}_{m,n}(\ell). \quad (12)$$

In particular, if $\mathfrak{Z}_{m,n}(\ell_0) \geq \mathfrak{C}_{m,n}(\ell_0)$, then no channel (ℓ, n) with $\ell \geq \ell_0$ can satisfy the strict proxy (1) at $K = k_m$.

Proof. Every old candidate in (8) gains exactly $(\ell+1)(\ell+2) - \ell(\ell+1) = 2(\ell+1)$ when ℓ is replaced by $\ell+1$; a new anchor can only increase the maximum. This proves (11). Formula (12) is direct.

If a channel were counted at k_m , its product part would give

$$(n^2 - 1)z^2 < M - \ell(\ell+1), \quad \text{hence} \quad z^2 < \mathfrak{C}_{m,n}(\ell),$$

whereas its propagated ball-zero part would give

$$\mathfrak{B}_n(\ell) < z^2 + M, \quad \text{hence} \quad z^2 > \mathfrak{Z}_{m,n}(\ell).$$

These strict inequalities are incompatible when $\mathfrak{Z} \geq \mathfrak{C}$. Equations (11) and (12) propagate the incompatibility to every larger angular index. $\square$

4 Packet B: moving-3z ordering

We first record rational radical enclosures. All six follow by positive squaring; the exact square margins are

lower comparison	margin	upper comparison	margin
$(183/10)^2 < 337$	211/100	$337 < (92/5)^2$	39/25
$(1051/100)^2 < 7073/64$	2221/40000	$7073/64 < (1052/100)^2$	6191/40000
$(1067/100)^2 < 114$	1511/10000	$114 < (107/10)^2$	49/100.

Thus

$$\frac{183}{10} < \sqrt{337} < \frac{92}{5}, \quad \frac{1051}{100} < \frac{\sqrt{7073}}{8} < \frac{1052}{100}, \quad \frac{1067}{100} < \sqrt{114} < \frac{107}{10}. \quad (13)$$

Lemma 4.1 (Endpoint and fixed-wall order). *With $d = \sqrt{337}/2$,*

$$d < 3z_{\rho_0} < 10 < \frac{81}{8} < \frac{21}{2} < \frac{\sqrt{7073}}{8} < \sqrt{114} < 3z_{\rho_c}. \quad (14)$$

Moreover $(81/8)^2 + 8 = 7073/64$.

Proof. The nontrivial endpoint estimates have the following rational margins. From (2) and (13),

$$3z_{\rho_0} > 3\pi > \frac{999}{106}, \quad d < \frac{46}{5}, \quad \frac{999}{106} - \frac{46}{5} = \frac{119}{530} > 0.$$

Also $\rho_0 < 10/183$, whence

$$3z_{\rho_0} < \frac{66}{7} \frac{183}{173} = \frac{12078}{1211}, \quad 10 - \frac{12078}{1211} = \frac{32}{1211} > 0.$$

At the other endpoint, $z_{\rho_c} = \pi + 1/2$; therefore

$$3z_{\rho_c} > \frac{579}{53}, \quad \sqrt{114} < \frac{107}{10}, \quad \frac{579}{53} - \frac{107}{10} = \frac{119}{530} > 0.$$

The rational middle comparisons are immediate. For the two radical steps one may square positive quantities:

$$\frac{7073}{64} - \left(\frac{21}{2}\right)^2 = \frac{17}{64} > 0, \quad 114 - \frac{7073}{64} = \frac{223}{64} > 0.$$

Finally, $(81/8)^2 + 8 = (6561 + 512)/64 = 7073/64$. $\square$

For a positive wall w , the unique solution of $3z_\rho = w$ is

$$r_w := 1 - \frac{3\pi}{w}.$$

The following table encloses each crossing between explicit rational numbers. The bounds use only Lemma 2.1 and (13); for instance r_w increases with w and decreases with π .

wall w	exact rational enclosure for w	exact enclosure for r_w
10	10	$\frac{2}{35} < r_{10} < \frac{61}{1060}$
81/8	81/8	$\frac{13}{189} < r_{81/8} < \frac{11}{159}$
21/2	21/2	$\frac{5}{49} < r_{21/2} < \frac{38}{371}$
$\sqrt{7073}/8$	$1051/100 < w < 1052/100$	$\frac{73}{757} < r_w < \frac{2903}{27878}$
$\sqrt{114}$	$1067/100 < w < 107/10$	$\frac{79}{679} < r_w < \frac{676}{5671}$

The four gaps between consecutive rational enclosures are, in order,

$$\frac{2251}{200340}, \quad \frac{256}{7791}, \quad \frac{183}{389921}, \quad \frac{231225}{18929162}, \quad (15)$$

and are all positive. For example, the first is $13/189 - 61/1060$, and similarly for the other three. The endpoint margins are also explicit:

$$\rho_0 < \frac{10}{183} < \frac{2}{35}, \quad \frac{2}{35} - \frac{10}{183} = \frac{16}{6405} > 0,$$

and, since $\rho_c > 7/51$,

$$\frac{676}{5671} < \frac{7}{51} < \rho_c, \quad \frac{7}{51} - \frac{676}{5671} = \frac{5221}{289221} > 0.$$

We have therefore proved the complete strict order

$$\rho_0 < r_{10} < r_{81/8} < r_{21/2} < r_{\sqrt{7073}/8} < r_{\sqrt{114}} < \rho_c. \quad (16)$$

This single table replaces the ten crossing-location predicates and the four crossing-order predicates of the executable registry.

5 Packet B: checkpoint boundary table and exhaustiveness

The ball-zero input consists of the following strict rational lower bounds:

$$\frac{n=2, \ell}{q_{\ell,2}^2} \mid \begin{array}{cc} 1 & 2 \\ (77/10)^2 & 9^2 \end{array} \quad \frac{n=3, \ell}{q_{\ell,3}^2} \mid \begin{array}{cc} 3 & 5 \\ (103/10)^2 & (129/10)^2 \end{array} \quad \frac{n=3, \ell}{q_{\ell,3}^2} \mid \begin{array}{cc} 1 & 2 \\ (21/2)^2 & (61/5)^2 \end{array} \quad (17)$$

Angular propagation gives the lower bound (8) at all larger angular indices. The seven first-excluded channels have the exact cuts shown below. Here $M = m(m+1)$, $B = \mathfrak{B}_n(\ell_0)$, $Z = B - M$, $C = (M - \ell_0(\ell_0 + 1))/(n^2 - 1)$, and the last column is the decisive incompatibility margin $Z - C$.

m	n	first excluded ℓ_0	B	Z	C	$Z - C$
7	2	2	81	25	50/3	25/3
8	2	4	11409/100	4209/100	52/3	7427/300
9	2	5	16641/100	7641/100	20	5641/100
10	2	6	17841/100	6841/100	68/3	13723/300
11	2	5	16641/100	3441/100	34	41/100
10	3	2	3721/25	971/25	13	646/25
11	3	2	3721/25	421/25	63/4	109/100

Every table entry is reconstructible from (17). The only propagated base values not already anchors are

$$\frac{11409}{100} = \left(\frac{103}{10}\right)^2 + (4 \cdot 5 - 3 \cdot 4), \quad \frac{17841}{100} = \left(\frac{129}{10}\right)^2 + (6 \cdot 7 - 5 \cdot 6).$$

All seven margins are positive; the smallest is $41/100$, so none of these exclusions relies on an equality convention. By Lemma 3.5, each row excludes not merely the displayed channel but its entire angular tail.

It remains to rule out the radial tails. On the high strip $\rho_c \leq \rho < 1$,

$$z \geq z_{\rho_c} = \pi + \frac{1}{2} > \frac{333}{106} + \frac{1}{2} = \frac{193}{53}, \quad z^2 > \frac{37249}{2809}.$$

The exact margins needed below are

$$\frac{37249}{2809} - \frac{90}{8} = \frac{22591}{11236} > 0, \quad \frac{37249}{2809} - \frac{132}{15} = \frac{62649}{14045} > 0. \quad (18)$$

At $m \leq 9$, an $n = 3$ channel would require $8z^2 < M - \ell(\ell + 1) \leq 90$, contradicting the first inequality. At $m \leq 11$, every $n \geq 4$ channel would require $15z^2 \leq (n^2 - 1)z^2 < M - \ell(\ell + 1) \leq 132$, contradicting the second inequality. These are the two base radial exclusions.

Theorem 5.1 (Exhaustive checkpoint inventories). *For $\rho_c \leq \rho < 1$, the channels that can satisfy the strict proxy (1) at the five checkpoints are contained in the following lists:*

checkpoint	possible modes
k_7	first 0:6, second 0:1, no $n \geq 3$
k_8	first 0:7, second 0:3, no $n \geq 3$
k_9	first 0:8, second 0:4, no $n \geq 3$
k_{10}	first 0:9, second 0:5, third 0:1, no $n \geq 4$
k_{11}	first 0:10, second 0:4, third 0:1, no $n \geq 4$.

Here n -th means radial index n , and $a:b$ denotes the angular indices $a, a + 1, \dots, b$. The word “possible” asserts an exhaustive upper inventory, not that every listed channel is present for every ratio.

Proof. For $n = 1$, the product part of the strict proxy at k_m is

$$z^2 + \ell(\ell + 1) < z^2 + m(m + 1).$$

Thus $\ell < m$. The equality channel $\ell = m$ and all larger indices are excluded, with the equality ownership fixed by Lemma 3.2. This gives precisely first 0: $m - 1$.

For $n = 2$, apply the first five rows of the checkpoint table and then Lemma 3.5. They give first-excluded angular indices 2, 4, 5, 6, 5 at $m = 7, 8, 9, 10, 11$, respectively. Hence the second blocks end at 1, 3, 4, 5, 4.

For $n = 3$, the first inequality in (18) removes all channels through k_9 . At k_{10} and k_{11} , the last two rows of the checkpoint table and cut monotonicity remove every $\ell \geq 2$, leaving only $\ell = 0, 1$. Finally, the second inequality in (18) removes all $n \geq 4$ through k_{11} . The five lists follow, without any bounded angular search. $\square$

6 Executable-row coverage contract

The former executable baseline contained 611 rows and had canonical digest

afd7e054f52aa9a3992fc8ef16d3e7551586c76bfb589053714f5f55a36792f1.

The digest is included only to identify the audited baseline. It is not a premise of any proof above. The exhibits above replace disjoint row groups as follows:

analytic exhibit	rows	exact ledger subfamilies
Lemma 2.1	4	one enclosure-nonemptiness row, two rational π -bound rows, and one reciprocal row
Lemma 3.2	24	sixteen integer/half-integer strict brackets, seven phase-proxy integer walls, and one optical radial equality owner
Lemma 3.3	97	twelve strict angular equality blocks, twelve optical angular blocks, 62 high-checkpoint cap identities, one six-cap lower tuple, and ten lower- d cap identities
Proposition 3.4	23	five general product-deficit predicates and eighteen sampled expansions
Lemma 3.5, the checkpoint boundary table, and Theorem 5.1	117	76 repeated cut-monotonicity rows, seven first-excluded comparisons, two radial exclusions, and 32 metadata/tail instances
Lemma 4.1 and the moving-crossing table	23	five fixed-wall orders, one propagation identity, three endpoint comparisons, ten crossing locations, and four crossing-order rows
total	288	

The following family table gives the same accounting against the executable entry points and makes the partial-family boundaries explicit. It maps the exact rows replaced by this note. “Collapsed” means that multiple sampled or bookkeeping rows are consequences of one displayed identity; it does not mean that an unprinted program enumeration is being used.

executable family	family rows	replaced here	analytic replacement
<code>verify_pi</code>	4	4	Lemma 2.1
<code>verify_thresholds</code>	34	23	five fixed-wall orders, one propagation identity, three endpoint comparisons, and fourteen moving-crossing predicates
<code>verify_strict_face_conventions</code>	35	35	Lemmas 3.2 and 3.3
<code>verify_lower_inventories_and_payments</code>	44	1	single six-cap tuple bookkeeping row
<code>verify_checkpoint_inventories</code>	117	117	boundary table, radial margins, and Theorem 5.1
<code>verify_cap_tables_and_payments</code>	98	62	all block-cap arithmetic: 61 cap-table rows plus the cap-74 block identity
<code>verify_optical_constants</code>	50	36	twelve angular blocks, one radial equality owner, five general product-deficit facts, and eighteen sampled expansions
supplemental k_6 /lower- d ledger	26	10	ten lower- d block-cap identities
total		288	

For the checkpoint family, the 117 rows decompose exactly as follows: 32 finite metadata/tail instances, seven first-excluded zero/channel comparisons, two base radial exclusions, and 76 repeated cut-monotonicity instances. Theorem 5.1 replaces all four groups. For the optical family, Proposition 3.4 replaces the 18 sampled (N, t) expansions by a symbolic identity valid for every integer $N \geq 1$ and every $0 < t \leq 1$.

The remaining 323 rows are outside Packets A and B. In particular, this note makes no claim to replace the high-event cap/payment implications, the Bessel sign registry, seam payments, optical screen constants, or the exact owner subtraction. Those require the other analytic replacement packets. This separation prevents a universal arithmetic lemma from being mistaken for proof of a geometric event assignment.

Connected High-Event Registry

Analytic ledger replacement, Packet C

Abstract

This note replaces the exact-executable arithmetic used in the high spherical-shell staircase by displayed rational witnesses. It contains the complete seventeen-face localization registry, the three algebraic splits, the three dashed (impossible) cap cells, the five global checkpoint payments, the seven conditional payments, and the corrected cap-74 path. It also contains the order-free closed-post-event theorem which maps every high-staircase state to a payment at its own event wall. That theorem, rather than the historical executable row census, is the finite-combinatorial closure used by the proof. All comparisons are strict rational comparisons after the two elementary bounds $333/106 < \pi < 22/7$. In particular, no decimal approximation, interval package, or external ledger is a logical input to this packet.

1 Definitions and strict proxy convention

For $0 < \rho < 1$, put

$$z = z_\rho := \frac{\pi}{1 - \rho}, \quad x = x_\rho := z_\rho^2, \quad k_m(\rho) := \sqrt{x_\rho + m(m + 1)}. \quad (1)$$

The three-dimensional Weyl term is

$$\mathcal{W}(\rho, K) := \frac{2}{9\pi}(1 - \rho^3)K^3. \quad (2)$$

At $K = k_m(\rho)$, the product part of the strict channel proxy for the mode (ℓ, n) , $n \geq 2$, is

$$n^2x + \ell(\ell + 1) < x + m(m + 1).$$

It is therefore governed by

$$\Delta_{m;n,\ell}(\rho) := m(m + 1) - \ell(\ell + 1) - (n^2 - 1)x_\rho. \quad (3)$$

The channel is product-possible only when $\Delta_{m;n,\ell} > 0$. Equality is excluded because the spectral proxy is strict. Since

$$x'_\rho = \frac{2\pi^2}{(1 - \rho)^3} > 0, \quad \Delta'_{m;n,\ell} = -(n^2 - 1)x'_\rho < 0, \quad (4)$$

a positive witness at r controls the side $\rho \leq r$, while a negative witness controls the side $\rho \geq r$.

We write H for the largest first-mode angular index retained in a cap cell and h for the largest second-mode angular index. A complete block through H has multiplicity $(H + 1)^2$, because $\sum_{\ell=0}^H (2\ell + 1) = (H + 1)^2$.

2 The only transcendental enclosure

Lemma 1 (Rational enclosure of π). *One has*

$$\pi_- := \frac{333}{106} < \pi < \frac{22}{7} =: \pi_+. \quad (5)$$

Proof. For $0 < u < 1$, the alternating expansion of $\arctan u$ lies between each pair of consecutive partial sums. The tangent addition formula gives

$$\tan\left(4 \arctan \frac{1}{5} - \arctan \frac{1}{239}\right) = 1;$$

indeed, $\tan(4 \arctan(1/5)) = 120/119$, so both the numerator and denominator in the last tangent quotient are $28561/(119 \cdot 239)$. The angle lies in $(0, \pi/2)$, and hence Machin's identity follows:

$$\pi = 16 \arctan \frac{1}{5} - 4 \arctan \frac{1}{239}.$$

Using an even partial sum for the positive term and an upper partial sum for the subtracted term gives

$$\begin{aligned} \pi &> 16 \left(\frac{1}{5} - \frac{1}{3 \cdot 5^3} + \frac{1}{5 \cdot 5^5} - \frac{1}{7 \cdot 5^7} \right) - \frac{4}{239}, \\ [\text{right side}] - \frac{333}{106} &= \frac{3418213}{41563593750} > 0. \end{aligned}$$

Likewise,

$$\begin{aligned} \pi &< 16 \left(\frac{1}{5} - \frac{1}{3 \cdot 5^3} + \frac{1}{5 \cdot 5^5} \right) - 4 \left(\frac{1}{239} - \frac{1}{3 \cdot 239^3} \right), \\ \frac{22}{7} - [\text{right side}] &= \frac{1845738322}{1493178640625} > 0. \end{aligned}$$

This proves (5). □

3 The seventeen rational localization faces

Let

$$\rho_c := \frac{1}{1 + 2\pi}.$$

Because $\pi > 3$, $\rho_c < 1/7$. The seventeen registry ratios are strictly ordered as follows:

$$\begin{aligned} \frac{4}{25} &< \frac{1}{5} < \frac{1}{4} < \frac{3}{10} < \frac{1}{3} < \frac{7}{20} < \frac{3}{8} < \frac{2}{5} < \frac{5}{12} < \frac{3}{7} < \frac{1}{2} \\ &< \frac{8}{15} < \frac{107}{200} < \frac{3}{5} < \frac{16}{25} < \frac{13}{20} < \frac{2}{3}. \end{aligned} \quad (6)$$

For a completely explicit distinctness check, the sixteen consecutive gaps in (6) are

$$\frac{1}{25}, \frac{1}{20}, \frac{1}{20}, \frac{1}{30}, \frac{1}{60}, \frac{1}{40}, \frac{1}{40}, \frac{1}{60}, \frac{1}{84}, \frac{1}{14}, \frac{1}{30}, \frac{1}{600}, \frac{13}{200}, \frac{1}{25}, \frac{1}{100}, \frac{1}{60}. \quad (7)$$

Moreover

$$\frac{4}{25} - \frac{1}{7} = \frac{3}{175} > 0, \quad \frac{7}{8} - \frac{2}{3} = \frac{5}{24} > 0.$$

Thus all seventeen ratios lie in the high strip $\rho_c < \rho < 7/8$.

For a positive row below, we use

$$\Delta_{m;n,\ell}(r) > m(m+1) - \ell(\ell+1) - (n^2 - 1) \left(\frac{\pi_+}{1-r} \right)^2.$$

For a negative row, we use

$$-\Delta_{m;n,\ell}(r) > (n^2 - 1) \left(\frac{\pi_-}{1-r} \right)^2 - m(m+1) + \ell(\ell+1).$$

The resulting exact witnesses are the following. “Retain” means only that the product wall does not permit deletion of the indicated block; the independent Bessel wall may still remove it.

no.	r	$(m; n, \ell)$	exact sign witness	one-sided cap/event consequence
1	1/5	(11; 3, 1)	$\Delta > 320/49$	retain the third block 0:1 for $\rho \leq 1/5$
2	3/10	(8; 2, 3)	$-\Delta > 58215/137641$	at k_8 , $h \leq 2$ for $\rho \geq 3/10$
3	1/3	(8; 2, 2)	$-\Delta > 27699/44944$	at k_8 , $h \leq 1$ for $\rho \geq 1/3$
4	3/8	(9; 2, 3)	$\Delta > 2622/1225$	retain the $h = 3$ column below this face
5	2/5	(9; 2, 2)	$\Delta > 248/147$	retain the $h = 2$ column below this face
6	5/12	(9; 2, 1)	$\Delta > 2200/2401$	retain the $h = 1$ column below this face
7	3/7	(9; 2, 0)	$-\Delta > 120843/179776$	no second mode at k_9 on the upper side
8	1/2	(10; 2, 0)	$-\Delta > 23677/2809$	no second mode at k_{10} on the upper side
9	107/200	(11; 2, 0)	$-\Delta > 13302732/2699449$	no second mode at k_{11} on the upper side
10	8/15	(11; 2, 0)	$-\Delta > 2175627/550564$	the clean stated cutoff: a second mode forces $\rho < 8/15$
11	3/5	(8; 2, 0)	$-\Delta > 5080707/44944$	select the no-second k_8 cap/payment side
12	16/25	(9; 2, 0)	$-\Delta > 1555635/11236$	select the no-second k_9 cap/payment side
13	13/20	(10; 2, 0)	$-\Delta > 18126190/137641$	select the no-second k_{10} cap/payment side
14	2/3	(11; 2, 0)	$-\Delta > 1510851/11236$	select the no-second k_{11} cap/payment side
15	4/25	(10; 3, 0)	$-\Delta > 260740/137641$	no third mode at k_{10} on the upper side
16	1/4	(11; 3, 0)	$-\Delta > 23484/2809$	a third mode at k_{11} forces $\rho < 1/4$
17	7/20	(9; 2, 4)	$-\Delta > 36230/474721$	the $H = 6, h = 4$ cap-74 cell can occur only below this face

Every entry is a displayed positive rational number. Together with the monotonicity (4), the table proves the asserted side on an interval, not merely at a sampled point.

4 Three algebraic splits and the five k_{11} implications

At $\rho = \rho_c$, $x = (\pi + 1/2)^2$, while at $\rho = 7/8$, $x = 64\pi^2$. Lemma 1 gives the exact chain

$$(\pi + \frac{1}{2})^2 < \left(\frac{51}{14}\right)^2 = \frac{2601}{196} < 16 < \frac{68}{3} < 34 < 64 \cdot 3^2 < 64\pi^2. \quad (8)$$

Consequently each of $x = 16, 68/3, 34$ is attained at a unique point in the high strip. The relevant channel arithmetic is

x	channel	exact value of Δ
16	$(m; n, \ell) = (11; 3, 1)$	$132 - 2 - 8 \cdot 16 = 2 > 0$,
68/3	$(10; 2, 6)$	$110 - 42 - 3(68/3) = 0$,
34	$(11; 2, 5)$	$132 - 30 - 3 \cdot 34 = 0$.

(9)

Thus the first split lies on the possible side. The latter two are exact channel walls, and their new modes are excluded by the strict convention.

The two localization consequences at k_{11} may also be read without the table:

$$x_{8/15} - 44 > \left(\frac{\pi_-}{1 - 8/15}\right)^2 - 44 = \frac{725209}{550564} > 0, \quad (10)$$

$$x_{1/4} - \frac{33}{2} > \left(\frac{\pi_-}{1 - 1/4}\right)^2 - \frac{33}{2} = \frac{5871}{5618} > 0. \quad (11)$$

Hence a second mode at k_{11} forces $\rho < 8/15$, and a third mode forces $\rho < 1/4$. Finally, the first-mode ball walls give

$$\left(\frac{69}{5}\right)^2 - 132 = \frac{1461}{25} = 44 + \frac{361}{25} = \frac{33}{2} + \frac{2097}{50}, \quad (12)$$

$$\left(\frac{64}{5}\right)^2 - 132 = \frac{796}{25} = \frac{33}{2} + \frac{767}{50}. \quad (13)$$

If $H = 10$ is present at k_{11} , then x is larger than both the second threshold 44 and the third threshold $33/2$; hence every second and third mode is excluded. If $H = 9$ is present, every third mode is excluded. Equations (10)–(13) are the five formerly separate k_{11} localization implications.

5 The three impossible cap cells

If the first mode H is counted at k_m , its strict ball wall $\beta_{H,1} < k_m$ forces $x > \beta_{H,1}^2 - m(m+1)$. If the second mode h is counted, its product wall forces $x < (m(m+1) - h(h+1))/3$. The three dashed cells are therefore empty by the following exact, incompatible bounds:

cell	required lower bound	required upper bound	gap
$k_8 : H = 6, h \geq 2$	$100 - 72 = 28$	$(72 - 6)/3 = 22$	6
$k_9 : H = 8, h \geq 0$	$(71/6)^2 - 90 = 1801/36$	$90/3 = 30$	$721/36$
$k_9 : H = 7, h \geq 3$	$(23/2)^2 - 90 = 169/4$	$(90 - 12)/3 = 26$	$65/4$.

(14)

No limiting equality can rescue a cell: both necessary modal inequalities are strict.

6 Checkpoint monotonicity and five global payments

Put $M = m(m+1)$ and $\mathcal{W}_m(\rho) := \mathcal{W}(\rho, k_m(\rho))$. Logarithmic differentiation shows that $\mathcal{W}'_m$ has the sign of

$$\pi^2(1+\rho) - M\rho^2(1-\rho)^2. \quad (15)$$

For $m \leq 11$,

$$\pi^2(1+\rho) - M\rho^2(1-\rho)^2 > 9 - \frac{132}{16} = \frac{3}{4} > 0.$$

Thus every $\mathcal{W}_m$, $m \leq 11$, is strictly increasing. This single argument replaces both executable derivative-bound predicates.

At ρ_c , one has $z = \pi + 1/2 > 193/53$, $\rho_c < 1/7$, and $1/\pi > 7/22$. Consequently

$$\mathcal{W}(\rho_c, K) > \frac{2}{9} \frac{7}{22} \left(1 - \frac{1}{7^3}\right) K^3 = \frac{38}{539} K^3. \quad (16)$$

For each row in the next table, the middle column proves $k_m(\rho_c) > t_m$, and the last column proves $\mathcal{W}_m(\rho_c) > C_m$.

m	cap C_m	t_m	$(193/53)^2 + M - t_m^2$	$(38/539)t_m^3 - C_m$
7	40	208/25	67049/1755625	5083656/8421875
8	53	923/100	1901439/28090000	656778873/269500000
9	73	254/25	61431/1755625	7911557/8421875
10	89	111/10	14211/280900	1999489/269500
11	121	241/20	65271/1123600	5076899/2156000

Every last-column entry is positive. Monotonicity therefore extends each global payment from ρ_c to the complete high strip.

7 Seven conditional first-block payments

We now give a general reduction which avoids any numerical enclosure of a radical. Suppose the presence of first mode H below the checkpoint k_m forces $K > t$. Set

$$a^2 := t^2 - m(m+1), \quad \rho_0 := 1 - \frac{\pi}{a}. \quad (17)$$

All seven rows below have $a > \pi$: the smallest listed radicand is

$$\frac{89}{4}, \quad \frac{89}{4} - \left(\frac{22}{7}\right)^2 = \frac{2425}{196} > 0.$$

If the mode is present for some $K \leq k_m(\rho)$, then $k_m(\rho) > t$, hence $x_\rho > a^2$ and $\rho > \rho_0$. By the strict increase of $\mathcal{W}_m$,

$$\mathcal{W}_m(\rho) > \mathcal{W}_m(\rho_0) = \mathcal{W}(\rho_0, t).$$

Choose a rational $r > \rho_0$. Since $\mathcal{W}(\rho, t)$ decreases with ρ at fixed t , Lemma 1 yields

$$\mathcal{W}(\rho_0, t) > \mathcal{W}(r, t) > \frac{7}{99}(1-r^3)t^3. \quad (18)$$

The condition $r > \rho_0$ follows, without taking a square root, from

$$L := \pi_-^2 - (1-r)^2 a^2 > 0. \quad (19)$$

Here is the complete calculation. The column P is the reserve in (18) above the stated cap.

m	H	t	cap	a^2	r	L	$P = \frac{7}{99}(1-r^3)t^3 - \text{cap}$
7	6	10	53	44	8/15	725209/2528100	18659/2673
8	7	23/2	80	241/4	3/5	64349/280900	213281/49500
9	8	71/6	81	1801/36	3/5	4713989/2528100	14506973/1336500
10	7	23/2	100	89/4	7/20	2105431/4494400	18539033/6336000
10	8	71/6	97	1081/36	3/7	317585/4955076	2865431/261954
10	9	64/5	100	1346/25	3/5	8811001/7022500	25143284/1546875
11	9	64/5	125	796/25	1/2	536261/280900	58757/12375

All L 's and P 's are positive. Each row therefore proves both the localization of its conditional branch and its Weyl payment.

8 The connected cap-74 path

The seventeenth localization row is tied to the exceptional $k_9, H = 6, h = 4$ cap as follows. Angular propagation from the strict $(3, 2)$ Bessel wall gives

$$j_{9/2,2}^2 > \left(\frac{103}{10}\right)^2 + 20 - 12 = \frac{11409}{100} = 108 + \frac{609}{100}. \quad (20)$$

Accordingly the strict proxy uses $\beta_{4,2}^2 = 11409/100$; the strict inequality belongs to the actual Bessel zero $j_{9/2,2}$, not to the proxy constant. At k_9 , the product wall for $(4, 2)$ is

$$3x < 90 - 20 = 70, \quad \text{i.e.} \quad x < \frac{70}{3}.$$

The ball wall (20), on the other hand, forces

$$x > \frac{11409}{100} - 90 = \frac{2409}{100}, \quad \frac{2409}{100} - \frac{70}{3} = \frac{227}{300} > 0.$$

Thus the fully sharpened cap-74 cell is empty.

For completeness, even the deliberately conservative cell is paid. At $\rho = 7/20$,

$$x_{7/20} - \frac{70}{3} > \frac{400}{169} \left(\frac{333}{106}\right)^2 - \frac{70}{3} = \frac{36230}{1424163} > 0, \quad (21)$$

so a counted $(4, 2)$ mode forces $\rho < 7/20$. Also

$$108 - \left(\frac{207}{20}\right)^2 = \frac{351}{400} > 0, \quad (22)$$

so its ball wall forces $K > 207/20$. Therefore

$$\begin{aligned} \mathcal{W}(\rho, K) &> \frac{7}{99} \left(1 - \left(\frac{7}{20}\right)^3\right) \left(\frac{207}{20}\right)^3 \\ &= \frac{52823261673}{704000000} = 74 + \frac{727261673}{704000000} > 74. \end{aligned} \quad (23)$$

The cap itself is exactly $(6+1)^2 + (4+1)^2 = 49 + 25 = 74$. Equations (20)–(23) cover both the actual incompatibility and every row of the conservative cap-74 fallback.

9 Order-free exhaustion and payment of every high event

This section supplies the finite-combinatorial implication which is not contained in a list of checkpoint payments alone. It is deliberately independent of the order in which two modal walls cross.

For $7 \leq m \leq 11$, set

$$\mathfrak{B}_m^+ := \{(\rho, K) : \rho_c \leq \rho \leq 39/50, k_{m-1}(\rho) \leq K < k_m(\rho)\}. \quad (24)$$

The final spectral band is closed on the right:

$$\mathfrak{B}_{11}^{\text{sp}} := \{(\rho, K) : \rho_c \leq \rho \leq 39/50, k_{10}(\rho) \leq K \leq k_{11}(\rho)\}.$$

At $K = k_{11}$ we use the original strict proxy. On the half-open event bands we use

$$b_{\ell,n}(z)^2 = \max\{n^2 z^2 + \ell(\ell+1), \beta_{\ell,n}^2\}, \quad C_z(K) = \sum_{\ell,n} (2\ell+1) \mathbf{1}_{\{b_{\ell,n}(z) < K\}},$$

and define the closed post-event count

$$\hat{C}_z(K) := \sum_{(\ell,n)} (2\ell+1) \mathbf{1}_{\{b_{\ell,n}(z) \leq K\}}. \quad (25)$$

The sum is over the checkpoint inventory appropriate to the upper endpoint of the band. Thus a simultaneous event is charged with all of its entering multiplicities before it is paid.

For a post-event state let H be the largest first-channel angular index, let $h = -1$ if no second channel is present and otherwise let h be its largest angular index, and put $\tau = 1$ if a third channel is present and $\tau = 0$ otherwise. Missing angular indices only lower the count, since

$$\hat{C}_z(K) \leq (H+1)^2 + \mathbf{1}_{\{h \geq 0\}}(h+1)^2 + 4\tau. \quad (26)$$

The exhaustive checkpoint inventories are

m	H	h	third channels
7	$H \leq 6$	$h \leq 1$	none
8	$H \leq 7$	$h \leq 3$	none
9	$H \leq 8$	$h \leq 4$	none
10	$H \leq 9$	$h \leq 5$	$\ell = 0, 1$
11	$H \leq 10$	$h \leq 4$	$\ell = 0, 1.$

(27)

First channels with $\ell \geq m$ are excluded at k_m by their product wall. For the first omitted second channel, the two necessary strict bounds are

m	7	8	9	10	11
first omitted ℓ	2	4	5	6	5
product wall	$x < 50/3$	$x < 52/3$	$x < 20$	$x < 68/3$	$x < 34$
ball wall	$x > 25$	$x > 4209/100$	$x > 7641/100$	$x > 6841/100$	$x > 3441/100.$

The $m = 10$ ball wall follows from angular propagation of the $\ell = 5, n = 2$ registry wall. In every column the bounds are incompatible; for larger ℓ , the product upper bound decreases and the propagated ball lower bound increases.

For $n = 3$, the product wall gives $x \leq 45/4 < (193/53)^2 < x$ through $m = 9$, and at $m = 10$ every $\ell \geq 2$ gives $x \leq 13 < (193/53)^2$. At $m = 11$, the first omitted third channel $\ell = 2$ would require

$$x < 63/4, \quad x > (61/5)^2 - 132 = 421/25.$$

Angular propagation removes the rest of that tail. For $n \geq 4$, the least restrictive omitted case is $m = 11, \ell = 0, n = 4$, which would require $x < 44/5 < (193/53)^2$. This proves (27) without an unprinted search.

The strict simultaneous restrictions used below are

m	restriction
8	$H = 6 \implies h \leq 1$, $H = 7$ unrestricted by this row,
9	$H = 8 \implies h = -1$, $H = 7 \implies h \leq 2$,
10	$H = 9 \implies h = -1, \tau = 0$, $H = 8 \implies h \leq 3, \tau = 0$, $H = 7 \implies \tau = 0$,
11	$H = 10 \implies h = -1, \tau = 0$, $H = 9 \implies \tau = 0$.

(28)

For $m = 8, 9$ these are exactly the incompatible cells in (14). For $m = 10$, the three first-wall lower bounds are $1346/25, 1081/36, 89/4$, which respectively exceed the applicable second or third thresholds $110/3, 30, 55/4$. For $m = 11$, (12)–(13) give the last two restrictions. All gaps are strict, so the same restrictions hold for the closed post-event count on the half-open bands.

9.1 The fourteen localization witnesses

If a closed post-event second mode h is present below k_m , then

$$4x + h(h + 1) \leq K^2 < x + m(m + 1), \quad 3x < m(m + 1) - h(h + 1).$$

At the strict spectral upper face the first inequality is strict and the same conclusion follows. Likewise, a third mode forces $8x < m(m + 1)$. The following table contains every localization used in the state table. Its last column is $(333/(106(1 - r)))^2 - A > 0$, where $x < A$ is the forced bound.

band	state implication	A	r	exact positive witness
$\mathfrak{B}_7^+$	some second channel	56/3	3/10	608779/412923
$\mathfrak{B}_8^+$	some second channel	24	3/8	88824/70225
$\mathfrak{B}_8^+$	$h \geq 2$	22	1/3	9233/44944
$\mathfrak{B}_8^+$	$h = 3$	20	3/10	19405/137641
$\mathfrak{B}_9^+$	some second channel	30	3/7	40281/179776
$\mathfrak{B}_9^+$	$h \geq 2$	28	5/12	138056/137641
$\mathfrak{B}_9^+$	$h \geq 3$	26	2/5	15889/11236
$\mathfrak{B}_9^+$	$h = 4$	70/3	7/20	36230/1424163
$\mathfrak{B}_{10}^+$	some second channel	110/3	1/2	23677/8427
$\mathfrak{B}_{10}^+$	$h \geq 4$	30	3/7	40281/179776
$\mathfrak{B}_{10}^+$	$h = 5$	80/3	2/5	25195/33708
$\mathfrak{B}_{10}^+$	$\tau = 1$	55/4	4/25	65185/275282
$\mathfrak{B}_{11}^+$	some second channel	44	8/15	725209/550564
$\mathfrak{B}_{11}^+$	$\tau = 1$	33/2	1/4	5871/5618

Every defining ratio face is therefore on the excluded side of its state; the directions in this table are interval implications, not sampled decisions.

9.2 Three payment rules

Put $F_j(\rho) := \mathcal{W}(\rho, k_j(\rho))$, and define

$$P(C; c, r) := \frac{7}{99}(1 - r^3)c^3 - C, \quad Q_j(c, r) := \left(\frac{333}{106(1 - r)} \right)^2 + j(j + 1) - c^2. \quad (29)$$

The derivative calculation in (15) gives, uniformly for $6 \leq j \leq 10$,

$$F'_j(\rho) > 0, \quad 9 - \frac{110}{16} = \frac{17}{8} > 0,$$

and $\mathcal{W}(\rho, c)$ decreases with ρ at fixed $c > 0$. Together with $2/(9\pi) > 7/99$, this gives three rules.

L. If a state forces $K \geq c$ and $\rho < r$, then $P(C; c, r) > 0$ implies $\mathcal{W}(\rho, K) > C$.

B. If a state forces $K \geq c$ in $\mathfrak{B}_m^+$, and $P(C; c, r) > 0$, $Q_{m-1}(c, r) > 0$, then it is paid on both sides of r . For $\rho \leq r$, use the fixed wall c . For $\rho \geq r$, use $K \geq k_{m-1}(\rho)$, the increase of F_{m-1} , and $k_{m-1}(r) > c$. In particular, $F_{m-1}(r) = \mathcal{W}(r, k_{m-1}(r)) > \mathcal{W}(r, c) > C$. Both arguments are strict at $\rho = r$.

G. If the state cap is at most a displayed global cap C , then $K \geq k_{m-1}(\rho)$ and the strict global lower-checkpoint payment pays it, including the lower checkpoint itself.

There is one initial checkpoint anchor of type B_0 . Since $\pi < 7/2$, one has $\rho_c > 1/8$. The increase of F_6 , together with the $B_{7,0}$ row below, gives

$$F_6(\rho) > F_6(1/8) > \mathcal{W}(1/8, 73/10) > 26.$$

9.3 The disjoint thirty-eight-state map

The following rows are literally disjoint. In a G -row the displayed state is the exact complement of the exceptional rows in its band, not a second overlapping estimate. The cap column is allowed to dominate more than one exact block sum, but every exact post-event cap is obtained from (26) and is one of the block sums represented here.

row	band	disjoint post-event state	cap C	c	r
$B_{7,0}$	7	$H \leq 4, -1 \leq h \leq 0$	26	73/10	1/8
$L_{7,1}$	7	$H \leq 4, h = 1$	29	77/10	3/10
$B_{7,1}$	7	$H = 5, h = -1$	36	87/10	1/2
$L_{7,2}$	7	$H = 5, 0 \leq h \leq 1$	40	87/10	3/10
$B_{7,2}$	7	$H = 6, h = -1$	49	10	3/5
$L_{7,3}$	7	$H = 6, 0 \leq h \leq 1$	53	10	3/10
G_8	8	$H \leq 4, h \leq 2$, or $H = 5, h \leq 1$	40	—	—
$L_{8,1}$	8	$H = 5, h = 2$	45	9	1/3
$L_{8,2}$	8	$H \leq 5, h = 3$	52	103/10	3/10
$B_{8,1}$	8	$H = 6, h = -1$	49	10	3/5
$L_{8,3}$	8	$H = 6, 0 \leq h \leq 1$	53	10	3/8
$B_{8,2}$	8	$H = 7, h = -1$	64	23/2	2/3
$L_{8,4}$	8	$H = 7, 0 \leq h \leq 3$	80	23/2	3/8
G_9	9	$H \leq 4$; or $H = 5, h \leq 3$; or $H = 6, h \leq 1$	53	—	—
$L_{9,1}$	9	$H = 6, h = 2$	58	10	5/12
$L_{9,2}$	9	$H = 6, h = 3$	65	103/10	2/5
$L_{9,3}$	9	$5 \leq H \leq 6, h = 4$	74	207/20	7/20
$B_{9,1}$	9	$H = 7, h = -1$	64	23/2	2/3
$L_{9,4}$	9	$H = 7, 0 \leq h \leq 2$	73	23/2	3/7
$B_{9,2}$	9	$H = 8, h = -1$	81	71/6	2/3
G_{10}	10	$H \leq 4$; or $H = 5$ except $(h, \tau) = (5, 1)$; or $H = 6, h \leq 3$; or $H = 7, h \leq 2$	73	—	—
$L_{10,1}$	10	$H = 6, h = 4, \tau = 0$	74	53/5	3/7
$L_{10,2}$	10	$H = 6, h = 4, \tau = 1$	78	53/5	4/25
$L_{10,3}$	10	$H = 6, h = 5, \tau = 0$	85	11	2/5
$L_{10,4}$	10	$5 \leq H \leq 6, h = 5, \tau = 1$	89	11	4/25

row	band	disjoint post-event state	cap C	c	r
$L_{10,5}$	10	$H = 7, h = 3$	80	23/2	1/2
$L_{10,6}$	10	$H = 7, h = 4$	89	23/2	3/7
$L_{10,7}$	10	$H = 7, h = 5$	100	23/2	2/5
$B_{10,1}$	10	$H = 8, h = -1$	81	71/6	3/5
$L_{10,8}$	10	$H = 8, 0 \leq h \leq 3$	97	71/6	1/2
$B_{10,2}$	10	$H = 9, h = -1$	100	64/5	16/25
G_{11}	11	$H \leq 6$; or $H = 7$ except $(h, \tau) = (4, 1)$; or $H = 8, h \leq 1$	89	–	–
$L_{11,1}$	11	$H = 7, h = 4, \tau = 1$	93	23/2	1/4
$B_{11,1}$	11	$H = 8, 2 \leq h \leq 4, \tau = 0$	106	71/6	3/7
$L_{11,2}$	11	$H = 8, 2 \leq h \leq 4, \tau = 1$	110	71/6	1/4
$B_{11,2}$	11	$H = 9, h = -1$	100	64/5	3/5
$L_{11,3}$	11	$H = 9, 0 \leq h \leq 4$	125	64/5	8/15
$B_{11,3}$	11	$H = 10, h = -1$	121	69/5	13/20

Apart from the checkpoint anchor $B_{7,0}$, the wall c in each exceptional row is forced by the corresponding fixed part of $b_{\ell,n}$. Besides the displayed first and second walls, the only weakenings are

$$\beta_{4,2}^2 - (207/20)^2 = 2787/400 > 0, \quad \beta_{4,2}^2 - (53/5)^2 = 173/100 > 0, \quad \beta_{5,2} - 11 = 19/10 > 0.$$

9.4 The twenty-two localized payments

For each L -row, the appropriate entry of the fourteen-row table gives $\rho < r$, and the fixed wall gives $K \geq c$. Direct substitution in (29) gives the following complete list.

row	$P(C; c, r)$	row	$P(C; c, r)$
$L_{7,1}$	21676933/90000000	$L_{7,2}$	58340437/11000000
$L_{7,3}$	1564/99	$L_{8,1}$	51/11
$L_{8,2}$	2294563597/99000000	$L_{8,3}$	88567/6336
$L_{8,4}$	8866645/405504	$L_{9,1}$	162353/21384
$L_{9,2}$	10063157/1375000	$L_{9,3}$	727261673/704000000
$L_{9,4}$	252947/9702	$L_{10,1}$	2173382/606375
$L_{10,2}$	126077081/21484375	$L_{10,3}$	386/125
$L_{10,4}$	73838/15625	$L_{10,5}$	89303/6336
$L_{10,6}$	97715/9702	$L_{10,7}$	7197/11000
$L_{10,8}$	943655/171072	$L_{11,1}$	72407/5632
$L_{11,2}$	810599/152064	$L_{11,3}$	32924779/41765625

All twenty-two fractions are positive, so rule L pays the dominant post-event cap in every localized row.

9.5 The twelve bridge pairs and four global rows

The eleven ordinary B -rows and the initial anchor $B_{7,0}$ have the following exact P, Q pairs. The subscript on Q is the lower checkpoint index.

row	$P(C; c, r)$	$Q_j(c, r)$
$B_{7,0}$	73625809/50688000	$Q_6 = 22025711/13764100$
$B_{7,1}$	417183/88000	$Q_6 = 1625379/280900$
$B_{7,2}$	637/99	$Q_6 = 165473/44944$
$B_{8,1}$	637/99	$Q_7 = 794689/44944$

row	$P(C; c, r)$	$Q_j(c, r)$
$B_{8,2}$	249635/21384	$Q_7 = 35314/2809$
$B_{9,1}$	249635/21384	$Q_8 = 80258/2809$
$B_{9,2}$	835355/577368	$Q_8 = 525692/25281$
$B_{10,1}$	14506973/1336500	$Q_9 = 4713989/404496$
$B_{10,2}$	202207748/21484375	$Q_9 = 648969/280900$
$B_{11,1}$	507845/261954	$Q_{10} = 317585/1617984$
$B_{11,2}$	25143284/1546875	$Q_{10} = 8811001/1123600$
$B_{11,3}$	151707121/11000000	$Q_{10} = 426449/3441025$

Every entry is positive. Rule B , and the preceding B_0 argument for the first row, therefore pay all bridge states at the event frequency, not at a later upper checkpoint.

The four G -rows use the already displayed global calculation (16). For clarity, here are the exact references needed by the state map; $S = (193/53)^2 + j(j+1) - t^2$ and $R = (38/539)t^3 - C$.

row	j	C	t	(S, R)
G_8	7	40	208/25	(67049/1755625, 5083656/8421875)
G_9	8	53	923/100	(1901439/28090000, 656778873/269500000)
G_{10}	9	73	254/25	(61431/1755625, 7911557/8421875)
G_{11}	10	89	111/10	(14211/280900, 1999489/269500).

Both components of every pair are positive.

Proposition 2 (Closed post-event domination). *For every $7 \leq m \leq 11$ and every $(\rho, K) \in \mathfrak{B}_m^+$,*

$$\widehat{C}_z(K) < \mathcal{W}(\rho, K).$$

At $K = k_{11}(\rho)$, the original strict proxy also satisfies $C_z(K) < \mathcal{W}(\rho, K)$. Consequently every fixed, moving, or coincident proxy event in $k_6 \leq K \leq k_{11}$ is paid at its own wall.

Proof. Fix a band. The upper checkpoint inventory (27) contains every possible event in that band; the algebraic splits in (9) cover the two higher radial tails at their equality faces. Choose H, h, τ only after all modes at the event have been inserted into (25). The simultaneous restrictions (28) and the mutually exclusive conditions in the thirty-eight-row table put the resulting state in exactly one row. Formula (26) proves its displayed cap, even if the modal set has gaps.

An L -row is paid by its localization witness and its positive P . A B -row is paid on both closed sides of its split by its positive P, Q . A G -row is paid by (30); the initial row is paid by $B_{7,0}$. These alternatives exhaust the table, proving the closed post-event inequality.

No ordering premise has entered the argument. Changing the order of two fixed or moving walls merely changes which disjoint state is selected. At a branch equality in the maximum defining $b_{\ell,n}$, both lower walls remain valid. At a ratio face the defining state is impossible by the strict positive localization witness. At a bridge face both payment arguments hold with positive reserve.

At an interior modal wall the original strict proxy assigns equality to the pre-event cap, whereas the stronger closed post-event cap has just been paid. At a lower checkpoint $K = k_{m-1}$, the closed post-event version is assigned to the following half-open band and is covered even when $K = k_{m-1}$; the original strict proxy omits every channel on its defining equality and remains owned by the preceding closed spectral band. At intermediate upper checkpoints the same convention passes the post-event state to the following band. Finally, at $K = k_{11}$ the strict inequalities $b_{\ell,n} < K$ give the same state restrictions and the $m = 11$ payment rows directly, while the equality channel (11, 1) is absent. Coincident walls are harmless because (25) inserts their combined multiplicity before the unique state row is chosen. $\square$

The proposition above is the final mathematical exhaustion theorem. It is not a row-count inference from the historical 611-predicate executable census. The map below is retained only as archival provenance for the older arithmetic implementation and is not a premise of Proposition 2.

10 Archival exact row map and scope audit

The following map uses the ordering of the two functions in the original exact registry. To keep the table readable, L denotes `verify_localizations`, and C denotes `verify_cap_tables_and_payments`. The map lets a referee match every replaced predicate to a displayed hand calculation.

registry family	row numbers	analytic replacement here
L	1–2	distinctness and connectedness: (6)–(7) and the seventeen-row table
L	3–19	all ratios in the high strip, by $\rho_c < 1/7 < 4/25$ and $2/3 < 7/8$
L	20–36	the seventeen exact sign witnesses in the long table
L	37–42	split location (8) and side/equality arithmetic (9)
L	43–47	the five k_{11} implications (10)–(13)
C	44–55	the twelve non-bookkeeping cap–74 predicates, (20)–(23); row $C56$, the block identity $49 + 25 = 74$, is allocated to Packet A
C	57–59	the three impossible cells, (14)
C	78–79	the general checkpoint monotonicity proof (15)
C	80–84	the five global-payment rows following (16)
C	85–98	the seven localization and seven conditional-payment rows in the final long table

The count reconciliation is

subfamily	connected mathematical objects	registry predicates
registry metadata	1	2
high-strip membership	17	17
localization signs	17	17
algebraic splits	3	6
special k_{11} implications	5	5
cap–74 path	1	12
impossible cells	3	3
checkpoint monotonicity	1	2
global payments	5	5
conditional payments	7	14
total	60	83.

In particular, the five headline Packet–C subfamilies contain exactly

$$17 + 3 + 3 + 5 + 7 = 35$$

connected objects (localization signs, algebraic splits, impossible cells, global payments, and conditional payments). Because each algebraic split has a location and a side predicate, and each conditional payment has a localization and a reserve predicate, these headline objects replace $17 + 6 + 3 + 5 + 14 = 45$ registry predicates. The remaining 38 predicates in the total are the explicitly listed auxiliary closure rows.

Thus this packet replaces all 47 predicates of L and 36 predicates of C . The elementary cap addition rows 1–43, row 56, and rows 60–77 of the latter family are outside the declared Packet C scope; they are instances of the general odd-sum/block-cap identity and are not claimed here. No row in the declared scope is unreconstructed.

Purely Analytic Ledger Replacement, Packet D: Seam Arithmetic and the Two-Sided Payments

Exact hand-checkable exhibit

Abstract

This packet replaces the twenty-two executable seam predicates and the eleven moving/fixed payment predicates used in the high side through k_6 . Every premise is an analytic inequality already derived in the seam or staircase argument, and every remaining decision is displayed as an exact rational identity or strict rational comparison. In particular, the derivative of the seam comparison function is derived explicitly and is bounded by an exact proxy with a positive margin to $16/5$. The six moving k_6 payments are printed with their full rational differences, not merely their numerators. No program execution, floating-point value, or interval calculation is a premise.

1 Rational constants used by both tables

For completeness, the needed bounds for π can be checked from the alternating arctangent series, rather than imported from an executable Machin enclosure. Machin's identity and the alternating remainder rule give

$$\begin{aligned}\pi &> 16 \left(\frac{1}{5} - \frac{1}{3 \cdot 5^3} + \frac{1}{5 \cdot 5^5} - \frac{1}{7 \cdot 5^7} \right) - \frac{4}{239} =: L_\pi, \\ \pi &< 16 \left(\frac{1}{5} - \frac{1}{3 \cdot 5^3} + \frac{1}{5 \cdot 5^5} \right) - 4 \left(\frac{1}{239} - \frac{1}{3 \cdot 239^3} \right) =: U_\pi.\end{aligned}$$

Indeed, the tangent-addition formula proves $\pi = 16 \arctan(1/5) - 4 \arctan(1/239)$, and the displayed truncations have the indicated directions. Direct rational subtraction gives

$$L_\pi - \frac{333}{106} = \frac{3418213}{41563593750} > 0, \quad \frac{1571}{500} - U_\pi = \frac{323354809}{853244937500} > 0. \quad (1)$$

Since

$$\frac{22}{7} - \frac{1571}{500} = \frac{3}{3500} > 0, \quad (2)$$

we may use throughout

$$\underline{\pi} := \frac{333}{106} < \pi < \bar{\pi} := \frac{1571}{500} < \frac{22}{7}. \quad (3)$$

All later occurrences of a rational π bound refer to (3).

2 The twenty-two seam rows

2.1 Analytic input and comparison function

We recall precisely the analytic data to which the finite arithmetic is applied. Put

$$\epsilon = 1 - \rho = y^2, \quad \kappa = K\epsilon^2, \quad 0 < y \leq \frac{1}{\sqrt{6}}, \quad \rho \geq \frac{5}{6}, \quad \kappa \geq 54.$$

For a first plateau of length p , and the quantities m, R in the seam decomposition, set

$$P = py, \quad r = Ry, \quad S = (p + m)y.$$

The angular estimate $d > 5/8$ and $R = p - dm$ imply, whenever $r > 0$,

$$0 < r \leq P, \quad S \leq S_*(P, r) := \frac{13P - 8r}{5}. \quad (4)$$

Equality in the second relation can occur when $m = 0$; no later step requires strictness at this comparison. The curvature-shelf calculation already proved in the seam argument gives the self-consistent inequality

$$P^2 < \frac{2\pi^2 Q}{(1 - \hat{q})^2}, \quad Q = 1 + \frac{y^2}{\kappa} + \frac{yS}{\kappa}, \quad \hat{q} = \frac{y^2}{\rho}(Q - 1). \quad (5)$$

Since the right-hand side increases with Q , define

$$Q_* = 1 + \frac{y^2}{\kappa} + \frac{yS_*}{\kappa}, \quad q_* = a(Q_* - 1), \quad a := \frac{y^2}{\rho} \leq \frac{1}{5}, \quad (6)$$

and

$$H(P, r) := \frac{2\pi^2 Q_*}{(1 - q_*)^2}. \quad (7)$$

Then (4) gives $Q \leq Q_*$ and $\hat{q} \leq q_*$. Since (5) is itself strict and its right-hand side increases with Q , it follows that $P^2 < H(P, r)$.

The remaining arithmetic is exactly the following table. Rows 1–5 are the five constant/shelf rows, rows 6–10 are the five displacement rows, rows 11–14 are the four derivative rows, rows 15–17 are the three endpoint rows, and rows 18–22 are the five terminal-payment rows. Thus the table reconciles one-for-one with all twenty-two seam predicates.

For clarity, the radical used in rows 7–9 follows directly from the analytic shelf bound. Namely,

$$p^2 < \frac{2\pi\rho K}{\epsilon}, \quad P = py, \quad K = \frac{\kappa}{y^4}$$

imply

$$\left(\frac{y^3 P}{\kappa \rho} \right)^2 < \frac{2\pi y^2}{\kappa \rho} \leq \frac{\pi}{135} < \frac{\bar{\pi}}{135}.$$

Together with $S_* \leq 13P/5$, this gives

$$q_* < \frac{1}{1620} + \frac{13}{5} \frac{763}{5000},$$

which is precisely the dependency recorded by rows 6–9.

#	quantity	exact rational check	role
1	seam π bound	$\bar{\pi} - U_\pi = 323354809/853244937500 > 0$	$\pi < \bar{\pi}; (1)$
2	lower $\sqrt{2}$ bound	$2 - (140/99)^2 = 2/9801 > 0$	$140/99 < \sqrt{2}$
3	upper $\sqrt{2}$ bound	$(99/70)^2 - 2 = 1/4900 > 0$	$\sqrt{2} < 99/70$
4	y ceiling	$(49/120)^2 - 1/6 = 1/14400 > 0$	$y < 49/120$
5	shelf ceiling	$(13/5)^2 - 2\bar{\pi} = 119/250 > 0$	$\sqrt{2\pi\rho} < 13/5;$ rows 1

#	quantity	exact rational check	role
6	first displacement term	$\frac{(1/6)^2}{54(5/6)} = \frac{1}{1620}$	$y^4/(\kappa\rho) \leq 1/1620$
7	radical ceiling	$(763/5000)^2 - \frac{\pi}{135} = \frac{8563}{675000000} > 0$	shelf radical; row 1
8	displacement proxy	$\frac{1}{1620} + \frac{13}{5} \frac{763}{5000} = \frac{804689}{2025000}$	rows 6–7
9	displacement margin	$\frac{1}{400} - \frac{804689}{2025000} = \frac{497}{4050000} > 0$	$q_*, \hat{q} < \bar{q} := 159/400$
10	outer-half localization	$\frac{5}{6}(1 - 159/400) = \frac{241}{480} = \frac{1}{2} + \frac{1}{480}$	$x_0/K > 1/2$; row 9
11	Q_* -derivative proxy	$D_Q := \frac{13}{5} \frac{49/120}{54} = \frac{637}{32400}$	$\partial_P Q_* < D_Q$; row 4
12	exact H -derivative proxy	$\frac{2\pi^2 D_Q (1 + 159/400 + 2/5)}{(1 - 159/400)^3} = \frac{2260740364246}{708624500625}$	rows 1, 9, 11 and $a \leq 1/5$
13	margin to $16/5$	$\frac{16}{5} - \frac{2260740364246}{708624500625} = \frac{6858037754}{708624500625} > 0$	$\partial_P H(P, B) < 16/5$
14	synthetic-slope margin	$2\frac{14}{3} - \frac{16}{5} = \frac{92}{15} > 0$	$\partial_P (P^2 - H) > 92/15$
15	endpoint Q_* proxy	$\bar{Q} := 1 + \frac{1/6}{54} + \frac{(49/120)(14/3)}{54} = \frac{10093}{9720}$, $\bar{Q} - 1 = \frac{373}{9720}$	$Q_*(B, B) < \bar{Q}$
16	endpoint q_* proxy	$\frac{1}{5}(\bar{Q} - 1) = \frac{373}{48600}$	$q_*(B, B) < 373/48600$
17	initial reserve	$\left(\frac{14}{3}\right)^2 - \frac{2\pi^2(10093/9720)}{(1 - 373/48600)^2} = \frac{2505132463469}{2616573970125} > 0$	$B^2 - H(B, B) > 0$; rows 1, 15–16
18	action-gain coefficient	$g := \frac{2(140/99)54}{3(1571/500)} = \frac{280000}{17281}$	$K\eta > g/y$; rows 1–2
19	gain after R loss	$g - \frac{14}{8} \frac{598066}{99} = \frac{51843}{132}$	use $R < 14/(3y)$
20	interface loss proxy	$\frac{8}{15} \frac{99}{70} = \frac{132}{175}$	$4\delta < 132/175$; row 3
21	dangerous-branch reserve	$\frac{598066}{51843} \frac{120}{49} - 1 - \frac{132}{175} = \frac{80132733}{3024175} > 0$	$R > 0$; rows 4, 19–20
22	safe-branch reserve	$\frac{280000}{17281} \frac{120}{49} - 1 - \frac{132}{175} = \frac{114694733}{3024175} > 0$	$R \leq 0$; rows 4, 18, 20

2.2 Why the table proves the seam implications

We spell out the two implications for which the table is more than a collection of arithmetic facts.

First, on the segment $r = B := 14/3$ and $P \geq B$,

$$\partial_P Q_* = \frac{13y}{5\kappa}, \quad \partial_P q_* = a \partial_P Q_*.$$

Differentiation of (7), followed only by the identity $aQ_* = q_* + a$, gives

$$\partial_P H = 2\pi^2(\partial_P Q_*) \frac{1 - q_* + 2aQ_*}{(1 - q_*)^3} = 2\pi^2(\partial_P Q_*) \frac{1 + q_* + 2a}{(1 - q_*)^3}. \quad (8)$$

Rows 1, 9, and 11 and $a \leq 1/5$ substitute monotonically into (8); rows 12–13 then prove $\partial_P H < 16/5$. Because $P \geq B$, row 14 proves

$$\frac{d}{dP}(P^2 - H(P, B)) = 2P - \partial_P H(P, B) > \frac{92}{15} > 0. \quad (9)$$

Rows 15–17 prove $B^2 - H(B, B) > 0$. Integration of (9) therefore yields $P^2 > H(P, B)$ for every $P \geq B$. If $r \geq B$, then $H(P, r) \leq H(P, B)$ because $S_*(P, r)$ decreases with r and H increases with Q_* . This contradicts $P^2 < H(P, r)$. Hence

$$r < B, \quad R < \frac{14}{3y}. \quad (10)$$

Second, the turning-point action bound gives

$$K\eta \geq \frac{2\sqrt{2}\kappa}{3\pi} \frac{1}{y} > \frac{g}{y},$$

where row 18 is the exact rational lower proxy. Also row 4 gives $1/y > 120/49$, and row 20 gives $4\delta < 8\sqrt{2}/15 < 132/175$. With $M = \lfloor K\eta \rfloor - R$ and the strict inequality $\lfloor x \rfloor > x - 1$, (10) and row 21 give, when $R > 0$,

$$M - 4\delta > \left(g - \frac{14}{3}\right) \frac{120}{49} - 1 - \frac{132}{175} > 0.$$

When $R \leq 0$, row 22 gives the still stronger estimate

$$M - 4\delta > g \frac{120}{49} - 1 - \frac{132}{175} > 0.$$

Thus the seam tail has the strict action payment claimed in the main argument, including the $R = 0$ and integer- $K\eta$ faces.

Remark 1 (Notation correction in the executable label). The supplemental verifier labels row 11 as an “exact dq/dP cap.” In the notation of the proof, the number 637/32400 is a proxy for $\partial_P Q_*$, not for $\partial_P q_*$. In fact $\partial_P q_* = a\partial_P Q_*$ and hence $\partial_P q_* < 637/162000$. Formula (8) and the verifier both use 637/32400 in its correct role as the Q_* -derivative proxy. The proof is therefore sound, but the old ledger label is inaccurate.

Remark 2 (Source wording for the π bound). One sentence in the seam passage groups $\pi < 1571/500$ with bounds said to be proved by positive rational squaring. Squaring proves rows 2–5, but not a bound for π . Equations (1)–(3) supply the required analytic alternating-series proof.

3 The eleven k_6 payment rows

Put

$$z_\rho = \frac{\pi}{1 - \rho}, \quad k_m(\rho)^2 = z_\rho^2 + m(m + 1), \quad p_1(\rho)^2 = 4z_\rho^2 + 2, \quad (11)$$

and

$$\mathcal{W}(\rho, K) = \frac{2}{9\pi}(1 - \rho^3)K^3. \quad (12)$$

For a moving wall $T^2 = az_\rho^2 + b$, direct differentiation shows that $\mathcal{W}(\rho, T)$ increases whenever

$$a\pi^2(1 + \rho) > b\rho^2(1 - \rho)^2. \quad (13)$$

Here $b/a \leq 42$. Since $\rho^2(1 - \rho)^2 \leq 1/16$ and $\pi > 3$, the left side of (13), after division by a , is greater than 9, whereas the right side is at most $42/16$; hence every moving payment in the table is increasing. At a fixed wall L , the function $\mathcal{W}(\rho, L)$ is decreasing in ρ .

For a rational split r , define

$$A(a, b; r) := a \frac{(333/106)^2}{(1 - r)^2} + b, \quad B(r) := \frac{7}{99}(1 - r^3). \quad (14)$$

The bounds (3) give, on a moving wall at $\rho = r$,

$$\mathcal{W}(r, T) > B(r)A(a, b; r)^{3/2}. \quad (15)$$

Thus, for a positive cap C , the rational inequality

$$\Delta_{\text{mov}} := B(r)^2 A(a, b; r)^3 - C^2 > 0 \quad (16)$$

implies the strict payment, because

$$\Delta_{\text{mov}} = (B(r)A^{3/2} - C)(B(r)A^{3/2} + C)$$

and the second factor is positive. At a fixed wall L , it is enough to check the ordinary rational reserve

$$\Delta_{\text{fix}} := B(r)L^3 - C > 0. \quad (17)$$

The following is the complete eleven-row registry: six moving rows and five fixed rows. The fractions in the last column are the full reduced differences in (16) or (17).

wall/branch	split r	wall data	cap C	full exact rational difference
k_1 moving	1/8	$(a, b) = (1, 2)$	4	$\Delta_{\text{mov}} = \frac{153864824754340538785}{348796101192617852928} > 0$
k_2 moving	3/10	$(a, b) = (1, 6)$	9	$\Delta_{\text{mov}} = \frac{1399810848073457853053}{394237491401523000000} > 0$
k_3 moving	1/2	$(a, b) = (1, 12)$	16	$\Delta_{\text{mov}} = \frac{137027505353736631}{514922437748928} > 0$
k_4 moving	1/2	$(a, b) = (1, 20)$	25	$\Delta_{\text{mov}} = \frac{2507119282010251109}{13902905819221056} > 0$
k_5 moving	13/25	$(a, b) = (1, 30)$	36	$\Delta_{\text{mov}} = \frac{92672404686738742434741}{709259556128000000000} > 0$
p_1 moving	1/5	$(a, b) = (4, 2)$	29	$\Delta_{\text{mov}} = \frac{373255772037478993002833}{868931613701316000000} > 0$
k_2 fixed	3/10	$L = 51/10$	9	$\Delta_{\text{fix}} = \frac{1387329}{11000000} > 0$
k_3 fixed	1/2	$L = 13/2$	16	$\Delta_{\text{fix}} = \frac{6277}{6336} > 0$
k_4 fixed	1/2	$L = 15/2$	25	$\Delta_{\text{fix}} = \frac{775}{6336} > 0$
k_5 fixed	13/25	$L = 17/2$	36	$\Delta_{\text{fix}} = \frac{704}{452843} > 0$
p_1 fixed	1/5	$L = 77/10$	29	$\Delta_{\text{fix}} = \frac{343750}{849901} > 0$

We now verify that the table pays every branch rather than merely listing positive numbers. For each maximum entry wall

$$\max\{k_2, 51/10\}, \quad \max\{k_3, 13/2\}, \quad \max\{k_4, 15/2\}, \quad \max\{k_5, 17/2\}, \quad \max\{p_1, 77/10\},$$

partition the ratio interval at the displayed r . On $\rho \geq r$, the maximum is at least the moving component and monotonicity together with the moving row pays C . On $\rho \leq r$, the maximum is at least the fixed component and the fixed row pays C . This argument does not assume that the two components are equal at the rational split. For k_1 , the applicable ratio interval starts at $\rho_c > 1/8$ (because $\pi < 22/7 < 7/2$), so the first moving row applies directly. Strict payments also cover coincident entry walls, while the wall itself remains assigned to the lower cap by the strict spectral convention.

4 Reconciliation with the lower- d rows

The two rational π predicates used by the two-sided ledger are already proved in (3). The eleven k_6 predicates are exactly the eleven rows in the preceding table. The remaining thirteen wall/cap predicates in that ledger are the ten transparent multiplicity sums

$$\begin{aligned}
& (1, 1+3, 1+3+5, 1+3+5+1, 1+3+5+7, \\
& \quad 1+3+5+7+1, 1+3+5+7+9+1, \\
& \quad 1+3+5+7+9+1+3, \\
& \quad 1+3+5+7+9+11+1+3, \\
& \quad 1+3+5+7+9+11+1+3+5) \\
& \quad = (1, 4, 9, 10, 16, 17, 26, 29, 40, 45),
\end{aligned} \tag{18}$$

and the endpoint ordering $13/2 < 2z_\rho < 15/2$. The latter follows without an executable check: using $\sqrt{337} < 19$ on the lower endpoint and (3),

$$2 \frac{333}{106} \frac{19}{18} - \frac{13}{2} = \frac{7}{53} > 0, \quad \frac{15}{2} - \left(2 \frac{22}{7} + 1\right) = \frac{3}{14} > 0. \tag{19}$$

Thus the cap-10 row is empty, and the three executable ordering rows collapse to (19) and its immediate empty-cell consequence.

The nine lower- d payment rows need no new table here: the main paper’s lemma “Lower side through d ” (label `cm:lem-lower-d`) already prints the seven fixed-wall differences

$$\frac{793292}{1546875}, \frac{486236661}{1375000000}, \frac{333100003}{99000000}, \frac{329797}{88000}, \frac{3590476297}{1125000000}, \frac{327078887}{99000000}, \frac{8805519}{1375000},$$

and states the initial and moving payments $\mathcal{W}(\rho_c, L(\rho_c)) > 19/6$, where $L(\rho) = (2\rho)^{-1}$ on this branch, and $\mathcal{W}(1/19, 2z) > 508/27$. If desired, their exact printed lower proxies close by the two one-line subtractions

$$\frac{57421}{16854} - \frac{19}{6} = \frac{675}{2809} > 0, \quad \frac{173863}{8427} - \frac{508}{27} = \frac{137795}{75843} > 0. \tag{20}$$

Consequently every seam and two-sided payment predicate assigned to Packet D is now visible as analytic reasoning plus printed integer arithmetic.

5 Dependency count and conclusion

The replacement count is exact:

$$\underbrace{22}_{\text{seam table}} + \underbrace{(2+6+5+10+3)}_{k_6/\text{lower-}d \text{ wall-cap ledger}} + \underbrace{9}_{\text{already printed lower-}d \text{ payments}}.$$

The two constant rows are supplied by (3); the six plus five payment rows are the eleven-row table; the ten bookkeeping rows are (18); the three ordering rows are (19); and the nine already printed payments are explicitly cross-referenced above. This removes program execution from the logical dependency of precisely the Packet-D portion of the exact ledger.

Upper-Owner and Optical Cleanup

Analytic ledger replacement, Packet E

Abstract

This note replaces the executable scope checks surrounding the upper owner of the final spherical-shell residual. It proves directly that the eligible upper wall is $K_0(\rho)$ and that $k_{11}(\rho) < K_0(\rho)$ on the required ratio interval. It also records the symbolic identities that make the repeated optical ledger samples unnecessary.

1 The upper owner

For $0 < \rho < 1$, set

$$a_\rho := \frac{2\pi\rho}{1-\rho}, \quad \eta_\rho := G_1(\max\{\rho, 1/2\}),$$

where

$$G_1(s) = \frac{\sqrt{1-s^2} - s \arccos s}{\pi}, \quad \omega_0 := G_1(1/2) = \frac{\sqrt{3}}{2\pi} - \frac{1}{6}.$$

Let $C_0 > 0$, and define

$$K_0(\rho) := \left(\frac{\sqrt{a_\rho} + \sqrt{a_\rho + 4\eta_\rho C_0}}{2\eta_\rho} \right)^2. \quad (1)$$

Also put

$$\rho_c := \frac{1}{1+2\pi}, \quad z_\rho := \frac{\pi}{1-\rho}, \quad k_{11}(\rho) := \sqrt{z_\rho^2 + 132}.$$

In the live owner graph of the main paper, $U(\rho)$ is the minimum of $K_0(\rho)$ and the small-hole wall $H_0(\rho)$ when the latter is eligible; the eligibility condition for H_0 is $\rho < \omega_0$.

Lemma 1 (The final upper wall is genuinely above the staircase). *For*

$$\rho_c \leq \rho < \frac{39}{50},$$

one has

$$k_{11}(\rho) < 64 < K_0(\rho) = U(\rho). \quad (2)$$

Proof. We use only

$$3 < \pi < \frac{22}{7}, \quad \sqrt{3} < \frac{7}{4}.$$

They imply

$$\omega_0 = \frac{\sqrt{3}}{2\pi} - \frac{1}{6} < \frac{7/4}{6} - \frac{1}{6} = \frac{1}{8}.$$

Since

$$G_1'(s) = -\frac{\arccos s}{\pi} < 0 \quad (0 \leq s < 1), \quad G_1(1) = 0,$$

one has $0 < \eta_\rho \leq \omega_0 < 1/8$. The function a_ρ is increasing, and its value at ρ_c is exactly one:

$$a_{\rho_c} = \frac{2\pi/(1+2\pi)}{2\pi/(1+2\pi)} = 1.$$

Thus $a_\rho \geq 1$. Formula (1) and $C_0 > 0$ give

$$\sqrt{K_0(\rho)} > \frac{\sqrt{a_\rho}}{\eta_\rho} > 8,$$

so $K_0(\rho) > 64$.

On the other hand, $\rho < 39/50 < 7/8$, and therefore

$$z_\rho < 8\pi < \frac{176}{7} < 26.$$

Consequently

$$k_{11}(\rho)^2 < 26^2 + 132 = 808 < 64^2,$$

which proves $k_{11}(\rho) < 64$.

The small-hole wall H_0 is eligible only when $\rho < \omega_0$. But

$$\rho_c > \frac{1}{8} > \omega_0,$$

because $\pi < 22/7 < 7/2$ implies $1 + 2\pi < 8$. Since $\rho \geq \rho_c$, the small-hole wall is therefore ineligible throughout the stated interval. The live upper-owner formula therefore gives $U(\rho) = K_0(\rho) > 64$. Combining this with $k_{11}(\rho) < 64$ proves (2). $\square$

2 Symbolic optical identities

The optical ledger evaluates the same algebraic statements at several sample values. The following identities replace those repetitions.

For $a/\pi = N + t$, with $N \geq 1$ and $0 < t \leq 1$, the exact continuous-minus-product deficit is

$$\frac{D(a)}{\pi^2} = \frac{N^2}{2} + N \left(t^2 + \frac{1}{6} \right) + \frac{2t^3}{3}. \quad (3)$$

If $C_D = 1382/3125$, direct subtraction gives

$$\frac{D(a)}{\pi^2} - C_D(N + t)^2 = \left(\frac{1}{2} - C_D \right) N^2 + N \left(t^2 - 2C_D t + \frac{1}{6} \right) + \frac{2t^3}{3} - C_D t^2. \quad (4)$$

The coefficient of N^2 is positive. The difference between the right sides at $N + 1$ and N is

$$3 \left(\frac{1}{2} - C_D \right) N + \left(\frac{1}{2} - C_D \right) + t^2 - 2C_D t + \frac{1}{6}.$$

The quadratic $t^2 - 2C_D t + 1/6$ has minimum $1/6 - C_D^2$ at $t = C_D$. Consequently, for $N \geq 1$, the displayed difference is at least

$$3 \left(\frac{1}{2} - C_D \right) + \frac{1}{6} - C_D^2 = \frac{4229603}{29296875} > 0.$$

Therefore the deficit increases in N , and it is enough to check $N = 1$. At $N = 1$, differentiation gives

$$\frac{d}{dt} \left(\frac{D(a)}{\pi^2} - C_D(1+t)^2 \right) = 2(t - C_D)(t+1).$$

Thus the unique minimum on $0 < t \leq 1$ occurs at $t = C_D$, where the exact value is

$$\frac{1822532}{91552734375} > 0. \quad (5)$$

This proves the product screen simultaneously for every N, t .

Likewise, the twelve repeated angular block checks are all instances of

$$\sum_{\ell=0}^{m-1} (2\ell+1) = m^2 \quad (m \in \mathbb{N}_0), \quad (6)$$

proved by subtracting consecutive squares. At a radial equality wall the strict convention excludes the newly proposed layer, so no boundary sample is added. These symbolic statements cover every repeated optical row; the two exact endpoint reserves remain those printed in the main optical lemma.

3 The two optical screens with all scaling shown

For completeness, we now print the algebra that was compressed in the main optical lemma. Set

$$\varepsilon_0 = \frac{11}{50}, \quad c = \frac{1126}{625}, \quad q = \frac{106}{333}, \quad C_D = \frac{1382}{3125}.$$

The inequalities

$$\pi > \frac{333}{106}, \quad \pi < \frac{22}{7}$$

give $q > 1/\pi$, and $0 < \varepsilon \leq \varepsilon_0 < 1/2$.

Lemma 2 (Complete optical screen arithmetic). *Let $a = \varepsilon K$.*

1. *If $\pi < a \leq c/\varepsilon$, then*

$$C_D > \frac{2cq}{3} + \frac{\varepsilon}{4} + \varepsilon^2 q^2. \quad (L)$$

2. *If $a \geq c/\varepsilon$, define*

$$L(\varepsilon) := \frac{(1-\varepsilon)^2}{4} - \frac{(22/7)(1-\varepsilon)\varepsilon}{4c},$$

$$A(\varepsilon) := q\varepsilon + \frac{\varepsilon^2}{4c} + \frac{q\varepsilon^3}{4c} + \frac{\varepsilon^4}{16c^2}.$$

Then $L(\varepsilon) > A(\varepsilon)$.

3. *The low and high domains are both closed at $a = c/\varepsilon$, so their union covers every $a > \pi$.*

Proof. For the low screen, $D(a) > C_D a^2$ was proved above. The sufficient product inequality obtained in the main optical argument is

$$D(a) \geq \varepsilon \left(\frac{2a^3}{3\pi} + \frac{a^2}{4} \right) + \frac{\varepsilon^2 a}{\pi}. \quad (1)$$

After division by a^2 , the three terms on the right are bounded by

$$\frac{2\varepsilon a}{3\pi} \leq \frac{2cq}{3}, \quad \frac{\varepsilon}{4} \leq \frac{\varepsilon_0}{4}, \quad \frac{\varepsilon^2}{\pi a} < \frac{\varepsilon^2}{\pi^2} < \varepsilon_0^2 q^2.$$

Exact subtraction gives

$$C_D - \left(\frac{2cq}{3} + \frac{11}{200} + \left(\frac{11}{50} \right)^2 q^2 \right) = \frac{39569}{2772225000} > 0. \quad (7)$$

This proves (L). For $0 \leq a \leq \pi$, the strict radial count is zero, including $a = \pi$, so no low-frequency gap remains.

For the high screen, the exact radial deficit and angular ceiling error from the main analytic argument are

$$\begin{aligned} \frac{D_{\text{rad}}}{a^2} &\geq \frac{(1-\varepsilon)^2}{4} - \frac{\pi(1-\varepsilon)}{4a}, \\ \frac{E}{a^2} &= \frac{\varepsilon}{\pi} + \frac{\varepsilon^2}{4\pi a} + \frac{\varepsilon}{4a} + \frac{\varepsilon^2}{16a^2}. \end{aligned}$$

Since $a \geq c/\varepsilon$, one has $1/a \leq \varepsilon/c$. Using the two rational bounds for π therefore gives

$$\frac{D_{\text{rad}}}{a^2} \geq L(\varepsilon), \quad \frac{E}{a^2} < A(\varepsilon).$$

Moreover,

$$L'(\varepsilon) = -\frac{1-\varepsilon}{2} - \frac{22/7}{4c}(1-2\varepsilon) < 0 \quad (0 < \varepsilon < 1/2),$$

whereas $A'(\varepsilon) > 0$ coefficientwise. Hence

$$L(\varepsilon) - A(\varepsilon) \geq L(\varepsilon_0) - A(\varepsilon_0).$$

The endpoint subtraction is

$$\begin{aligned} L(\varepsilon_0) - A(\varepsilon_0) &= \frac{1}{4} \left(\frac{39}{50} \right)^2 - \frac{22}{7} \frac{(39/50)(11/50)}{4(1126/625)} \\ &\quad - \left[\frac{11}{50} q + \frac{(11/50)^2}{4(1126/625)} + \frac{(11/50)^3 q}{4(1126/625)} + \frac{(11/50)^4}{16(1126/625)^2} \right] \\ &= \frac{14817541}{472867032960000} > 0. \end{aligned} \quad (8)$$

Thus the radial deficit strictly exceeds the angular error.

Finally,

$$\frac{c}{\varepsilon} \geq \frac{c}{\varepsilon_0} = \frac{2252}{275} > \frac{22}{7} > \pi, \quad \frac{2252}{275} - \frac{22}{7} = \frac{9714}{1925} > 0.$$

Therefore the common face lies in the low domain as well as the high domain, and

$$\{a : \pi < a \leq c/\varepsilon\} \cup \{a : a \geq c/\varepsilon\} = \{a : a > \pi\},$$

and equality $a = c/\varepsilon$ belongs to both pieces. Together with the zero-count interval $0 \leq a \leq \pi$, this covers every $a \geq 0$. At $K = a = 0$, both the spectral count and the Weyl term vanish. $\square$

Ledger rows replaced. Lemma 1 replaces the eight executable `verify_u_and_k11` scope checks. Equations (3)–(6) replace the repeated product samples and odd-block identities. These are rows of `verify_optical_constants`. Lemma 2 prints the remaining scaling, monotonicity, common-face, and exact-reserve implications. Thus all fifty optical rows are consequences of displayed identities rather than executable predicates.

Exact Owner Subtraction and Direct Analytic Closure

Analytic ledger replacement, Packet F

Abstract

The executable exact ledger checks a 9×7 truth table when the accepted lower and staircase regions are subtracted from the Round-19 residual. This note replaces all sixty-three state checks and the five additional interface checks by one set-theoretic calculation. The surviving set is then placed directly inside the closed domain of the all-frequency retained-remainder theorem. Every open and closed face is retained explicitly, and no artificial frequency interface is introduced.

1 Sets and inherited owners

Let

$$0 < \rho_* < \rho_0 < \rho_c < \rho_{\text{opt}}, \quad \rho_{\text{opt}} := \frac{39}{50},$$

and let $U(\rho)$, $k_6(\rho)$, and $k_{11}(\rho)$ be the upper owner and the two staircase walls from the main proof. Put

$$D_{\text{low}} := \left\{ \rho_* < \rho \leq \rho_0, (2\rho)^{-1} < K < U(\rho) \right\} \cup \left\{ \rho_0 < \rho < \rho_c, d < K < U(\rho) \right\}, \quad (1)$$

$$D_{\text{high}} := \left\{ \rho_c \leq \rho < \rho_{\text{opt}}, k_6(\rho) < K < U(\rho) \right\}, \quad (2)$$

where $d = (2\rho_0)^{-1}$, and define

$$D_{19} := D_{\text{low}} \dot{\cup} D_{\text{high}}. \quad (3)$$

The dot is justified by the disjoint ratio ranges.

The already-proved inclusive owners used in the subtraction are:

$$L_{\text{own}} := D_{\text{low}}, \quad (4)$$

$$S_{\text{own}} := \left\{ \rho_c \leq \rho < \rho_{\text{opt}}, k_6(\rho) < K \leq k_{11}(\rho) \right\}, \quad (5)$$

The lower owner includes the complete $\rho = \rho_0$ fibre. The staircase includes its upper wall $K = k_{11}(\rho)$. The optical owner from the main proof has already restricted the high component of D_{19} to $\rho < \rho_{\text{opt}}$. The residual also excludes $K = U(\rho)$.

2 The subtraction

Proposition 1 (Exact residual identity). *One has*

$$\boxed{D_{19} \setminus (L_{\text{own}} \cup S_{\text{own}}) = D_{20}}, \quad (6)$$

where

$$D_{20} := \left\{ \rho_c \leq \rho < \rho_{\text{opt}}, k_{11}(\rho) < K < U(\rho) \right\}. \quad (7)$$

Proof. Since $L_{\text{own}} = D_{\text{low}}$ and the union in (3) is disjoint,

$$D_{19} \setminus L_{\text{own}} = D_{\text{high}}. \quad (8)$$

Inside D_{high} , the lower frequency inequality is strict: $k_6(\rho) < K$. Removing the inclusive staircase (5) therefore gives exactly

$$D_{\text{high}} \setminus S_{\text{own}} = \{\rho_c \leq \rho < \rho_{\text{opt}}, k_{11}(\rho) < K < U(\rho)\}. \quad (9)$$

There is no missing equality case: $K = k_{11}(\rho)$ belongs to S_{own} , while $K = U(\rho)$ never belonged to D_{19} . The face $\rho = \rho_{\text{opt}}$ was already removed when the optical theorem was applied in forming D_{19} , whereas $\rho = \rho_c$ remains in D_{high} . Thus (9) is precisely (7). $\square$

3 The five delicate interfaces

For clarity, the five extra assertions formerly checked separately by the executable ledger are consequences of the displayed sets:

1. At $\rho = \rho_0$, the identity $(2\rho_0)^{-1} = d$ makes the prospective second lower fibre empty; the first set in (1) owns the full equality fibre, and L_{own} removes it exactly once.
2. The face $\rho = \rho_c$ belongs to D_{high} . It is not an optical face, so points with $k_{11}(\rho_c) < K < U(\rho_c)$ remain in D_{20} .
3. The face $\rho = \rho_{\text{opt}} = 39/50$ belongs to the optical theorem and is already absent from D_{19} , hence from D_{20} .
4. The frequency face $K = k_{11}(\rho)$ belongs to S_{own} , whereas $K = U(\rho)$ is excluded already by the strict inequality in D_{19} .
5. If $\rho_c < \rho < \rho_{\text{opt}}$ and $k_{11}(\rho) < K < U(\rho)$, the point is in D_{high} but in neither L_{own} nor S_{own} ; hence it survives, and every surviving point has exactly these properties.

4 Direct retained-remainder ownership

Let

$$\mathcal{R}_{\text{tr}} := \{(\rho, K) : \rho_c \leq \rho \leq \rho_{\text{opt}}, \quad K \geq k_{11}(\rho)\}. \quad (10)$$

This is the closed domain of the all-frequency retained-remainder theorem proved by the structural, angular-block, and scalar-positivity modules. In the notation of the main proof, that theorem states

$$(\rho, K) \in \mathcal{R}_{\text{tr}} \implies N_D(A_{\rho,1}, K^2) < W(\rho, K). \quad (11)$$

Proposition 2 (Direct ownership of the exact residual). *One has*

$$\boxed{D_{20} \subset \mathcal{R}_{\text{tr}}}, \quad D_{20} \setminus \mathcal{R}_{\text{tr}} = \emptyset. \quad (12)$$

Consequently the strict Pólya inequality holds at every point of D_{20} .

Proof. If $(\rho, K) \in D_{20}$, then

$$\rho_c \leq \rho < \rho_{\text{opt}} \quad \text{and} \quad k_{11}(\rho) < K < U(\rho).$$

The first pair of inequalities implies $\rho_c \leq \rho \leq \rho_{\text{opt}}$, and the second implies $K \geq k_{11}(\rho)$. Hence $(\rho, K) \in \mathcal{R}_{\text{rr}}$, which proves (12); the final assertion follows from (11).

The inclusion does not change any inherited owner. The face $\rho = \rho_{\text{opt}}$ and the frequency face $K = k_{11}(\rho)$ are outside D_{20} , although the stronger theorem also proves them. The face $K = U(\rho)$ remains excluded from D_{20} , while $\rho = \rho_c$ remains included subject to the two strict frequency inequalities. There is no additional frequency face. $\square$

Ledger rows replaced. Proposition 1 replaces all $9 \cdot 7 = 63$ state evaluations in the executable family `verify_exact_subtraction`. The five items above replace its five separately asserted interface predicates. Proposition 2 supplies the final residual closure by one set inclusion, without a frequency classifier or executable truth table.

A Global Structural Reduction for the Spherical-Shell Proxy

Global analytic structural module

Abstract

This note gives the global structural component of the purely analytic middle-ratio argument for the spherical-shell proxy. The weighted proxy is first written as an exact layer-cake sum. Its deficit from the action integral is then decomposed into a radial shifted-quadrature deficit and an angular rounding defect. A Stieltjes calculation retains two nonnegative remainders that are discarded by the optical estimate, and one of those remainders admits a smooth explicit lower bound. The result is the direct lower bound (36). All structural identities hold for every $0 < \rho < 1$ and $K > 0$; in particular they apply on the upward-closed middle-ratio region $\rho_c \leq \rho \leq 39/50$, $K \geq k_{11}(\rho)$.

1 Action and the global target

For $a > 0$ and $z \geq 0$, put

$$G_a(z) = \begin{cases} \frac{1}{\pi} \left(\sqrt{a^2 - z^2} - z \arccos \frac{z}{a} \right), & 0 \leq z < a, \\ 0, & z \geq a. \end{cases} \quad (1)$$

Fix $0 < \rho < 1$ and $K > 0$, and define

$$A(z) = A_{\rho,K}(z) := G_K(z) - G_{\rho K}(z), \quad T := A(0) = \frac{(1-\rho)K}{\pi}. \quad (2)$$

The strict positive-integer count is

$$[x]_{<} := \#\{n \in \mathbb{N} : n < x\}.$$

The spherical-shell proxy and its continuous payment are

$$P(\rho, K) := \sum_{\substack{\nu=\ell+1/2 < K \\ \ell \in \mathbb{N}_0}} 2\nu \left[A(\nu) + \frac{1}{4} \right]_{<}, \quad (3)$$

$$W(\rho, K) := \int_0^K 2zA(z) dz = \frac{2(1-\rho^3)K^3}{9\pi}. \quad (4)$$

The upward-closed middle-ratio region used below is

$$\mathcal{R}_c := \left\{ (\rho, K) : \rho_c \leq \rho \leq \frac{39}{50}, \quad K \geq k_{11}(\rho) \right\}, \quad \rho_c := \frac{1}{1+2\pi}, \quad k_{11}(\rho) := \sqrt{\frac{\pi^2}{(1-\rho)^2} + 132}. \quad (5)$$

The middle-ratio target is $P < W$ on $\mathcal{R}_c$.

For $0 < z < K$, direct differentiation gives

$$-A'(z) = \frac{1}{\pi} \begin{cases} \arccos \frac{z}{K} - \arccos \frac{z}{\rho K}, & 0 < z < \rho K, \\ \arccos \frac{z}{K}, & \rho K \leq z < K. \end{cases} \quad (6)$$

In particular, A is strictly decreasing from T to 0. Let

$$R : [0, T] \longrightarrow [0, K]$$

be its decreasing inverse, and set

$$F(t) := R(t)^2, \quad g(t) := -F'(t), \quad \tau := A(\rho K) = KG_1(\rho). \quad (7)$$

The derivative in (6) increases in magnitude on $(0, \rho K)$ and decreases in magnitude on $(\rho K, K)$. The inverse curvature has the related, but slightly stronger, property

$$g \text{ decreases on } (0, \tau), \quad g \text{ increases on } (\tau, T). \quad (8)$$

Indeed, on the outer branch, if $z = R(t)$ then

$$g(t) = \frac{2\pi z}{\arccos(z/K)};$$

the quotient increases with z , while z decreases with t . On the inner branch, put

$$\delta(z) := \arccos \frac{z}{K} - \arccos \frac{z}{\rho K}.$$

Differentiation with respect to the intermediate ratio gives

$$\frac{\delta(z)}{z} = \int_{\rho}^1 \frac{ds}{s\sqrt{s^2 K^2 - z^2}}.$$

This integral increases with z . Thus $z/\delta(z)$ decreases with z , and $g(t) = 2\pi z/\delta(z)$ increases as t increases and z decreases. The endpoint and interface data are

$$F(\tau) = \rho^2 K^2, \quad g(\tau) = \frac{2\pi \rho K}{\arccos \rho}, \quad g(T) = \frac{2\pi \rho K}{1 - \rho}. \quad (9)$$

Here and below $g(T)$ is the one-sided limit. Near $t = 0$, one has $g(t) = O(t^{-1/3})$, so all improper Stieltjes boundary terms used below vanish.

2 Exact layer cake and angular rounding

Let

$$x_n := n - \frac{1}{4}, \quad M(r) := \#\left\{\ell \in \mathbb{N}_0 : \ell + \frac{1}{2} < r\right\} = \max\left\{0, \left\lceil r - \frac{1}{2} \right\rceil\right\}. \quad (10)$$

Lemma 1 (Exact layer cake). *For every $0 < \rho < 1$ and $K > 0$,*

$$P(\rho, K) = \sum_{\substack{n \geq 1 \\ x_n < T}} M(R(x_n))^2, \quad W(\rho, K) = \int_0^T F(t) dt. \quad (11)$$

Both identities retain the strict convention at every radial and angular wall.

Proof. For a fixed half-integer ν ,

$$\left[A(\nu) + \frac{1}{4} \right]_{<} = \#\{n \geq 1 : x_n < A(\nu)\}.$$

Since A and R are decreasing, $x_n < A(\nu)$ is equivalent to $\nu < R(x_n)$. Interchanging the two finite sums and using

$$\sum_{\ell=0}^{m-1} (2\ell + 1) = m^2$$

proves the first identity. At $R(x_n) = m + 1/2$, the strict angular count is m , exactly as specified by M .

For the second identity, the area formula for a decreasing inverse gives

$$\int_0^T R(t)^2 dt = \int_0^K 2zA(z) dz.$$

The last integral is (4). □

For every active n , abbreviate

$$r_n := R(x_n), \quad m_n := M(r_n), \quad \alpha_n := r_n - m_n + \frac{1}{2}. \quad (12)$$

The strict convention implies

$$m_n - \frac{1}{2} < r_n \leq m_n + \frac{1}{2}, \quad 0 < \alpha_n \leq 1. \quad (13)$$

This includes $m_n = 0$. Define the angular rounding defect

$$\mathcal{E}_{\text{ang}}(\rho, K) := \sum_{x_n < T} (m_n^2 - r_n^2). \quad (14)$$

The defect of a single radial layer is exactly

$$m_n^2 - r_n^2 = (1 - 2\alpha_n)r_n + \left(\alpha_n - \frac{1}{2}\right)^2. \quad (15)$$

Unlike the one-sided estimate $m_n^2 - r_n^2 < r_n + 1/4$, formula (15) retains the cancellation when $\alpha_n > 1/2$.

Combining (11) and (14) gives the first exact two-term decomposition:

$$W - P = \mathcal{D}_{\text{rad}} - \mathcal{E}_{\text{ang}}, \quad \mathcal{D}_{\text{rad}} := \int_0^T F(t) dt - \sum_{x_n < T} F(x_n). \quad (16)$$

3 The radial deficit with retained positive remainders

Define the shifted counting function and its two primitives by

$$C(t) := \#\{n \geq 1 : x_n < t\}, \quad w(t) := t - C(t), \quad \mathfrak{W}(t) := \int_0^t w(s) ds, \quad \Psi(t) := \mathfrak{W}(t) - \frac{t}{4}. \quad (17)$$

On $0 \leq t \leq 3/4$,

$$\Psi(t) = \frac{1}{2} \left(t - \frac{1}{4} \right)^2 - \frac{1}{32}. \quad (18)$$

For $t = x_n + u$, $0 \leq u \leq 1$,

$$\Psi(t) = \frac{1}{2} \left(u - \frac{1}{2} \right)^2 - \frac{1}{32}. \quad (19)$$

Consequently

$$\mathfrak{W}(t) \geq 0, \quad -\frac{1}{32} \leq \Psi(t) \leq \frac{3}{32} \quad (t \geq 0). \quad (20)$$

Proposition 2 (Exact retained-remainder identity). *For every $0 < \rho < 1$ and $K > 0$, put*

$$\mathcal{R}_{\text{dec}} := - \int_{(0,\tau)} \mathfrak{W}(t) dg(t), \quad (21)$$

$$\mathcal{R}_{\text{osc}} := \left(\Psi(T) + \frac{1}{32} \right) g(T) + \int_{(\tau,T)} \left(\frac{3}{32} - \Psi(t) \right) dg(t). \quad (22)$$

Then both remainders are nonnegative, and

$$\begin{aligned} \mathcal{D}_{\text{rad}} &= \frac{F(\tau)}{4} - \frac{g(T)}{8} + g(\tau) \left(\frac{\tau}{4} + \frac{3}{32} \right) \\ &\quad + \mathcal{R}_{\text{dec}} + \mathcal{R}_{\text{osc}}. \end{aligned} \quad (23)$$

Therefore

$$W - P = \mathcal{B}(\rho, K) + \mathcal{R}_{\text{dec}} + \mathcal{R}_{\text{osc}} - \mathcal{E}_{\text{ang}}, \quad (24)$$

where

$$\mathcal{B}(\rho, K) := \frac{\rho^2 K^2}{4} - \frac{\pi \rho K}{4(1-\rho)} + \frac{2\pi \rho K}{\arccos \rho} \left(\frac{KG_1(\rho)}{4} + \frac{3}{32} \right). \quad (25)$$

Proof. Distributionally, $dw = dt - dC$. Stieltjes integration by parts, using $F(T) = 0$ and $w(0) = 0$, gives

$$\mathcal{D}_{\text{rad}} = \int_0^T w(t) g(t) dt. \quad (26)$$

The improper boundary term at 0 vanishes because $\mathfrak{W}(t) = t^2/2$ there and $g(t) = O(t^{-1/3})$.

On $(0, \tau)$, integrate with $d\mathfrak{W} = w dt$:

$$\int_0^\tau wg = \mathfrak{W}(\tau)g(\tau) - \int_{(0,\tau)} \mathfrak{W} dg. \quad (27)$$

On (τ, T) , use $w = 1/4 + \Psi'$ and $\int_\tau^T g = F(\tau)$:

$$\int_\tau^T wg = \frac{F(\tau)}{4} + \Psi(T)g(T) - \Psi(\tau)g(\tau) - \int_{(\tau,T)} \Psi dg. \quad (28)$$

Since $\mathfrak{W}(\tau) = \tau/4 + \Psi(\tau)$, adding (27) and (28) cancels the uncontrolled interface value $\Psi(\tau)$.

Now add and subtract the extremal sawtooth values in (20). The exact identity

$$\begin{aligned} &\Psi(T)g(T) - \int_{(\tau,T)} \Psi dg \\ &= -\frac{g(T)}{8} + \frac{3g(\tau)}{32} + \left(\Psi(T) + \frac{1}{32} \right) g(T) + \int_{(\tau,T)} \left(\frac{3}{32} - \Psi \right) dg \end{aligned}$$

gives (23). By (8), $-dg$ is a positive measure on $(0, \tau)$ and dg is a positive measure on (τ, T) . Equations (20), (21), and (22) therefore prove nonnegativity of both remainders. Finally substitute (9) and $\tau = KG_1(\rho)$, and then use (16). $\square$

4 A smooth lower bound for the decreasing-branch remainder

The term $\mathcal{R}_{\text{dec}}$ is quantitatively important on the middle-ratio region; discarding it loses too much. It has the following elementary smooth minorant.

Lemma 3 (A global primitive minorant). *For $t \geq 0$, define*

$$L(t) := \frac{t^2}{2(1+2t)}, \quad L'(t) = \frac{t(1+t)}{(1+2t)^2}. \quad (29)$$

Then

$$0 \leq L(t) \leq \mathfrak{W}(t) \quad (t \geq 0). \quad (30)$$

Proof. For $0 \leq t \leq 3/4$, one has $\mathfrak{W}(t) = t^2/2 \geq L(t)$. For $t \geq 3/4$, (20) gives

$$\mathfrak{W}(t) \geq \frac{t}{4} - \frac{1}{32}.$$

Moreover,

$$\frac{t}{4} - \frac{1}{32} - \frac{t^2}{2(1+2t)} = \frac{6t-1}{32(1+2t)} \geq 0.$$

This proves (30). □

Define

$$\mathcal{R}_*(\rho, K) := - \int_{(0, \tau)} L(t) dg(t). \quad (31)$$

Because $-dg$ is positive on the decreasing branch,

$$0 \leq \mathcal{R}_* \leq \mathcal{R}_{\text{dec}}. \quad (32)$$

Integration by parts is legitimate at 0 because $L(t)g(t) = O(t^{5/3})$. Hence

$$\mathcal{R}_* = -L(\tau)g(\tau) + \int_0^\tau L'(t)g(t) dt. \quad (33)$$

On the outer branch $t = A(z)$, one has $g(t) dt = -2z dz$. Therefore

$$\mathcal{R}_* = -\frac{\tau^2}{2(1+2\tau)}g(\tau) + \int_{\rho K}^K 2z \frac{A(z)(1+A(z))}{(1+2A(z))^2} dz. \quad (34)$$

This expression is smooth at every point with $0 < \rho < 1$ and $K > 0$ and has no floor or wall discontinuity.

Corollary 4 (Direct retained-remainder bound). *For every $0 < \rho < 1$ and $K > 0$, define*

$$\begin{aligned} \mathcal{H}(\rho, K) := & \frac{\rho^2 K^2}{4} - \frac{\pi \rho K}{4(1-\rho)} \\ & + \frac{2\pi \rho K}{\arccos \rho} \left(\frac{\tau}{4(1+2\tau)} + \frac{3}{32} \right) \\ & + \int_{\rho K}^K 2z \frac{A(z)(1+A(z))}{(1+2A(z))^2} dz, \quad \tau = KG_1(\rho). \end{aligned} \quad (35)$$

Then

$$W(\rho, K) - P(\rho, K) \geq \mathcal{H}(\rho, K) - \mathcal{E}_{\text{ang}}(\rho, K). \quad (36)$$

In particular, $\mathcal{E}_{\text{ang}}(\rho, K) < \mathcal{H}(\rho, K)$ implies $P(\rho, K) < W(\rho, K)$.

Proof. By (25), (34), and

$$\frac{\tau}{4} - \frac{\tau^2}{2(1+2\tau)} = \frac{\tau}{4(1+2\tau)},$$

one has $\mathcal{H} = \mathcal{B} + \mathcal{R}_*$. Now use (24), (32), and $\mathcal{R}_{\text{osc}} \geq 0$. □

Remark 5 (Handoff). The companion one-layer estimate proves, whenever at least one shifted radial layer is active,

$$\mathcal{E}_{\text{ang}} < \frac{(1-\rho^2)K^2}{6} - \frac{N}{2}, \quad N = \#\{n \geq 1 : n - 1/4 < T\}.$$

Combining this with (36) gives one direct smooth lower bound for $W - P$. The separate double-sawtooth expansion is not needed for this implication and is kept outside the live proof as a research reformulation.

Global Angular Blocks for the Middle-Ratio Shell Proxy

Global analytic angular module

Abstract

The global structural reduction for the spherical-shell proxy leaves an angular rounding defect $\mathcal{E}_{\text{ang}}$ and a smooth positive payment $\mathcal{H}$. This note controls the defect for every $0 < \rho < 1$ and $K > 0$ with at least one active radial layer. A single paired radial-block argument, using only the sharp slope cap $-A' \leq 1/2$ and the exact strict angular rounding, gives

$$\mathcal{E}_{\text{ang}} < \frac{(1 - \rho^2)K^2}{6} - \frac{N}{2},$$

where N is the number of active shifted radial layers. On the upward-closed middle-ratio region $\rho_c \leq \rho \leq 39/50$, $K \geq k_{11}(\rho)$, one has $N \geq 1$. The global middle-ratio theorem is therefore reduced to the smooth scalar inequality (19), which is established in the companion scalar-positivity module.

1 Definitions and the target payment

For $a > 0$ and $z \geq 0$, define

$$G_a(z) = \begin{cases} \frac{1}{\pi} \left(\sqrt{a^2 - z^2} - z \arccos \frac{z}{a} \right), & 0 \leq z < a, \\ 0, & z \geq a. \end{cases} \quad (1)$$

Fix $0 < \rho < 1$ and $K > 0$, and put

$$A(z) := G_K(z) - G_{\rho K}(z), \quad T := A(0) = \frac{(1 - \rho)K}{\pi}. \quad (2)$$

The function A decreases strictly from T to 0. Denote its decreasing inverse by $R : [0, T] \rightarrow [0, K]$, and set

$$F(t) := R(t)^2, \quad q(z) := -A'(z), \quad \tau := A(\rho K) = KG_1(\rho). \quad (3)$$

Direct differentiation gives

$$q(z) = \frac{1}{\pi} \begin{cases} \arccos \frac{z}{K} - \arccos \frac{z}{\rho K}, & 0 < z < \rho K, \\ \arccos \frac{z}{K}, & \rho K \leq z < K. \end{cases} \quad (4)$$

Consequently

$$0 \leq q(z) \leq \frac{1}{2}, \quad q \text{ increases on } (0, \rho K), \quad q \text{ decreases on } (\rho K, K). \quad (5)$$

Let

$$x_n := n - \frac{1}{4}, \quad M(r) := \max \left\{ 0, \left\lceil r - \frac{1}{2} \right\rceil \right\}, \quad N := \#\{n \geq 1 : x_n < T\}. \quad (6)$$

For $1 \leq n \leq N$, write

$$r_n := R(x_n), \quad m_n := M(r_n), \quad (7)$$

and define the exact angular rounding defect

$$\mathcal{E}_{\text{ang}}(\rho, K) := \sum_{n=1}^N (m_n^2 - r_n^2). \quad (8)$$

The strict convention is built into M : at $r = m + 1/2$ one has $M(r) = m$, not $m + 1$.

The smooth payment obtained from the retained radial Stieltjes remainder is

$$\begin{aligned} \mathcal{H}(\rho, K) &:= \frac{\rho^2 K^2}{4} - \frac{\pi \rho K}{4(1-\rho)} \\ &\quad + \frac{2\pi \rho K}{\arccos \rho} \left(\frac{\tau}{4(1+2\tau)} + \frac{3}{32} \right) \\ &\quad + \int_{\rho K}^K 2z \frac{A(z)(1+A(z))}{(1+2A(z))^2} dz. \end{aligned} \quad (9)$$

The preceding structural reduction proves the direct lower bound

$$W(\rho, K) - P(\rho, K) \geq \mathcal{H}(\rho, K) - \mathcal{E}_{\text{ang}}(\rho, K). \quad (10)$$

This note derives a smooth upper bound for the left side.

2 A one-layer paired-block estimate

The cap $q \leq 1/2$ in (5) forces a useful separation: one unit of radial action spans at least two units of angular radius. The shift $x_n = n - 1/4$ leaves a left block of length $3/4$, enough to pay the worst angular rounding in that layer.

Lemma 1 (One-layer paired block). *For every active n ,*

$$r_n \leq \frac{4}{3} \int_{n-1}^{x_n} R(t) dt - \frac{3}{4}, \quad (11)$$

and

$$m_n^2 - r_n^2 < r_n + \frac{1}{4}. \quad (12)$$

Both statements remain valid, with the displayed strictness, at every radial-angular common wall.

Proof. Fix $t \in [n-1, x_n]$ and put $z = R(t) \geq r_n$. Since $A(r_n) = x_n$, (5) gives

$$x_n - t = A(r_n) - A(z) = \int_{r_n}^z q(s) ds \leq \frac{z - r_n}{2}.$$

Hence

$$R(t) \geq r_n + 2(x_n - t).$$

Integration over the interval of length $3/4$ yields

$$\int_{n-1}^{x_n} R(t) dt \geq \frac{3}{4}r_n + \frac{9}{16},$$

which is (11).

By strict counting, $m_n < r_n + 1/2$ even when r_n is a half-integer: at $r_n = m + 1/2$ the value is $m_n = m$. Squaring gives

$$m_n^2 < (r_n + 1/2)^2 = r_n^2 + r_n + 1/4,$$

and proves (12). $\square$

Proposition 2 (Global angular block bound). *For every $0 < \rho < 1$ and every $K > 0$ with $N \geq 1$,*

$$\boxed{\mathcal{E}_{\text{ang}}(\rho, K) < \frac{(1 - \rho^2)K^2}{6} - \frac{N}{2}.} \quad (13)$$

In particular,

$$\mathcal{E}_{\text{ang}}(\rho, K) < \frac{(1 - \rho^2)K^2}{6} - \frac{1}{2}. \quad (14)$$

Proof. The left blocks $[n - 1, x_n]$, $1 \leq n \leq N$, are disjoint subsets of $[0, T]$. Summing (11) and then (12) gives

$$\begin{aligned} \mathcal{E}_{\text{ang}} &< \sum_{n=1}^N \left(r_n + \frac{1}{4} \right) \\ &\leq \frac{4}{3} \sum_{n=1}^N \int_{n-1}^{x_n} R(t) dt - \frac{N}{2} \\ &\leq \frac{4}{3} \int_0^T R(t) dt - \frac{N}{2}. \end{aligned}$$

The area identity for decreasing inverses gives

$$\int_0^T R(t) dt = \int_0^K A(z) dz.$$

Scaling $z = as$ in (1) and using

$$\int_0^1 \sqrt{1 - s^2} ds = \frac{\pi}{4}, \quad \int_0^1 s \arccos s ds = \frac{\pi}{8},$$

shows that

$$\int_0^a G_a(z) dz = \frac{a^2}{8}.$$

Consequently

$$\int_0^T R(t) dt = \frac{(1 - \rho^2)K^2}{8},$$

which proves (13). Equation (14) follows from $N \geq 1$. $\square$

Combining the structural and angular estimates removes the staircase defect from the live handoff altogether.

Proposition 3 (Direct analytic proxy gap). *For every $0 < \rho < 1$ and $K > 0$ with $N \geq 1$,*

$$\boxed{W(\rho, K) - P(\rho, K) > \mathcal{H}(\rho, K) - \frac{(1 - \rho^2)K^2}{6} + \frac{N}{2}.} \quad (15)$$

This one inequality includes the first and terminal radial pieces and every strict angular wall; no separate common-jump estimate is a premise.

Proof. Insert (13) into (10). □

3 The upward-closed middle-ratio scalar inequality

Set

$$\rho_c := \frac{1}{1 + 2\pi}, \quad k_{11}(\rho) := \sqrt{\frac{\pi^2}{(1 - \rho)^2} + 132}, \quad (16)$$

and consider the upward-closed middle-ratio region

$$\mathcal{R}_c := \left\{ (\rho, K) : \rho_c \leq \rho \leq \frac{39}{50}, \quad K \geq k_{11}(\rho) \right\}. \quad (17)$$

On this set,

$$T \geq \frac{(1 - \rho)k_{11}(\rho)}{\pi} = \sqrt{1 + \frac{132(1 - \rho)^2}{\pi^2}} > 1, \quad (18)$$

so $N \geq 1$. Proposition 3 reduces the global middle-ratio theorem to the single inequality

$$\boxed{\mathcal{S}(\rho, K) := \mathcal{H}(\rho, K) - \frac{(1 - \rho^2)K^2}{6} + \frac{1}{2} > 0} \quad ((\rho, K) \in \mathcal{R}_c). \quad (19)$$

Everything in $\mathcal{S}$ is smooth at every finite point of $\mathcal{R}_c$; the floor functions and all radial-angular common jumps have disappeared. Substituting (9) gives the explicit form

$$\begin{aligned} \mathcal{S}(\rho, K) &= \frac{(5\rho^2 - 2)K^2}{12} - \frac{\pi\rho K}{4(1 - \rho)} + \frac{1}{2} \\ &\quad + \frac{2\pi\rho K}{\arccos \rho} \left(\frac{KG_1(\rho)}{4(1 + 2KG_1(\rho))} + \frac{3}{32} \right) \\ &\quad + \int_{\rho K}^K 2z \frac{A(z)(1 + A(z))}{(1 + 2A(z))^2} dz. \end{aligned} \quad (20)$$

The elementary identity

$$\frac{u(1 + u)}{(1 + 2u)^2} = \frac{1}{4} - \frac{1}{4(1 + 2u)^2} \quad (21)$$

turns the same scalar gap into the more useful loss form

$$\begin{aligned} \mathcal{S}(\rho, K) &= \frac{(1 + 2\rho^2)K^2}{12} - \frac{1}{2} \int_{\rho K}^K \frac{z}{(1 + 2A(z))^2} dz \\ &\quad - \frac{\pi\rho K}{4(1 - \rho)} + \frac{1}{2} + \frac{2\pi\rho K}{\arccos \rho} \left(\frac{KG_1(\rho)}{4(1 + 2KG_1(\rho))} + \frac{3}{32} \right). \end{aligned} \quad (22)$$

Thus the only non-endpoint quantity still requiring control is the positive boundary-layer loss integral in (22).

Remark 4 (Status and next step). The direct proxy gap (15) is proved analytically from one retained-radial estimate and one one-layer angular estimate. The first remaining step within this module is the smooth scalar inequality (19). The companion scalar-positivity module proves it on all of $\mathcal{R}_c$ by a uniform upper bound for the boundary-layer loss in (22) and strict positivity on the base curve. No upper cutoff in K occurs in the angular estimate or in the verification that $N \geq 1$.

Purely Analytic Positivity of the Upward-Closed Scalar Gap

Global middle-ratio analytic module

Abstract

The angular-block reduction for the middle-ratio spherical-shell proxy leaves one smooth scalar inequality. This note proves that inequality on the upward-closed region $\rho_c \leq \rho \leq 39/50$, $K \geq k_{11}(\rho)$ without interval arithmetic or a finite numerical certificate. The loss integral is first differentiated with respect to the spectral cutoff. A turning-point lower bound and an exact beta integral show that the scalar gap is strictly increasing in that cutoff for every $K > 12$. At the base curve, a rational secant majorant gives a uniform analytic estimate for the loss. The remaining one-variable inequality is then reduced to positivity of a degree-nine polynomial; all of its Bernstein coefficients on one rational interval are displayed and are positive. Thus there is no upper-frequency cutoff in the scalar-positivity theorem.

1 The scalar gap

For $0 \leq s \leq 1$, put

$$G_1(s) := \frac{1}{\pi} \left(\sqrt{1 - s^2} - s \arccos s \right). \quad (1)$$

Fix

$$\rho_c := \frac{1}{1 + 2\pi}, \quad k_{11}(\rho) := \sqrt{\frac{\pi^2}{(1 - \rho)^2} + 132}, \quad (2)$$

and consider

$$\mathcal{R}_c := \left\{ (\rho, K) : \rho_c \leq \rho \leq \frac{39}{50}, \quad K \geq k_{11}(\rho) \right\}. \quad (3)$$

The scalar gap left by the angular-block estimate is

$$\begin{aligned} \mathcal{S}(\rho, K) := & \frac{(1 + 2\rho^2)K^2}{12} - \mathcal{J}(\rho, K) - \frac{\pi\rho K}{4(1 - \rho)} + \frac{1}{2} \\ & + \frac{2\pi\rho K}{\arccos \rho} \left(\frac{KG_1(\rho)}{4(1 + 2KG_1(\rho))} + \frac{3}{32} \right), \end{aligned} \quad (4)$$

where

$$\mathcal{J}(\rho, K) := \frac{K^2}{2} \int_{\rho}^1 \frac{s}{(1 + 2KG_1(s))^2} ds. \quad (5)$$

Theorem 1 (Upward-closed scalar positivity). *For every $\rho_c \leq \rho \leq 39/50$ and every $K \geq k_{11}(\rho)$,*

$$\boxed{\mathcal{S}(\rho, K) > 0}. \quad (6)$$

We use only the elementary rational estimates

$$\frac{157}{50} < \pi < \frac{22}{7}, \quad \frac{140}{99} < \sqrt{2}, \quad \frac{5}{3} < \sqrt{3}. \quad (7)$$

The two square-root estimates follow immediately by squaring; for the closer one we record the exact margin

$$2 \cdot 99^2 - 140^2 = 2 > 0.$$

The displayed bounds for π are the classical Archimedean bounds (with the lower one weakened to $157/50$).

2 Strict monotonicity in the spectral cutoff

The identity

$$G_1(s) = \frac{1}{\pi} \int_s^1 \arccos u \, du \quad (8)$$

and $1 - \cos v \leq v^2/2$ imply

$$G_1(s) \geq \frac{2\sqrt{2}}{3\pi} (1-s)^{3/2}. \quad (9)$$

Indeed, $\arccos u \geq \sqrt{2(1-u)}$, and integration gives (9).

Lemma 2 (Derivative of the boundary-layer loss). *For every fixed $0 < \rho < 1$ and every $K > 0$,*

$$\partial_K \mathcal{J}(\rho, K) = K \int_\rho^1 \frac{s}{(1 + 2KG_1(s))^3} ds. \quad (10)$$

Moreover, if $K > 12$, then

$$\int_\rho^1 \frac{s}{(1 + 2KG_1(s))^3} ds < \frac{88}{567}. \quad (11)$$

Proof. Differentiating (5) under the integral sign gives

$$\begin{aligned} \partial_K \mathcal{J} &= K \int_\rho^1 \frac{s}{(1 + 2KG_1(s))^2} ds - 2K^2 \int_\rho^1 \frac{sG_1(s)}{(1 + 2KG_1(s))^3} ds \\ &= K \int_\rho^1 \frac{s}{(1 + 2KG_1(s))^3} ds, \end{aligned}$$

which proves (10).

Set $a = 4\sqrt{2}/(3\pi)$. By (9), the change of variable $t = 1 - s$, and $s \leq 1$,

$$\begin{aligned} \int_\rho^1 \frac{s \, ds}{(1 + 2KG_1(s))^3} &\leq \int_0^\infty \frac{dt}{(1 + aKt^{3/2})^3} \\ &= \frac{8\pi}{27\sqrt{3}} (aK)^{-2/3}. \end{aligned} \quad (12)$$

The last equality is the beta integral

$$\frac{2}{3} (aK)^{-2/3} B\left(\frac{2}{3}, \frac{7}{3}\right) = \frac{8\pi}{27\sqrt{3}} (aK)^{-2/3}.$$

At $K = 12$, the cube of the last member of (12) is

$$\frac{\pi^5}{59049\sqrt{3}}.$$

Using (7),

$$\frac{\pi^5}{59049\sqrt{3}} < \frac{5153632}{1654060905} < \left(\frac{88}{567}\right)^3,$$

where the second strict inequality is the exact identity

$$\left(\frac{88}{567}\right)^3 - \frac{5153632}{1654060905} = \frac{27812576}{44659644435} > 0.$$

The right side of (12) decreases with K , proving (11). $\square$

Proposition 3 (Monotonicity of the scalar gap). *For every $\rho_c \leq \rho \leq 39/50$ and every $K \geq k_{11}(\rho)$,*

$$\partial_K \mathcal{S}(\rho, K) > 0. \quad (13)$$

Consequently,

$$\mathcal{S}(\rho, K) \geq \mathcal{S}(\rho, k_{11}(\rho)). \quad (14)$$

Proof. Write

$$\theta := \arccos \rho, \quad y := KG_1(\rho).$$

The derivative of the final line of (4) is

$$\frac{2\pi\rho}{\theta} \left(\frac{y(1+y)}{2(1+2y)^2} + \frac{3}{32} \right) \geq \frac{3\rho}{8}, \quad (15)$$

because $\theta \leq \pi/2$. Also

$$-\frac{\pi\rho}{4(1-\rho)} > -\frac{11\rho}{14(1-\rho)}.$$

Every $K \geq k_{11}(\rho)$ in the stated ratio interval is greater than 12; this also follows from the stronger estimate established in Lemma 4 below. Lemma 2 therefore gives

$$\begin{aligned} \partial_K \mathcal{S} &> K \left(\frac{1+2\rho^2}{6} - \frac{88}{567} \right) + \frac{3\rho}{8} - \frac{11\rho}{14(1-\rho)} \\ &= K \left(\frac{13}{1134} + \frac{\rho^2}{3} \right) + \frac{3\rho}{8} - \frac{11\rho}{14(1-\rho)} \\ &> \frac{26}{189} + 4\rho^2 + \frac{3\rho}{8} - \frac{11\rho}{14(1-\rho)}. \end{aligned} \quad (16)$$

It remains to check the sign of the last expression. Multiplication by $1-\rho$ produces

$$p(\rho) := (1-\rho) \left(\frac{26}{189} + 4\rho^2 + \frac{3\rho}{8} \right) - \frac{11\rho}{14}. \quad (17)$$

The bound $\pi < 22/7$ gives $\rho_c > 7/51$. On the larger rational interval $[7/51, 39/50]$, set

$$u := \frac{\rho - 7/51}{39/50 - 7/51} \in [0, 1].$$

In the degree-three Bernstein basis $B_{j,3}(u) = \binom{3}{j}u^j(1-u)^{3-j}$, exact expansion gives

$$p(\rho) = \sum_{j=0}^3 b_j B_{j,3}(u), \quad (18)$$

with

$$\begin{array}{c|c} j & b_j \\ \hline 0 & \frac{335005}{32947387} \\ 1 & \frac{2785671}{281783183} \\ 2 & \frac{196635600}{578340000} \\ 3 & \frac{231517}{13500000} \end{array} \quad (19)$$

All four coefficients are positive. Since the Bernstein basis functions are nonnegative and sum to one, $p(\rho) > 0$. Equation (16) proves (13), and (14) follows by integration in K . $\square$

3 An exact rational upper bound for the loss

Lemma 4 (Uniform loss bound). *For every $\rho_c \leq \rho \leq 39/50$ and every $K \geq k_{11}(\rho)$,*

$$K > \frac{241}{20} \quad (20)$$

and

$$\int_{\rho}^1 \frac{x}{(1 + 2KG_1(x))^2} dx < \frac{17}{100}. \quad (21)$$

Proof. Since $\rho \geq \rho_c$,

$$\frac{\pi}{1-\rho} \geq \frac{\pi}{1-\rho_c} = \pi + \frac{1}{2} > \frac{91}{25}.$$

Consequently

$$K^2 \geq k_{11}(\rho)^2 > 132 + \left(\frac{91}{25}\right)^2 = \left(\frac{241}{20}\right)^2 + \frac{471}{10000},$$

which proves (20). The rational bounds in (7) now give

$$\frac{4\sqrt{2}K}{3\pi} > \frac{4(140/99)(241/20)}{3(22/7)} = \frac{23618}{3267} = \frac{36}{5} + \frac{478}{16335}. \quad (22)$$

Put $y = 1 - x$. Equations (9) and (22) yield

$$\begin{aligned} \int_{\rho}^1 \frac{x dx}{(1 + 2KG_1(x))^2} &< \int_0^{1-\rho} \frac{1-y}{(1 + (36/5)y^{3/2})^2} dy \\ &\leq \int_0^1 \frac{1-y}{(1 + (36/5)y^{3/2})^2} dy. \end{aligned} \quad (23)$$

We bound the last integral by exact rational arithmetic. Set

$$c := \frac{36}{5}, \quad \phi(z) := \frac{1}{(1+z)^2},$$

and use the rational grid

$$(t_i)_{i=0}^{10} = \left(0, \frac{1}{4}, \frac{5}{16}, \frac{3}{8}, \frac{7}{16}, \frac{1}{2}, \frac{9}{16}, \frac{5}{8}, \frac{3}{4}, \frac{7}{8}, 1\right). \quad (24)$$

For $z_i = ct_i^3$, let

$$\beta_i := \frac{\phi(z_{i+1}) - \phi(z_i)}{z_{i+1} - z_i}, \quad \alpha_i := \phi(z_i) - \beta_i z_i.$$

Because $\phi''(z) = 6(1+z)^{-4} > 0$, its secant satisfies

$$\phi(z) \leq \alpha_i + \beta_i z \quad (z_i \leq z \leq z_{i+1}).$$

After $y = t^2$, the integral in (23) is

$$\int_0^1 2t(1-t^2)\phi(ct^3) dt.$$

Thus it is at most

$$Q := \sum_{i=0}^9 (F_i(t_{i+1}) - F_i(t_i)), \quad (25)$$

where the exact polynomial antiderivative is

$$F_i(t) := \alpha_i \left(t^2 - \frac{t^4}{2}\right) + 2c\beta_i \left(\frac{t^5}{5} - \frac{t^7}{7}\right). \quad (26)$$

Every number in (25) is rational. Substitution of the grid (24) gives

$$Q = \frac{1305767238061446154064110434835938779253983373178558326887}{7706446260271137290663804190970969288616921855152043745280}, \quad (27)$$

and direct subtraction gives

$$\frac{17}{100} - Q = \frac{21643130923235926743681388145629999054466710986445549053}{38532231301355686453319020954854846443084609275760218726400} > 0. \quad (28)$$

Equations (23)–(28) prove (21). $\square$

4 The base curve

Fix ρ and now set

$$t := \sqrt{1-\rho}, \quad k := k_{11}(\rho) = \sqrt{132 + \frac{\pi^2}{t^4}}, \quad \rho = 1 - t^2. \quad (29)$$

The middle-ratio range lies strictly inside the rational interval

$$\frac{7}{15} < t < \frac{14}{15}. \quad (30)$$

Indeed,

$$t^2 \geq \frac{11}{50}, \quad \frac{11}{50} - \left(\frac{7}{15}\right)^2 = \frac{1}{450} > 0,$$

while $\rho \geq \rho_c > 7/51$ gives

$$t^2 = 1 - \rho < \frac{44}{51} < \left(\frac{14}{15}\right)^2, \quad \left(\frac{14}{15}\right)^2 - \frac{44}{51} = \frac{32}{3825} > 0.$$

Lemma 5 (Analytic endpoint payment). *For every $\rho_c \leq \rho \leq 39/50$, let t and k be as in (29), and let*

$$C(\rho) := \frac{100\rho^2 - 1}{600} \quad (31)$$

and

$$E(t) := \frac{11}{14t^2} - \frac{14}{15(1+2t)} - \frac{3}{8t}. \quad (32)$$

Then $C(\rho) > 0$, $E(t) > 0$, and

$$\mathcal{S}(\rho, k) > C(\rho)k^2 - \rho E(t)k + \frac{1}{2}. \quad (33)$$

Proof. First, $\rho \geq \rho_c > 7/51 > 1/10$, so $C(\rho) > 0$. Lemma 4 and

$$\frac{1+2\rho^2}{12} - \frac{17}{200} = \frac{100\rho^2 - 1}{600}$$

show that the first two terms in (4) are strictly greater than $C(\rho)k^2$.

Set $\theta = \arccos \rho$ and $Y = kG_1(\rho)$. For $0 \leq t \leq 1$,

$$\theta = \arccos(1 - t^2) = 2 \arcsin \frac{t}{\sqrt{2}} \leq \frac{\pi t}{2}. \quad (34)$$

To see the last inequality, use convexity of $\arcsin$ on $[0, 1/\sqrt{2}]$: its graph lies below the chord joining $(0, 0)$ to $(1/\sqrt{2}, \pi/4)$. Furthermore, (9) and $k > \pi/t^2$ give

$$Y > \frac{2\sqrt{2}}{3}t > \frac{14}{15}t. \quad (35)$$

Since $Y \mapsto Y/(1+2Y)$ is increasing, (34)–(35) imply

$$\begin{aligned} & \frac{2\pi\rho k}{\theta} \left(\frac{Y}{4(1+2Y)} + \frac{3}{32} \right) \\ & > \rho k \left(\frac{14}{15(1+2t)} + \frac{3}{8t} \right). \end{aligned} \quad (36)$$

Also

$$-\frac{\pi\rho k}{4t^2} > -\frac{11\rho k}{14t^2}.$$

Combining these inequalities with (4) proves (33).

Finally,

$$840t^2(1+2t)E(t) = 660 + 1005t - 1414t^2. \quad (37)$$

The quadratic on the right is concave, so its minimum on $[7/15, 14/15]$ occurs at an endpoint. Its endpoint values are

$$\frac{184739}{225}, \quad \frac{82406}{225},$$

respectively. Both are positive, proving $E(t) > 0$. $\square$

Proposition 6 (Rational positivity on the base curve). *For every $\rho_c \leq \rho \leq 39/50$,*

$$\mathcal{S}(\rho, k_{11}(\rho)) > 0. \quad (38)$$

Proof. The elementary inequality

$$\sqrt{x^2 + 132} < x + \frac{66}{x} \quad (x > 0)$$

and (7) give

$$k^2 > 132 + \frac{(157/50)^2}{t^4}, \quad (39)$$

$$k < \frac{22}{7t^2} + \frac{3300t^2}{157}. \quad (40)$$

Since $C(\rho) > 0$ and $\rho E(t) > 0$, Lemma 5 yields

$$\mathcal{S}(\rho, k) > R(t), \quad (41)$$

where

$$\begin{aligned} R(t) := & C(1 - t^2) \left(132 + \frac{(157/50)^2}{t^4} \right) \\ & - (1 - t^2)E(t) \left(\frac{22}{7t^2} + \frac{3300t^2}{157} \right) + \frac{1}{2}. \end{aligned} \quad (42)$$

It remains only exact polynomial algebra. Put

$$H(t) := t^4(1 + 2t)R(t).$$

Using (37), its factorized form is

$$\begin{aligned} H(t) = & C(1 - t^2)(1 + 2t) \left(132t^4 + \left(\frac{157}{50} \right)^2 \right) \\ & - \frac{(1 - t^2)(660 + 1005t - 1414t^2)}{840} \left(\frac{22}{7} + \frac{3300}{157}t^4 \right) \\ & + \frac{1}{2}t^4(1 + 2t). \end{aligned} \quad (43)$$

For an independently checkable expansion,

$$\begin{aligned} H(t) = & -\frac{20642567}{24500000} - \frac{6205067}{12250000}t + \frac{547983}{122500}t^2 - \frac{1033727}{367500}t^3 \\ & + \frac{34911551}{16485000}t^4 + \frac{187093801}{8242500}t^5 + \frac{8679}{1099}t^6 \\ & - \frac{138149}{2198}t^7 - \frac{2101}{157}t^8 + 44t^9. \end{aligned} \quad (44)$$

Set

$$u := \frac{t - 7/15}{7/15} \in [0, 1].$$

In the degree-nine Bernstein basis,

$$H(t) = \sum_{j=0}^9 d_j B_{j,9}(u), \quad B_{j,9}(u) = \binom{9}{j} u^j (1 - u)^{9-j}, \quad (45)$$

where exact conversion of (44) gives

j	d_j
0	$\frac{65408801807437}{9463832437500000}$
1	$\frac{299549381760793}{1135659892500000}$
2	$\frac{15904699805720773}{28391497312500000}$
3	$\frac{2765605256590021}{3154610812500000}$
4	$\frac{33203943051258761}{28391497312500000}$
5	$\frac{38636221171331747}{28391497312500000}$
6	$\frac{2538003206646073}{1892766487500000}$
7	$\frac{28758289199580211}{28391497312500000}$
8	$\frac{12920786461478803}{28391497312500000}$
9	$\frac{3428732855971679}{9463832437500000}$

(46)

Every d_j is positive. The Bernstein basis functions are nonnegative and sum to one; hence $H(t) > 0$ throughout $[7/15, 14/15]$. Since $t^4(1+2t) > 0$, this proves $R(t) > 0$. Equation (41) proves (38). $\square$

Proof of Theorem 1. Proposition 3 reduces every point of the upward-closed middle-ratio region to the base curve $K = k_{11}(\rho)$. Proposition 6 proves strict positivity on that curve. Therefore $\mathcal{S}(\rho, K) > 0$ throughout $\mathcal{R}_c$, with no upper restriction on K . $\square$

Remark 7 (Nature of the certificate). The rational secants and Bernstein expansions above are finite exact algebra, not interval arithmetic: convexity proves the functional majorant on each secant interval, and positivity follows from explicitly displayed rational coefficients. No floating-point evaluation, sampled minimum, or numerical root isolation is used as a premise.

Purely Analytic Closure of the Final Spherical-Shell Residual

Abstract

This note assembles the all-frequency retained-remainder estimate after the exact owner subtraction. It proves that the final spherical-shell residual is empty without invoking an interval certificate, an executable truth table, an aggregate tail argument, or a frequency split. The structural, angular-block, and scalar-positivity modules cover one closed superset of the entire residual.

1 Notation and analytic inputs

Let

$$A_{\rho,1} := \{x \in \mathbb{R}^3 : \rho < |x| < 1\}, \quad W(\rho, K) := \frac{2(1-\rho^3)K^3}{9\pi},$$

and let $N_D(A_{\rho,1}, K^2)$ count Dirichlet eigenvalues strictly below K^2 . Put

$$\rho_c := \frac{1}{1+2\pi}, \quad k_{11}(\rho) := \sqrt{\frac{\pi^2}{(1-\rho)^2} + 132}.$$

The preceding analytic owner lemmas and the exact set subtraction give

$$\mathcal{D}_{20} = \left\{(\rho, K) : \rho_c \leq \rho < \frac{39}{50}, \quad k_{11}(\rho) < K < U(\rho)\right\}. \quad (1)$$

On this interval $U(\rho) = K_0(\rho)$, although the formula for the upper wall will not be needed below.

We use the shell action

$$G_a(z) := \frac{1}{\pi} \left(\sqrt{a^2 - z^2} - z \arccos \frac{z}{a} \right) \quad (0 \leq z < a), \quad G_a(z) := 0 \quad (z \geq a),$$

$$A_{\rho,K}(z) := G_K(z) - G_{\rho K}(z),$$

and the strict lattice proxy

$$P(\rho, K) := \sum_{\substack{\nu=\ell+1/2 < K \\ \ell \in \mathbb{N}_0}} 2\nu \# \left\{ n \in \mathbb{N} : n < A_{\rho,K}(\nu) + \frac{1}{4} \right\}. \quad (2)$$

The analytic Bessel-phase reduction proved earlier in the main argument is

$$N_D(A_{\rho,1}, K^2) \leq P(\rho, K). \quad (3)$$

Put

$$M_c(\rho, K) := \frac{(1-\rho^2)K^2}{6} - \frac{1}{2}.$$

Parts VII–IX of the detailed analytic supplement establish, respectively, the following three inputs. The retained-remainder module gives the structural implication

$$\mathcal{E}_{\text{ang}}(\rho, K) < \mathcal{H}(\rho, K) \implies P(\rho, K) < W(\rho, K). \quad (4)$$

The paired angular-block module gives, on $\rho_c \leq \rho \leq 39/50$ and $K \geq k_{11}(\rho)$,

$$\mathcal{E}_{\text{ang}}(\rho, K) < M_c(\rho, K). \quad (5)$$

The scalar-positivity module gives on the same unbounded domain

$$\mathcal{H}(\rho, K) - M_c(\rho, K) > 0. \quad (6)$$

The all-frequency scope in (6) is part of that module: its scalar gap has positive K -derivative whenever $K > 12$, one has $k_{11}(\rho) > 241/20 > 12$, and the gap is strictly positive on the base curve $K = k_{11}(\rho)$. Thus the proof integrates upward from the base curve to every finite $K \geq k_{11}(\rho)$; it uses no upper-frequency bound.

2 All-frequency retained-remainder closure

Theorem 1 (All-frequency middle-ratio theorem). *For*

$$\rho_c \leq \rho \leq \frac{39}{50}, \quad K \geq k_{11}(\rho),$$

one has

$$N_D(A_{\rho,1}, K^2) < W(\rho, K).$$

Proof. Equations (5) and (6) give the strict chain

$$\mathcal{E}_{\text{ang}}(\rho, K) < M_c(\rho, K) < \mathcal{H}(\rho, K).$$

The structural implication (4) therefore gives $P(\rho, K) < W(\rho, K)$, and the phase reduction (3) completes the proof. Strictness includes both closed ratio faces, the base face $K = k_{11}(\rho)$, and every common radial–angular proxy wall. $\square$

3 Direct ownership of the exact residual

Proposition 2 (Analytic closure of $\mathcal{D}_{20}$). *Define the closed all-frequency theorem domain*

$$\mathcal{R}_{\text{tr}} := \left\{ (\rho, K) : \rho_c \leq \rho \leq \frac{39}{50}, \quad K \geq k_{11}(\rho) \right\}.$$

Then $\mathcal{D}_{20} \subset \mathcal{R}_{\text{tr}}$, the strict Pólya inequality holds throughout $\mathcal{D}_{20}$, and the successor residual is empty.

Proof. If $(\rho, K) \in \mathcal{D}_{20}$, then (1) gives

$$\rho_c \leq \rho < \frac{39}{50}, \quad k_{11}(\rho) < K < U(\rho).$$

After weakening the strict upper ratio and lower frequency inequalities, these conditions place (ρ, K) in $\mathcal{R}_{\text{tr}}$. Therefore Theorem 1 proves every point of $\mathcal{D}_{20}$, and

$$\mathcal{D}_{20} \setminus \mathcal{R}_{\text{tr}} = \emptyset.$$

The already owned faces $\rho = 39/50$, $K = k_{11}(\rho)$, and $K = U(\rho)$ remain outside $\mathcal{D}_{20}$, even though the stronger theorem also covers the first two. The face $\rho = \rho_c$ remains in $\mathcal{D}_{20}$ whenever both strict frequency inequalities hold. No artificial frequency face is present. $\square$

Logical dependency. After Parts I–VI identify $\mathcal{D}_{20}$, the closure uses only the structural, angular-block, and scalar-positivity modules in Parts VII–IX. It does not use an aggregate-tail module, a frequency split, the former 10,580-box compact certificate, the 1,286-box aggregate certificate, or the 63-state executable subtraction table.