

Probability

April 22, 2016

1 Probability Spaces

1.1 Classical Probability Spaces

Probability theory [5, 7, 16] is defined using the notions of a *sample space* Ω , a space of *events* \mathcal{E} , and a *probability measure* μ . In this paper, we will only consider *finite* sample spaces: we therefore define a sample space Ω as an arbitrary non-empty finite set, the space of events \mathcal{E} as 2^Ω , the powerset of Ω , and the *probability measure* as a function $\mu : \mathcal{E} \rightarrow [0, 1]$ such that:

- $\mu(\Omega) = 1$, and
- for a collection of pairwise disjoint events E_i , the probability measures are additive $\mu(\bigcup E_i) = \sum \mu(E_i)$.

Example of a problem on a finite sample space (Two coin experiment) Consider an experiment that tosses two coins. We have four possible outcomes that constitute the sample space $\Omega = \{HH, HT, TH, TT\}$. The event that the first coin is “heads” is $\{HH, HT\}$; the event that the two coins land on opposite sides is $\{HT, TH\}$; the event that at least one coin is tails is $\{HT, TH, TT\}$. Depending on the assumptions regarding the coins, we can define several probability measures. Here is a possible one:

$\mu(\emptyset)$	$= 0$	$\mu(\{HT, TH\})$	$= 2/3$
$\mu(\{HH\})$	$= 1/3$	$\mu(\{HT, TT\})$	$= 0$
$\mu(\{HT\})$	$= 0$	$\mu(\{TH, TT\})$	$= 2/3$
$\mu(\{TH\})$	$= 2/3$	$\mu(\{HH, HT, TH\})$	$= 1$
$\mu(\{TT\})$	$= 0$	$\mu(\{HH, HT, TT\})$	$= 1/3$
$\mu(\{HH, HT\})$	$= 1/3$	$\mu(\{HH, TH, TT\})$	$= 1$
$\mu(\{HH, TH\})$	$= 1$	$\mu(\{HT, TH, TT\})$	$= 2/3$
$\mu(\{HH, TT\})$	$= 1/3$	$\mu(\{HH, HT, TH, TT\})$	$= 1$

Note that the probability measure for disjoint events such as $\{HT\}$ and $\{TH\}$ do indeed add.

1.2 Quantum Probability Spaces

The mathematical framework above assumes that one has complete knowledge of the events and their relationships. But even in many classical situations, the structure of the event space is only partially known and the precise dependence of two events on each other cannot be determined with certainty. In the quantum case, this partial knowledge is compounded by the fact that not all quantum events can be observed simultaneously. Indeed, in the quantum world, there are non-commuting events which cannot even happen simultaneously. To accommodate these more complex situations, we abandon the sample space Ω and define and reason directly about events. A quantum probability space consist of just two components: a set of events \mathcal{E} and a probability measure $\mu : \mathcal{E} \rightarrow [0, 1]$. We give an example before giving the formal definition.

Consider the two-qubit Hilbert space with computational basis $|0\rangle$ and $|1\rangle$ and states:

$$|+\rangle = \frac{|0\rangle + |1\rangle}{\sqrt{2}}, \quad |-\rangle = \frac{|0\rangle - |1\rangle}{\sqrt{2}}.$$

The set of events associated with this Hilbert space consists of all projections including the empty projection \emptyset and the unit projection $\mathbb{1} = |0\rangle\langle 0| + |1\rangle\langle 1|$:

$$\{\emptyset, |0\rangle\langle 0|, |1\rangle\langle 1|, |+\rangle\langle +|, |-\rangle\langle -|, \dots, \mathbb{1}\}$$

Each event is interpreted as a possible post post-measurement state of a quantum system as follows: given some arbitrary current quantum state $|\psi\rangle$ to be measured, the event $|0\rangle\langle 0|$ states that the post-measurement state will be $|0\rangle$; the event $|1\rangle\langle 1|$ states that the post-measurement state will be $|1\rangle$; the event $|+\rangle\langle +|$ states that the post-measurement state will be $|+\rangle$; the event $|-\rangle\langle -|$ states that the post-measurement state will be $|-\rangle$; the event $\mathbb{1}$ states that the post-measurement state will be a linear combination of $|0\rangle$ and $|1\rangle$; and the event \emptyset states that the post-measurement state will be the empty state.

Irrespective of the current state $|\psi\rangle$ and irrespective of the particular experiment, the probability of event \emptyset will always be 0 (it is an impossible event) and the probability of event $\mathbb{1}$ will always be 1 (it is a certain event). The probabilities attached to other events will depend on the particular state in question. If the state is $|0\rangle$, the probability of event $|0\rangle\langle 0|$ is 1; the probability of event $|1\rangle\langle 1|$ is 0; the probability of event $|+\rangle\langle +|$ is $\frac{1}{2}$; and the probability of event $|-\rangle\langle -|$ is $\frac{1}{2}$. If the state is $|+\rangle$, the probability of each event $|0\rangle\langle 0|$ and $|1\rangle\langle 1|$ will be $\frac{1}{2}$; the probability of event $|+\rangle\langle +|$ is 1; and the probability of event $|-\rangle\langle -|$ is 0.

We now formalize a *quantum probability space* as follows [3, 6, 15, 1, 11]. We first assume an ambient Hilbert space \mathcal{H} and define the set of events \mathcal{E} as all *projections* on \mathcal{H} . Each quantum state $|\psi\rangle \in \mathcal{H} \setminus \{0\}$ induces a probability measure $\mu_\psi : \mathcal{E} \rightarrow [0, 1]$ on the space of events defined for any event $E \in \mathcal{E}$ as follows¹:

$$\mu_\psi(E) = \frac{\langle \psi | E \psi \rangle}{\langle \psi | \psi \rangle} \quad (1)$$

Similarly to the classical case, this probability measure must satisfy:

- $\mu(\mathbb{1}) = 1$, and
- for a collection of pairwise orthogonal E_i , we have $\mu(\sum_i E_i) = \sum_i \mu(E_i)$.

Yu-Tsung says: Claim: If $\sum_i |\psi_i\rangle\langle \psi_i|$ is a projection, then they are orthogonal.
Proof: If $\sum_i |\psi_i\rangle\langle \psi_i|$ is a projection, then

$$\begin{aligned} |\psi_j\rangle &= \left(\sum_i |\psi_i\rangle\langle \psi_i| \right) |\psi_j\rangle \\ &= \sum_i |\psi_i\rangle\langle \psi_i | \psi_j \rangle \end{aligned}$$

Therefore, we have

$$0 = \sum_{i \neq j} |\psi_i\rangle\langle \psi_i | \psi_j \rangle$$

1.3 Plan

Several assumptions are woven in the definition of a quantum probability space:

¹Recently, people extend the domain of μ_ψ to all operators \mathcal{A} on \mathcal{H} and consider $\mu_\psi : \mathcal{A} \rightarrow \mathbb{C}$ [11, 17]. When an operator $A \in \mathcal{A}$ is Hermitian, $\mu_\psi(A)$ is the expectation value of A . We do not take this approach because we want to focus only on probability.

- the Hilbert space \mathcal{H} ;
- the real interval $[0, 1]$;
- the fact that each state induces a probability measure, i.e., the Born rule (1) [4, 12];
- the fact that every probability measure is induced by a state, i.e., Gleason's theorem [6, 13, 15].

In the remainder of the paper, we examine each of these assumptions and consider variations motivated by computation of numerical quantities in a world with limited resources. In particular, we will consider a variant of the Hilbert space over finite fields $\mathbb{F}_{p^2}^d$ [10, 9, 8].

For the probability values, we will consider set-valued probability measures [2, 14], in particular $\mathcal{L}_2 = \{\text{impossible}, \text{possible}\}$, where impossible is a singleton set, $\{0\}$, and possible is an open interval, $(0, \infty)$.

If there may be more than one probability measure, we will discuss whether we will keep using the Born rule (1) or there is another formula $\tilde{\mu}$ such that $\tilde{\mu}_{|\psi\rangle}$ is a probability measure for all $|\psi\rangle \in \mathcal{H} \setminus \{0\}$.

Then, Gleason's theorem states given a probability measure μ there is a mixed state $|\psi\rangle$ such that $\mu = \mu_{|\psi\rangle}$, i.e., the Born rule is surjective². For any other formula $\tilde{\mu}$, we can ask whether $\tilde{\mu}$ is surjective as well.

Obviously, we don't want arbitrarily assign a state $|\psi\rangle \in \mathcal{H} \setminus \{0\}$ with a probability measure $\tilde{\mu}_{|\psi\rangle}$, for example, assigning every state to the same probability measure. We want $\tilde{\mu}_{|\psi\rangle}$ satisfying the following properties with some physical meaning:

- $\langle \psi | \phi \rangle = 0 \Leftrightarrow \tilde{\mu}_{|\psi\rangle}(|\phi\rangle\langle\phi|) = \tilde{0}$, where $\tilde{0}$ is 0 for $[0, 1]$ and $\tilde{0}$ is impossible for $\mathcal{L}_2 = \{\text{impossible}, \text{possible}\}$.
- $\tilde{\mu}_{|\psi\rangle}(|\phi\rangle\langle\phi|) = \tilde{\mu}_{|\mathbf{U}|\psi\rangle}(\mathbf{U}|\phi\rangle\langle\phi|\mathbf{U}^\dagger)$, where $|\psi\rangle, |\phi\rangle$ are states and \mathbf{U} is any unitary map, i.e., $\mathbf{U}^\dagger \mathbf{U} = \mathbb{1}$.

And the results can be summarized in the following table:

State space \mathcal{H}	Probability values	Is there a $\tilde{\mu}$ satisfying the given conditions?	Is the $\tilde{\mu}$ surjective?
\mathbb{C}^d for $d \geq 3$	$[0, 1]$	Yes	Yes
\mathbb{C}^d	\mathcal{L}_2	Yes	No
$\mathbb{F}_{p^2}^{d*}$ for $d \geq 3$ except $d = p = 3$	$[0, 1]$	No	
$\mathbb{F}_{p^2}^{d*}$	\mathcal{L}_2	Yes	No

References

- [1] Samson Abramsky. Big toy models: Representing physical systems as Chu spaces. *CoRR*, abs/0910.2393, 2009.
- [2] Zvi Artstein. Set-valued measures. *Transactions of the American Mathematical Society*, 165:103–125, 1972.
- [3] Garrett Birkhoff and John Von Neumann. The logic of quantum mechanics. *Annals of mathematics*, pages 823–843, 1936.
- [4] Max Born. Zur quantenmechanik der stoßvorgänge (1926). In *Die Deutungen der Quantentheorie*, pages 48–52. Springer, 1984.
- [5] William G. Faris. Appendix: Probability in quantum mechanics. In *The infamous boundary : seven decades of controversy in quantum physics*. Boston : Birkhauser, 1995.

²If we extend the domain of $\mu_{|\psi\rangle}$ including the mixed states.

- [6] Andrew Gleason. Measures on the closed subspaces of a hilbert space. *Indiana Univ. Math. J.*, 6:885–893, 1957.
- [7] R.L. Graham, D.E. Knuth, and O. Patashnik. *Concrete Mathematics: A Foundation for Computer Science*. Addison-Wesley, 1994.
- [8] Andrew J Hanson, Gerardo Ortiz, Amr Sabry, and Yu-Tsung Tai. Geometry of discrete quantum computing. *Journal of Physics A: Mathematical and Theoretical*, 46(18):185301, 2013.
- [9] Andrew J Hanson, Gerardo Ortiz, Amr Sabry, and Yu-Tsung Tai. Discrete quantum theories. *Journal of Physics A: Mathematical and Theoretical*, 47(11):115305, 2014.
- [10] Andrew J Hanson, Gerardo Ortiz, Amr Sabry, and Yu-Tsung Tai. Corrigendum: Geometry of discrete quantum computing. *Journal of Physics A: Mathematical and Theoretical*, 49(3):039501, 2015.
- [11] Hans Maassen. Quantum probability and quantum information theory. In *Quantum information, computation and cryptography*, pages 65–108. Springer, 2010.
- [12] N. D. Mermin. *Quantum Computer Science*. Cambridge University Press, 2007.
- [13] A. Peres. *Quantum Theory: Concepts and Methods*. Fundamental Theories of Physics. Springer, 1995.
- [14] Madan L Puri and Dan A Ralescu. Strong law of large numbers with respect to a set-valued probability measure. *The Annals of Probability*, pages 1051–1054, 1983.
- [15] Michael Redhead. *Incompleteness, Nonlocality, and Realism: A Prolegomenon to the Philosophy of Quantum Mechanics*. Oxford University Press, 1987.
- [16] V.K. Rohatgi and A.K.M.E. Saleh. *An Introduction to Probability and Statistics*. Wiley Series in Probability and Statistics. Wiley, 2011.
- [17] Jan Swart. Introduction to quantum probability. *Lecture Notes*, 2013.