

# Summary: Calculating Derivatives

## Derivatives of constant multiples

If  $g(x) = k f(x)$  for some constant  $k$ , then

$$g'(x) = k f'(x)$$

at all points where  $f$  is differentiable.

## Derivatives of sums

If  $h(x) = f(x) + g(x)$ , then

$$h'(x) = f'(x) + g'(x)$$

at all points where  $f$  and  $g$  are differentiable.

## Derivatives of differences

Similarly, if  $j(x) = f(x) - g(x)$ , then

$$j'(x) = f'(x) - g'(x)$$

at all points where  $f$  and  $g$  are differentiable.

## Derivatives of constant multiples - proof

Suppose that  $g(x) = kf(x)$  for all  $x$ , where  $k$  is a constant. We want to prove that  $g'(x) = kf'(x)$  at any point  $x$  where  $f$  is differentiable.

We know that

$$\begin{aligned}
g'(x) &= \lim_{\Delta x \rightarrow 0} \frac{g(x + \Delta x) - g(x)}{\Delta x} \\
&= \lim_{\Delta x \rightarrow 0} \frac{kf(x + \Delta x) - kf(x)}{\Delta x} \\
&= \lim_{\Delta x \rightarrow 0} k \frac{f(x + \Delta x) - f(x)}{\Delta x} \\
&= \lim_{\Delta x \rightarrow 0} k \lim_{\Delta x \rightarrow 0} \frac{f(x + \Delta x) - f(x)}{\Delta x}.
\end{aligned}$$

The first limit is just  $k$ , and the second limit is the definition of  $f'(x)$ . So we get  $g'(x) = kf'(x)$ .

## What is linearity?

We've seen that differentiation "respects" addition and multiplication by a constant. That is, if you take a derivative of a sum of functions, you get the same thing as if you differentiated each part, and then added the derivatives. Similarly, if you take the derivative of  $k$  times a function, where  $k$  is a constant, then you get  $k$  times the derivative of the original function.

Respecting addition and constant multiplication in this way is called "linearity," and it is an important property of the derivative operation!

## The Power Rule

If  $n$  is any fixed number, and  $f(x) = x^n$ , then  $f'(x) = nx^{n-1}$ .