

Assignment 1: Characterising a NAC AMOC Time Series

BBF0104yuxiangluan


Data choice

This analysis uses the North Atlantic Current (NAC) dataset from AMOCatlas and the TRANS_NAC_PROXY variable. This series is a six-monthly transport proxy in Sverdrups (Sv), based on satellite altimetry and float observations. I chose the NAC dataset because it provides an AMOC-relevant North Atlantic transport time series and does not duplicate the datasets selected by other students.

The record spans 1993-01-01 to 2025-07-02. It contains 66 samples, with a median sampling interval of 182.625 days. There are no missing values in the proxy series, so no interpolation was needed. The gap-filling step in the script is kept for reproducibility, but it leaves this particular series unchanged.


Time-domain statistics

The NAC transport proxy has a mean of 27.16 Sv and a standard deviation of 1.74 Sv. The median is 27.21 Sv. The minimum and maximum are 23.30 Sv and 30.63 Sv, giving a total range of 7.33 Sv. The histogram shows that the values are concentrated around the high-20s Sv, with variability of a few Sverdrups around the mean.



NAC transport proxy distribution

The raw time series shows interannual to decadal variability in the NAC transport proxy. Because the data are sampled every six months, I applied a 5-year Tukey-window low-pass filter using an 11-sample centred rolling window. This filter suppresses shorter interannual fluctuations and highlights slower multi-year changes in the transport proxy.




Raw and filtered NAC transport proxy

Frequency-domain statistics

I estimated the power spectral density using Welch's method with a Hann window, a 64-sample segment length, and no overlap. The record is short, so this configuration is close to a tapered periodogram, but it keeps nearly the full record length and gives a stable variance budget. The segment length corresponds to about 32 years because the sampling interval is approximately half a year.

The integrated spectrum satisfies the Parseval variance check. The ratio between integrated PSD and time-domain variance is 0.996 for the raw series, which is very close to 1. This indicates that the spectral normalisation is consistent with the variance of the time series. The largest spectral peak in this estimate corresponds to a timescale of about 32 years. Given the short record, this peak should be interpreted cautiously as low-frequency variability over the full record rather than as a precisely resolved oscillation.

The 5-year Tukey low-pass filter reduces power at higher frequencies while retaining the lower-frequency part of the spectrum. The filtered spectrum lies below the raw spectrum at shorter timescales, which is the expected behaviour of a low-pass filter designed to isolate slower AMOC-related transport variability.



Raw and filtered NAC transport proxy spectrum

Reproducibility

The analysis can be reproduced with:

```
python assignment_analysis.py  
pytest -q
```

The script downloads/loads the NAC dataset through AMOCatlas, writes the figures to figures/, and writes the numerical summary to outputs/assignment1_summary.json.