



mid_Egamma TxTx: max C_{py} regions



TxTx characteristic max $C_{p\gamma}$ configuration

3D momenta




c_p_gamma_mid_egamma_txtx_region_0 C_{p\gamma}=0.999461
 region: mid_Egamma TxTx_region_0 final-pair delta_xy=0 rad
 outgoing-state purity: 1
 kinematic point: high_s_high_theta_in_mid_Egamma
 incoming state: $|T_{x,e}T_{x,p}\rangle$

$s=21.5158$, $\sqrt{s}=4.63852$, $|\vec{P}|=2.22442$, $|\vec{P}'|=1.15253$
 $\theta_{in}=1.57079$, $\phi_l=4.71239$, $\phi_p=1.5708$
 $E_\gamma=1$, $\phi_\gamma=1.5708$
 $Q^2=2.90272e-10$, $x_B=4.35241e-10$, $t=-0.287559$, $W^2=1.54677$, $y=0.0323184$
 $\theta(l',\gamma)=180$ deg

four-momenta $[E, p_x, p_y, p_z]$ GeV:
 l $E= 2.2244$ $p_x= -0.0000$ $p_y= -2.2244$ $p_z= -0.0000$
 P $E= 2.4141$ $p_x= 0.0000$ $p_y= 2.2244$ $p_z= 0.0000$
 l' $E= 2.1525$ $p_x= -0.0000$ $p_y= -2.1525$ $p_z= 0.0000$
 P' $E= 1.4860$ $p_x= 0.0000$ $p_y= 1.1525$ $p_z= 0.0000$
 q_γ $E= 1.0000$ $p_x= 0.0000$ $p_y= 1.0000$ $p_z= 0.0000$

Transverse plane




$h_e=+1, h_p=-1, h_\gamma=+1$
 electron Tx, proton Tx

$h_e=-1, h_p=+1, h_\gamma=-1$
 electron Tx, proton Tx

$h_e=-1, h_p=-1, h_\gamma=+1$
 electron Tx, proton Tx


$h_e=+1, h_p=+1, h_\gamma=-1$
 electron Tx, proton Tx

Leading initial-ensemble/final-state components (N=4, retained=100.0%)



TxTx characteristic max $C_{p\gamma}$ configuration

3D momenta




c_p_gamma_mid_egamma_txtx_region_1 C_{p\gamma}=0.963626
 region: mid_Egamma TxTx_region_1 final-pair delta_xy=1.37445 rad
 outgoing-state purity: 1
 kinematic point: high_s_high_theta_in_mid_Egamma
 incoming state: $|T_{X,e}T_{X,p}\rangle$

$s=21.5158$, $\sqrt{s}=4.63852$, $|\vec{P}|=2.22442$, $|\vec{P}'|=1.44782$
 $\theta_{in}=1.57079$, $\phi_l=4.71239$, $\phi_P=1.5708$
 $E_\gamma=1$, $\phi_\gamma=0.19635$
 $Q^2=1.20336$, $x_B=0.294314$, $t=-0.128405$, $W^2=3.76517$, $y=0.198133$
 $\theta(l', \gamma)=132.086$ deg

four-momenta $[E, p_x, p_y, p_z]$ GeV:
 l $E= 2.2244$ $p_x= -0.0000$ $p_y= -2.2244$ $p_z= -0.0000$
 P $E= 2.4141$ $p_x= 0.0000$ $p_y= 2.2244$ $p_z= 0.0000$
 l' $E= 1.9134$ $p_x= -0.9808$ $p_y= -1.6429$ $p_z= 0.0000$
 P' $E= 1.7251$ $p_x= 0.0000$ $p_y= 1.4478$ $p_z= 0.0000$
 q_γ $E= 1.0000$ $p_x= 0.9808$ $p_y= 0.1951$ $p_z= 0.0000$

Transverse plane




$h_e=-1, h_p=+1, h_\gamma=-1$
 electron Tx, proton Tx

$h_e=+1, h_p=-1, h_\gamma=+1$
 electron Tx, proton Tx

$h_e=-1, h_p=-1, h_\gamma=+1$
 electron Tx, proton Tx


$h_e=+1, h_p=+1, h_\gamma=-1$
 electron Tx, proton Tx

Leading initial-ensemble/final-state components (N=4, retained=99.4%)



TxTx characteristic max $C_{p\gamma}$ configuration

3D momenta




c_p_gamma_mid_egamma_txtx_region_2 C_{p\gamma}=0.963626
 region: mid_Egamma TxTx_region_2 final-pair delta_xy=1.37445 rad
 outgoing-state purity: 1
 kinematic point: high_s_high_theta_in_mid_Egamma
 incoming state: $|T_{X,e}T_{X,p}\rangle$

$s=21.5158$, $\sqrt{s}=4.63852$, $|\vec{P}|=2.22442$, $|\vec{P}'|=1.44782$
 $\theta_{in}=1.57079$, $\phi_l=4.71239$, $\phi_P=1.5708$
 $E_\gamma=1$, $\phi_\gamma=2.94524$
 $Q^2=1.20336$, $x_B=0.294314$, $t=-0.128405$, $W^2=3.76517$, $y=0.198133$
 $\theta(l', \gamma)=132.086$ deg

four-momenta $[E, p_x, p_y, p_z]$ GeV:

l $E= 2.2244$ $p_x= -0.0000$ $p_y= -2.2244$ $p_z= -0.0000$
 P $E= 2.4141$ $p_x= 0.0000$ $p_y= 2.2244$ $p_z= 0.0000$
 l' $E= 1.9134$ $p_x= 0.9808$ $p_y= -1.6429$ $p_z= 0.0000$
 P' $E= 1.7251$ $p_x= 0.0000$ $p_y= 1.4478$ $p_z= 0.0000$
 q_γ $E= 1.0000$ $p_x= -0.9808$ $p_y= 0.1951$ $p_z= 0.0000$

Transverse plane



$h_e=+1, h_p=-1, h_\gamma=+1$
 electron Tx, proton Tx

$h_e=-1, h_p=+1, h_\gamma=-1$
 electron Tx, proton Tx

$h_e=+1, h_p=+1, h_\gamma=-1$
 electron Tx, proton Tx

$h_e=-1, h_p=-1, h_\gamma=+1$
 electron Tx, proton Tx

Leading initial-ensemble/final-state components (N=4, retained=99.4%)

