


# Ty electron: $C_{ep}$ two-angle scans at characteristic kinematics

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_\gamma = 0.25 \text{ GeV}$ ,  $\max C_{ep} = 0.000$

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_\gamma = 1 \text{ GeV}$ ,  $\max C_{ep} = 0.000$

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_\gamma = 1.8 \text{ GeV}$ ,  $\max C_{ep} = 0.000$




# Ty electron: $C_{e\gamma}$ two-angle scans at characteristic kinematics

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_\gamma = 0.25 \text{ GeV}$ ,  $\max C_{e\gamma} = 1.000$

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_\gamma = 1 \text{ GeV}$ ,  $\max C_{e\gamma} = 1.000$

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_\gamma = 1.8 \text{ GeV}$ ,  $\max C_{e\gamma} = 1.000$




# Ty electron: $C_{p\gamma}$ two-angle scans at characteristic kinematics

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_\gamma = 0.25 \text{ GeV}$ ,  $\max C_{p\gamma} = 0.000$

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_\gamma = 1 \text{ GeV}$ ,  $\max C_{p\gamma} = 0.000$

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_\gamma = 1.8 \text{ GeV}$ ,  $\max C_{p\gamma} = 0.000$




# Ty electron: $C_{e|p\gamma}$ two-angle scans at characteristic kinematics

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_\gamma = 0.25 \text{ GeV}$ ,  $\max C_{e|p\gamma} = 0.000$

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_\gamma = 1 \text{ GeV}$ ,  $\max C_{e|p\gamma} = 0.000$

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_\gamma = 1.8 \text{ GeV}$ ,  $\max C_{e|p\gamma} = 0.000$




# Ty electron: $C_{p|e\gamma}$ two-angle scans at characteristic kinematics

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_\gamma = 0.25 \text{ GeV}$ ,  $\max C_{p|e\gamma} = 0.000$

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_\gamma = 1 \text{ GeV}$ ,  $\max C_{p|e\gamma} = 0.000$

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_\gamma = 1.8 \text{ GeV}$ ,  $\max C_{p|e\gamma} = 0.000$




# Ty electron: $C_{\gamma|ep}$ two-angle scans at characteristic kinematics

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_\gamma = 0.25 \text{ GeV}$ ,  $\max C_{\gamma|ep} = 0.000$

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_\gamma = 1 \text{ GeV}$ ,  $\max C_{\gamma|ep} = 0.000$

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_\gamma = 1.8 \text{ GeV}$ ,  $\max C_{\gamma|ep} = 0.000$




# Ty electron: $M_e$ two-angle scans at characteristic kinematics

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_Y = 0.25 \text{ GeV}$ ,  $\max M_e = -0.000$

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_Y = 1 \text{ GeV}$ ,  $\max M_e = -0.000$

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_Y = 1.8 \text{ GeV}$ ,  $\max M_e = -0.000$




# Ty electron: $M_p$ two-angle scans at characteristic kinematics

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_\gamma = 0.25 \text{ GeV}$ ,  $\max M_p = -0.000$

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_\gamma = 1 \text{ GeV}$ ,  $\max M_p = -0.000$

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_\gamma = 1.8 \text{ GeV}$ ,  $\max M_p = -0.000$






# Ty electron: $M_Y$ two-angle scans at characteristic kinematics

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_Y = 0.25 \text{ GeV}$ ,  $\max M_Y = -0.000$

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_Y = 1 \text{ GeV}$ ,  $\max M_Y = -0.000$

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_Y = 1.8 \text{ GeV}$ ,  $\max M_Y = -0.000$




# Ty electron: $F_3$ two-angle scans at characteristic kinematics

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_\gamma = 0.25 \text{ GeV}$ ,  $\max F_3 = 0.000$

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_\gamma = 1 \text{ GeV}$ ,  $\max F_3 = 0.000$

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_\gamma = 1.8 \text{ GeV}$ ,  $\max F_3 = 0.000$




# Ty electron: $P_{\text{GHZ}+}$ + two-angle scans at characteristic kinematics

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_Y = 0.25 \text{ GeV}$ ,  $\max P_{\text{GHZ}+} = 0.280$

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_Y = 1 \text{ GeV}$ ,  $\max P_{\text{GHZ}+} = 0.384$

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_Y = 1.8 \text{ GeV}$ ,  $\max P_{\text{GHZ}+} = 0.483$



# Ty electron: $P_W$ two-angle scans at characteristic kinematics

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_Y = 0.25 \text{ GeV}$ ,  $\max P_W = 0.307$

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_Y = 1 \text{ GeV}$ ,  $\max P_W = 0.307$

$s = 21.5 \text{ GeV}^2$ ,  $\theta_{\text{in}} = 1.57 \text{ rad}$   
 $E_Y = 1.8 \text{ GeV}$ ,  $\max P_W = 0.307$

