



mid\_Egamma L proton: max  $C_{p\gamma}$  regions



# L proton characteristic max $C_{p\gamma}$ configuration

3D momenta




c\_p\_gamma\_mid\_egamma\_l\_proton\_region\_0 C\_{p\gamma}=0.617601  
 region: mid\_Egamma\_L\_proton\_region\_0 final-pair delta\_xy=2.94524 rad  
 outgoing-state purity: 0.531619  
 kinematic point: high\_s\_high\_theta\_in\_mid\_Egamma  
 incoming state:  $\frac{1}{2}\sum_{h_e} \rho(h_e, L_p)$


$s=21.5158$ ,  $\sqrt{s}=4.63852$ ,  $|\vec{P}|=2.22442$ ,  $|\vec{P}'|=2.20638$   
 $\theta_{in}=1.57079$ ,  $\phi_i=4.71239$ ,  $\phi_p=1.5708$   
 $E_\gamma=1$ ,  $\phi_\gamma=4.51604$   
 $Q^2=0.0686461$ ,  $x_B=0.00746835$ ,  $t=-4.94517e-05$ ,  $W^2=10.0028$ ,  $y=0.445416$   
 $\theta(l', \gamma)=20.2945$  deg

four-momenta  $[E, p_x, p_y, p_z]$  GeV:  
 $l$   $E= 2.2244$   $p_x= -0.0000$   $p_y= -2.2244$   $p_z= -0.0000$   
 $P$   $E= 2.4141$   $p_x= 0.0000$   $p_y= 2.2244$   $p_z= 0.0000$   
 $l'$   $E= 1.2410$   $p_x= 0.1951$   $p_y= -1.2256$   $p_z= 0.0000$   
 $P'$   $E= 2.3975$   $p_x= 0.0000$   $p_y= 2.2064$   $p_z= 0.0000$   
 $q_\gamma$   $E= 1.0000$   $p_x= -0.1951$   $p_y= -0.9808$   $p_z= 0.0000$

Transverse plane




Leading initial-ensemble/final-state components (N=4, retained=100.0%)



# L proton characteristic max $C_{p\gamma}$ configuration

3D momenta




c\_p\_gamma\_mid\_egamma\_l\_proton\_region\_1 C\_{p\gamma}=0.313279  
 region: mid\_Egamma\_L\_proton\_region\_1 final-pair delta\_xy=1.5708 rad  
 outgoing-state purity: 0.528905  
 kinematic point: high\_s\_high\_theta\_in\_mid\_Egamma  
 incoming state:  $\frac{1}{2}\sum_{h_e}\rho(h_e, L_p)$

$s=21.5158$ ,  $\sqrt{s}=4.63852$ ,  $|\vec{P}|=2.22442$ ,  $|\vec{P}'|=1.5395$   
 $\theta_{in}=1.57079$ ,  $\phi_i=4.61421$ ,  $\phi_p=1.47262$   
 $E_\gamma=1$ ,  $\phi_\gamma=3.14159$   
 $Q^2=1.78711$ ,  $x_B=0.3314$ ,  $t=-0.128343$ ,  $W^2=4.48534$ ,  $y=0.26132$   
 $\theta(\ell', \gamma)=123.006$  deg


four-momenta  $[E, p_x, p_y, p_z]$  GeV:

$\ell$   $E= 2.2244$   $p_x= -0.2180$   $p_y= -2.2137$   $p_z= -0.0000$   
 $P$   $E= 2.4141$   $p_x= 0.2180$   $p_y= 2.2137$   $p_z= 0.0000$   
 $\ell'$   $E= 1.8358$   $p_x= 1.0000$   $p_y= -1.5395$   $p_z= 0.0000$   
 $P'$   $E= 1.8027$   $p_x= 0.0000$   $p_y= 1.5395$   $p_z= 0.0000$   
 $q_\gamma$   $E= 1.0000$   $p_x= -1.0000$   $p_y= 0.0000$   $p_z= 0.0000$

Transverse plane




Leading initial-ensemble/final-state components (N=6, retained=99.5%)



# L proton characteristic max $C_{p\gamma}$ configuration

3D momenta




c\_p\_gamma\_mid\_egamma\_l\_proton\_region\_2 C\_{p\gamma}=0.313279  
 region: mid\_Egamma\_L\_proton\_region\_2 final-pair delta\_xy=1.5708 rad  
 outgoing-state purity: 0.528905  
 kinematic point: high\_s\_high\_theta\_in\_mid\_Egamma  
 incoming state:  $\frac{1}{2}\sum_{h_e} \rho(h_e, L_p)$

$s=21.5158$ ,  $\sqrt{s}=4.63852$ ,  $|\vec{P}|=2.22442$ ,  $|\vec{P}'|=1.5395$   
 $\theta_{in}=1.57079$ ,  $\phi_l=4.81056$ ,  $\phi_p=1.66897$   
 $E_\gamma=1$ ,  $\phi_\gamma=0$   
 $Q^2=1.78711$ ,  $x_B=0.3314$ ,  $t=-0.128343$ ,  $W^2=4.48534$ ,  $y=0.26132$   
 $\theta(l', \gamma)=123.006$  deg

four-momenta  $[E, p_x, p_y, p_z]$  GeV:

$l$   $E= 2.2244$   $p_x= 0.2180$   $p_y= -2.2137$   $p_z= -0.0000$   
 $P$   $E= 2.4141$   $p_x= -0.2180$   $p_y= 2.2137$   $p_z= 0.0000$   
 $l'$   $E= 1.8358$   $p_x= -1.0000$   $p_y= -1.5395$   $p_z= 0.0000$   
 $P'$   $E= 1.8027$   $p_x= 0.0000$   $p_y= 1.5395$   $p_z= 0.0000$   
 $q_\gamma$   $E= 1.0000$   $p_x= 1.0000$   $p_y= 0.0000$   $p_z= 0.0000$

Transverse plane



Leading initial-ensemble/final-state components (N=6, retained=99.5%)

