



low_Egamma L massless lepton + L proton: max $C_{p\gamma}$ regions



L massless lepton + L proton characteristic max $C_{p\gamma}$ configuration

3D momenta




c_p_gamma_low_egamma_ll_region_0 C_{p\gamma}=0.981704
region: low_Egamma_LL_region_0 final-pair delta_xy=2.84707 rad
outgoing-state purity: 1
kinematic point: high_s_high_theta_in_low_Egamma
incoming state: $|L_{\ell_0}\rangle \otimes |L_p\rangle$

$s=21.5158$, $\sqrt{s}=4.63852$, $|\vec{P}|=2.22442$, $|\vec{P}'|=2.21812$
 $\theta_{in}=1.57079$, $\phi_{\ell}=4.71239$, $\phi_P=1.5708$
 $E_{\gamma}=0.25$, $\phi_{\gamma}=5.00691$
 $Q^2=0.00591804$, $x_B=0.00260549$, $t=-5.99798e-06$, $W^2=3.1453$, $y=0.110068$
 $\theta(\ell', \gamma)=18.9753$ deg

four-momenta $[E, p_x, p_y, p_z]$ GeV:
 ℓ $E= 2.2244$ $p_x= -0.0000$ $p_y= -2.2244$ $p_z= -0.0000$
 P $E= 2.4141$ $p_x= 0.0000$ $p_y= 2.2244$ $p_z= 0.0000$
 ℓ' $E= 1.9802$ $p_x= -0.0726$ $p_y= -1.9789$ $p_z= 0.0000$
 P' $E= 2.4083$ $p_x= 0.0000$ $p_y= 2.2181$ $p_z= 0.0000$
 q_{γ} $E= 0.2500$ $p_x= 0.0726$ $p_y= -0.2392$ $p_z= 0.0000$


Transverse plane



$h_e=+1, h_p=-1, h_{\gamma}=+1$
massless lepton L, proton L


$h_e=+1, h_p=+1, h_{\gamma}=-1$
massless lepton L, proton L

Leading initial-ensemble/final-state components (N=2, retained=98.5%)



L massless lepton + L proton characteristic max $C_{p\gamma}$ configuration

3D momenta




c_p_gamma_low_egamma_ll_region_1 C_{p\gamma}=0.438613
 region: low_Egamma_LL_region_1 final-pair delta_xy=2.06167 rad
 outgoing-state purity: 1
 kinematic point: high_s_high_theta_in_low_Egamma
 incoming state: $|L_{t_0}\rangle \otimes |L_p\rangle$

$s=21.5158$, $\sqrt{s}=4.63852$, $|\vec{P}|=2.22442$, $|\vec{P}'|=2.14934$
 $\theta_{in}=1.57079$, $\phi_l=4.71239$, $\phi_p=1.5708$
 $E_\gamma=0.25$, $\phi_\gamma=3.63247$
 $Q^2=0.0530724$, $x_B=0.0306384$, $t=-0.000876114$, $W^2=2.55899$, $y=0.0839415$
 $\theta(l', \gamma)=68.0691$ deg

four-momenta $[E, p_x, p_y, p_z]$ GeV:
 ℓ $E= 2.2244$ $p_x= -0.0000$ $p_y= -2.2244$ $p_z= -0.0000$
 P $E= 2.4141$ $p_x= 0.0000$ $p_y= 2.2244$ $p_z= 0.0000$
 ℓ' $E= 2.0434$ $p_x= 0.2205$ $p_y= -2.0315$ $p_z= 0.0000$
 P' $E= 2.3451$ $p_x= 0.0000$ $p_y= 2.1493$ $p_z= 0.0000$
 q_γ $E= 0.2500$ $p_x= -0.2205$ $p_y= -0.1178$ $p_z= 0.0000$


Transverse plane



$h_e=+1, h_p=-1, h_\gamma=+1$
 massless lepton L, proton L


$h_e=+1, h_p=+1, h_\gamma=-1$
 massless lepton L, proton L

Leading initial-ensemble/final-state components (N=2, retained=100.0%)



L massless lepton + L proton characteristic max $C_{p\gamma}$ configuration

3D momenta




c_p_gamma_low_egamma_ll_region_2 C_{p\gamma}=0.255296
 region: low_Egamma_LL_region_2 final-pair delta_xy=1.47262 rad
 outgoing-state purity: 1
 kinematic point: high_s_high_theta_in_low_Egamma
 incoming state: $|L_{\ell_0}\rangle \otimes |L_p\rangle$

$s=21.5158$, $\sqrt{s}=4.63852$, $|\vec{P}|=2.22442$, $|\vec{P}'|=2.07351$
 $\theta_{in}=1.57079$, $\phi_{\ell}=4.71239$, $\phi_P=1.5708$
 $E_{\gamma}=0.25$, $\phi_{\gamma}=0.0981748$
 $Q^2=0.0653998$, $x_B=0.0593636$, $t=-0.00364768$, $W^2=1.91613$, $y=0.0533864$
 $\theta(\ell', \gamma)=102.388$ deg


four-momenta $[E, p_x, p_y, p_z]$ GeV:
 ℓ $E= 2.2244$ $p_x= -0.0000$ $p_y= -2.2244$ $p_z= -0.0000$
 P $E= 2.4141$ $p_x= 0.0000$ $p_y= 2.2244$ $p_z= 0.0000$
 ℓ' $E= 2.1127$ $p_x= -0.2488$ $p_y= -2.0980$ $p_z= 0.0000$
 P' $E= 2.2758$ $p_x= 0.0000$ $p_y= 2.0735$ $p_z= 0.0000$
 q_{γ} $E= 0.2500$ $p_x= 0.2488$ $p_y= 0.0245$ $p_z= 0.0000$

Transverse plane




$h_e=+1$, $h_p=-1$, $h_{\gamma}=+1$
 massless lepton L, proton L

Leading initial-ensemble/final-state components (N=1, retained=98.3%)



L massless lepton + L proton characteristic max $C_{p\gamma}$ configuration

3D momenta




c_p_gamma_low_egamma_ll_region_3 C_{p\gamma}=0.106191
 region: low_Egamma_LL_region_3 final-pair delta_xy=0.687223 rad
 outgoing-state purity: 1
 kinematic point: high_s_high_theta_in_low_Egamma
 incoming state: $|L_{t_0}\rangle \otimes |L_p\rangle$

$s=21.5158$, $\sqrt{s}=4.63852$, $|\vec{P}|=2.22442$, $|\vec{P}'|=1.98975$
 $\theta_{in}=1.57079$, $\phi_l=4.71239$, $\phi_p=1.5708$
 $E_\gamma=0.25$, $\phi_\gamma=2.25802$
 $Q^2=0.0255969$, $x_B=0.0718149$, $t=-0.00912784$, $W^2=1.21068$, $y=0.0172722$
 $\theta(l', \gamma)=144.78$ deg

four-momenta $[E, p_x, p_y, p_z]$ GeV:
 l $E= 2.2244$ $p_x= -0.0000$ $p_y= -2.2244$ $p_z= -0.0000$
 P $E= 2.4141$ $p_x= 0.0000$ $p_y= 2.2244$ $p_z= 0.0000$
 l' $E= 2.1888$ $p_x= 0.1586$ $p_y= -2.1830$ $p_z= 0.0000$
 P' $E= 2.1998$ $p_x= 0.0000$ $p_y= 1.9898$ $p_z= 0.0000$
 q_γ $E= 0.2500$ $p_x= -0.1586$ $p_y= 0.1933$ $p_z= 0.0000$

Transverse plane



$h_e=+1$, $h_p=-1$, $h_\gamma=+1$
 massless lepton L, proton L

Leading initial-ensemble/final-state components (N=1, retained=99.7%)

