



low_Egamma Tx massless lepton + Tx proton: max C_{py} regions



Tx massless lepton + Tx proton characteristic max $C_{p\gamma}$ configuration

3D momenta




c_p_gamma_low_egamma_txtx_region_0 C_{p\gamma}=0.999987
 region: low_Egamma_TxTx_region_0 final-pair delta_xy=0 rad
 outgoing-state purity: 1
 kinematic point: high_s_high_theta_in_low_Egamma
 incoming state: $|T_{x,\ell_0}\rangle \otimes |\bar{T}_{x,p}\rangle$

$s=21.5158$, $\sqrt{s}=4.63852$, $|\vec{P}|=2.22442$, $|\vec{P}'|=1.96296$
 $\theta_{in}=1.57079$, $\phi_\ell=4.71239$, $\phi_P=1.5708$
 $E_\gamma=0.25$, $\phi_\gamma=1.5708$
 $Q^2=8.66557e-12$, $x_B=8.15204e-11$, $t=-0.0114583$, $W^2=0.986143$, $y=0.00515116$
 $\theta(\ell', \gamma)=180$ deg

four-momenta $[E, p_x, p_y, p_z]$ GeV:
 ℓ $E= 2.2244$ $p_x= -0.0000$ $p_y= -2.2244$ $p_z= -0.0000$
 P $E= 2.4141$ $p_x= 0.0000$ $p_y= 2.2244$ $p_z= 0.0000$
 ℓ' $E= 2.2130$ $p_x= -0.0000$ $p_y= -2.2130$ $p_z= 0.0000$
 P' $E= 2.1756$ $p_x= 0.0000$ $p_y= 1.9630$ $p_z= 0.0000$
 q_γ $E= 0.2500$ $p_x= 0.0000$ $p_y= 0.2500$ $p_z= 0.0000$

Transverse plane




$h_e=-1, h_p=+1, h_\gamma=-1$
 massless lepton Tx, proton Tx

$h_e=+1, h_p=-1, h_\gamma=+1$
 massless lepton Tx, proton Tx

$h_e=+1, h_p=+1, h_\gamma=-1$
 massless lepton Tx, proton Tx


$h_e=-1, h_p=-1, h_\gamma=+1$
 massless lepton Tx, proton Tx

Leading initial-ensemble/final-state components (N=4, retained=100.0%)



Tx massless lepton + Tx proton characteristic max $C_{p\gamma}$ configuration

3D momenta




c_p_gamma_low_egamma_txtx_region_1 $C_{p\gamma}=0.984344$
 region: low_Egamma_TxTx_region_1 final-pair delta_xy=1.37445 rad
 outgoing-state purity: 1
 kinematic point: high_s_high_theta_in_low_Egamma
 incoming state: $|T_{x,\ell_0}\rangle \otimes |T_{x,p}\rangle$


$s=21.5158$, $\sqrt{s}=4.63852$, $|\vec{P}|=2.22442$, $|\vec{P}'|=2.06107$
 $\theta_{in}=1.57079$, $\phi_\ell=4.71239$, $\phi_P=1.5708$
 $E_\gamma=0.25$, $\phi_\gamma=0.19635$
 $Q^2=0.0631736$, $x_B=0.0635316$, $t=-0.00429535$, $W^2=1.81103$, $y=0.0481859$
 $\theta(\ell', \gamma)=107.879$ deg

four-momenta $[E, p_x, p_y, p_z]$ GeV:
 ℓ $E= 2.2244$ $p_x= -0.0000$ $p_y= -2.2244$ $p_z= -0.0000$
 P $E= 2.4141$ $p_x= 0.0000$ $p_y= 2.2244$ $p_z= 0.0000$
 ℓ' $E= 2.1240$ $p_x= -0.2452$ $p_y= -2.1098$ $p_z= 0.0000$
 P' $E= 2.2645$ $p_x= 0.0000$ $p_y= 2.0611$ $p_z= 0.0000$
 q_γ $E= 0.2500$ $p_x= 0.2452$ $p_y= 0.0488$ $p_z= 0.0000$

Transverse plane




Leading initial-ensemble/final-state components (N=4, retained=99.3%)



Tx massless lepton + Tx proton characteristic max $C_{p\gamma}$ configuration

3D momenta




c_p_gamma_low_egamma_txtx_region_2 $C_{p\gamma}=0.984344$
 region: low_Egamma_TxTx_region_2 final-pair delta_xy=1.37445 rad
 outgoing-state purity: 1
 kinematic point: high_s_high_theta_in_low_Egamma
 incoming state: $|T_{x,\ell_0}\rangle \otimes |\bar{T}_{x,p}\rangle$


$s=21.5158$, $\sqrt{s}=4.63852$, $|\vec{P}|=2.22442$, $|\vec{P}'|=2.06107$
 $\theta_{in}=1.57079$, $\phi_\ell=4.71239$, $\phi_P=1.5708$
 $E_\gamma=0.25$, $\phi_\gamma=2.94524$
 $Q^2=0.0631736$, $x_B=0.0635316$, $t=-0.00429535$, $W^2=1.81103$, $y=0.0481859$
 $\theta(\ell', \gamma)=107.879$ deg

four-momenta $[E, p_x, p_y, p_z]$ GeV:
 ℓ $E= 2.2244$ $p_x= -0.0000$ $p_y= -2.2244$ $p_z= -0.0000$
 P $E= 2.4141$ $p_x= 0.0000$ $p_y= 2.2244$ $p_z= 0.0000$
 ℓ' $E= 2.1240$ $p_x= 0.2452$ $p_y= -2.1098$ $p_z= 0.0000$
 P' $E= 2.2645$ $p_x= 0.0000$ $p_y= 2.0611$ $p_z= 0.0000$
 q_γ $E= 0.2500$ $p_x= -0.2452$ $p_y= 0.0488$ $p_z= 0.0000$

Transverse plane




Leading initial-ensemble/final-state components (N=4, retained=99.3%)



Tx massless lepton + Tx proton characteristic max $C_{p\gamma}$ configuration

3D momenta




c_p_gamma_low_egamma_txtx_region_3 $C_{p\gamma}=0.658932$
 region: low_Egamma_TxTx_region_3 final-pair delta_xy=2.74889 rad
 outgoing-state purity: 1
 kinematic point: high_s_high_theta_in_low_Egamma
 incoming state: $|T_{x,\ell_0}\rangle \otimes |T_{x,p}\rangle$

$s=21.5158$, $\sqrt{s}=4.63852$, $|\vec{P}|=2.22442$, $|\vec{P}'|=2.21331$
 $\theta_{in}=1.57079$, $\phi_\ell=4.71239$, $\phi_P=1.5708$
 $E_\gamma=0.25$, $\phi_\gamma=5.10509$
 $Q^2=0.0102647$, $x_B=0.00459348$, $t=-1.87039e-05$, $W^2=3.10419$, $y=0.108287$
 $\theta(\ell', \gamma)=25.263$ deg

four-momenta $[E, p_x, p_y, p_z]$ GeV:

ℓ	$E= 2.2244$	$p_x= -0.0000$	$p_y= -2.2244$	$p_z= -0.0000$
P	$E= 2.4141$	$p_x= 0.0000$	$p_y= 2.2244$	$p_z= 0.0000$
ℓ'	$E= 1.9846$	$p_x= -0.0957$	$p_y= -1.9823$	$p_z= 0.0000$
P'	$E= 2.4039$	$p_x= 0.0000$	$p_y= 2.2133$	$p_z= 0.0000$
q_γ	$E= 0.2500$	$p_x= 0.0957$	$p_y= -0.2310$	$p_z= 0.0000$

Transverse plane



Leading initial-ensemble/final-state components (N=5, retained=99.0%)

